Retrospective Methodology to Estimate Daily Infections from

Deaths (REMEDID) in COVID-19: the Spain case study

David García-García[1], María Isabel Vigo[1], Eva S. Fonfría[2], Zaida Herrador[3],

Miriam Navarro[2], Cesar Bordehore [2,4*]

[1] Department of Applied Mathematics, University of Alicante, Alicante, Spain.

[2] Multidisciplinary Institute for Environmental Studies "Ramon Margalef", University of Alicante,

Alicante, Spain.

[3] National Centre of Epidemiology, Instituto de Salud Carlos III, Madrid, Spain.

[4] Department of Ecology, University of Alicante, Alicante, Spain.

Corresponding author:

Dr. Cesar Bordehore

Multidisciplinary Institute for Environmental Studies "Ramon Margalef" & Department

of Ecology, University of Alicante, Campus San Vicente del Raspeig, 03690, Spain

Tel: +34 5903400 ext 8755, Fax: +34 965903625

cesar.bordehore@ua.es

Keywords: COVID-19, MoMo, mortality, deaths, epidemiology, Case Fatality Ratio,

first infection, infections inference

Abstract

The number of new daily infections is one of the main parameters to understand the dynamics of an epidemic. During the COVID-19 pandemic in 2020, however, such information has been underestimated. Here, we propose a retrospective methodology to estimate daily infections from daily deaths, because those are usually more accurately documented. This methodology is applied to Spain and its 19 administrative regions. Our results showed that probable infections during the first wave were between 35 and 42 times more than those officially documented on 14 March, when the national government decreed a national lockdown and 9 times more than those documented by the updated version of the official data. The national lockdown had a strong effect on the growth rate of virus transmission, which began to decrease immediately. Finally, the first inferred infection in Spain is about 43 days before it was officially reported during the first wave. The current official data show delays of 15-30 days in the first infection relative to the inferred infections in 63% of the regions. In summary, we propose a methodology that allows reinterpretation of official daily infections, improving data accuracy in infection magnitude and dates because it assimilates valuable information from the National Seroprevalence Studies.

1. Introduction

The key parameter to understand and model the evolution of the COVID-19 pandemic

is the number of daily infections. Despite its importance, however, it is difficult to

estimate reliable data due to the bias in official figures[1]. For symptomatic patients, there

are delays of 5.79 days from infection to symptom onset and 5.82 days from symptom

onset to diagnosis[2]. In addition, the presence of asymptomatic people and those with

mild symptoms hinder their detection. For example, 86% of infections were undetected

in Wuhan, China, prior to 23 January 2020, when travel restrictions were imposed[3].

Additionally, not all patients with COVID-19 compatible symptoms are tested,

especially at the beginning of the pandemic. So, it is widely assumed that the reported

infections are just a fraction of the actual ones. In Spain, a National Seroprevalence

Study based on ~55,000 random participants estimated that only 10.7% (CI 95%:

10.1%-11.3%) of the actual infections had been detected during the first wave (before

22 June 2020) in Spain[4, 5].

Another problem is the inconsistency of daily infection time series and the variability of

the case definitions. For example, on 13 January 2020, China changed the parameters to

confirm new cases and ~15,000 new infections were counted in a single day, while

daily infections during the previous week were less than 3,500[6]. In Spain, the method of

counting official numbers of infections has been modified several times since the

pandemic was decreed. On 22 April 2020, a total of 213,024 cases were reported from

positive PCR and antibody tests[7]; however, total cases decreased to 202,990 the next

day and thereafter, because only positive PCRs were included in the total cases[8].

Accumulated cases on a given day are obtained by adding the new cases that day to the

accumulated cases of the previous day. That has not always been the method, however,

because some Spanish regions reviewed and modified the cases in previous days[9,10].

This inconsistency of time series hampers any kind of accurate analysis. Additionally,

data are continuously being updated. In Spain, official daily infection data available

during the first wave reported 1,832 new infected on 14 March 2020, but current official

data (as downloaded in February 2021) reported 7,478 new cases for the same date.

High fidelity time series of each parameter of an epidemic are crucial to run reliable

epidemiological models. The number of new infections per unit time is one of the

essential parameters. The official data do not reflect the actual day of infection nor the

real number of infections. To overcome this limitation, we propose a retrospective

methodology to infer daily infections from daily deaths, believing the number of deaths

to be a more reliable parameter than the official number of daily infections[1]. These

estimated infections are closer to the date of infection, avoiding delays from the

incubation period and symptom onset to diagnosis. Then, the time series of daily

infections would help to better understand the pandemic dynamics and to quantify the

effectiveness of the different control strategies, avoiding the bias of the official data. In

addition, those numbers can be used to feed epidemiological models with more realistic

data. Finally, we will apply this methodology in Spain as a whole and each of its 19

administrative regions (17 autonomous communities and 2 autonomous cities).

2. Methods

2.1 REMEDID

We named this methodology REMEDID, **RE**trospective **M**ethodology to **E**stimate

Daily **I**nfections from **D**eaths and it is described as follows. Date of infection (DI) of an

individual may be estimated from the death date (DD) by subtracting the incubation

period (IP) and illness onset to death (IOD) periods, hence

$$DI + IP + IOD = DD.$$

Therefore, DI can be estimated from DD as far as IP and DD are known; however,

neither IP nor IOD are fixed values. On the contrary, they are random variables that can

be approximated by probability distributions. From dozens of cases in Wuhan, Linton et

al.[11] approximated IP for COVID-19 with a lognormal distribution, X_{IP}, with a mean of

5.6 days and median of 5 days, and IOD with a lognormal distribution, X_{IOD}, with a

mean of 14.5 days and median of 13.2 days. Therefore, the infection to death period is a

random variable, X_{IP+IOD}, that follows the distribution $X_{IP} + X_{IOD}$.

Although the addition of two lognormal distributions does not follow any commonly

used probability distribution, its probability density function (PDF) can be estimated

convolving the PDF of the two variables[12]. If $g(t)$ and $h(t)$ are the PDF of IP and IOD,

respectively, then their convolution defines the PDF of X_{IP+IOD},

$$f(t_0) = \int\limits_{-\infty}^{+\infty} g(t_0 - t) \cdot h(t)\, dt,$$

125   where t_0 is a positive real number representing days from infection. Then, *f(t)* is the PDF

126   of the time from infection to death, with a mean of 20.1 days and a median of 18.8 days.

127   Figure 1 shows *g(t)*, *h(t)*, and *f(t)*, as well as a lognormal PDF with the same mean and

128   median as *f(t)* for comparison. Note that the probability that X_{IP+IOD} is less than or equal

129   to 33 days is 0.95.

131   [Figure 1]

Given a time series of deaths produced by the illness, *x(t)*, we estimate the infection

time series that produced such deaths, *y(t)*. If we assume a case fatality ratio (CFR) of

100%, *y(t)* would represent all daily infections. To calculate *y(t)*, we use the following

likelihood-based estimation procedure.

Because the relative likelihood that a given infection will produce a death *t* days later is

f(t), the infections at a given time t_0 can be estimated as

$$y(t_0) = \int\limits_{t_0}^{+\infty} x(t) \cdot f(t - t_0)\, dt.$$

142   In practice, *x(t)* is a discrete time series and could be written as *x(n)*, where *n* is an

143   integer representing entire days. Let *F(n)* be a discrete approximation to *f(t)* as follows:

$$F(n) = \int_{n}^{n+1} f(t)\, dt.$$

146   Then,

$$y(n_0) = \sum_{n=n_0}^{+\infty} x(n) \cdot F(n - n_0). \qquad (1)$$

149   A similar analysis was used previously to estimate a time-variable effective

150   reproduction number for the severe acute respiratory syndrome (SARS) in 2003[13].

151   For a different CFR, the associated/estimated infections are estimated from *y(n)* in

152   Equation 1 as

$$\text{Inferred infections}(n) = \frac{y(n)}{CFR} \times 100. \qquad (2)$$

Equations 1 and 2 constitute the kernel of the REMEDID algorithm. Validation of the

capacity of REMEDID to reproduce the number of new daily infections is in Appendix

A.

2.2. Data

2.2.a. Official numbers of COVID-19 infections and deaths

COVID-19 daily infections and deaths for Spain are reported by the *Centro de*

Coordinación de Alertas y Emergencias Sanitarias (CCAES), which depends on the

Ministry of Health, Social Services and Equality. Time series for the 19 regions of

Spain can be downloaded from the national organization that compiles and publishes

the data from all regions, the Instituto de Salud Carlos III (ISCIII;

https://covid19.isciii.es/). For consistency, time series for Spain were estimated by

adding the time series of the 19 regions. The final access to infections from official data

was on 16 February 2021 (hereafter referred as IO_{21}). Those data have been greatly

improved with respect to the daily situation reports published by CCAES during the

first wave in the first semester of 2020 (hereafter referred as IO_{20};

https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov-

China/situacionActual.htm). For example, IO_{20} cases were given the date of diagnosis,

whereas IO_{21} cases were given the date at symptoms onset. When the latter was not

available, it was estimated as follows: infections prior to 10 May 2020 were given the

diagnosis date minus 6 days, whereas since 11 May 2020 only 3 days were subtracted

from the date of diagnosis. Asymptomatic cases were given the diagnosis date. These

corrections brought reported cases closer to the real infection dates, although they were

still delayed by the length of the incubation period. In addition, re-evaluation of the

suspicious case of the first infectee on 20 February 2020 in IO_{20} gave it the date of 1

January 2020 in IO_{21}. The first official death was reported on 13 February 2020.

However, no deaths were reported until 3 March 2020 and afterwards deaths were

continuously reported. Then, time spans we considered for IO_{21} is from 1 January 2020

to 29 November 2020 and for official daily deaths from 3 March 2020 to 1 January

2021, specifically with 33 extra days as required to apply the REMEDID methodology.

When applying REMEDID the use of a reliable CFR is essential. Thus, to determine a

realistic CFR in Spain, we used the total infections data from the longitudinal

seroprevalence study conducted by ISCIII in 4 phases. The first three phases were

completed from 27 April to 22 June and the results were published on 6 July 2020[4, 5]. In

this longitudinal cohort study, ~68,000 people were randomly selected and ~55,000 of

them agreed to be interviewed and tested in the three phases. The study found that 5.2%

(CI 95%: 4.9%-5.5%) of the participants presented SARS-Cov2 IgG antibodies.

Extrapolation of this result to the ~47 million of the population in Spain yielded 2.45

million infections until 22 June, which, with 29,674 accumulated COVID-19 deaths on

22 June, would give a CFR of 1.21% (CI 95%: 1.15%-1.29%). We assumed this value

in our analysis for Spain until 22 June. The fourth phase of the study was completed

from 16 to 29 November 2020 and the results published on 15 December 2020[14]. In that

study, ~51,000 people agreed to participate. The study found that 9.9% (CI 95%: 9.4%-

10.4%) of the participants presented SARS-Cov2 IgG antibodies at any of the four

phases. Then, there were 4.75 millions of accumulated infections in Spain, meaning

2.30 million new infections from 23 June to 29 November. Because there were 47,113

accumulated COVID-19 deaths on 29 November (17,439 new deaths from 23 June), it

would give a CFR of 0.76% (CI 95%: 0.71%-0.81%). Thus, we assumed different CFRs

for the different periods, in order to smooth the discontinuity effects, we propose a CFR

time series segmentation as follows: i) a constant value of 1.21% from 1 January to 22

June 2020, ii) a constant value of 0.76% from 30 June to 29 November 2020, and iii) a

linear function from 23 June to 28 June that would correspond to the interpolation of the

CFR values for 22 June and 29 June. The CFRs of each region were estimated

analogously (summarized in Tables 1 and 2). Prior to 22 June, Asturias, Castilla – La

Mancha, Castilla y León, Extremadura, País Vasco, and La Rioja had CFRs greater than

1.5%. In contrast, Andalucía, Canarias, Melilla, and Murcia had CFRs less than 0.7%.

After 22 June, the CFR for Spain diminished in 0.45%, probably due to improved

medical treatments and experience gained during the first wave. In this period, the

number of regions with a CFR below 0.7% increased to 10 and those with a CFR

greater than 1.5% decreased to 3. Most of the regions revealed certain improvement,

particularly remarkable was La Rioja, with a CFR decrease from 3.13% to 1.12%.

2.2.b. Excess of expected deaths from MoMo

Expected deaths from any cause, with an error band representing a 99% confidence

interval, are modelled by the Mortality Monitoring (MoMo) surveillance system for 22

countries in Europe (www.euromomo.eu). In Spain, MoMo depends on the ISCIII[15] and

data are available at

https://momo.isciii.es/public/momo/dashboard/momo_dashboard.html#nacional. MoMo

also collects daily deaths from any cause from the Spanish *Instituto Nacional de*

Estadística (INE) and the notaries and civil registries. Daily deaths within the error

estimates of the expected deaths are considered usual. Any deviation allows

identification of unusual mortality patterns, as for the COVID-19 pandemic. The useful

variable here is the excess deaths (*ED*), estimated as the difference between actual and

expected deaths. The error band for *ED* can be estimated from the error band of the

expected deaths by subtracting the latter from the actual deaths. Note that *ED* can show

negative values (Figure 2). By cautiously assuming that all significant *ED* is caused by

COVID-19, *ED* can be used to estimate non-reported COVID-19 deaths. In this case,

negative values for *ED* make no sense and are set to zero. The MoMo time series was

downloaded on 16 February 2021 and we considered the period from 3 March 2020

(first non-isolated official COVID-19 death) to 1 January 2021.

Given that MoMo *ED* in Spain was 45,119 (CI 99%: 37,593-55,098) deaths from 3

March to 22 June, and 23,612 (CI 99%: 11,356-39,690) from 23 June to 29 November,

the CFR for the country estimated similarly as in Section 2.1 was 1.85% (CI 95%:

1.74%-1.96%) prior to 22 June, and 1.02% (CI 95%: 0.97%-1.09%) after 23 June. For

each region, the CFRs were estimated similarly and are provided in Tables 1 and 2. In

all cases, we used a piece wise CFR time series segmented as described in Section 2.2a.

3. Results

3.1. COVID-19 vs. MoMo

The COVID-19 official deaths and MoMo *ED* time series overlaped for the period from

3 March 2020 to 1 January 2021 for Spain and its 19 regions (Figure 2). In general,

there was good agreement between both datasets, meaning that most of MoMo *ED* were

related to COVID-19 deaths. During the first wave, the most important differences were

observed in Spain, Madrid, Cataluña, Castilla-La Mancha, and Castilla y León. Before

22 June in Spain, MoMo *ED* showed 15,445 accumulated deaths more than the official

COVID-19 deaths, which is beyond the error band. That difference comes basically

from the four regions with the largest numbers of deaths (Madrid, Cataluña, Castilla-La

Mancha, and Castilla y León). Table 1 shows the accumulated values before 22 June,

which were used to estimate the CFR for Spain and its 19 regions according to the third

phase of the National Seroprevalence Study[4,5]. For all regions, the CFR estimated from

MoMo *ED* was larger than the CFR estimated from COVID-19 deaths. In particular,

Asturias, Canarias, and Murcia were twice as large. Ceuta and Melilla dramatically

increased their CFR from MoMo *ED*, although that may be biased due to their small

populations and numbers of deaths.

Similarly, the same variables for the period from 23 June to 29 November 2020 are

reported in Table 2. In Spain, MoMo *ED* showed 6,173 accumulated deaths more than

the official COVID-19 deaths. This difference is a third of the difference observed prior

to 22 June; because this is within the error band, there was a significant improvement in

the detection of COVID-19 deaths in this period. Figure 2 also shows a general

agreement between MoMo *ED* and official COVID-19 deaths time series after the first

wave, with the exception of late July and early August. These differences were due to

two heat waves that were responsible for at least 25% of the MoMo *ED*[16].

276   [Table 1]

277   [Table 2]

278   [Figure 2]

3.2. Infections estimated from COVID-19 deaths

To illustrate the delay between official daily infections data and REMEDID estimated

daily infections, we applied REMEDID from COVID-19 deaths assuming CFR= 100%.

Figure 3 shows the current IO_{21} and the infections associated with COVID-19 deaths

for the first wave. The latter in Spain reached a maximum on 13 March 2020 (Table 3),

the day before the national government decreed a state of emergency and national

lockdown. Thus, the adopted measures had an immediate effect, which was observed in

the official data IO_{21} 7 days later (20 March). This delay is similar to the incubation

period (mean 5.78 days[2]), which could be explained because official infections were

reported when symptoms appeared. This delay reached 16 days when we compared with

earlier version IO_{20} (not shown), which highlights the usefulness of the methodology to

reinterpret official data from very early stages of the pandemic. On the other hand, the

maximum number of deaths was reached on 1 April, which was 19 days after the

inferred infection maximum, bringing this delay close to the 20 days expected between

infection and death (Figures 1 and 3).

296   [Figure 3]

We applied REMEDID to the official COVID-19 death with the corresponding

estimated CFRs (see section 2.2) to obtain the time series of estimated daily infections,

hereafter referred to as IR_O. Figure 4 shows IR_O and the accumulated infections for

Spain and its 19 regions. Note that in Spain, IR_O are amplified versions of inferred

infections in Figure 3. In Spain, the first infection, according to IR_O, is on 8 January

2020 (Table 3), 43 days before the first infection was officially reported on 20 February

2020 according to IO_{20}. By contrast, IO_{21} places the first infection on 1 January 2020.

Spain reached the maximum number of IR_O on 13 March, a day before the state of

emergency and lockdown were enforced (Table 4). On 14 March, $IO_{20} = 1,832$, and IO_{21}

$= 7,478$; however, $IR_O = 63,727$ (CI 95%: 60,050-67,403), 35 and 9 times IO_{20} and IO_{21},

respectively (Table 5). This implies that on that day, IO_{20} and IO_{21} only reported 2.9%

(CI 95%: 2.7%-3.1%) and 11.7% (CI 95%: 11.1%-12.5%) of new infections,

respectively. Although detection of infections clearly improved from IO_{20} to IO_{21},

almost 90% of the infections are still not documented in the peak of the first wave. The

situation is similar for the accumulated infections before 22 June 2020, as reported by

the National Seroprevalence Study[4,5].

In almost all regions, IO_{20} showed a delay of 1 month or more between the first

infection and IR_O (Table 3). No delay in IO_{21} occurred in Islas Baleares, Castilla-León,

and Galicia, while in three regions (Cataluña, Madrid, and La Rioja), the first case

occurred earlier than the first case of IR_O. However, 6 regions had delays of 15 days and

other 6 regions had delays of 1 month. According to IR_O, all regions except Ceuta and

Melilla had some infections in January, but in IO_{21} only 6 regions had infections in that

month. In all scenarios, the first infections were in Madrid and Cataluña.

During the first wave, according to IR_O most of the regions had maximum daily

infections around 14 March. In Madrid, the maximum was reached on 11 March,

coinciding with the educational centres closing and an official warning by the regional

government (Table 4). Asturias was the last region to reach peak infections (25-26

March). The maximum percentage of documented cases (12.6%,CI 95%: 9.2%-18.4%)

occurred in Asturias on 14 March, but in the other regions, only between 1.2 to 8 % of

the infections were documented (Table 5).

Figure 4 shows how the IO_{21} and IR_O curves of Spain and the 19 different regions

fluctuated following the same pattern until the middle of June 2020, but thereafter, they

showed different patterns. This reflects the fact that the Spanish government had

decreed the control measures for the whole nation until June, but thereafter, each

regional government implemented its own control measures. For example, some regions

(e.g., Aragón, Islas Baleares, Cantabria, Comunidad Valenciana, Extremadura, Galicia,

Murcia, País Vasco, and La Rioja) had two peaks, but others had only one. An apparent

maximum on 22 June in Islas Baleares is an artifact produced by the interpolation for

transition from the two CFRs. Although beyond the scope of this work, it would be very

interesting to investigate the effects of the different control measures implemented on

the corresponding IR_O for the 19 regions.

The Spanish COVID-19 second wave reached a maximum of daily infections on 22

October from IR_O and on 26 October from IO_{21}. The delay of 4 days is similar to the

mean incubation period (5.78 days[2]). The estimated number of new infections is still

larger than the documented cases, but the shapes of the two curves are more similar in

the second wave than in the first wave (Figure 1a). The same is true for the 19 regions,

most of which had the largest peak around 22-26 October, with the exceptions of

Canarias and Madrid, which reached maxima in late August and early September,

respectively.

352   [Table 2]

353   [Table 3]

354   [Figure 4]

3.3. Infections from MoMo Excess Deaths

Assuming that MoMo *ED* accounts for both recorded and non-recorded COVID-19

deaths, negative deaths are meaningless, and they were set to zero. Then, the associated

daily infections can be estimated, as in Section 3.2, with a CFR of 100% from MoMo

ED for Spain (Figure 5). Note two main differences between this time series and that

estimated from official COVID-19 deaths: (1) MoMo data present an error band that

was inherited by the estimated infections; (2) MoMo *ED* estimated infections reached a

maximum of 1,443 (CI 99%: 1,329-1,547), doubling the 776 inferred daily infections

from official COVID-19 deaths in Figure 3. This is because maximum MoMo *ED* was

1,584 (CI 99%: 1,468-1,686%) and maximum COVID-19 official deaths was 828, both

estimated from the 14-day running mean time series. The maximum of inferred

infections was reached on 13 March, just one day prior to the state of emergency and

lockdown. The expected and observed delays with respect to official infections and

MoMo *ED* were similar to those observed for estimated infections from official

COVID-19 deaths. Error bounds of the estimated infections in Figure 5 were computed

from the MoMo *ED* error bounds. However, it should be highlighted that the

combination of the error bounds from MoMo *ED* and the estimated CFRs might lead to

unrealistic error estimates. To avoid this, the error estimates in Figure 6 were estimated

from the MoMo *ED* time series (no error bounds) and the error bounds of the estimated

CFRs.

380   [Figure 5]

The REMEDID was applied to the MoMo *ED* with the corresponding CFRs (see

section 2.2) to obtain the estimated daily infections, which will be referred hereafter as

IR_M. The IR_M were calculated for Spain and its 19 regions and are depicted in Figure 6,

as well as the accumulated IR_M. In Spain, the first infection shown by IR_M happened on

9 January, with an error estimate from 9 to 10 January, 41 to 42 days before the first

documented infection of IO_{20} on 20 February 2020 (Table 3). The maximum IR_M was

77,855 (CI 95%: 73,364 - 82,347) reached on 13 March. On 14 March, IR_M showed

14,128 infections more than IR_O (Table 5). Notice that the CFR used with MoMo *ED*

data was larger than the one used with official COVID-19 deaths data, which makes this

difference even more remarkable, because the larger the CFR the lower the estimated

infections. Therefore, if the true CFRs, which are unknown, were used in both cases,

IR_M would double IR_O on 14 March, as happened when a CFR of 100% was used

(Figures 3 and 5). Notice that with the CFRs used, the IR_M and IR_O resulted in the same

accumulated infections on 22 June and 29 November, matching the results of the

seroprevalence study. Nevertheless, IR_M showed 42 times more cases than IO_{20} and 10

times more than IO_{21} on 14 March, detection of official cases of only 2.4% (2.2%-2.5%)

and 9.6% (9.1%-10.2%), respectively.

Table 3 shows the estimated date of first infection for Spain and by region. Note that the

first cases of IR_M in Spain were on 9 January and in Aragón, Canarias, and Navarra on 8

January, which is possible because significant excess deaths in a region may not

become significant for the whole country. In general, the maxima of daily infections

were closer to those on 14 March in IR_M than in IR_O. During the first wave, all regions

showed a single maximum, except for Ceuta, Melilla, and Murcia, which showed two

maxima (Figure 6). In general, the IR_M time series in all regions were similar during that

period. The official data clearly under-detected infections during the first wave. On 14

March, IR_M were comparable to IR_O, overlapping CI in all regions, but not in Spain as a

whole (Table 5). During the second wave, there was improved detection of cases with

differences among regions.

412   [Figure 6]

4. Discussion

Infection dynamics is key to understand and model the COVID-19 pandemic.

Nevertheless, documented infections in Spain (and nearly worldwide) are of limited

quality qualitatively and quantitatively. Specifically, reported infections were

underestimated and delayed 16 days compared to the estimated date of infection during

the first months of the pandemic (according to IO_{20}) and 6 days in the current version of

the data (IO_{21}). According to the National Seroprevalence Study, only 10.7% of

infections were documented in Spain prior to 22 June 2020[4,5] and 64.3% from 23 June

to 29 November 2020[14]. The huge underreporting during the first wave is mainly due to:

(1) the lack of testing for asymptomatic and mildly symptomatic people[3], which could

have been detected with a "test, track and trace" strategy, as done in South Korea,

China, and Singapore[17]; (2) deaths outside of the hospitals, with many cases from

nursing homes for the elderly, where 6,664 deaths with COVID-19 and 7,082 with

similar symptoms, but not officially diagnosed with COVID-19, occurred from January

to May 2020[18]. This poor quality of data hinders any options to study the infection

dynamics based on official data.

To overcome these difficulties, we propose the REMEDID algorithm, which allows

calculation of time series for daily infections from daily death time series, a more

reliable source, which can be applied in early stages of any pandemic with only 33 days

delay. The REMEDID algorithm needs only three inputs:

(1) PDF of the period from date of infection to date of death. In this study, that was

calculated by combining the PDF of the Incubation Period and the Illness Onset to

Death period, as estimated by Linton et al.[11] for cases in Wuhan, China. The incubation

period can be assumed to be similar in China and elsewhere, because it is a

characteristic of the virus, but the illness onset to death period could be affected by the

type of health care and, thus, by each national health system[2] or even ethnicity[19]. No

such information is available for Spain to date, but it would be desirable to use a local

distribution of Illness Onset to Death when available for Spain. The calculations could

be redone easily by applying the MATLAB code available in the Supplementary

Material.

(2) Time series of daily deaths. We used two sources, the official COVID-19 deaths and

the MoMo *ED*, because official COVID-19 deaths have been underestimated, especially

during the first wave in the regions of Castilla–La Mancha, Castilla y León, Cataluña,

and Madrid, because non-hospital deaths and cases during the early pandemic were

somewhat undertested (Table 1 and Figure 2). We expect that undocumented COVID-

19 deaths appeared in the excess of expected deaths from any cause calculated from

MoMo[15]. Interpretation of those results must be made carefully because *ED* is a

statistical variable calculated from what happened in previous years and some expected

deaths, such as those produced by traffic and work accidents, may have been reduced

due to the national lockdown that began on 14 March 2020 and gradually has been

eased. Nonetheless, expected fatalities were very low (traffic and work accidents less

than 5 deaths daily) compared to the daily death toll from COVID-19 (300 deaths per

day average from March to May). Some non-COVID-19 deaths may have been

influenced by other pandemic-related factors[22], such as breakdown in medical follow-up

(especially chronic conditions and among older patients), delays in attending to the

healthcare appointments and/or being attended due to the collapse of the Health System,

transplant delay and reduction in donors' numbers from 7.2 to 1.2 per day[23], and the

decrease of the interventional cardiology activity between 40% and 81%[24]. Thus,

MoMo *ED* do not exactly correspond to COVID-19 deaths and must be regarded with

caution. A detailed study of the changes in other sources of mortality is recommended.

Nevertheless, MoMo *ED* is a valuable source of data that must be explored considering

the enormous bias in official data of daily infections.

(3) CFR estimate. CFR was defined as the ratio of deaths among total infections. Thus,

the lower the documented infections, the larger the CFR and under-detection of

infections leads to overestimation of the CFR. For example, in Spain on 22 June,

official data showed 261,111 accumulated infections and 29,676 deaths, a CFR of

11.37%, which was clearly overestimated because the number of infections was

underestimated. The National Seroprevalence Study in Spain with random testing

showed that 5.2% of the population (~47 million) had already been infected[4,5]. Because

CFR also depends on death records, we calculated a CFR of 1.21% (CI 95%: 1.15%-

1.29%) from official COVID-19 deaths and a CFR of 1.85% (CI 95%: 1.74%-1.96%)

from MoMo *ED*. However, the CFR is not constant in time nor in space. Different

regions showed different CFRs and the fourth phase of the seroprevalence study showed

a general decrease in the CFRs for the period 23 June – 29 November. In Spain, the

CFR decreased from 1.21% to 0.76% (CI 95%: 0.71%-0.81%) from official COVID-19

deaths and from 1.85% to 1.02% (CI 95%: 0.97%-1.09%) from MoMo *ED*. This decline

is probably due to the improvement of COVID-19 treatments and experience gained

since the beginning of the pandemic[25]. Similar seroprevalence studies are desirable in

other countries, because CFRs from official data are so variable. For example, on 22

June 2020, CFRs of 0.6, 5.2, and 14.5% were reported for Iceland, the world, and Italy,

respectively[26]. In addition, we assumed a constant CFR prior to 22 June and from 23

June to 29 November 2020, although introducing time dependence on the CFR would

enhance the infected estimates. A constant CFR does not consider who is affected by

saturation of the health systems due to limited human and material resources. However,

to estimate a time-variable CFR requires accurate infection records among other

variables, which are not currently available. Moreover, CFRs among severe cases

(mostly detected) are much greater than those among asymptomatic and mild cases

(mostly undetected). Consequently, the CFR calculation has a high degree of

uncertainty. Nevertheless, because information on the CFRs is expected to improve,

calculations in this study can be updated easily using the MATLAB code for

REMEDID algorithm provided as Supplementary Material.

We ran the REMEDID algorithm to provide the estimated time series of infections for

Spain and for its 19 regions for the period from 8 January to 29 November 2020. These

time series provided valuable information to understand the time evolution of the

pandemic. Our main findings for Spain are:

(1) On 14 March, when the national government decreed the state of emergency and

national lockdown, estimated infections were between 35 and 42 times larger than

officially documented ones at that time, and 9-10 times more than current official data.

(2) The national lockdown had a strong effect on the transmission of the virus, as shown

by a rapid slope decrease around day 13 March.

(3) The first infection was better determined from inferred infections than from official

records during the first months of the pandemic.

The REMEDID algorithm has strengths and limitations. First, it uses the number of

deaths, which are more accurately recorded than infections. Second, it allows

elucidation of the date of the infections estimated during the seroprevalence studies,

which only determines the total number of infections. Thus, the REMEDID algorithm

complements the seroprevalence studies. Third, the estimated daily infections place the

probable date of infection more accurately than the official numbers. Note that official

infections are delayed by the incubation period (from infection to illness onset).

Therefore, although the maxima of infections theoretically should have coincided with

the state of emergency and the national lockdown on 14 March, official infections in

Spain were maximum 6 days later. Infections estimated from death data do not show

such delay, however. Determination of the actual day of infection is very important for

evaluation of the immediate effect of the implemented countermeasures. In Madrid the

estimated infections reached their maximum on 11 March when regional government

warned the population to stay at home and schools and universities were closed, forcing

1.2 million students to stay at home. Moreover, overall recommendations on disease

control and social distancing were given by the Ministry of Health to the general public

since de beginning of March (see

https://www.mscbs.gob.es/en/profesionales/saludPublica/ccayes/alertasActual/nCov-

China/ciudadania.htm).

Four, the REMEDID algorithm can be applied to any region, regardless of population

size, but being cautious with small populations. For example, in this study we applied

REMEDID methodology to a small city (Ceuta with 84,777 inhabitants), a medium size

region (Madrid with more than 6.6 million inhabitants), and a country (Spain with 47

million inhabitants). This versatility allows study of different epidemic dynamics as

reflected in the variety of shapes and slopes in the curves depicting death and infection

records, that show the heterogeneity of the epidemic at each region with different

population size and density, social behaviour and transmission pattern, especially during

the second wave, showing unique dynamics that must be addressed individually.

Knowledge of the real epidemic dynamics of different population nodes is key to

succeed in modelling attempts, because we could calculate the sole effect of group

size[27]. This spatially-explicit information, in combination with population size per node

and mobility would allow us to use a metapopulation approach in future models[28-30].

Among the methodology limitations, the most important is that it can only be

implemented retrospectively. Thus, these estimates cannot be used to control the

pandemic in real time. But considering that different regions or countries are at different

stages at the same time, results of this methodology for the first communities could be

applied elsewhere. Furthermore, it is useful to enhance models and improve our

knowledge of the pandemic dynamics, including the effectiveness of the different

measures adopted to flatten the curve and design safe post-lockdown measures. In

addition, more realistic and accurate models could be obtained by use of the more

realistic daily number of infections provided by REMEDID, which, in turn, would

improve the models outcome and enhance comparisons of different lockdown and post-

lockdown measures[31].

We believe that REMEDID methodology applied to daily COVID-19 deaths (if

accurately reported) or to MoMo *ED* could be useful to analyse the dynamics of the

pandemic retrospectively and more accurately quantify the real daily infections with

respect to the official numbers. We only need the CFR and, for greater precision, the

PDF of Infections to Death or, alternatively, the PDF of Incubation Period and Illness

Onset to Death. This approach could be implemented anywhere, improving our

knowledge and understanding of the dynamics of the pandemic and the effectiveness of

the confinement measures. This is of high importance to develop successful strategies to

face and reduce the effects of future epidemic episodes.

Appendix A. Validation test

In order to validate REMEDID, it would be desirable to have an accurate time series of

daily infections and deaths. Then, the REMEDID infections inferred from deaths could

be compared to the recorded infections. However, such datasets are not available

because many infections in any disease or plague are usually undetected. On the other

hand, REMEDID-estimated infections cannot be compared to those of dynamical

models, which also need accurate data to be tuned. In fact, the main motivation for

developing REMEDID has been to provide (more) reliable data to feed dynamical

models.

As an alternative of validation, we propose the following experiment as a proof of

concept. In the official COVID-19 time series, IO_{21}, infections are given the date of

symptoms onset. Then, REMEDID applied to official deaths with the PDF of illness

onset to death, $h(t)$, instead of the PDF of infection to death, $f(t)$, should produce a time

series comparable to IO_{21}. Official accumulated cases and deaths on 22 June 2020 were

261,111 and 29,676, respectively, which gives a CFR of 11.37%. This CFR is

overestimated due to an under detection of cases in that period. On the other hand,

official accumulated cases and deaths from 23 June to 29 November 2020 were

1,482,293 and 17,439, which gives a CFR of 1.18%. Although closer to reality, this

value is still overestimated. In any case, if we use these CFR values to construct the

segmented timeseries of CFRs, as in Section 2.2, when applying REMEDID the

resulting estimated daily infections should be comparable to IO_{21}. The results are shown

in Figure A.1 where the REMEDID estimated daily infections is in very good

agreement with the smoothed IO_{21} (14-day running mean), with a correlation between

the two series of 0.98 % (p-value<0.001). Because smoothed time series present a high

degree of serial correlation, the correlation and the significance level was estimated

according to a specific Monte Carlo analysis based on the randomization of phases in

the frequency domain[32]. Notice that (i) application of the 14-day running mean to the

IO_{21} series removes the weekly variability due to the weekend infections misreport; (ii)

the agreement of the two series would improve with an improved estimate of the CFR

time series. Thus, the good match between the estimated and the original daily

infections timeseries proves the theoretical concept that supports REMEDID.

Note the differences with the results presented in section 3.2, where the PDF includes

the incubation period and a more realistic CFR, as estimated from the National

Seroprevalence Study data.

References

1. Holmdahl, S. M. I. & Buckee, D. P. C. Wrong but Useful — What Covid-19

epidemiologic models can and cannot tell us. *N Eng J Med*. DOI:

10.1056/NEJMp2016822 (2020)

2. Fonfría, E.S. et al. Essential epidemiological parameters of COVID-19 for clinical

and mathematical modeling purposes: a rapid review and meta-analysis. medRxiv

2020.06.17.20133587; doi: https://doi.org/10.1101/2020.06.17.20133587 (2020)

3. Li, R. et al. Substantial undocumented infection facilitates the rapid dissemination of

novel coronavirus (SARS-CoV-2). *Science* **368**, Issue 6490, 489-493,

DOI: 10.1126/science.abb3221 (2020).

4. Instituto de Salud Carlos III. Estudio ENE-COVID19: Estudio Nacional de sero-

Epidemiología de la infección por SARS-COV-2 en España.

Informe final, 6 July 2020. Available online at

https://www.mscbs.gob.es/ciudadanos/ene-covid/docs/ESTUDIO_ENE-

COVID19_INFORME_FINAL.pdf

5. Pollán, M., et al. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide,

population-based seroepidemiological study. *The Lancet* **396**, Issue 10250, 535-544,

https://doi.org/10.1016/ S0140-6736(20)31483-5 (2020).

6. World Health Organization. Coronavirus disease 2019 (COVID-19) Situation Report

– 24, published online on February 2020, https://www.who.int/docs/default-

source/coronaviruse/situation-reports/20200213-sitrep-24-covid-

19.pdf?sfvrsn=9a7406a4_4 (WHO, 2020)

7. Centro de Coordinación de Alertas y Emergencias Sanitarias. Actualización nº 84.

Enfermedad por el coronavirus (COVID-19), published online on 23 April 2020,

https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov-

China/documentos/Actualizacion_84_COVID-19.pdf (CCAES, 2020a).

8. Centro de Coordinación de Alertas y Emergencias Sanitarias. Actualización nº 85.

Enfermedad por el coronavirus (COVID-19), published online on 24 April 2020,

https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov-

China/documentos/Actualizacion_85_COVID-19.pdf (CCAES, 2020b).

9. Centro de Coordinación de Alertas y Emergencias Sanitarias. Actualización nº 111.

Enfermedad por el coronavirus (COVID-19), published online on 20 May 2020,

https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov-

China/documentos/Actualizacion_111_COVID-19.pdf (CCAES, 2020c).

10. Centro de Coordinación de Alertas y Emergencias Sanitarias. Actualización nº 112.

Enfermedad por el coronavirus (COVID-19), published online on 21 May 2020,

https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov-

China/documentos/Actualizacion_112_COVID-19.pdf (CCAES, 2020d).

11. Linton et al. Incubation period and other epidemiological characteristics of 2019

novel coronavirus infections with right truncation: a statistical analysis of publicly

available case data. *Journal of Clinical Medicine* **9** (2), 538, DOI: 10.3390/jcm9020538

(2020).

12. Uspensky, J. V. Introduction to mathematical probability. Ed. McGraw-Hill Book

Company, New York and London (1937).

13. Wallinga, J. & Teunis, P. Different epidemic curves for severe acute respiratory

syndrome reveal similar impacts of control measures. *American Journal of*

Epidemiology **160**, 509-516 (2004).

14. Instituto de Salud Carlos III. Estudio ENE-COVID19: Cuarta ronda. Estudio

Nacional de sero-Epidemiología de la infección por SARS-COV-2 en España.

Cuarta ronda, 6 July 2020. Available online at

https://portalcne.isciii.es/enecovid19/informes/informe_cuarta_ronda.pdf

15. León-Gómez, I. et al. Exceso de mortalidad relacionado con la gripe en España en el

invierno de 2012, *Gac Sanit* **29**, 258-65 (2015).

16. YY. Gil Bellosta, C. J., Luz Frías, Ludovica Verrieri, Concha Delgado, Inmaculada

León y Amparo Larrauri. Informe MOMOCalor, Estimaciones de la mortalidad

atribuible a excesos de temperatura en España, 1 de junio a 15 de septiembre de 2020.

Centro Nacional de Epidemiología. Ciber de Epidemiología y Salud Pública

(CIBERESP). Instituto de Salud Carlos III. Octubre 2020.

https://www.isciii.es/QueHacemos/Servicios/VigilanciaSaludPublicaRENAVE/Enferme

dadesTransmisibles/MoMo/Paginas/MOMOcalor.aspx [Last access: 16-02-2021]

17. Fang Yong, S. E., et al., Connecting clusters of COVID-19: an epidemiological and

serological investigation. *The Lancet*, published online April 21, 2020

https://doi.org/10.1016/S1473-3099(20)30273-5 (2020).

18. Defunciones según la Causa de Muerte - Avance enero-mayo de 2019 y de 2020.

Notas de prensa del Instituto Nacional de Estadística. 10 diciembre 2020.

https://www.ine.es/dyngs/INEbase/es/operacion.htm?c=Estadistica_C&cid=125473617

6780&menu=ultiDatos&idp=1254735573175 [Last access on 16 February 2021].

19. Risk for COVID-19 Infection, Hospitalization, and Death By Race/Ethnicity.

https://www.cdc.gov/coronavirus/2019-ncov/covid-data/investigations-

discovery/hospitalization-death-by-race-ethnicity.html [last access on 5 March 2021]

20. Dirección General de Tráfico, Ministry of Interior. Cuadro comparativo de

accidentes mortales y fallecidos a 24 h en vías interurbanas, published online on 13 May

2020, http://www.dgt.es/Galerias/seguridad-vial/estadisticas-e-indicadores/accidentes-

24-horas/2020/Mayo/CC_-2020_05_13.pdf (DGT, 2020).

21. Ministerio de Trabajo y Economía Social. Estadística de Accidentes de Trabajo,

Avance enero-marzo 2020, published online on 18 May 2020,

http://www.mitramiss.gob.es/estadisticas/eat/eat20_03/ATR_03_2020_Resumen.pdf

(MTES, 2020).

22. Vandoros, S. Excess Mortality during the Covid-19 pandemic: Early evidence from

England and Wales. medRxiv 2020.04.14.20065706; doi:

https://doi.org/10.1101/2020.04.14.20065706 (2020).

23. Domínguez☐Gil, B. et al. COVID-19 in Spain: Transplantation in the Midst of the

Pandemic. *Am J Transplant*. Accepted Author Manuscript. doi:10.1111/ajt.15983

(2020).

24. Rodríguez-Leor, O. et al. Impacto de la pandemia de COVID-19 sobre la actividad

asistencial en cardiología intervencionista en España. *REC Interv Cardiol* **2** (2), 82-89

(2020).

25. Boudourakis L, Uppal A. Decreased COVID-19 Mortality—A Cause for Optimism.

JAMA Intern Med. Published online December 22, 2020.

doi:10.1001/jamainternmed.2020.8438

26. Roser, M., Ritchie, H., Ortiz-Ospina, E. & Hasell, J. Mortality Risk of COVID-19.

Published online in Our World in Data: https://ourworldindata.org/mortality-risk-

covid#the-case-fatality-rate (accessed on 23rd May 2020).

27. Caillaud, D., Craft, M.E. & Meyers, A. Epidemiological effects of group size

variation in social species. *J. R. Soc. Interface* 10: 20130206, DOI:

10.1098/rsif.2013.0206 (2013).

28. Levins, R. Some demographic and genetic consequences of environmental

heterogeneity for biological control. *Bull. Entomol. Soc. Am. Am.* **15** (3), 237–240

(1969).

29. Hanski, I. Metapopulation Ecology. Oxford University Press (1999).

30. Ball, F. et al. Seven challenges for metapopulation models of epidemics, including

households models. *Epidemics* **10**, 63-67.

http://dx.doi.org/10.1016/j.epidem.2014.08.001 (2015)

31. Bordehore, C. et al. Understanding COVID-19 spreading through simulation

modeling and scenarios comparison: preliminary results. medRxiv

2020.03.30.20047043, doi: https://doi.org/10.1101/2020.03.30.20047043 (2020)

32. Ebisuzaki, W. A method to estimate the statistical significance of a correlation when

the data are serially correlated, J. Clim., 10, 2147–2153, (1997).

746
747

753 **Table 1.** Accumulated COVID-19 deaths and MoMo Excess Deaths (*ED*) for Spain and

754 its 19 regions. Values from 3 March to 22 June 2020. Population data from INE. *:

755 differences beyond the error estimation.

Region	2019 Population	Percentage of infections[4,5]	Official data		MoMo data			*ED* minus COVID-19 deaths
		% (CI 95%)	COVID-19 deaths	CFR (CI 99%)	*ED* (CI 95%).	Days with lower limit >0	CFR (CI 95%)	
Spain	47,026,208	5.2 (4.9; 5.5)	29674	1.21 (1.15; 1.29)	45,119 (37,593; 55,098)	57	1.85 (1.74; 1.96)	15445*
Andalucía	8,414,240	3.0 (2.6; 3.6)	1441	0.57 (0.48; 0.66)	2,123 (651; 5,168)	23	0.84 (0.70; 0.97)	682
Aragón	1,319,291	4.8 (3.7; 6.2)	915	1.44 (1.12; 1.87)	1,073 (408; 2,165)	30	1.69 (1.31; 2.20)	158
Asturias	1,022,800	1.9 (1.3; 2.6)	336	1.73 (1.26; 2.53)	689 (73; 1,885)	14	3.54 (2.59; 5.18)	353
Islas Baleares	1,149,460	1.4 (0.8; 2.4)	226	1.4 (0.82; 2.46)	279 (11; 1,090)	6	1.73 (1.01; 3.03)	53
Canarias	2,153,389	2.3 (1.5; 3.5)	166	0.33 (0.22; 0.51)	337 (13; 1,585)	5	0.68 (0.45; 1.04)	171
Cantabria	581,078	3.7 (2.5; 5.4)	212	0.99 (0.68; 1.46)	296 (34; 932)	10	1.38 (0.94; 2.04)	84
Castilla - La Mancha	2,032,863	9.6 (7.8; 11.6)	2951	1.51 (1.25; 1.86)	5,468 (4,464; 7,003)	47	2.80 (2.32; 3.45)	2517*
Castilla y León	2,399,548	7.8 (6.8; 9.0)	2829	1.51 (1.31; 1.73)	3,813 (2,763; 5,452)	45	2.04 (1.77; 2.34)	984
Cataluña	7,675,217	5.9 (5.0; 7.1)	6507	1.44 (1.19; 1.7)	12,182 (10,034; 15,537)	54	2.69 (2.24; 3.17)	5675*
Ceuta	84,777	0.7 (0.3; 1.8)	5	0.84 (0.33; 1.97)	85 (5; 170)	3	14.24 (5.54; 33.22)	80
C. Valenciana	5,003,769	2.4 (1.9; 3.1)	1450	1.21 (0.94; 1.53)	2,078 (753; 4,391)	29	1.73 (1.34; 2.19)	628
Extremadura	1,067,710	3.1 (2.2; 4.3)	513	1.55 (1.12; 2.18)	919 (298; 2,044)	25	2.78 (2.00; 3.91)	406
Galicia	2,699,499	1.9 (1.5; 2.4)	618	1.20 (0.95; 1.53)	755 (122; 2,472)	13	1.47 (1.16; 1.86)	137

Madrid	6,663,394	11.7 (.3; 13.3)	8720	1.12 (0.98; 1.27)	14,443 (12,766; 16,612)	62	1.85 (1.63; 2.10)	5723*
Melilla	86,487	3.4 (2.3; 5.1)	2	0.07 (0.05; 0.10)	63 (0; 144)	0	2.14 (1.43; 3.17)	61
Murcia	1,493,898	1.6 (1.0; 2.7)	147	0.62 (0.36; 0.98)	387 (37; 1,437)	9	1.62 (0.96; 2.59)	240
Navarra	654,214	6.6 (5.1; 8.5)	528	1.22 (0.95; 1.58)	776 (400; 1,494)	29	1.80 (1.39; 2.32)	248
País Vasco	2,207,776	3.6 (2.9; 4.7)	1555	1.96 (1.50; 2.43)	1,866 (838; 3,388)	32	2.35 (1.80; 2.91)	311
La Rioja	316,798	3.7 (2.8; 4.9)	367	3.13 (2.36; 4.14)	386 (59; 849)	21	3.29 (2.49; 4.35)	19

760 **Table 2.** Accumulated COVID-19 deaths and MoMo Excess Deaths (*ED*) for Spain

761 and its 19 regions. Values from 23 June to 29 November 2020. The percentages for this

762 period were estimated as differences between the accumulated percentages on 29

763 November and on 22 June.

Region	Percentage of infections[14]	Official data		MoMo data			*ED* minus COVID-19 deaths
	% (CI 95%)	COVID-19 deaths	CFR (CI 95%)	*ED* (CI 99%).	Days with lower limit >0	CFR (CI 95%))	
Spain	4.9 (4.6; 5.2)	17439	0.76 (0.71; 0.81)	23,612 (11,356; 39,690)	104	1.02 (0.97; 1.09)	6173
Andalucía	4.1 (3.5; 4.7)	3007	0.87 (0.76; 1.02)	5,814 (2,173; 10,738)	77	1.69 (1.47; 1.97)	2807
Aragón	7.1 (5.9; 8.4)	1416	1.51 (1.28; 1.82)	1,,666 (421; 3,263)	59	1.78 (1.50; 2.14)	250
Asturias	3.7 (2.8; 5.1)	709	1.87 (1.36; 2.48)	1,198 (320; 2,933)	31	3.17 (2.30; 4.18)	489
Islas Baleares	4.7 (3.4; 6.3)	218	0.40 (0.30; 0.56)	470 (18; 1,666)	6	0.87 (0.65; 1.20)	252
Canarias	1.4 (1.0; 2.0)	178	0.59 (0.41; 0.83)	592 (31; 2,227)	9	1.96 (1.37; 2.75)	414
Cantabria	2.7 (2.0; 3.6)	109	0.70 (0.52; 0.94)	287 (6; 1,127)	4	1.83 (1.37; 2.47)	178
Castilla - La Mancha	7.2 (6.6; 8.0)	1020	0.70 (0.63; 0.76)	1,601 (214; 3,748)	36	1.09 (0.98; 1.19)	581
Castilla y León	4.6 (4.1; 5.1)	1843	1.67 (1.51; 1.87)	2,053 (481; 4,397)	47	1.86 (1.68; 2.09)	210
Cataluña	6.1 (5.1; 7.1)	1914	0.41 (0.35; 0.49)	3,677 (777; 8,125)	37	0.79 (0.67; 0.94)	1763
Ceuta	8.9 (6.2; 12.2)	51	0.68 (0.49; 0.97)	144 (6; 276)	4	1.90 (1.39; 2.73)	93
C. Valenciana	3.4 (2.7; 4.2)	970	0.57 (0.46; 0.72)	2,235 (197; 5,772)	29	1.31 (1.06; 1.65)	1265
Extremadura	4.3 (3.7; 5.0)	443	0.97 (0.83; 1.12)	965 (72; 2,480)	16	2.10 (1.81; 2.44)	522
Galicia	2.4 (1.8; 3.2)	600	0.93 (0.70; 1.23)	1,148 (64; 3,972)	10	1.77 (1.33; 2.36)	548
Madrid	8.2 (7.3; 9.1)	2757	0.51 (0.46; 0.57)	2,930 (793; 6,373)	61	0.54 (0.48; 0.60)	173
Melilla	8.4 (6.1; 11.3)	39	0.54 (0.40; 0.74)	135 (6; 260)	5	1.86 (1.38; 2.56)	96

Murcia	4.0 (2.8; 5.4)	478	0.80 (0.60; 1.14)	785 (58; 2,212)	13	1.31 (0.97; 1.88)	307
Navarra	7.9 (6.8; 9.1)	361	0.70 (0.61; 0.81)	472 (24; 1,474)	7	0.91 (0.79; 1.06)	111
País Vasco	4.4 (3.6; 5.3)	1038	1.07 (0.89; 1.31)	1,311 (174; 3,494)	25	1.35 (1.12; 1.65)	273
La Rioja	5.2 (3.4; 7.6)	183	1.12 (0.76; 1.71)	341 (24; 956)	12	2.07 (1.42; 3.17)	158

771   Table 3. Date of first infection for REMEDID estimated daily infections from COVID-

772   19 deaths (IR_O) and from MoMo Excess Deaths (ED) (IR_M), and for official COVID-19

773   daily infections released on June 2020 (IO_{20}) and on February 2021 (IO_{21}). All dates

774   refer to year 2020.

	First infection			
	IR_O (CI 95%)	IR_M (CI 95%)	Official COVID-19 data	
			IO_{20}	IO_{21}
Spain	8 Jan (8-8 Jan)	9 Jan (9–10 Jan)	20 Feb	1 Jan
Andalucía	18 Jan (17-18 Jan)	12 Jan (12-13 Jan)	26 Feb	2 Feb
Aragón	21 Jan (20-22 Jan)	8 Jan (8 Jan-8 Jan)	4 March	11 Feb
Asturias	28 Jan (26-29 Jan)	20 Jan (18 Jan-21 Jan)	29 Feb	23 Feb
Islas Baleares	27 Jan (24-29 Jan)	14 Jan (11 Jan-17 Jan)	20 Feb	29 Jan
Canarias	22 Jan (20-23 Jan)	8 Jan (8 Jan-8 Jan)	20 Feb	29 Jan
Cantabria	28 Jan (26-30 Jan)	11 Jan (9 Jan-13 Jan)	29 Feb	23 Feb
Castilla - La Mancha	17 Jan (16-17 Jan)	14 Jan (13 Jan-14 Feb)	29 Feb	24 Feb
Castilla y León	19 Jan (19-20 Jan)	17 Jan (16-17 Jan)	28 Feb	20 Jan
Cataluña	16 Jan (16-17 Jan)	10 Jan (10 Jan-11 Feb)	25 Feb	3 Jan
Ceuta	16 Feb (11-21 Feb)	21 Jan (16 Jan-26 Jan)	14 March	3 March
C. Valenciana	9 Jan (8-11 Jan)	12 Jan (11 Jan-13 Jan)	25 Feb	1 Feb
Extremadura	25 Jan (24-26 Jan)	12 Jan (10 Jan-13 Jan)	29 Feb	8 Feb
Galicia	25 Jan (24-26 Jan)	14 Jan (13 Jan-15 Jan)	3 March	1 Feb
Madrid	11 Jan (10-11 Jan)	9 Jan (8 Jan-9 Jan)	25 Feb	1 Jan
Melilla	2 Feb (31 Jan-4 Feb)	17 Jan (15 Jan-19 Jan)	12 March	5 March
Murcia	28 Jan (26-29 Jan)	13 Jan (10 Jan-15 Jan)	7 March	22 Feb
Navarra	24 Jan (23-25 Jan)	8 Jan (8 Jan-8 Jan)	29 Feb	10 Feb
País Vasco	18 Jan (17-18 Jan)	9 Jan (9 Jan-11 Feb)	28 Feb	12 Feb
La Rioja	27 Jan (25-28 Jan)	14 Jan (12 Jan-15 Jan)	1 March	5 Jan

779 Table 4. Date of the most prominent relative maxima, for Spain and the 19 regions, of

780 the REMEDID estimated daily infections from COVID-19 deaths (IR_O) and from

781 MoMo Excess Deaths (*ED*) (IR_M), and official COVID-19 daily infections (IO_{21}).

782 Maxima of IO_{21} were estimated from the 14-day running mean time series.

	Relative Maxima in 2020		
	IR_O	IR_M	IO_{21}
Spain	13 March, 22 October	13 March, 18 July, 30 Aug, 19 Oct	20 March, 16-17 Sept, 26 Oct
Andalucía	15 March, 21 October	12 March, 19 October	20 March, 26 October
Aragón	19 March, 27 July, 24 Oct	16 March, 25 July, 21 Oct	23 March, 2 Aug, 26 Oct
Asturias	25-26 March, 2-3 Nov	12-14 March, 22-23 March, 25 July, 24 Oct	20 March, 9 Nov
Islas Baleares	17 March, 16 Aug, 5 Sept, 20-21 Oct	12 March, 30-31 July, 3 Oct	20 March, 17 Aug, 8 Nov, 20-21 Nov
Canarias	11 March, 26-27 Aug	14 March, 12 Aug	19 March, 29 Aug
Cantabria	17 March, 2 Sept, 26 Oct, 16 Nov	17 March	22 March, 4 Sept, 8 Nov
Castilla - La Mancha	12 March, 19 Oct	12 March, 18 July, 15 Oct	20 March, 17 Sept, 29 Oct
Castilla y León	15 March, 25 Oct	14 March, 21-22 Oct	21 March, 26 Oct
Cataluña	14 March, 12 Oct	15 March, 22 July, 19 Oct	20 March, 26 Oct
Ceuta	17-19 March, 16-17 May, 25-26 Aug, 15-16 Oct	22-24 July, 24-26 Aug, 14-15 Oct	23 March, 29 Oct
C. Valenciana	15 March, 1 Sept, 31 Oct	13 March, 18 July, 24 Nov	19 March, 24 Aug, 2 Nov
Extremadura	14 March, 12-13 Sept, 19 Oct	14 March, 3 July, 13 Oct	22 March, 18 Sept, 17 Oct
Galicia	18 March, 28-29 Aug, 25 Oct	17 March, 8-9 Oct	22 March, 29 Aug, 27 Oct
Madrid	11 March, 9 Sept	11 March, 18 July, 8 Sept	28 March, 16 Sept
Melilla	11 March, 20 Aug, 16 Oct	9-10 March, 17-18 July, 15 Oct	17 March, 8 Sept, 22 Oct
Murcia	14-15 March, 28-29 Aug, 19-20 Oct	11-12 March, 17 April, 18-19 Aug, 17-18 Oct	19 March, 18 Sept, 30 Oct

Navarra	16 March, 21 Oct	14 March, 23 Sept, 17 Oct	19 March, 17 Sept, 20 Oct
País Vasco	18 March, 31 Aug, 30-31 Oct	15-16 March, 30 Aug, 16 Oct	20 March, 30 Aug, 1 Nov
La Rioja	18 March, 3-4 Sept, 22-23 Oct	13-14 March, 21-22 Aug, 10-11 Oct	21 March, 14 Sept, 26 Oct

 Table 5. REMEDID estimated daily infections from COVID-19 deaths (IR_O) and from

788 MoMo Excess Deaths (ED) (IR_M) on 14 March, and for official COVID-19 daily

789 infections released on June 2020 (IO_{20}) and released on February 2021 (IO_{21}). Infections

790 are inferred from COVID-19 deaths and MoMo ED with their corresponding CFR and

791 95% confidence interval detailed in Table 1. Percentages are official COVID-19

792 infections as proportions of the estimated infections.

	Daily infections on 14 March 2020				Percentage of detected daily infections (%)			
	IR_D (CI 95%)	IR_M (CI 95%)	Official COVID-19		IO_{20}		IO_{21}	
			IO_{20}	IO_{21}	IR_D (CI 95%)	IR_M (CI 95%)	IR_D (CI 95%)	IR_M (CI 95%)
Spain	63,727 (60,050; 67,403)	77855 (73364; 82347)	1832	7478	2.9 (2.7; 3.1)	2.4 (2.2; 2.5)	11.7 (11.1; 12.5)	9.6 (9.1; 10.2)
Andalucía	7,144 (6,192; 8,573)	6788 (5883; 8145)	168	396	2.4 (2.0; 2.7)	2.5 (2.1; 2.9)	5.5 (4.6; 6.4)	5.8 (4.9; 6.7)
Aragón	1,599 (1,232; 2,065)	1804 (1391; 2330)	26	107	1.6 (1.3; 2.1)	1.4 (1.1; 1.9)	6.7 (5.2; 8.7)	5.9 (4.6; 7.7)
Asturias	358 (245; 490)	346 (237; 473)	45	81	12.6 (9.2; 18.4)	13 (9.5; 19)	22.6 (16.5; 33.1)	23.4 (17.1; 34.2)
Islas Baleares	412 (235; 706)	307 (175; 526)	14	69	3.4 (2.0; 6.0)	4.6 (2.7; 8)	16.7 (9.8; 29.4)	22.5 (13.1; 39.4)
Canarias	1,385 (903; 2,108)	1086 (708; 1653)	19	96	1.4 (0.9; 2.1)	1.7 (1.1; 2.7)	6.9 (4.6; 10.6)	8.8 (5.8; 13.6)
Cantabria	597 (403; 871)	501 (338; 730)	20	59	3.4 (2.3; 5.0)	4 (2.7; 5.9)	9.9 (6.8; 14.6)	11.8 (8.1; 17.5)
Castilla - La Mancha	5,303 (4,309; 6,408)	6458 (5247; 7804)	112	545	2.1 (1.7; 2.6)	1.7 (1.4; 2.1)	10.3 (8.5; 12.6)	8.4 (7; 10.4)
Castilla y León	4,688 (4,087; 5,410)	5871 (5119; 6775)	54	474	1.2 (1.0; 1.3)	0.9 (0.8; 1.1)	10.1 (8.8; 11.6)	8.1 (7; 9.3)
Cataluña	11,795 (9,996; 14,194)	13854 (11740; 16671)	184	1482	1.6 (1.3; 1.8)	1.3 (1.1; 1.6)	12.6 (10.4; 14.8)	10.7 (8.9; 12.6)

Ceuta	23 (10; 59)	9 (4; 23)	1	3	4.3 (1.7; 10.0)	11.1 (4.3; 25)	13.0 (5.1; 30)	33.3 (13; 75)
C. Valenciana	3,476 (2,752; 4,490)	3555 (2814; 4592)	279	463	8.0 (6.2; 10.1)	7.8 (6.1; 9.9)	13.3 (10.3; 16.8)	13 (10.1; 16.5)
Extremadura	981 (696; 1,361)	883 (627; 1225)	29	71	3.0 (2.1; 4.2)	3.3 (2.4; 4.6)	7.2 (5.2; 10.2)	8 (5.8; 11.3)
Galicia	1,476 (1,165; 1,865)	1443 (1139; 1823)	80	244	5.4 (4.3; 6.9)	5.5 (4.4; 7)	16.5 (13.1; 20.9)	16.9 (13.4; 21.4)
Madrid	19,565 (17,224; 22,241)	24315 (21405; 27640)	604	2622	3.1 (2.7; 3.5)	2.5 (2.2; 2.8)	13.4 (11.8; 15.2)	10.8 (9.5; 12.2)
Melilla	139 (94; 208)	46 (31; 68)	5	3	3.6 (2.4; 5.3)	10.9 (7.4; 16.1)	2.2 (1.4; 3.2)	6.5 (4.4; 9.7)
Murcia	867 (542; 1,464)	446 (279; 752)	24	84	2.8 (1.6; 4.4)	5.4 (3.2; 8.6)	9.7 (5.7; 15.5)	18.8 (11.2; 30.1)
Navarra	1,165 (900; 1,500)	1447 (1118; 1863)	37	121	3.2 (2.5; 4.1)	2.6 (2; 3.3)	10.4 (8.1; 13.4)	8.4 (6.5; 10.8)
País Vasco	1,965 (1,583; 2,566)	2269 (1828; 2962)	109	421	5.5 (4.2; 6.9)	4.8 (3.7; 6)	21.4 (16.4; 26.6)	18.6 (14.2; 23)
La Rioja	314 (238; 416)	266 (201; 352)	22	137	7.0 (5.3; 9.2)	8.3 (6.3; 10.9)	43.6 (32.9; 57.6)	51.5 (38.9; 68.2)

800   **Figure 1.** Probability density functions (PDF) of incubation period (*IP*, blue line) and

801   illness onset to death period (*IOD*, red line) from Linton et al. (2020). Green line, *f(t)*, is

802   the convolution of both and represents the PDF of infection to death. Black dotted line

803   is a lognormal PDF with the same mean and median as *f(t)*.

813 Figure 2. COVID-19 deaths and MoMo Excess Deaths (*ED*). a) Spain; b-t) Spanish

814 regions. Red lines are official COVID-19 deaths, solid black lines are MoMo deaths,

815 and thin black lines delimit MoMo death error bands (99% confidence interval).

821

Figure 3. Official COVID-19 infections and deaths, and estimated infections with case

fatality ratio (CFR) of 100% in Spain during the first wave. Left y-axis: COVID-19

daily infections IO_{21} (blue curve). Right y-axis: COVID-19 deaths (orange curve) and

its REMEDID-estimated infections with CFR=100% (red curve). All curves are for

Spain. Thin blue and orange curves are daily data, and thick curves are smoothed by 14-

day running mean. Arrows show delays between the maximum of inferred infections

and maxima from COVID-19 deaths (orange arrows) and COVID-19 infections (blue

arrows). Solid arrows are expected delays, dotted while arrows are observed delays.

Figure 4. Daily and accumulated infections for official COVID-19 daily infections

(IO_{21}), and daily infections estimated from COVID-19 deaths (IR_O). Lines are daily

infections and refer to the y-axis on the right; bars are accumulated infections and refer

to the y-axis on the left. Red lines and cyan bars are official COVID-19 data; orange

lines and blue bars are inferred infections with case fatality ratio (CFR) in Table 1. Thin

orange lines correspond to the CFR confidence interval.

Figure 5. Official COVID-19 infections, MoMo Excess Deaths (*ED*), and estimated

infections with case fatality ratio (CFR) of 100% in Spain during the first wave. Left y-

axis: COVID-19 daily infections IO_{21} (blue curve). Right y-axis: MoMo *ED* (orange

curve) and its REMEDID-estimated infections with CFR=100% (red curve). All curves

are for Spain. Thin blue and orange curves are daily data, and thick curves are smoothed

by 14-days running mean. Dashed curves represent the error estimate of MoMo *ED*

(orange) and inferred infections (red). Arrows show delays between the maximum of

inferred infections and maxima from MoMo *ED* (orange arrows) and COVID-19

infections (blue arrows). Solid arrows are expected delays, dotted while arrows are

observed delays.

Figure 6. Daily and accumulated infections for official COVID-19 daily infections

(IO_{21}), and daily infections estimated from MoMo Excess Deaths (*ED*) (IR_M). Lines are

daily infections and refer to the y-axis on the right; and bars are accumulated infections

and refer to the y-axis on the left. Red lines and cyan bars are for official COVID-19

data; and orange lines and blue bars are for inferred infections with case fatality ratio

(CFR) in Table 1. Thin orange lines represent the error estimate of inferred infections.

886   Figure A.1. REMEDID validation test. Thin blue line is official infections and thick

887   blue line is its 14-day-running mean; green line is official deaths; and red line is

888   REMEDID applied with the PDF of illness onset to death, *h(t)*.