

Hydroxychloroquine serum concentrations in non-critical care patients infected with SARS-CoV-2

Alasdair MacGowan^{1*}, Fergus Hamilton¹, Mark Bayliss¹, Liam Read¹, Marie Attwood¹, Alan Noel¹, Sally Grier¹, Anna Morley², David Arnold², Nicholas Maskell²

1. Bristol Centre for Antimicrobial Research and Evaluation (BCARE), Infection Sciences, Pathology Sciences Building – Phase 2, Science Quarter, Southmead Hospital, Westbury-on-Trym, Bristol, BS10 5NB, UK

2. Academic Respiratory Unit, Second Floor, Learning & Research Building, Science Quarter, Southmead Hospital, Westbury-on-Trym, Bristol, BS10 5NB, UK

*Corresponding Author

e-mail: alasdair.macgowan@nbt.nhs.uk

Tel: +44 (0) 117 414 6215

26 **Abstract**

27

28 Hydroxychloroquine(HCQ) has been widely used to treat SARS-CoV-2 infection however
29 HCQ pharmacokinetics in this condition have not been studied in non-critical care patient
30 groups. Here we report the serum concentrations of HCQ in a small cohort of patients
31 treated with HCQ as part of the RECOVERY trial.

32

33 **Introduction**

34

35 Hydroxychloroquine (HCQ) has been widely used to treat infection with SARS-CoV-2 but
36 most recently has been shown to be of no benefit in a large randomised clinical trial
37 (RECOVERY 2020). Detailed information on the pharmacodynamics of HCQ is lacking but
38 measures of the *in vitro* activity of HCQ against SARS-CoV-2 are available as well as
39 pharmacokinetic data derived from analysis of patients with conditions other than
40 SARS-CoV-2 infection who received HCQ. HCQ is much less active against SARS-CoV-2 *in*
41 *vitro* than it is against Plasmodium species. Pharmacokinetic data from patients with
42 SARS-CoV-2 is sparse and absent for those outside critical care (Perinel et al, 2020). Here we
43 report data on the serum concentrations of HCQ in patients who were recruited into the
44 Randomised Evaluation of COVID-19 Therapy (RECOVERY) Trial. Permission to take and use
45 stored serum was ethically approved as part of the Diagnostic and Severity markers in
46 COVID-19 to Enable Rapid triage (DISCOVER) Study. Patients received HCQ at a dose of
47 800mg 6 hrly apart, then 400mg 6 hours later followed by 400mg 12 hrly subsequently.

48

49

50 **Methods and Results**

51

52 HCQ was assayed using a recently developed LC-MS/MS method. Serum was deproteinised
53 with acetonitrile after addition of the internal standard, chloroquine. The method was
54 validated over a working range of 0.005 to 1 mg/L. There was no evidence of drug loss in gel
55 separated tubes compared to plain tubes or after 24 hr at room temperature. The mean
56 blood/plasma partitioning ratio was 1.64 ± 0.25 .

57

58 Seven patients were recruited into the HCQ treatment arm. Subsequently one patient
59 (male, aged 58 years) had no blood samples drawn and a second (male, aged 84 years) only
60 received a single dose of 800mg. Of the five remaining, two were male, three female with a
61 mean \pm std (min-max) age of 57.4 ± 20.7 (26-83) and weight of 70.6 ± 4.2 (65.4-75.0) kgs. All
62 had eGFRs >80 ml/min on day 0 and none had serum bilirubin, alkaline phosphatase or
63 alamine aminotransferase enzyme levels of more than three times normal. Three of the
64 five had co-morbidities – that is, cardiac, neurological or neoplastic disease. All HCQ serum
65 concentrations were ≤ 0.005 mg/L either before or on the first day of dosing. Serum
66 concentrations on subsequent days after starting treatment are shown on Figure 1.

67

68 Concentrations on day 1, 2, 3 and 4 were 0.4 ± 0.35 mg/L (n=3); 0.43 ± 0.14 mg/L (n=3), 0.46
69 mg/L (n=2), and 0.62 mg/L (n=2) respectively.

70

71 The timing of sampling after oral dose was variable – 2.9 ± 3.5 (0.12-8.5 hours), though
72 three quarters were taken within 4 hours of the dose being administered.

73

74 **Discussion**

75

76 Data on the pharmacokinetics of HCQ has been accumulated in patients with conditions
 77 other than SARS-CoV-2 infection. The European SmPC suggest that after a single 400mg
 78 dose, the peak serum concentration was 0.105mg/L at 1.8 hr post dose (SmPC, 2017).
 79 Pharmacokinetic and modelling data exists for SARS-CoV-2 infected patients . Yoa et al,
 80 2020 simulated several potential doses of HCQ suggesting that in Chinese subjects (usually
 81 smaller and lighter than European or North American patients) that a dose of 400mg BD and
 82 200mg BD may be optimal. Interestingly, mean serum HCQ concentrations were not
 83 predicted to be >1mg/L until 4-5 days therapy with this regimen. Downes et al, 2020
 84 simulated doses more akin to those used in the RECOVERY trial –that is, 1600mg on day 1
 85 and 400mg/day thereafter. The predicted Cmax (IQR) was 2.7mg/L (2.2-3.3) and AUC
 86 109.9mg/L.h. Perinel et al, 2020 performed a study of HCQ serum concentrations in French
 87 patients in critical care receiving 200mg TDS; most were mechanically ventilated. It took 2-7
 88 days for serum concentration to reach 1mg/L. They also suggested a reference range of
 89 >1mg/L and <2mg/L but only 8 out of 13 patients reached these concentrations. HCQ
 90 dosing in the RECOVERY trial is higher than those in these studies and, despite this, all our
 91 patients, none of whom were receiving critical care, had HCQ concentration of <1mg/L. It
 92 may be expected that patients not receiving critical care may have better absorption than
 93 those in ICU.

94

95 There are a number of limitations of our study – most obviously, the small number of
 96 patients and samples available. However, our data indicates that HCQ concentrations are
 97 lower than might be expected, even with a high dose regimen.

98

99 Given that HCQ is likely to have marginal pharmacodynamics in SARS-CoV-2 infection, this

100 data may in part help explain the lack of benefit in randomised clinical trials.

101

102 **Funding**

103 This study was supported by the Severn Infection Sciences Partnership, Southmead

104 Research Foundation, and North Bristol NHS Trust.

105

106 **Transparency declarations**

107 Bristol Centre for Antimicrobial Research & Evaluation (BCARE) receives grant funding for

108 research on antimicrobials from Paratek, Wockhardt, InfectoPharm, Venatorx, Merck,

109 AiCURIS, NosoPharm, Evotec, IMI JU, COMBACTE MAGNET and GNA NOW, MRC(UK) and

110 NIHR(UK).

111

References

- Statement from the Chief Investigators of the Randomised Evaluation of COVID-19 thERapY (RECOVERY) Trial on hydroxychloroquine, 5 June 2020.
<https://www.recoverytrial.net/files/hcq-recovery-statement-050620-final-002.pdf>
- Perinel S, Launay M, Botelho-Nevers E, Diconne E, Louf-durier A, Lachand R, Murgier M, Page D, Vermesch R, Thierry G, Delavanne X. Towards Optimization of Hydroxychloroquine Dosing in Intensive Care Unit COVID-19 Patients. *Clin Inf Dis* 2020 published online.
<https://doi:10.1093/cid/ciaa394/5816960>
- Summary of Product Characteristics (SmPC). Quinoric 200mg Film-Coated Tablets 2017.
<https://www.medicines.org.uk/emc/product/477/smpc/print>
- Yao X, Ye F, Zhang M, Cui C, Huang B, Niu P, Liu X, Zhao L, Dong E, Song C, Zhan S, Lu R, Li H, Tan W, Liu D. In vitro Antiviral Activity and Projection of Optimized Dosing Design of hydroxychloroquine for the Treatment of Severe Acute Respiratory Syndrome Corona virus 2 (SARS-CoV-2). *Clin Inf Dis* 2020. <https://doi/10.1093/cid/ciaa237/5801998>
- Downes KJ, Chiotos K, Fitzgerald JC, Scheetz MH, Zuppa AF. Rational Dosing of Hydroxychloroquine for the Treatment of COVID-19. *Open Science Framework PrePrints*
<https://osf.io/py3kv/>

FIGURE 1 Serum concentrations of hydroxyl chloroquine

