1 **Only a combination of social distancing and massive testing can**

2 **effectively stop COVID-19 progression in densely populated urban areas**

Mario Moisés Alvarez and Grissel Trujillo-de Santiago[1,3]

[1] Centro de Biotecnología-FEMSA, Tecnologico de Monterrey, Monterrey 64849, NL, México

[2] Departamento de Bioingeniería, Escuela de Ingeniería y Ciencias, Tecnologico de Monterrey,

Monterrey 64849, NL, México

[3] Departamento de Ingeniería Mecatrónica y Eléctrica, Escuela de Ingeniería y Ciencias,

Tecnologico de Monterrey, Monterrey 64849, NL, México

(*) corresponding author: *mario.alvarez@tec.mx*

13 **Abstract**

We present a simple epidemiological model that includes demographic density, social
distancing, and efficacy of massive testing and quarantine as main parameters to model the
progression of COVID-19 pandemics in densely populated urban areas (i.e., above 10,000
hab km^2). Our model demonstrates that effective containment of pandemic progression in
densely populated cities is achieved only by combining social distancing, widespread
testing, and quarantining of infected subjects. This finding has profound epidemiological
significance, and sheds light on the controversy regarding the relative effectiveness of
widespread testing and social distancing. Our simple epidemiological simulator can also be
used to assess the efficacy of a governmental/societal response to an outbreak.
This study has also relevant implications on the concept of smart cities; densely populated
areas are hot spots highly vulnerable to epidemic crisis.

Keywords: *COVID-19, SARS-CoV2, mathematical modeling, social distancing, massive*

testing, pandemic, demographic density

MedRxiv

Introduction

COVID-19 has clearly illustrated that we were unprepared for the second pandemic of the 21st century. By the fourth week of June 2020, the official cumulative number of infected persons worldwide was more than 9,000,000, with a death toll higher than 450,000. COVID-19 has not recognized frontiers; it started in China, migrated to Iran, and formed a later epicenter in Italy and Spain. It then extended to France, England, Germany, and other European Countries. Now, COVID-19 has a strong presence in Las Americas, mainly in the USA[1], Brazil, and Mexico. Among all affected territories, pandemic COVID-19 has encountered containment responses ranging from aggressive to mediocre. Some of these responses, namely the ones that appear to be more successful, have been based on a combination of social distancing and massive testing and selective quarantine[2]. Social distancing effectively decreases the progression of the disease essentially by decreasing the demographic density, while massive testing enables the identification of infected subjects and opportune quarantining (which is technically a version of selective social distancing). However, discussion persists regarding the relative effectiveness of countermeasures such as social distancing or massive testing. Countries like South Korea have based their strategy of containment on strict social distancing and massive testing in their open populations. Spain initially based its strategy on gradual social distancing. Later, it significantly increased its diagnostic efforts to plateau the pandemic. England started by using testing mainly for epidemiological recording, but later significantly scaled up its efforts to aggressively diagnose its population. By contrast, Mexican officials have openly declared that massive testing and quarantine have no significant value as a countermeasure for COVID-19 progression and that diagnostics is merely informative. Consequently, Mexico

is one of the countries with the fewest diagnostic tests conducted per 100,000 inhabitants,

according to OCDE (Organisation de coopération et de développement économiques).

The benefit of social distancing and massive diagnostics has not only been a subject of

controversy(*3*), but assessing the potential benefit has also been challenging. Here, we

introduce a simple mathematical model with a formulation that explicitly considers

demographic variables (i.e., the demographic density and the total population) to calculate

the progression of COVID-19 in urban areas. The model also considers the effectiveness of

social distancing measures and of massive testing for expeditious identification and

quarantining of infective subjects as inputs. We present a wide range of simulation

scenarios for "representative" urban areas and show that both social distancing and

widespread and effective testing should be combined for effective containment of the

pandemic advance in densely populated urban areas.

Rationale of the model formulation

We developed a very simple epidemiological model for the propagation of COVID-19 in

urban areas. The model considers two variable populations of individuals, infected (X) and

retrieved (R). The cumulative number of infected patients (X) is the total number of

subjects among the population that have been infected by SARS-CoV-2. The number of

retrieved patients should be interpreted as the number of individuals that have been

retrieved from the general population and are not contributing to the propagation of

COVID-19. Retrieved subjects include subjects who have recovered from the infection and

do not shed virus, quarantined individuals, and deceased patients. Importantly, the model

assumes that infection results in (at least) short-term immunity upon recovery. This

assumption is based in experimental evidence that suggests that rhesus macaques that

recovered from SARS-CoV-2 infection could not be reinfected(*4*). However, the

acquisition of full immunity to reinfection has not been confirmed in humans, although it is

well documented for other coronavirus infections, such as SARS and MERS(*5*, *6*).

Two set of parameters, demographic and clinical/epidemiological, determine the interplay

between these two main populations and other subpopulations that include asymptomatic

infected (A), symptomatic infected (S), and deceased (D). Clinical parameters include an

intrinsic infection rate constant (μ_o) that is calculated from the initial stage of the pandemic

in that particular region, the fraction of asymptomatic patients (a), the delay between the

period of viral shedding by an infected patient (delay_r), the period from the onset of

shedding to the result of first diagnosis and quarantine in the fraction of patients effectively

diagnosed (delay_q), and the fraction of infected patients effectively diagnosed and

retrieved from the population (α). Demographic parameters include the population of the

region (P_o), the relative demographic density (δ), the social distancing (σ), and the fraction

of infected individuals retrieved from the population due to massive and effective testing

(α). The model is based on a set of two simple differential equations.

95               $dX/dt = \mu_o\, \delta\, (1-\sigma)\, (X-R)\, (P_o-X)/P_o$ equation (1)

96               $dR/dt = \alpha \int_{t=0}^{t=t-delay\_q} dX/dt - (1-\alpha) \int_{t=0}^{t=t-delay\_r} dX/dt$ equation (2)

The first equation of the set (equation 1) states that the rate of accumulation of infected

habitants (symptomatic and asymptomatic) in an urban area (assumed to be a closed

system) is proportional to the number of infective subjects (X-R) present in that population

at a given point and the fraction of the population susceptible to infection ((P_o-X)/P_o). Note

that the number of infective subjects is given by the difference between the accumulated

number of infected subjects (X) and the number of retrieved subjects (R). The fraction of

the susceptible population decreases over time as more inhabitants in the community get

infected. The proportionality constant in equation 1 (μ_o) is an intrinsic rate of infection that

is weighted by the population density (δ) in that urban area, and the effective fractional

reduction of social distancing on the population density (1- σ).

The second equation (equation 2) describes the rate at which infected patients are retrieved

from the infective population. Eventually, all infected subjects are retrieved from the

population of infected individuals, but this occurs at distinctive rates. A fraction of infected

individuals (α) is effectively retrieved from the general population soon after the onset of

symptoms or after a positive diagnosis. Another fraction of infected subjects (1- α) is not

effectively retrieved from the population until they have recovered or died from the disease.

Therefore, in our formulation, the overall rate of retrieval (dR/dt) has two distinctive

contributions, each one associated with different terms on the right-hand side of equation 2.

The first term accounts for the active rate of retrieving infected patients through the

diagnosis and quarantine of SARS-CoV-2 positive subjects. For this term, the delay from

the onset of virus shedding to positive diagnosis and quarantine (delay_q) is considered

short (*i.e.*, between two or three days), to account for a reasonable time between the

positive diagnosis and the action of quarantine. In our model formulation, this term is

represented by (1-α). A second term relates to the recovery or death of infected patients

(symptomatic or asymptomatic) and is represented by the integral of all infected subjects

recovered or deceased from the onset of the epidemic episode in the region, considering a

delay of 21 days (delay_r), which accounts for the average time of recovery of an infected

individual.

Note that the simultaneous solution of equations 1 and 2 is sufficient to describe the

evolution of the number of asymptomatic individuals (A), symptomatic individuals (S), and

deceased patients (D) through the specification of several constants and simple relations.

$$a\ dX/dt = dA/dt \qquad \text{equation (3)}$$

$$(1-a)\ dX/dt = dS/dt \qquad \text{equation (4)}$$

$$m\ [(1-a)\ dX/dt\ ] = dD/dt \qquad \text{equation (5)}$$

Here, **a** is the fraction of asymptomatic subjects among the infected population, (1-**a**) is the

fraction of infected individuals that exhibit symptoms, and **m** is the mortality rate expressed

as a fraction of symptomatic individuals.

136 Selection of relevant epidemiological parameters for COVID-19

As with any epidemiological model, this model relies on some basic assumptions that must

be sustained in clinical or epidemiological data. We now briefly discuss the assumptions

that were made and the rationale behind the relevant values of the parameters of the model:

the fraction of asymptomatic infected, the average time to recover, the fraction of

symptomatic patients that would require hospitalization, and the average time of bed

occupancy per hospitalized patient.

The fraction of asymptomatic infected is one of the critical inputs to the model; it

determines the final and maximum feasible threshold of symptomatic infected. However,

the current evidence is not yet sufficient to support a conclusive value for this parameter.

Nevertheless, a recent serological study conducted in New York City (NYC) found anti-

SARS-CoV-2 IGGs among 21.2% of the population (www.360dx.com; www.cnn.com),

and this result is consistent with previous information related to massive epidemic episodes.

For instance, in the context of the pandemic influenza A/H1N1/2009, up to 20–40% of the

population in urban areas (i.e., Monterrey, México, and Pittsburgh, USA) (*7*, *8*) exhibited

specific antibodies regardless of experiencing symptoms, while the fraction of confirmed

symptomatic infections was lower than less than 10%. This serological result based

exclusively on information from NYC suggests that more than 90% of exposed New

Yorkers (~91.4%) were asymptomatic or exhibited minor symptoms. Based on this (still

unpublished) data, we assumed a symptomatic fraction of only 10% in the calculations and

forecasts presented here.

In addition, the average time of sickness was set at 21 days in our simulations, as this is

within the reported range of 14 to 32 days(*9*), with a median time to recovery of 21

days(*10*). Viral shedding can last for three to four weeks after the onset of symptoms, with

a peak at day 10-11.(*11*) Therefore, we assume that all those infected not quarantined

could continue to transmit the virus until full recovery (21 days). Similarly, asymptomatic

patients are only removed from the pool of susceptible persons after full virus clearance.

Note that, in the current version of our model, asymptomatic patients are considered part of

the population capable of transmitting COVID-19; reported evidence that suggests that

asymptomatic subjects (or minimally symptomatic patients) may exhibit similar viral

loads(*12*) to those of symptomatic patients and may be active transmitters of the disease(*13*,

14).

The average fraction of deceased patients worldwide is estimated as 0.064 of those infected

21 days before. This mortality percentage (case fatality rate) lies within the range reported

in the recent COVID-19 literature(*15–18*). The time lapse of 14 days between the onset of

disease and death was statistically estimated by Linton et al. in a recent report (*19*). We also

consider that the average time for bed occupancy of hospitalized patients is 14 days. The

estimated average hospitalization stays range from 9.3 to 13 days in the United States (*20*)

and China(*21*, *22*), but much longer stays have been reported in intensive care units in Italy

(20 to 25 days)(*23*). Anecdotal data collected in México suggests that hospitalization stays

of at least two weeks are a more accurate figure for Latin American societies.

Table 1. Specific infection rates (μ_o) and the associated doubling times (t_d) for COVID-19
infection in different geographic regions.
180

Territory	Demographic density (hab km^2)	μ (day^{-1})	t_d (day)
Spain (Madrid)	5,185	0.358	1.937
NYC	10,194	0.720	0.968

Definition of representative scenarios

We aimed to reproduce representative settings for COVID-19 progression; therefore, we

selected two hypothetical but realistic urban scenarios. Figure 1 shows the most densely

populated cities in the world (according to information concentrated in

www.citymayors.com) represented by dots in a bidimensional plot of demographic density

versus population. Two clusters of cities can be readily identified. The first group of 5 to7

important cities in the world (i.e, Porto Alegre, Ankara, Athens, Guadalajara, Monterrey,

Barcelona) have populations between 3.0 and 3.7 million inhabitants and a demographic

density of ~5,000 inhabitants per square kilometer (hab km^2). A second group is centered

on the coordinates of a population of 10 million citizens and a demographic density of

10,000 hab km^2 (i.e., Baghdad, Ho Chi Minh, Bangalore, Tianjin, Kinshasa, Hyderabad).

Based on this analysis, we centered our estimates of COVID-19 progression in these two

classes of "representative" cities (3.5 million inhabitants, 5,000 hab km^2; and 10 million

citizens, 10,000 hab km^2).

Figure 1. Population and demographic density of the world largest cities. (A) Two clusters of cities, referred here as Type I (green cluster) and Type II (red cluster) are indicated in a plot of population versus demographic density. The 51 largest cities in the world, according to www.citymayors.com/, are included. Madrid Spain is indicated in yellow (•), New York City in red (•), and Mexico City in green (•). (B) Natural log of the number of symptomatic versus time in Madrid from March 2 to March 17 (initial stage of the pandemics in Spain). The initial rate of infection (μ_o) can be calculated from the slope of this straight line.

We also based our estimates of μ_o, the intrinsic rate of COVID_19 propagation, on data extracted from the local dynamics of the pandemics in Madrid (a city in our first category) and New York City (a city in our second category). These two cities also belong to a limited number of cities that have generated reliable datasets on the local progression of the

number of COVID-19 positive cases over time. We calculate the value of μ_o (*i.e.*, the

intrinsic rate of infectivity of SARS-CoV2 before interventions) by assuming that the initial

rate of propagation is $d(X)/dt = \mu_o$ [X], where [X] is the number of initially infected

subjects. We then simply calculate the intrinsic rate of infection from the initial slope of a

plot of ln [X] *vs* time, which is a usual procedure for calculation of intrinsic growth rates in

cell culture scenarios under the assumption of first order rate growth dependence.

Following this rationale, we set μ_o for all our simulations. Consistently, we noted that the

initial rate of propagation observed in NYC, a city with twice the demographic density of

Madrid, doubles this μ_o value (Table 1). Therefore, we take the demographic density of

Madrid as a reference (5,185 hab km^2) and assume that multiplying μ_o by the normalized

demographic density (with respect to that in Madrid) is a reasonable procedure for adapting

the model to any urban area.

Effect of social distancing and massive testing

Social distancing has been regarded as the one of the most effective buffering measures for

containment of local COVID-19 epidemics(*24*, *25*). However, the effectiveness of massive

testing, either alone or in combination with social distancing, has not been evaluated

formally.

First, we conducted simulations in which we evaluated the independent impact of different

degrees of social distancing and massive testing in a Type I city (3.5 million inhabitants and

medium demographic density). Figure 2a shows the impact of different degrees of social

distancing at a fixed and basal value of massive testing. In this simulation, a basal value of

social distancing (α=0.1) means that only 10% of the infected patients are diagnosed and

quarantined, while the rest of the infected subjects continue active until recovery. This

strategy is consistent with that adopted by countries that diagnosed essentially only those

subjects who were symptomatic and asked for medical assistance (i.e., México, Chile, and

Bolivia, with fewer than 2 tests per confirmed case). The pandemic progression (number of

cumulative symptomatic cases, new infections per day, and bed occupancy per day) is

indicated with grey curves for a reference scenario with no social distancing and basal level

of testing. Higher degrees of enforcement of social distancing (i.e., such that the

demographic density is effectively reduced by 20, 40, and 60%) are presented with blue,

green, and red lines, respectively. Levels of social distancing of 20% and 40% delay the

pandemic curve by 15 and 30 days, whereas the pandemic progression is successfully

buffered only when social distancing effectively reduces demographic density (and

therefore activity) by 60% for extended times (i.e., for 6 months).

Figure 2b presents predictions for scenarios where no social distancing is implemented, but

where the level of diagnostic testing is increased from a basal value (10% of infected are

diagnosed and quarantined) to situations were 20, 30, and 40% of infected patients

(symptomatic and asymptomatic) are diagnosed within the first 3 days of viral shedding

and quarantined. These levels of massive testing, in the absence of social distancing

enforcement, are not sufficiently effective to buffer the pandemic.

Figure 2c shows a scenario in which social distancing is set at 20% and different testing

emphasis is applied. As before, trends related to the reference scenario of low testing and

no social distance enforcement are indicated in grey. The combination of moderate social

distancing and testing renders better results than any one of the two strategies

independently applied. When social distancing is elevated to a 40% and combined with

more intensive testing efforts, the epidemic peak is dramatically delayed.

258
259
Figure 2. Pandemic progression scenarios for a Type I city (3.5×10^6 citizens) with a medium

demographic density (5,500 inhabitants km^{-2}). The number of new infections per day (thin and

light line curves), the bed occupancy (thick line curves) and the cumulative number of infections

(thin cumulative curve) are presented for each scenario. (a) Scenario I: basal level of testing (α=0.1)

and increasingly higher levels of social distancing: σ=0.0 (gray), σ=0.2 (blue), σ=0.4 (green), and

σ=0.6 (red). (b) Scenario II: social distance not enforced (σ=0.0), and increasing values of testing

α=0.1 (gray), α=0.2 (blue), α=0.3 (green), and α=0.4 (red). (c) Scenario III: social distance modestly

enforced (σ=0.2), and increasing values of testing, α=0.1 (blue), α=0.2 (green), and α=0.3 (red). (d)

Scenario IV: social distance moderately enforced (σ=0.4), and increasing values of testing α=0.2

(blue), α=0.3 (green), and α=0.4 (red). Gray curves correspond to the pandemic progression with

minimum intervention (no social distancing and α=0.1).

A value of 40% social distancing in combination with an effort to identify and quarantine

30% of newly infected subjects delays the peak of maximum bed occupancy from day 25 to

day 75, and lowers the highest demand of beds from 75,000 to fewer than 30,000. Increased

testing at 40% social distancing further contributes to extinguish the epidemic peak.

We also estimated the pandemic progression under the same set of scenarios for a densely

populated city (Type II: 5×10^6 citizens and 10,000 hab km^2). The results similarly indicate

that only a combined strategy of social distancing and scaled-up testing and quarantine may

effectively control the pandemic progression due to "selective social distancing". However,

the higher population leads to increases in the number of cases in the same time frame (let

us say, the first 120 days from pandemic onset) and in the maximum threshold of

symptomatic cases. The higher demographic density also causes a higher rate of transition.

Therefore, the containment strategies need to be stronger than those required for Type I

cities. For example, in the absence of intensified testing, the degree of social distancing

required to buffer the pandemic is much higher in a Type II city ($\alpha=0.75$) than in a type I

city ($\alpha=0.60$). Similarly, only aggressive social distancing interventions combined with

intensified testing, *i.e.*, ($\sigma=0.6$, $\alpha=0.3$) or ($\sigma=0.7$, $\alpha=0.5$) can mitigate the pandemic

progression in larger or denser cities (Type II). A summary of the results for different

combinations of scenarios is presented in Tables 2 and 3. We present scenarios for both

moderately and densely populated cities (Type I and II). Four different indicators are

calculated for each scenario, including the day of the epidemic peak, the number of new

infection cases at the epidemic peak, the cumulative number of symptomatic infections

after 120 days of the local pandemic onset (4 months), and the maximum bed occupancy.

Figure 3. Pandemic progression scenarios for a Type II city (5.0×10^6 citizens) with a high demographic density (10,000 inhabitants km^{-2}). The number of new infections per day (thin and ligth line curves), the bed occupancy (thick line curves) and the cumulative number of infections (thin cumulative curve) are presented for each scenario. (a) Scenario I: basal level of testing (α=0.1) and increasingly higher levels of social distancing: σ=0.00 (gray), σ=0.25 (blue), σ=0.50 (green), and σ=0.75 (red). (b) Scenario II: social distance not enforced (σ=0.0), and increasing values of testing α=0.1 (gray), α=0.3 (blue), α=0.5 (green), and α=0.7 (red). (c) Scenario III: social distance enforced (σ=0.6), and increasing values of testing: α=0.1 (blue), α=0.3 (green), and α=0.5 (red). (d) Scenario IV: social distance moderately enforced (σ=0.7), and increasing values of testing: α=0.1 (blue), α=0.3 (green), and α=0.5 (red). Gray curves correspond to the pandemic progression with minimum intervention (no social distancing and α=0.1).

307 **Table 2.** Effect of social distancing and testing in a city with a demographic density of 5,500 inhabitants/km^2 and a population of 3.5×10^6

308 persons).

σ:			α								
			0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90
	0.00	peak @ day:	38.00	39.00	42.00	45.00	50.00	56.00	64.00	70.00	100.00
		infected after 120 days:	349037.52	349072.31	349410.58	349731.45	349801.94	349008.93	323718.47	134600.03	5963.75
		peak of infection (hab/day)	29155.11	26837.36	24343.10	21633.49	18652.11	15311.81	11469.93	2849.59	90.84
		maximum oc beds:	43076.11	41297.81	39084.22	36297.24	32740.85	28128.49	20667.05	5774.46	190.73
	0.25	peak @ day:	51.00	54.00	59.00	64.00	65.00	70.00	81.00	90.00	90.00
		infected after 120 days:	349994.09	349980.90	349723.77	342776.43	274280.90	122542.87	24923.94	2614.45	248.17
		peak of infection (hab/day)	21114.08	19067.32	16862.56	14462.14	8280.49	2397.63	398.56	37.26	2.59
		maximum oc beds:	35889.86	33437.60	30474.06	25109.72	14935.49	4900.12	833.91	78.27	5.45
	0.5	peak @ day:	67.00	67.00	76.00	81.00	82.00	86.00	88.00		
		infected after 120 days:	175936.83	91893.77	38845.99	13861.23	4332.07	1227.77	341.99		
		peak of infection (hab/day)	3803.14	1649.97	632.51	215.77	64.49	17.03	4.18		
		maximum oc beds:	7523.66	3411.07	1325.73	452.39	135.43	35.79	8.78		
	0.6	peak @ day:	77.00	81.00	83.00	86.00					
		infected after 120 days:	18919.36	8432.49	3534.87	1408.84					
		peak of infection (hab/day)	298.44	129.46	52.56	20.02					
		maximum oc beds:	625.81	271.66	110.40	42.07					
	0.7	peak @ day:	88.00								
		infected after 120 days:	1147.06								
		peak of infection (hab/day)	16.38								
		maximum oc beds:	34.42								

309 (*) Conditions at which the progression can be controlled or extinguished are indicated in light and intense green, respectively.

312 **Table 3.** Effect of social distancing and testing in a city with a demographic density of 10,000 inhabitants/km^2 and a population of 5.0×10^6

313 persons).

			α								
			0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90
σ:	0.00	peak @ day:	19.00	19.00	19.00	20.00	21.00	21.00	22.00	24.00	25.00
		infected after 120 days:	499989.80	499971.06	499917.90	499766.95	499336.94	498102.78	494503.52	483634.61	446996.48
		peak of infection (hab/day)	86423.60	82575.85	78404.48	73841.74	68794.07	63125.83	56629.06	48953.93	39422.43
		maximum oc beds:	73747.96	73345.63	72776.58	71953.49	70731.66	68861.46	65889.88	60938.49	52151.75
	0.25	peak @ day:	25.00	26.00	27.00	28.00	30.00	32.00	35.00	39.00	47.00
		infected after 120 days:	499846.93	499665.38	499268.89	498409.15	496596.85	493100.98	487611.16	479682.50	443985.07
		peak of infection (hab/day)	63930.77	60111.56	55971.34	51440.99	46422.90	40772.27	34255.89	26450.93	16419.13
		maximum oc beds:	70822.64	69733.12	68276.26	66293.98	63540.74	59619.71	53853.41	44986.81	30352.40
	0.50	peak @ day:	38.00	41.00	44.00	46.00	51.00	57.00	64.00	72.00	96.00
		infected after 120 days:	498882.25	499096.74	499540.23	499809.64	499807.31	498253.90	446201.67	158799.47	6043.61
		peak of infection (hab/day)	41649.98	38338.88	34775.60	30904.68	26645.47	21873.49	15986.09	3184.80	92.64
		maximum oc beds:	61537.12	58996.65	55834.29	51852.80	46772.08	40182.73	27909.42	6544.93	194.56
	0.60	peak @ day:	49.00	52.00	56.00	61.00	64.00	67.00	80.00	90.00	90.00
		infected after 120 days:	499990.74	499988.72	499925.44	498000.70	455117.57	259517.01	62505.34	6072.10	390.58
		peak of infection (hab/day)	32518.26	29514.18	26280.30	22762.02	17097.07	6020.73	1047.72	91.03	4.37
		maximum oc beds:	53795.88	50486.51	46503.25	41012.07	29013.61	11822.72	2187.91	191.19	9.19
	0.70	peak @ day:	63.00	65.00	67.00	67.00	76.00	82.00	94.00		
		infected after 120 days:	486459.87	414489.06	263372.49	115959.71	36130.22	8585.78	1674.53		
		peak of infection (hab/day)	21722.22	13239.99	5938.16	2056.64	581.04	130.80	23.44		
		maximum oc beds:	35466.46	23463.24	11668.79	4276.82	1216.45	274.64	49.25		

314 (*) Conditions at which the progression can be controlled or extinguished are indicated in light and intense green, respectively

Next, we investigated the fidelity of the model prediction by simulating the advance of the pandemic in Madrid, NYC, and Mexico City. The number of new infections per day in these three cities has been made available by government officials. Interestingly, the demographic characteristics of these major cities exhibit important differences, as well as the types of counter-measures adopted to contain COVID-19.

Madrid (a Type-I city) and NYC (nearly a Type-II city) have practically extinguished the pandemic. Therefore, the position and magnitude of the pandemic peak offers valuable information to estimate values of the magnitude of social distancing (σ) and diagnostic effort (α). In contrast, Mexico City has not reached the pandemic peak yet, and struggles to contain COVID-19 in a challenging demographic situation. We found sets of parameters that properly describe the evolution of COVID-19 in each of these cities, despite the obvious differences in the behavior of each progression curve (Figure 4). For Madrid, our simulations suggest that social distancing measures have achieved a degree of 58% of reduction in population density (α=0.58). However, even considering these relatively higher values of social distancing, an aggressive testing program (α=0.50) was needed to bend the curve at a cumulative number of ~70,000 cases and reduce the emergence of new cases to the current levels (less than 10 per day). This implies that overall approximately 50% of the active infected subjects were found and quarantined through testing.

We also fitted the model to the pandemic progression observed in NYC. NYC is a densely populated area with a population mark of 8,400,000 inhabitants (i.e., nearly a Type-II city). Therefore, only a combination of social distancing and intensified testing can stop progression effectively (Table II). Our results suggest that NYC has applied a strategy mostly based on aggressive testing.

Figure 4. Progression of the COVID-19 Pandemic in (a) Madrid, (b) New York City (NYC), and (c) Mexico City. Actual data points, as officially reported, are shown using circles. Model predictions of new COVID-19 cases (yellow line) and simulation predictions of the cumulative number of symptomatic patients (blue line). The values of α and σ used in the simulations for each city are indicated within the corresponding pannels.

Indeed, a scenario based on a sustained level of social distancing of 27% efficacy ($\sigma=0.27$;

effective reduction of 30% in demographic density) and a massive testing effort ($\alpha=0.98$) is

the combination that better recapitulates the actual pandemic evolution in NYC. NYC is,

therefore, a remarkable example of the efficacy of massive diagnosis and quarantine, as a

city that was able to stop COVID-19 progression at a cumulative count of ~250,000

symptomatic subjects, which is about 25% of the maximum expected. In perspective, with

an effective distancing of 27% and no aggressive testing (i.e., $\alpha=0.50$ instead of $\alpha=0.98$),

the peak of bed occupancy in the city would have been three-fold higher, causing a total

collapse of the hospital system. Our results also show that demographic density is a key

factor in COVID-19 progression. Note that the testing effort in NYC was more intense than

in Madrid. However, NYC has double the demographic density of Madrid, resulting in a

higher number of those infected than in Madrid (250,000 versus 70,000).

While Madrid and NYC have transitioned through the pandemic peak and successfully

extinguished the pandemic, Mexico City is still in the middle of it. Several mathematical

models have been used to forecast pandemic scenarios for Mexico City (*26*, *27*). However,

the predictions of the progression of COVID-19 and the occurrence of the pandemic peak

in the capital of Mexico have not been accurate. Mexico City has a demographic density of

1.69 times that of Madrid and a population of more than 15,000 within the city limits.

In terms of demographic density, Mexico City lies between a Type-I and Type-II city,

possessing a demographic density 1.69 times greater than Madrid. However, in terms of

population, Mexico City is double that of NYC. The Mexico City Metropolitan Area,

which comprehends the city and its surrounding municipalities in Estado de México,

houses 25,000,000 inhabitants. From the scenarios simulated before (Figure 3), we may

readily infer that only a combination of aggressive social distancing and massive testing

effort can stop pandemic progression in Mexico City. However, Mexican public health

officials have publicly stated their decision to deemphasize diagnostics, instead privileging

social distancing in the entire Mexican territory. Figure 4c compares the actual evolution of

the number of cases in Mexico City versus the prediction of our model for a scenario where

social distancing is set at σ=0.75 (75% effective reduction on demographic density) and the

testing effort at α=0.10% (only 10% of infected individuals are effectively quarantined after

being diagnosed). The Mexican strategy has been successful in delaying the pandemic's

progression, but not in bending the slope of the epidemic curve. As a result, Mexico City is

steadily positioned in a pandemic plateau, with a nearly constant number of cases per day

(600 cases), which is consistent with journalistic information and official data

(https://coronavirus.gob.mx/datos/).

Concluding remaks

Here we introduce a mathematical model, based on demographic and clinical data that

enables evaluating the relative benefit of social distancing and massive testing. Using this

simple model, we investigate scenarios of COVID-19 evolution in two types of

representative cities (i.e., 3,500,000 inhabitants and 5,000 hab km^2, and 5.0 X 10^6

inhabitants and 10,000 hab km^2).

Our modeling simulations show that for Type-I cities, extreme and sustained social

distancing (i.e., effectively decreasing the demographic density by 60% or more) may

extinguish the pandemic progression. However, in Type-II cities (or larger), only

combinations of social distancing and aggressive testing can effectively control the

pandemic evolution. This finding has relevant implications for the planning of

countermeasures for the effective containment of pandemic progression and for the

design/redesign of urban areas. The concept of sustainability and cost-effectiveness of

densely populated urban areas has to be revisited. Our results make explicit that large cities

are highly vulnerable to epidemic crisis.

In principle, this model can be adapted to any urban area by setting the population and the

demographic density. The predictions on the evolution of COVID-19 based on this

mathematical model could represent important tools for designing and/or evaluating

countermeasures.

Acknowledgments

401 MMA and GTdS acknowledge the funding received from CONACyT (Consejo Nacional de

402 Ciencia y Tecnología, México) and Tecnológico de Monterrey.

Author contributions

405 MMA and GTdS collected and analyzed epidemiology data. MMA formulated the model

406 and ran the simulations. MMA and GTdS wrote the manuscript. Both authors reviewed and

407 approved the manuscript.

Competing interest

The authors declare no competing interests.

Supplementary Material:

File S1: Implementation of a mathematical model for COVID-19 epidemic

progression in Excel. Spreadsheet for the calculation of the progression of the number of

COVID-19 infected subjects in a particular geographic region (urban area). Calculation is

based on a simple differential model, epidemiological data reported in literature (as detailed

in the text of the manuscript) and the local demographical density.

References:

1. M. L. Holshue *et al.*, First Case of 2019 Novel Coronavirus in the United States. *N.*

Engl. J. Med. (2020), doi:10.1056/nejmoa2001191.

2. J. Cohen, K. Kupferschmidt, Countries test tactics in "war" against COVID-19.

Science. **367**, 1287–1288 (2020).

3. A. Terriau, J. Albertini, A. Poirier, Q. Le Bastard, Impact of virus testing on

COVID-19 case fatality rate: estimate using a fixed-effects model,

doi:10.1101/2020.04.26.20080531.

4. L. Bao *et al.*, *bioRxiv*, in press, doi:10.1101/2020.03.13.990226.

5. E. Prompetchara, C. Ketloy, T. Palaga, Allergy and Immunology Immune responses

in COVID-19 and potential vaccines: Lessons learned from SARS and MERS

epidemic, doi:10.12932/AP-200220-0772.

6. W. Liu *et al.*, Two-Year Prospective Study of the Humoral Immune Response of

Patients with Severe Acute Respiratory Syndrome. *J. Infect. Dis.* **193**, 792–795

(2006).

7. L. Elizondo-Montemayor *et al.*, Seroprevalence of antibodies to influenza

A/H1N1/2009 among transmission risk groups after the second wave in Mexico, by

a virus-free ELISA method. *Int. J. Infect. Dis.* **15**, e781–e786 (2011).

8. S. M. Zimmer *et al.*, Seroprevalence Following the Second Wave of Pandemic 2009

H1N1 Influenza in Pittsburgh, PA, USA, doi:10.1371/journal.pone.0011601.

9.   L. Lan *et al.*, Positive RT-PCR Test Results in Patients Recovered From COVID-19.

JAMA (2020), doi:10.1001/jama.2020.2783.

10.   Q. Bi *et al.*, *medRxiv*, in press, doi:10.1101/2020.03.03.20028423.

11.   R. Wölfel *et al.*, Virological assessment of hospitalized patients with COVID-2019.

Nature. **581**, 465–469 (2020).

12.   L. Zou *et al.*, SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected

Patients. *N. Engl. J. Med.* **382**, 1177–1179 (2020).

13.   Y. Bai *et al.*, Presumed Asymptomatic Carrier Transmission of COVID-19. *JAMA*

(2020), doi:10.1001/jama.2020.2565.

14.   C. R. MacIntyre, Global spread of COVID-19 and pandemic potential. *Glob.*

Biosecurity. **1** (2020), doi:10.31646/gbio.55.

15.   C. C. Lai, T. P. Shih, W. C. Ko, H. J. Tang, P. R. Hsueh, Severe acute respiratory

syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19):

The epidemic and the challenges. *Int. J. Antimicrob. Agents*. **55** (2020), p. 105924.

16.   S. Jung *et al.*, Real-Time Estimation of the Risk of Death from Novel Coronavirus

(COVID-19) Infection: Inference Using Exported Cases. *J. Clin. Med.* **9**, 523 (2020).

17.   Z. Xu *et al.*, Pathological findings of COVID-19 associated with acute respiratory

distress syndrome. *Lancet Respir. Med.* **0** (2020), doi:10.1016/S2213-

2600(20)30076-X.

18.   R. Porcheddu, C. Serra, D. Kelvin, N. Kelvin, S. Rubino, Similarity in Case Fatality

Rates (CFR) of COVID-19/SARS-COV-2 in Italy and China. *J. Infect. Dev. Ctries.*

14, 125–128 (2020).

19.   N. M. Linton *et al.*, Epidemiological characteristics of novel coronavirus infection:

A statistical analysis of publicly available case data,

doi:10.1101/2020.01.26.20018754.

20.   J. A. Lewnard *et al.*, Incidence, clinical outcomes, and transmission dynamics of

severe coronavirus disease 2019 in California and Washington: Prospective cohort

study. *BMJ*. **369** (2020), doi:10.1136/bmj.m1923.

21.   W. Guan *et al.*, Clinical Characteristics of Coronavirus Disease 2019 in China. *N.*

Engl. J. Med. **382**, 1708–1720 (2020).

22.   L. Pan *et al.*, Clinical characteristics of COVID-19 patients with digestive symptoms

in Hubei, China: A descriptive, cross-sectional, multicenter study. *Am. J.*

Gastroenterol. **115**, 766–773 (2020).

23.   L. Rosenbaum, Facing Covid-19 in Italy - Ethics, Logistics, and Therapeutics on the

Epidemic's Front Line. *N. Engl. J. Med.* **382**, 1873–1875 (2020).

24.   R. M. Anderson, H. Heesterbeek, D. Klinkenberg, T. D. Hollingsworth, How will

country-based mitigation measures influence the course of the COVID-19 epidemic?

Lancet. **395** (2020), pp. 931–934.

25.   Isolation, quarantine, social distancing and community containment: pivotal role for

old-style public health measures in the novel coronavirus (2019-nCoV) outbreak |

Journal of Travel Medicine | Oxford Academic, (available at

https://academic.oup.com/jtm/article/27/2/taaa020/5735321).

26.   R. H. Mena *et al.*, Using posterior predictive distributions to analyse epidemic

models: COVID-19 in Mexico City (2020) (available at

http://arxiv.org/abs/2005.02294).

27.   U. Avila-Ponce de León, Á. G. C. Pérez, E. Avila-Vales, *medRxiv*, in press,

doi:10.1101/2020.05.11.20098517.

References (websites):

• (http://www.citymayors.com/statistics/largest-cities-population-125.html).

• https://www.360dx.com/infectious-disease/new-york-california-serology-studies-

give-early-estimates-covid-19-prevalence#.Xu0aJmgzaUk

• https://www.cnbc.com/2020/04/23/new-york-antibody-study-estimates-

13point9percent-of-residents-have-had-the-coronavirus-cuomo-says.html

• https://coronavirus.gob.mx/datos/