

1 **Article Summary Line:** In one of the areas of first expansion of the epidemic in Italy IgG
2 seroprevalence for SARS-CoV-2 increases with age, possibly suggesting age related differences in
3 susceptibility to infection.

4 **Running Title:** Seroprevalence screening in Castiglione D'Adda.

5 **Keywords:** mass screening, seroprevalence, SARS-CoV-2, COVID-19

6 **Title:** Seroprevalence of SARS-CoV-2 IgG significantly varies with age: results from a mass
7 population screening (SARS-2-SCREEN-CdA)

8

9 **Authors:** Gabriele Pagani^{1,2}, Federico Conti^{1,2}, Andrea Giacomelli^{1,2}, Dario Bernacchia^{1,2}, Rossana
10 Rondanin³, Andrea Prina³, Vittore Scolari⁴, Cecilia Eugenia Gandolfi¹, Silvana Castaldi^{1,5},
11 Giuseppe Marano¹, Cosimo Ottomano⁶, Patrizia Boracchi¹, Elia Biganzoli^{1*}, Massimo Galli^{1,2*}.

12

13 **Affiliations:**

14 1: Università degli Studi di Milano, Italy

15 2: ASST FBF-Sacco, Milano, Italy

16 3: Medispa s.r.l.

17 4: Institut Curie-PSL Research University, CNRS, Sorbonne Université, UMR3664, F-75005, Paris,
18 France.

19 5: Fondazione IRCCS Ca' Granda Ospedale Maggiore policlinico di Milano, Italy

20 6: Synlab s.p.a.

21 *Equally contributing Authors

22

23

24

25

26

27 **Abstract – 48 words**

28 Castiglione D’Adda is one of the towns earlier and more severely affected by the SARS-CoV-2
29 epidemic in Lombardy. In a mass screening involving 4174 out of about 4550 total inhabitants,
30 significant age-related differences in anti SARS-CoV-2 IgG seroprevalence were found, with the
31 lowest prevalence in the youngest inhabitants.

32

33 **Text – 691 words**

34 ***Background and Methods***

35 Coronavirus Induced Disease 2019 (COVID-19) is caused by a novel *betacoronavirus* which was
36 first identified in China and denominated SARS-CoV-2 (Severe Acute Respiratory Syndrome
37 Coronavirus 2) [1]. Italy was the first European country that suffered a wide spread of the infection,
38 which caused hundreds of thousands of cases [2].

39 The municipality of Castiglione d’Adda, a town of about 4550 inhabitants, has been heavily
40 affected by SARS-CoV-2 infection since the earliest stages of the epidemic and subjected to
41 movement restrictions since February 23rd, 2020. As of June 21, 2020, 184 confirmed cases of
42 COVID-19 were reported, the large majority of which requiring hospitalization, accounting for
43 about 4% of the total population. At the same time, 76 deaths were officially attributed to COVID-
44 19, with an overall population mortality of 1.7%. During the epidemic, testing was restricted to
45 severely symptomatic cases. Consequently, the true extent of the SARS-CoV-2 infection remains
46 unknown. In this study, the entire population of Castiglione D’Adda was invited to perform a
47 lateral-flow immunocromatographic tests on capillary blood (Prima Lab, Switzerland). Moreover, a
48 random sample of 562 subjects (stratified per sex and age) and all subjects tested positive to the
49 rapid test were invited to undergo confirmatory tests by chemiluminescent method on venipuncture
50 drawn blood (CLIA, IgG anti-SARS-CoV-2, Abbott, USA) and SARS-CoV-2 PCR on NPS [3].

51

52 The analysis of IgG prevalence in the different age groups was performed by logistic regression
53 models with response variable equal to 1 for positive IgG results, and 0 for negative IgG results.
54 Age and gender were included as independent variables. Results were reported in terms of
55 estimated probabilities of being positive to IgG test as a function of age, with respective 95%
56 confidence intervals.

57

58 ***Results and Discussion***

59 The analysis was based on 509 subjects of the random sample with available results. The overall
60 seroprevalence found in the tested sample was 22.6% (95% confidence interval 17.2- 29.1). A
61 significant effect of age was found ($p < 0.0001$) while no significant association emerged between
62 IgG results and gender ($p = 0.2560$). The possible existence of a non-linear effect of age was tested
63 by including spline polynomials, without significant results ($p = 0.9078$). Furthermore, an
64 age/gender interaction effect did not result significant ($p = 0.5199$). Estimates of probabilities of
65 being positive to IgG test, from a model including only age as independent variable, are reported in
66 Fig.1 and Tab. 1.

67 Since the early phases of the pandemic, advanced age was identified as an independent predictor for
68 severe disease and worse outcomes [4]. Beside this, it remains unclear if the limited number of
69 cases reported in children [5] is due to a milder course of disease, with a larger percentage of
70 asymptomatic cases, or to a lower susceptibility to infection, as our results seem to suggest.

71 Different ACE2 expression according to age have been postulated to explain clinical expression and
72 susceptibility to the infection. In particular, a higher expression of ACE2 in lung tissues in
73 advanced age groups had been speculated [6, 7]. Moreover, a variable susceptibility to other
74 coronavirus such as HCoV-NL63, which also use ACE2 as cell receptor in humans, in different age
75 groups, has been also reported in different age groups [8].

76 Another possible explanation may be that an asymptomatic/pauci-symptomatic infection, more
77 common in younger subjects, could elicit a less marked, or transient, antibody response, as already
78 found in the closely related Middle East Respiratory Syndrome Coronavirus (MERS-CoV) [9].

79 A possible confounding factor in our findings could be related to social distancing measures:
80 schools of any grade were among the first institutions to be closed in Italy, starting from the 5th of
81 March. This could have led to a lower exposure to the infection in children in pre-scholar and
82 scholar age groups.

83 In conclusion, our findings suggest that IgG seroprevalence for SARS-CoV-2 increases with
84 increasing age and these data suggest a lower susceptibility to infection in the lower age groups.

85 These findings have important implications in epidemiology and public health, particularly in
86 designing future population screenings, and could be an important contribution in the re-opening
87 process, especially considered that more than three-fourths of the population could be still
88 susceptible to SARS-CoV-2 infection, even in an area of initially unrestricted viral circulation.

89

90 **Conflict of interests**

91 The authors declare no conflicts of interest.

92

93 **Acknowledgments**

94 MG, GP, FC, DB, AG, RR and EB defined the study protocol. GP drafted a first version of the
95 manuscript, which was then revised and integrated by MG, EB, AG, CEG and SC. EB, PB and GM
96 analyzed the data. All authors approved the final version of the manuscript.

97 We also acknowledge the non-conditioning economical support from CISOM (Corpo Italiano di
98 Soccorso dell'Ordine di Malta), Banca Mediolanum s.p.a., Fondazione Rava and FC Internazionale
99 Milano.

100 The study was approved by University of Milan's Ethical Committee.

101

102 **Author Bio**

103 Gabriele Pagani is an Infectious Diseases resident at Luigi Sacco Hospital (Milano, Italy), currently

104 ending his last year of residency. His primary interests are tropical and traveler diseases and

105 anthropo-zoonoses, particularly in One Health and Global Health perspectives.

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128 **References:**

- 129 [1] Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with Pneumonia in
130 China, 2019. *N Engl J Med* 2020; 382: 727–733.
- 131 [2] EpiCentro. Coronavirus | Istituto Superiore di Sanità, <https://www.epicentro.iss.it/coronavirus/>
132 (accessed 29 May 2020).
- 133 [3] Pagani G, Bernacchia D, Conti F, et al. Dynamics of the SARS-CoV-2 epidemic in the
134 earliest-affected areas in Italy: 1 Mass screening for SARS-CoV-2 serological positivity
135 (SARS-2-SCREEN). *medRxiv* 2020; 2020.06.06.20124081.
- 136 [4] Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of
137 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *The Lancet* 2020;
138 395: 507–513.
- 139 [5] Lee P-I, Hu Y-L, Chen P-Y, et al. Are children less susceptible to COVID-19? *Journal of*
140 *Microbiology, Immunology and Infection* 2020; 53: 371–372.
- 141 [6] Li Y, Zhou W, Yang L, et al. Physiological and pathological regulation of ACE2, the SARS-
142 CoV-2 receptor. *Pharmacological Research* 2020; 157: 104833.
- 143 [7] Bourgonje AR, Abdulle AE, Timens W, et al. Angiotensin-converting enzyme-2 (ACE2),
144 SARS-CoV-2 and pathophysiology of coronavirus disease 2019 (COVID-19). *The Journal of*
145 *Pathology*; n/a. DOI: 10.1002/path.5471.
- 146 [8] Huang S-H, Su M-C, Tien N, et al. Epidemiology of human coronavirus NL63 infection
147 among hospitalized patients with pneumonia in Taiwan. *Journal of Microbiology, Immunology*
148 *and Infection* 2017; 50: 763–770.

149 [9] Choe PG, Perera RAPM, Park WB, et al. MERS-CoV Antibody Responses 1 Year after
150 Symptom Onset, South Korea, 2015. *Emerg Infect Dis* 2017; 23: 1079–1084.

151

152 **Corresponding author:**

153 Gabriele Pagani

154 Infectious Diseases department, III Division

155 ASST FBF-Sacco

156 Via G.B. Grassi 74, 20157, Milano

157 gabriele.pagani@unimi.it

158 02 3904 2591

159

160 **Tab.1 Estimates of the probability of IgG positivity for age interval of five years**

161 C.I. = confidence interval

162

163 **Fig.1: Estimated probability of IgG positivity as a function of age**

164 Solid line: estimates, dashed lines: 95% confidence intervals.

AGE	PROBABILITY (95% C.I.)	AGE	PROBABILITY (95% C.I.)
5	9.1% (4.4%, 13.8%)	50	22.3% (18.5%, 26.0%)
10	10.1% (5.4%, 14.8%)	55	24.3% (20.4%, 28.3%)
15	11.2% (6.6%, 15.9%)	60	26.6% (22.2%, 30.9%)
20	12.5% (7.9%, 17.0%)	65	28.9% (23.8%, 33.9%)
25	13.8% (9.3%, 18.2%)	70	31.3% (25.4%, 37.3%)
30	15.2% (11.0%, 19.5%)	75	33.9% (26.8%, 41.0%)
35	16.8% (12.7%, 20.9%)	80	36.6% (28.3%, 44.9%)
40	18.5% (14.6%, 22.4%)	85	39.3% (29.7%, 48.9%)
45	20.3% (16.6%, 24.0%)	90	42.1% (31.1%, 53.1%)

