

1 **Title: „The transcriptome-wide association search for genes and genetic variants which**  
2 **associate with BMI and gestational weight gain in women with type 1 diabetes”**

3 **Short title: “TWAS and variant association analysis for BMI and GWG in T1DM**  
4 **patients”**

5  
6 Agnieszka H. Ludwig-Słomczyńska,<sup>1</sup>; Michał T. Seweryn,<sup>1,2</sup>; Przemysław Kapusta,<sup>1</sup>; Ewelina  
7 Pitera,<sup>1</sup>; Urszula Mantaj,<sup>3</sup>; Katarzyna Cyganek,<sup>4,5</sup>; Paweł Gutaj,<sup>3</sup>; Łucja Dobrucka,<sup>4</sup>; Ewa  
8 Wender-Ożegowska,<sup>3</sup>; Maciej T. Małecki,<sup>4,5</sup>; Paweł P. Wołkow,<sup>1\*</sup>

9  
10 <sup>1</sup>Center for Medical Genomics OMICRON, Jagiellonian University Medical College,  
11 Kraków, Poland;

12 <sup>2</sup>The Ohio State University Wexner Medical Center, Department of Cancer Biology and  
13 Genetics, Columbus, Ohio, United States of America;

14 <sup>3</sup>Poznan University of Medical Sciences, Department of Reproduction, Poznań, Poland;

15 <sup>4</sup>University Hospital Kraków, Department of Metabolic Diseases, Kraków, Poland;

16 <sup>5</sup>Department of Metabolic Diseases, Jagiellonian University Medical College, Kraków,  
17 Poland.

18  
19 Agnieszka H. Ludwig-Słomczyńska – ORCID ID 0000-0002-0548-4767

20 Michał T. Seweryn – ORCID ID 0000-0002-9090-3435

21 Przemysław Kapusta – ORCID ID 0000-0003-4467-1175

22 Ewelina Pitera - ORCID ID 0000-0001-6208-5976

23 Urszula Mantaj – ORCID ID 0000-0002-0042-778X

24 Katarzyna Cyganek – ORCID ID 0000-0003-2673-3512

25 Paweł Gutaj – ORCID ID 0000-0003-2885-9792

26 Łucja Dobrucka – ORCID ID 0000-0002-8056-8805

27 Ewa Wender-Ożegowska – ORCID ID 0000-0002-5492-8651

28 Maciej T. Małecki – ORCID ID 0000-0002-4794-0820

29 Paweł P. Wołkow – ORCID ID 0000-0002-9322-5545

30

31 **\*Corresponding author:**

32 e-mail: [pawel.wolkow@uj.edu.pl](mailto:pawel.wolkow@uj.edu.pl) (PPW), [agnieszka.ludwig@uj.edu.pl](mailto:agnieszka.ludwig@uj.edu.pl) (ALS)

33

## 34 **ABSTRACT**

35 **Background:** Clinical data suggest that BMI and gestational weight gain (GWG) are strongly  
36 interconnected phenotypes, however the genetic basis of the latter is rather unclear. Here we  
37 aim to find genes and genetic variants which influence BMI and/or GWG. **Methods:** We have  
38 genotyped 316 type 1 diabetics using Illumina Infinium Omni Express Exome-8 v1.4 arrays.  
39 The GIANT, ARIC and T2D-GENES summary statistics were used for TWAS (performed  
40 with PrediXcan) in adipose tissue. Next, the analysis of association of imputed expression  
41 with BMI in the general and diabetic cohorts (Analysis 1 and 2) or GWG (Analysis 3 and  
42 4) was performed, followed by variant association analysis (1Mb around identified loci) with  
43 the mentioned phenotypes. **Results:** In Analysis 1 we have found  
44 175 BMI associated genes and 19 variants ( $p < 10^{-4}$ ) which influenced GWG, with the

45 strongest association for rs11465293 in *CCL24* ( $p=3.18E-06$ ). Analysis 2, with diabetes  
46 included in the model, led to discovery of 1812 BMI associated loci and 207 variants ( $p<10^{-4}$ )  
47 influencing GWG, with the strongest association for rs9690213 in *PODXL* ( $p=9.86E-07$ ). In  
48 Analysis 3, among 648 GWG associated loci, 2091 variants were associated with BMI  
49 ( $FDR<0.05$ ). In Analysis 4, 7 variants in GWG associated loci influenced BMI in the ARIC  
50 cohort. **Conclusions:** Here, we have shown that loci influencing BMI might have an impact  
51 on GWG and GWG associated loci might influence BMI, both in the general and T1DM  
52 cohorts. The results suggest that both phenotypes are related to insulin signaling, glucose  
53 homeostasis, mitochondrial metabolism, ubiquitinylation and inflammatory responses.

54

55 **Keywords:** obesity, BMI, gestational weight gain (GWG), Transcriptomic-wide association  
56 study (TWAS), association analysis

57

## 58 **DECLARATIONS**

59 **Funding:** The study was funded by the National Science Centre in Poland through the Sonata  
60 Grant “Search for genetic variants influencing gestational weight gain in type 1 diabetes  
61 patients by genome wide association method” to ALS (Nr 2013/11/D/NZ5/03219).

62 **Conflict of interests:** The authors declare they have no competing interests.

63 **Ethics approval and consent to participate:** This study was approved by the Bioethical  
64 Committees of the Jagiellonian University and Poznan University of Medical Sciences and  
65 performed according to the Helsinki Declaration.

66 **Consent to participate:** Informed consent was obtained from all individual participants  
67 included in the study.

68 **Consent to publication:** Not applicable

69 **Availability of data and materials:** Sequence data has been deposited at the European Genome-  
70 phenome Archive (EGA), which is hosted by the EBI and the CRG, under accession number  
71 EGAS00001004408. Further information about EGA can be found on [https://ega-](https://ega-archive.org)  
72 [archive.org](https://ega-archive.org) "The European Genome-phenome Archive of human data consented for  
73 biomedical research" (<http://www.nature.com/ng/journal/v47/n7/full/ng.3312.html>).

74

## 75 **ACKNOWLEDGEMENTS**

76 The authors thank prof. Katarzyna Cypryk and dr Monika Żurawska-Kliś for their help in  
77 recruiting patients.

78 This research was supported in part by PLGrid Infrastructure. The bioinformatic analysis was  
79 performed using Prometheus (AGH, Krakow, Poland), Michigan Imputaton Server  
80 (Michigan, MI, USA) and The Ohio Supercomputer Center (Columbus, OH, USA).

81 For manuscripts with ARIC GWAS data (including HapMap and 1000G imputed data): The  
82 Atherosclerosis Risk in Communities study has been funded in whole or in part with Federal  
83 funds from the National Heart, Lung, and Blood Institute, National Institutes of Health,  
84 Department of Health and Human Services (contract numbers HHSN268201700001I,  
85 HHSN268 201700002I, HHSN268201700003I, HHSN268201700004I and  
86 HHSN268201700005I), R0 1HL087641, R01HL059367 and R01HL086694; National Human  
87 Genome Research Institute contract U01HG004402; and National Institutes of Health contract  
88 HHSN268200625226C. The authors thank the staff and participants of the ARIC study for  
89 their important contributions. Infrastructure was partly supported by Grant Number  
90 UL1RR025005, a component of the National Institutes of Health and NIH Roadmap for  
91 Medical Research.

92 The BioMe Biobank Program T2D-GENES Exome Sequencing Study was conducted by the  
93 BioMe Biobank Program T2D-GENES Exome Sequencing Study Investigators and supported  
94 by the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). The data  
95 from the BioMe Biobank Program T2D-GENES Exome Sequencing Study reported here were  
96 supplied by the Broad Institute and Icahn School of Medicine at Mount Sinai. This manuscript  
97 was not prepared in collaboration with Investigators of the BioMe Biobank Program T2D-  
98 GENES Exome Sequencing study and does not necessarily reflect the opinions or views of  
99 the BioMe Biobank Program T2D-GENES Exome Sequencing study, or the NIDDK.

100 **Authorship statement:** All authors whose names appear on the submission made substantial  
101 contributions to the conception or design of the work; or the acquisition, analysis, or  
102 interpretation of data; or the creation of new software used in the work; drafted the work or  
103 revised it critically for important intellectual content; approved the version to be published;  
104 agree to be accountable for all aspects of the work in ensuring that questions related to the  
105 accuracy or integrity of any part of the work are appropriately investigated and resolved.

## 106 INTRODUCTION

107 In the recent years, a significant attention was paid to transcriptomic-wide association  
108 studies (TWAS), which enable the prediction of gene-level associations between the 'imputed  
109 expression based on genetic data' and complex phenotypes (Gamazon et al., 2015). This is  
110 possible, since it was shown that a proportion of GWAS risk variants co-localize with genetic  
111 variants, which regulate gene expression (i.e. expression quantitative trait loci, eQTL)  
112 (Hormozdiari et al., 2016). The prediction of gene expression based on genotype removes the  
113 noise created by environmental factors as well as the potential reverse causation (when the  
114 trait affects gene expression), therefore the TWAS analysis not only increases the statistical

115 power, but also enables to focus on regions for which finding a functional interpretation of the  
116 associations is often easier (B. Li et al., 2018; Mancuso et al., 2017).

117 Gestational weight gain (GWG) has been extensively studied over the last several  
118 years as it was suggested that it's inadequacy may lead to adverse both maternal  
119 (preeclampsia, hypertension, obesity later in life, cesarean section) and neonatal (preterm  
120 birth, stillbirth, inadequate neonatal weight – small or large for gestational age, obesity later in  
121 life) outcomes (*Influ. Pregnancy Weight Matern. Child Heal.*, 2007; Kominiarek &  
122 Peaceman, 2017; Voerman et al., 2019). Simultaneously, clinicians tried to optimize  
123 pregnancy care for women with type 1 diabetes whose pregnancy outcomes (including  
124 developmental abnormalities, spontaneous abortions, neonatal hyperglycemia, neonatal  
125 hyperinsulinemia, maternal retinopathy, maternal nephropathy, preeclampsia) are far worse  
126 than in women from the general population (Celia et al., 2016). For a long time it was  
127 believed that it stemmed from maternal hyperglycemia, however, even though glycemic goals  
128 for these women have been achieved (HbA1c < 6.0%) still large for gestational age (LGA) or  
129 macrosomic neonates are born more frequent than in general population (Bashir, Naem, Taha,  
130 Konje, & Abou-Samra, 2019; Dori-Dayan et al., 2020; Scifres, Feghali, Althouse, Caritis, &  
131 Catov, 2014). Thus, there must be other potential contributors to adverse maternal and fetal  
132 outcomes (Mastella et al., 2018; Rys, Ludwig-Slomczynska, Cyganek, & Malecki, 2018;  
133 Secher et al., 2014). Among them are also those which affect the general population -  
134 maternal body lipids, pre-pregnancy BMI and GWG (McWhorter et al., 2018).

135 Even though GWG can have an impact on the health of future generations, to date,  
136 most of the performed studies are retrospective or observational and their main goal was to  
137 assess the relationship between GWG and environmental factors (Nunnery, Ammerman, &

138 Dharod, 2018; Siega-Riz, Bodnar, Stotland, & Stang, 2020), while the genetic risks for  
139 inadequate GWG have been scarcely analyzed. In a recent paper, the authors show that 43%  
140 and 26% of the variation in GWG can be explained by genetic factors in the first pregnancy  
141 and second pregnancy, respectively (Andersson et al., 2015). The first GWAS study on GWG  
142 in the general, multiethnic cohort was performed in 2018. The study has shown that 20% of  
143 the variability in GWG can be explained by maternal genetic variants. Unfortunately, it did  
144 not find significant associations of GWG with any genomic loci (Warrington et al., 2018).

145         Since, both environmental and genetic data show that there is a correlation between  
146 pre-pregnancy BMI and GWG (Luecke et al., 2018) we took the TWAS approach to study the  
147 genetic correlation between BMI and GWG - in particular, to find loci and variants which are  
148 associated with GWG and BMI, or (on the contrary) only with GWG. We find this  
149 particularly important, as the reports on the overlap of genes and/or variants which influence  
150 both phenotypes appear to be conflicting (Kawai, Nwosu, Kurnik, Harrell, & Stein, 2019;  
151 Lawlor et al., 2011).

152         In this study, we aimed to investigate the genetic factors associated with GWG from  
153 the perspective of the genetics of obesity. To this aim, we performed TWAS analyses on the  
154 GIANT cohort representing the general population and two diabetic cohorts - T2D-GENES  
155 and ARIC to find genes associated with BMI. The ARIC and T2D-GENES cohorts were used  
156 since diabetes mellitus itself might influence BMI and creates a special metabolic context,  
157 while no cohort with patients with type 1 diabetes was available. Only regions which were  
158 associated with BMI in these cohorts were subjected to the analysis of association with GWG  
159 in T1DM patients. Secondly, we also searched for reverse associations and checked whether  
160 genes in TWAS associated with GWG in T1DM cohort might affect BMI in ARIC or GIANT

161 cohorts. If so, we decided to search for genes which imputed expression correlated with BMI  
162 in order to restrict the analysis of genetic association with GWG to these genes only and  
163 conversely, to check whether the genes which correlate with GWG (at the imputed expression  
164 level) do influence BMI. Thus, this work comprises 4 different analyses, as presented in  
165 Figure 1.

166

## 167 **METHODS**

168 **Patients:** Patients were recruited either in Department of Metabolic Diseases University  
169 Hospital in Krakow or in Division of Reproduction Department of Obstetrics, Gynecology  
170 and Gynecological Oncology, Poznan University of Medical Sciences. All patients enrolled in  
171 the study were women with type 1 diabetes (T1D) and treated with insulin. Pre-pregnancy  
172 weight was based either on women self-report or weight measurement (if the women  
173 were/was under antenatal care). Only singleton pregnancies were included. Pregnancies that  
174 resulted in a miscarriage or stillbirth were excluded. Total GWG was defined as the difference  
175 between last gestational weight before delivery and pre-pregnancy weight. Whole blood  
176 samples were drawn and stored at -80C. This study was approved by the Bioethical  
177 Committees of the Jagiellonian University and Poznan University of Medical Sciences and  
178 performed according to the Helsinki Declaration. Written informed consent was collected  
179 from all patients.

180 **Genotyping:** DNA was extracted from whole blood with the use of automated nucleic acid  
181 extraction system Maxwell (Promega). Five hundred twenty-seven samples were genotyped  
182 on Illumina Infinium Omni Express Exome-8 v1.4 arrays. Only in term life births with

183 information regarding age, pre-pregnancy BMI, GWG, diabetes duration, treatment method  
184 and daily insulin dose available were included in the final analysis.

185 **Data processing and imputation:** The detailed protocol of the data processing, QC analysis  
186 and imputation is presented in (Ludwig-Slomczynska et al., 2018).

### 187 **Genotype Data and the analysis of genetic variants**

188 The GWAS analysis on the T1DM cohort was performed on a group of 316 females with  
189 complete data on: age, parity, insulin dose prior pregnancy, pre-pregnancy BMI and GWG.

190 The mixed-effects model approach was used as implemented in the package GENESIS in R.

191 The random effect was associated with the individual ID and the genetic relatedness matrix  
192 was estimated via the PCARelate method in package GENESIS in R. The 'null' mixed effects  
193 model was considered as:

194  $GWG \sim age + parity + pre-pregnancy\_BMI + pre-pregnancy\_InsDose + OriginOfSample + (1|ID)$ ,

195 where the OriginOfSample is an indicator of the sample being collected in Division of  
196 Reproduction Department of Obstetrics, Gynecology and Gynecological Oncology, Poznan  
197 University of Medical Sciences. To the aim of testing significance of the (additive) effect of  
198 the genotype the Wald's test statistics was used.

199 The summary statistics for BMI tested in the GIANT Consortium cohort were downloaded  
200 from [https://portals.broadinstitute.org/collaboration/giant/images/1/15/SNP\\_gwas\\_mc\\_merge](https://portals.broadinstitute.org/collaboration/giant/images/1/15/SNP_gwas_mc_merge_nogc.tbl.uniq.gz)  
201 [\\_nogc.tbl.uniq.gz](https://portals.broadinstitute.org/collaboration/giant/images/1/15/SNP_gwas_mc_merge_nogc.tbl.uniq.gz)

202 The genotype and phenotype data for the ARIC cohort (GENEVA study) were accessed via  
203 dbGaP. As far as the genotype data is concerned, the imputed data for participants of  
204 European ancestry were used as deposited under the phg000248.v1 code. Per-chromosome

205 genotype probability data were transformed to dosages. From these dosages, genotype  
206 relatedness matrices were estimated on a per-chromosome basis and merged with the aid of  
207 the SNPRelate package in R. For the GWAS analysis the GENESIS package was used. As far  
208 as the phenotype files are concerned, these were accessed under the phs000090.v3 study code.  
209 The 'null' mixed effects model was defined as:

210  $BMI \sim age + sex + diabetes\_status + (1|ID)$

211 As GENEVA is a longitudinal study, for diabetic participants, we used the BMI and age  
212 measurement at the first time when the status of the participant was 'diabetic', whereas for the  
213 non-diabetic participants we used the measurements at the study entry.

214 The T2Dgenes genotype files were accessed via dbGaP under the phg000573.v1 code. The  
215 raw vcf files were transformed to dosages prior analysis. The phenotype data were accessed  
216 from the phs000462.v2 study code. Similarly to the data analysis in the ARIC cohort, we  
217 recorder BMI and age at the first time when the diabetic status was positive or at the study  
218 entry, otherwise.

219 The TWAS analysis was performed via the PrediXcan and MetaXcan software with Adipose  
220 as the tissue of interest. For the GIANT Cohort data, the MetaXcan framework was  
221 implemented based on the GTEx.v7 models. For the ARIC, T2Dgenes and T1DM cohorts the  
222 PrediXcan software was used also with GTEx.v7 models. In the TWAS analyses in the ARIC  
223 and T1DM cohorts the same independent variables were used as with the GWAS models  
224 defined above – with the exception that the classical linear models were used as implemented  
225 in the limma package (with no random effects). For the T2Dgenes TWAS analysis, from the  
226 'predicted expression' matrix the principal components were first estimated and the first  
227 component was added to the models as it did not correlate with any independent variable –

228 and it differ between families (as tested with the Kruskal-wallis test). As before the target  
229 analysis in the T2Dgenes cohort was performed with the aid of linear models as implemented  
230 in the limma package.

231 The FUMA analysis was performed through the webserver at [fuma.ctglab.nl](http://fuma.ctglab.nl)

232 The COJO and fast-BAT software were used as implemented by in the GTCA framework.

233 The LD structure was estimated via the 1000 genomes data.

234 GO enrichment analysis was performed as implemented in the topGO package in R.

235

## 236 **RESULTS**

### 237 *Gestational weight gain analysis in T1DM cohort*

238 Our analysis comprised 316 women with T1DM for whom full phenotype data were  
239 available. The basic characterization of patients included in the analysis is presented in Table  
240 1.

241 *Table 1. The clinical characteristics of the T1DM cohort.*

	<b>Med; IQR</b>
<b>Age [years]</b>	29; (25.75, 32)
<b>Pre-pregnancy BMI [kg/m<sup>2</sup>]</b>	23.41; (21.09, 26.2)
<b>GWG [kg]</b>	14; (10.5, 18)
<b>Diabetes duration [years]</b>	12; (6, 18)

<b>Treatment method – CSII [%]</b>	0.55
<b>Daily insulin dose before pregnancy [U]</b>	38; (28, 50)
<b>Parity</b>	1; (1, 2)

242

243         PrediXcan was used to impute gene expression in subcutaneous and visceral adipose  
244 tissue based on imputed genotype data and the gene-based search for associations with GWG  
245 and BMI was performed using a linear model approach. Due to small cohort size, genes  
246 nominally significantly associated with GWG (442 genes in subcutaneous and 328 in visceral  
247 adipose tissue) and BMI (416 genes in subcutaneous and 326 in visceral adipose tissue) were  
248 considered for further analysis. It is worth noting that only sixty genes overlapped between  
249 the two phenotypes (Table S1). We further explored this phenomenon by looking at the  
250 strength of the association between gene expression for these two traits. We found a negative  
251 correlation between logFCs of genes associated with GWG and BMI. This is consistent with  
252 clinical recommendations as patients with high pregestational BMI are advised to restrict their  
253 GWG, however, we believe that the genetic background might also play a role.

254

255 ***PrediXcan analysis on BMI in the general population (GIANT cohort) and variant***  
256 ***associations with GWG in T1DM cohort - Analysis 1***

257         Gene expression prediction in visceral and subcutaneous adipose tissue in the GIANT  
258 cohort (234069 patients, (Locke, Kahali, Berndt, Justice, & Pers, 2015)) as performed using  
259 PrediXcan software. One hundred seventy-five genes significantly associated ( $p < 1E-04$ )  
260 with BMI in the adipose (subcutaneous and visceral combined) tissue were found (Table S2).

261 The Venn diagram of genes which affected BMI, GWG or both phenotypes is presented in  
 262 Table S3. Only 15 genes influenced both phenotypes, 160 influenced only BMI, while 633  
 263 had an impact on GWG only (Figure 2).

264 We further analyzed the loci of interest and searched for variants (within the 1Mb  
 265 window around the gene) which influence BMI and/or GWG. We first mapped variants in 175  
 266 BMI associated genes (+/- 500000 bp) and applied these restricted variant panels to the  
 267 analysis of the association with GWG in the T1DM cohort. We filtered the results by MAF <  
 268 5% and p-value < 1E-04 and found an association with GWG for 19 variants in the T1DM  
 269 cohort (Table 2).

270 **Table 2. The list of variants localized to genes associated with BMI in the GIANT cohort**  
 271 **which associate with GWG.**

SNP ID	chr	MAF [%]	B	p-value	Gene/ nearest gene	localization/type
rs11465293	7	5.7	4.46	3.18E-06	<i>CCL24</i>	missense
rs1978202	7	8.7	3.51	4.53E-06	<i>CCL26</i>	intergenic
rs9347707	6	35.1	1.94	4.74E-05	<i>PACRG</i>	intron
rs72849841	17	17.1	-2.19	1.25E-04	<i>RNF213</i>	missense
rs11807240	1	10.4	2.58	1.61E-04	<i>FUBP1</i>	intron
rs9659938	1	10.4	2.58	1.61E-04	<i>NEXN204</i>	intron
rs13340504	7	14.4	2.32	2.29E-04	<i>CCL24</i>	intergenic
rs1541725	2	46.0	1.61	3.95E-04	<i>AC009502.4</i>	intergenic
rs2136682	1	9.3	2.54	4.25E-04	<i>GIPC2</i>	intron

rs4329088	6	46.2	1.63	4.33E-04	<i>PACRG</i>	intron
rs886131	12	41.0	-1.60	4.78E-04	<i>RPL29P25</i>	noncoding transcript
rs11037234	11	33.1	1.66	5.34E-04	<i>RP11-111A24.2</i>	intergenic
rs8122282	20	45.7	-1.49	6.44E-04	<i>C20orf166</i>	intron
rs4808209	17	6.9	3.00	7.13E-04	<i>ZNF101</i>	start loss
rs9352745	6	26.1	1.70	7.27E-04	<i>LCA5</i>	intron
rs876373	12	43.4	1.50	7.29E-04	<i>RPL29P25</i>	intron
rs6062267	20	45.3	-1.49	7.94E-04	<i>C20orf166</i>	intron
rs7961894	12	8.5	2.75	8.88E-04	<i>WDR66</i>	intron
rs7564856	2	34.5	-1.59	9.62E-04	<i>SPEG</i>	intergenic

272

273 The FUMA GWAS method further prioritized the variants obtained in the analysis.  
 274 Twelve leading SNPs were detected. Three of them (rs7564856, rs11465293 and rs7961894)  
 275 were associated with several traits in GWAS Catalog, mainly blood parameters, but, none  
 276 significantly with metabolic outcome. At the same time however, these were also associated  
 277 at the level of significance  $p\text{-value} < 1E-03$  with LDL cholesterol levels, nonalcoholic fatty  
 278 liver disease, body mass index, mean arterial pressure or metabolite levels. These variants  
 279 were eQTLs for 40 genes in several tissues, while rs9659938, rs11807240, rs2136682 (*NEXN*,  
 280 *NEXN-AS1*), rs7564856 (*SPEG*), rs9352745 (*SH3BGRL2*), rs13340504 (*CCL24*), rs11465293  
 281 (*CCL24*), rs886131 (*RPL29P25*) influenced gene expression in both subcutaneous and  
 282 visceral adipose. Thus, we conclude that there is a subset of genes which influence both BMI  
 283 and GWG as well as genetic variants associated with GWG which co-localize to these loci.

284           Next, we tried to determine, whether the signals for the two traits in the genes of interest  
285 are correlated, thus we checked whether the genetic variants associated with BMI and GWG  
286 are in linkage disequilibrium or they belong to different haplotypes. We performed the  
287 analysis of LD for all genes with variants significantly associated with GWG, however, below  
288 we present only the most interesting examples. In the gene *GPN3*, we have found 2 bins of  
289 eQTLs which are not in LD either with BMI or GWG variants. Variants associated with BMI  
290 also create a cluster, which is separate from a small cluster of 2 variants associated with  
291 GWG. One of GWG associated variants (rs876373) is in moderate LD with the BMI cluster  
292 (Figure S1a). In another gene, *PMS2P3*, we found a cluster of 3 variants which are associated  
293 with GWG, all in LD with one eQTL (rs707395). At the same time, variants associated with  
294 BMI form a separate, bin which is in moderate LD with a subset of eQTLs for this gene. This  
295 cluster is not in LD with the GWG bin or rs707395 (Figure S1b). The third example is the  
296 *STAG3L1* gene in which three clusters - GWG associated variants, BMI associated variants  
297 and eQTLs can be seen ( $r^2$ ), however, in each, few members of any cluster are in LD with  
298 members of other clusters ( $D'$ ) making them rather dependent on each other (Figure S1c).

299

300 ***PrediXcan analysis on BMI in the diabetic cohorts (ARIC and T2D-GENES) and variant***  
301 ***associations with GWG in T1DM cohort - Analysis 2***

302           Since it is known that diabetes and its treatment might impact BMI, we searched for  
303 variants that might affect BMI in the context of diabetes status. Since no T1DM cohort was  
304 available, we used T2D patients' cohorts – T2D-GENES (590 patients) and ARIC (8746  
305 patients) (being aware that the two phenotypes obesity and diabetes are interconnected and  
306 T2D patients are predisposed to obesity, while T1DM patients not). After adjusting for  
307 diabetes, we found that 1812 genes were associated with BMI  $p < 0.05$  (Table S4). Among

308 them 135 influenced both phenotypes, 1677 influenced only BMI and 513 influenced GWG  
 309 (Table S5, Figure 3).

310 Again, we mapped variants to 1812 BMI associated genes (+/- 500000bp) and used this  
 311 panel to look for associations with GWG in T1DM cohort. When filtered by MAF < 5% and  
 312 p-value < 1E-04 the analysis has shown that GWG correlates with 207 variants (Table S6).  
 313 Table 3 shows 20 variants which were significant at the level of p-value < 1E-05.

314 **Table 3. The list of variants localized to genes associated with BMI in T2D-GENES and**  
 315 **ARIC cohorts which associate with GWG.**

SNP ID	chr	MAF [%]	B	p-value	Gene/ nearest gene	localization/type
rs9690213	7	18.4	2.66	9.86E-07	<i>PODXL</i>	regulatory region variant
rs11465293	7	5.7	4.46	3.17E-06	<i>CCL24</i>	missense
rs9393623	6	31.0	2.18	4.11E-06	<i>CMAHP</i>	intron
rs4796675	17	22.2	2.38	4.40E-06	<i>LINC00974</i>	noncoding transcript exon variant
rs1978202	7	8.7	3.51	4.53E-06	<i>CCL26</i>	intergenic
rs8080053	17	23.1	2.33	5.57E-06	<i>LINC00974</i>	noncoding transcript exon variant
rs12534221	7	17.7	2.55	5.85E-06	<i>PODXL</i>	intergenic
rs6747327	2	35.6	2.04	1.35E-05	<i>CAPN13</i>	intron
rs4742339	9	37.8	1.97	2.97E-05	<i>KANK1</i>	intergenic

rs11714248	3	43.0	-1.86	4.40E-05	<i>TRANK1</i>	intron
rs9347707	6	35.1	1.94	4.74E-05	<i>PACRG</i>	intron
rs1398095	3	25.6	2.15	4.85E-05	<i>CAPDS</i>	intron
rs9940705	16	6.2	3.61	6.91E-05	<i>FAM92B</i>	intron
rs2421052	5	43.1	-1.86	7.63E-05	<i>AC008691</i>	intron
rs6815557	4	39.2	1.82	8.24E-05	<i>LOC107986225</i>	intergenic
rs12788347	11	10.8	2.92	8.72E-05	<i>AL137224</i>	intron
rs11235519	11	24.2	-2.02	8.93E-05	<i>AP005019.4</i>	intergenic
rs6546217	2	11.7	2.64	9.89E-05	<i>AC118345.1</i>	intergenic
rs7609526	2	11.1	2.68	9.97E-05	<i>AC118345.1</i>	intergenic

316

317 Most of the variants which influenced both GWG and BMI (p-value < 1E-05) were  
 318 eQTLs associated with the expression of 13 genes (*LRRFIP2*, *MLH1*, *GOLGA4*, *CCL26*,  
 319 *RHBDD2*, *AC004980*, *POR*, *CCL26*, *GTF2IRD2*, *UPK3B*, *ART2P*, *LINC00974*, *KTP15*,  
 320 *KRT17*) in 14 tissues (Thyroid, Skeletal Muscle, Whole Blood, Artery, Cultured Fibroblasts,  
 321 Tibial Nerve, Testis, Skin (Unexposed), Visceral Adipose Tissue, Minor Salivary Gland,  
 322 Small Intestine, Colon, Brain (several regions).

323 FUMA analysis showed that among variants significant for association with GWG (p-value <  
 324 1E-04), 18 were listed for associations with multiple phenotypes in the GWAS Catalog. The  
 325 KEGG analysis showed enrichment for autoimmune disease systemic lupus  
 326 erythematosus (adj. p-value = 3.32E-7) and taste transduction (adj. p-value = 2.18E-2). In the  
 327 molecular functions GO enrichment analysis we found these variants to be responsible,  
 328 among others, for carbohydrate binding (adj. p-value = 3.15E-8), bitter taste receptor activity

329 (adj. p-value = 4.51E-5), trace amine receptor activity (adj. p-value = 9.69E-5) and taste  
330 receptor activity (adj. p-value = 9.04E-4) (FigureS2).

331 Finally, when comparing the lists of variants obtained in Analyses 1 and 2, we found a  
332 significant overlap, what supports our hypothesis that loci associated with BMI, both in the  
333 general population and diabetic cohorts may also associate with GWG. Several SNPs -  
334 rs7564856 (*SPEG*), rs1541725 (*AC009502.4*), rs9347707 (*PCARG*), rs4329088 (*PCARG*),  
335 rs1978202 (*CCL26*), rs13340504 (*CCL24*), rs11465293 (*CCL24*), rs11037234 (*RP11-*  
336 *111A24.2*), rs876373 (*RPL29P25*), rs886131 (*RPL29P25*), rs7961894 (*WDR66*), rs4808209  
337 (*ZNF101*) were found to influence GWG in T1DM, ARIC, T2D-GENES and GIANT cohorts.  
338 We conclude, that despite the relatively large overlap between traits at the gene level, the  
339 genetic variants which localize with these loci most often are linked to a single phenotype  
340 only.

341 Next, we aimed to study the “reverse association” - i.e. we asked whether genes  
342 associated with GWG influence BMI as well.

343

344 ***PrediXcan analysis on GWG in T1DM cohort and variant associations with BMI in the***  
345 ***general population (GIANT cohort) - Analysis 3***

346 Variants were mapped to 648 GWG associated genes (+/- 500000 bp) (Table S1) and a  
347 search for associations with BMI in the GIANT cohort was performed. Among 394149  
348 variants analyzed, 2091 were significantly associated with BMI (at FDR < 0.05). These  
349 significant SNPs encompassed 0.53% of the whole list of variants analyzed. In comparison,  
350 the significant SNPs in the analysis of BMI in the GIANT cohort make up 0.41%, what means  
351 that genes which influence GWG are significantly enriched in the association test for BMI.

352 We used the FUMA tool to prioritize these 2091 variants. Twelve lead SNPs and 12  
 353 genomic risk loci were found. Not surprisingly, the GWAS Catalog analysis has shown a  
 354 significant enrichment for genes associated with BMI, body fat distribution, attendance to  
 355 gym or sport groups, sleep duration, height, lipoprotein levels, low HDL-level cholesterol etc.  
 356 as presented in Figure 4 and Table S7.

357 Thus, we find a significant enrichment of variants associated with BMI in the general  
 358 population within loci associated with GWG.

359

360 ***PrediXcan analysis on GWG in T1DM cohort and variant associations with BMI in the***  
 361 ***diabetic cohort (ARIC) - Analysis 4***

362 A GWAS analysis on BMI in the ARIC cohort was performed. Due to a large number of  
 363 variants in the analysis we searched for lead SNPs using LD score regression. COJO analysis  
 364 has returned 15 variants statistically significantly associated with the trait of interest (Table  
 365 4A).

366 ***Table 4. The list of variants localized to genes associated with BMI (A) and GWG (B) in***  
 367 ***ARIC cohort.***

SNP ID	chr	MAF [%]	B	p-value	Gene/ nearest gene	localization/type
A.						
rs138445632	2	7.92	-2.88	4.54E-12	AC021851,2	intron
rs73646471	9	11.40	-5.59	2.60E-10	LOC105375974	intergenic
rs72703233	9	8.47	-1.57	1.31E-07	DMRT2	noncoding transcript

						exon variant
rs78231573	2	6.14	-6.02	2.62E-07	<i>LOC105373643</i>	intergenic
rs144768710	2	5.40	2.77	3.30E-07	<i>CHN1</i>	intron
rs73419031	11	5.72	-6.34	6.96E-07	<i>SOX6</i>	intron
rs6456353	6	90.25	42.14	7.10E-07	<i>E2F3</i>	intron
rs1486942	5	8.15	-2.66	1.61E-06	<i>LOC112267929</i>	intergenic
rs74036632	14	7.05	-2.85	3.69E-06	<i>NYNRIN</i>	3'UTR
rs77346003	1	10.58	-17.55	4.12E-06	<i>ADIPOR1</i>	intergenic
rs73382735	9	5.68	4.49	4.28E-06	<i>LOC105375955</i>	intergenic
rs12685009	9	8.42	-8.94	7.74E-06	<i>SLC24A2</i>	intergenic
rs34427781	22	19.46	-2.54	8.47E-06	<i>TCF20</i>	intron
rs13359900	5	15.98	-27.84	8.95E-06	<i>MCTP1</i>	intron
rs4131847	12	5.86	44.27	9.31E-06	<i>CCDC60</i>	intergenic
B.						
rs7875240	9	10.53	-3.08	1.21E-06	<i>PTPRD</i>	intron
rs1486942	5	8.15	-2.66	1.61E-06	<i>LOC112267929</i>	intergenic
rs74036632	14	7.05	-2.85	3.69E-06	<i>NYNRIN</i>	3'UTR
rs34427781	22	19.46	-2.54	8.47E-06	<i>TCF20</i>	intron

368

369 Only 4 variants from this list were in GWG associated genes in PrediXcan, three of  
 370 which overlapped with those associated with BMI (Table 4B). None of the variants were  
 371 reported in GWAS Catalog or PheWas for the association with any trait. 6 variants were  
 372 eQTLs - two for genes in which they were localized rs13359900 and rs73419031 for *MCTP1*

373 and *SOX6*, respectively; rs144768710 for *ATP5G3*, rs74036632 for *ZFHX2-AS1*, rs4131847  
 374 for *HSPB8*, while rs34427781 for several genes all localized 200000bp upstream or  
 375 downstream from the variant.

376 Since the COJO analysis was performed on loci associated with GWG (+/- 500000bp  
 377 around the gene), it was possible that the most important variants influencing BMI were  
 378 located outside those windows. Thus, we used a different tool, fastBAT, to find genes  
 379 associated with BMI. The analysis revealed 8 genes which nominally associate with the trait  
 380 of interest (Table 5).

381 **Table 5. The list of variants localized to genes associated with BMI in ARIC cohort and**  
 382 **associated with GWG as shown by fast-BAT.**

Gene/nearest gene	chr	localization	# of SNPs	p-value	Top SNP p-value	Top SNP
<i>HLD-DQB</i>	6	intergenic	4654	0.016	1.02E-06	rs115258523
<i>GATAD2</i>	19	intron	1256	0.017	6.23E-05	rs145702982
<i>AC099791.3</i>	1	intron	1800	0.020	8.85E-05	rs2485748
<i>LINC01141</i>	1	intron	1501	0.027	9.95E-05	rs12027661
<i>HLA-DRB</i>	6	intergenic	9354	0.029	1.89E-05	rs139964305
<i>HLA-B</i>	6	stop gained	4017	0.044	2.11E-04	rs149512147
<i>LHFPL6</i>	13	intron	1648	0.047	3.34E-06	rs6563700

383

384 Again, none the of variants was previously mentioned to be significantly associated with  
 385 any trait in GWAS Catalog or PheWas. Two were eQTLs - rs6563700 for *LHFPL6*, while  
 386 rs12027661 for *CAMK2N1*.

387

## 388 **DISCUSSION**

389           In this work we performed TWAS as well as single variant associations to determine  
390 the loci related to GWG and/or BMI. About 15% of loci overlap between BMI and GWG at  
391 the TWAS level. In the general population (GIANT cohort) the pathway analysis has shown  
392 that loci which contribute to both phenotypes affect mainly energy metabolism. Among the  
393 enriched pathways are regulation of mitochondrion (GO:1903749, GO:1903747,  
394 GO:0010822, GO:0070585, GO:0010821, GO:0006839) and Golgi apparatus (GO:0051683,  
395 GO:0048313, GO:0051645). There are also few pathways which affect transcription,  
396 polyadenylation or methylation of RNA (GO:1903311, GO:0016071, GO:1900363,  
397 GO:0080009). The analysis of the loci which TWAS associates with GWG only leads to  
398 necrosis (GO:0070266, GO:0097300, GO:0070265), protein kinases signaling (GO:0046330,  
399 GO:0043507, GO:0032874, GO:0070304, GO:0043506, GO:0007256) and metabolism of  
400 sugars (GO:0034033, GO:0034030, GO:0033866,GO:0009226). The BMI associated loci  
401 enrich cell cycle progression pathways (GO:0010972,GO:1902750) host to pathogen  
402 signaling (GO:0043921, GO:0052312, GO:0052472) and macromolecule metabolism  
403 (GO:0010604, GO:0034641, GO:0006139, GO:0009308, GO:0046483) and insulin signaling  
404 (GO:0046626, GO:1900076). The results of the GO enrichment analysis are presented in  
405 Supplementary Table S8 and Figures S3a, b.

406           The analysis of the overlap between BMI and GWG associated loci in the ARIC  
407 cohort and T1DM cohort respectively showed enrichment in the TGF $\beta$  signaling pathway  
408 (GO:0071559, GO:0071560), regulation of stem cell differentiation (GO:1901532,  
409 GO:1902036, GO:0060218, GO:2000736) and polyol pathway (GO:0019751, GO:0046173).  
410 Pathways associated with GWG were similar to those enriched in general population,

411 however the important contribution of genes which affect antigen presentation could be seen  
412 (GO:0002495, GO:0002504, GO:0019886). The pathways characteristic for BMI associated  
413 loci were very much involved into response to environmental stimuli i.e. temperature  
414 (GO:0009266, GO:0009408), oxygen levels (GO:0070482, GO:0001666, GO:0036293) and  
415 sensory perception (GO:0050954, GO:0007600). The results of the GO enrichment analysis  
416 are presented in Supplementary Table S9 and Figures S4a, b.

417 At the genetic variant level, we detected SNPs in BMI associated loci which are  
418 related to GWG, and variants associated with both GWG and BMI (Analysis 1). At the same  
419 time, we performed a similar analysis with diabetes adjusted BMI and GWG, which led us to  
420 congruent findings (Analysis 2). These results point us towards inflammatory response, TGF $\beta$   
421 signaling, ER stress and glucose homeostasis. Subsequently, we investigated the association  
422 of SNPs in loci which influenced GWG (based on TWAS) with BMI and found 2091 variants  
423 in the GIANT cohort (Analysis 3). A parallel analysis was done on diabetes adjusted BMI (in  
424 the ARIC cohort) – resulting in 15 variants localized to GWG associated loci (Analysis 4).  
425 The results of these analyses show an impact of lipid biosynthesis, appetite regulation, Ca<sup>2+</sup>  
426 homeostasis (and ER stress) and inflammatory response on obesity. Our results point to the  
427 source of the genetic correlation between GWG and BMI and confirms the interconnection of  
428 the phenotypes. We compared the results of our gene focused analysis with GWAS on GWG  
429 published in 2018 (Warrington et al., 2018); even though variants from the discovery cohort  
430 did not replicate, their potential functional roles overlap with those found in our study.  
431 *TMEM163* is one of the best known genes associated with obesity, *LCORL* was shown to  
432 associate with height, *UGDH* with TGF $\beta$  signaling, *HLA-C* with autoimmune response,  
433 *HSD17B3* with fatty acids, *GLRX3* with oxidative stress, *RBM19* with ribosomal biogenesis,  
434 *SYT4* with Ca<sup>2+</sup> binding and pancreatic functioning, *PSG5* with pregnancy development,

435 while *NTS4* with neutrophilin signaling pathway. Our results are also congruent with those  
436 obtained in the computational search for functional annotation of 445 loci associated with  
437 obesity (Cheng et al., 2018). Therefore, we hypothesize that a further search for clinical  
438 phenotypes which affect GWG might allow identification of other loci that associate with it.  
439 Below we shed more light on the biological interpretation of our results and links between  
440 them (Fig. 5, 6).

441 In analyses 1 and 2, the strongest association with GWG was observed for SNPs in  
442 chemokine receptor ligands - rs1978202 – *CCL26* and rs13340504, rs11465293 - *CCL24*.  
443 Apart from variants located in these genes we identified several other (rs4742339, rs4796675,  
444 rs11235519) which serve as eQTLs for them. Chemokines are chemoattractant with the  
445 proinflammatory role. Their receptor expression was shown to correlate with BMI (being  
446 higher in obese individuals) (Ignacio, Gibbs, Lee, & Son, 2016) and BMI changes (decreased  
447 after bariatric surgery) (Gentili et al., 2016). *CCL26* is also involved in adipose tissue being  
448 in response to cold (Finlin et al., 2017). At the same time, we find three variants rs9659938,  
449 rs11807240, rs2136682 associated with GWG in the *NEXN* gene (or serving as eQTL for it)  
450 also involved in the inflammatory response. Of note, the augmentation of *NEXN* antisense  
451 RNA (*NEXN-AS1*) inhibits TLR4 oligomerization and NF $\kappa$ B activity and leads to suppression  
452 of proinflammatory response (Hu et al., 2019). *NEXN* is also known for its abundant  
453 expression in striated muscles (Zhu et al., 2018) and cardiomyocytes; its expression is higher  
454 under high glucose conditions (Barbati et al., 2017) (Figure 5).

455 Interestingly, we find GWG associated variants in genes involved in TGF $\beta$  signaling:  
456 *SPEG*, *LINC000974*, *GIPC2*. The *GIPC2* gene (rs2136682 of which associates with GWG),  
457 belongs to the GIPC family known for regulating proliferation, cytokinesis or migration and  
458 involved in the trafficking of various transmembrane proteins. *GIPC1* is necessary for cell-

459 surface expression of transmembrane receptors, such as IGF1R and TGF $\beta$ R3 (Katoh, 2013;  
460 Song et al., 2016). The inhibition of *LINC000974* leads to TGF $\beta$  secretion and repression of  
461 phosphorylated SMAD2 expression (Fang et al., 2019; Tang et al., 2014). Both TGF $\beta$  and  
462 SMADs are involved in insulin expression and signaling; SMAD4 deficiency was shown to  
463 improve glucose tolerance and glucose induced insulin release (Lin et al., 2009). Of note,  
464 three GWG associated variants (rs9690213, rs9393623, rs1978202) are eQTLs for  
465 *LINC000974* and four (rs8080053, rs1978202, rs4796675, rs8080053) are located in the gene  
466 itself (Figure 6).

467 The rs7564856, negatively associated with GWG is located in *SPEG*, and its  
468 expression is significantly greater in endurance than power athletes (Kusić et al., 2020). Also,  
469 at the protein level, SPEG differs between muscles of high-response vs low response rat  
470 trainers. The coimmunoprecipitation experiment with SPEG in the former showed a  
471 significant number of proteins involved in JNK and TGF $\beta$  signaling (Kusić et al., 2020),  
472 which, as said before, is known to influence insulin signaling and ER stress response (Z. Liu  
473 et al., 2019) (Figure 4, 5).

474 Two other variants - rs9347707 and rs4329088 - located in *PACRG1* gene, known to  
475 be involved in the insulin pathway. In *C. elegans* PACRG mutants have reduced insulin  
476 signaling. Also, PCARG plays a key role in the regulation of mitophagy (Stephenson et al.,  
477 2018) and ubiquitynylation (Loucks et al., 2016), its knockout leads to defective NF $\kappa$ B  
478 signaling (Meschede et al., 2020) and CHOP upregulation (Han et al., 2017), which in adipose  
479 tissue result in increased proinflammatory macrophage polarization (M1) and insulin  
480 resistance (Suzuki et al., 2017). Thus, we note that PACRG is also involved in ER stress. The  
481 *RHBDD2* gene is also involved in this biological process, with six eQTLs (rs4742339,  
482 rs11235519, rs1978202, rs13340504, rs4796675 and rs13340504) associated with GWG. It is

483 worth noting that silencing of *RHBDD2* leads to increased expression of ATF6, IRE1, PERK,  
484 CRT, BiP, ATF4 and CHOP (32) (Figure 4).

485 The association of rs72849841 (in *RNF213*) with GWG draws our particular attention.

486 Namely, in Akita mice (which develop diabetes spontaneously) the knockout of *RNF213*

487 lowers glucose tolerance by 20% and leads to increased insulin contents in pancreases (by

488 150%) compared to wild type animals. This is obtained via the inhibition of ubiquitinylation

489 (*RNF213* is a ubiquitin ligase) as shown by the 30% lower percentage of CHOP positive B-

490 cells (Kobayashi et al., 2013). The depletion of *RNF213* protects cells from lipotoxicity

491 (Piccolis et al., 2019), reduces palmitate-induced cell death and modifies non-mitochondrial

492 oxygen consumption rate. Moreover, *RNF213* is targeted to lipid droplets, which are

493 specialized for neutral lipid storage and increases their abundance and stability in cells

494 through the elimination of adipose triglyceride lipase (ATGL) (Sugihara et al., 2019). It is

495 interesting to note, that the latter is also negatively regulated by insulin (Kershaw et al., 2006).

496 *RNF213* variants in GWAS analyses suggestively associated (p-value ~E-03) with diabetic

497 polyneuropathy, portal hypertension, diabetic retinopathy and abnormal glucose (Figure 4). Its

498 expression differed between cyclic and pregnant heifers (Forde et al., 2012).

499 We identified GWG associated loci involved in glucose homeostasis or pancreatic

500 functioning (Figure 5) - e.g. rs11807240 in *FUBP1*, which upregulates the mRNA levels of

501 the two hexokinase genes *Hk1* and *Hk2* - the rate limiting enzymes of glycolysis. A positive

502 correlation between *FUBP1* mRNA and both of hexokinases was found in several types of

503 cancers (Kang, Lee, Kim, Lee, & Kim, 2019). Variants in *LCA5* gene were shown to modify

504 glucose response in a clinical trial of insulin and potassium (GIK) infusion in acute coronary

505 syndromes (Ellis et al., 2015). The rs9352745 associated with GWG is an eQTL for

506 *SH3BRGL2*, whose expression is associated with diabetes (type 1, type 2 and gestational) in a

507 meta-analysis study. An interaction between *SH3BRGL2* SNP and total fat intake was found  
508 to affect LDL-PPD (Rudkowska et al., 2015). We identified GWG associated SNP rs7609526  
509 in *KANK1* related to fasting proinsulin (Huyghe et al., 2013) and fasting plasma glucose  
510 (Hwang et al., 2015) in GWAS. *CADPS* (with GWG associated SNP rs1398095) plays a role  
511 in insulin secretion (Speidel et al., 2008). Cg14451730 located near *CADPS* is differentially  
512 methylated with HbA1c levels in the Taiwan Biobank cohort (Hsiung et al., 2019). The  
513 suggestive association with fasting plasma glucose of rs487321 in *CADPS* gene was found in  
514 Arabian population of T2D patients (Hebbar et al., 2020). We also found a variant correlating  
515 with GWG in *RPL29P25* gene. The presence of anti-60S ribosomal protein L29 (RPL29)  
516 antibody in human serum was shown to inhibit the proliferation of pancreatic cancer cells in  
517 various cancers and it is believed to be a novel candidate for a prognostic marker for  
518 unresectable pancreatic cancer (Muro, Miyake, Kato, Tsutsumi, & Yamamoto, 2015). The  
519 knockdown of *RPL29* leads to suppression of cell proliferation, induces cell arrest (at  
520 G0/G1phase), enhances cell apoptosis and decreases intracellular ROS generation (C. Li, Ge,  
521 Yin, Luo, & Chen, 2012). The rs4808209 (in *ZNFI* gene) is an eQTL for *LPAR2* - the  
522 receptor for a signaling lipid LPA. It is a part of the LPAR2/Gab1/PI3K/Akt pathway, which  
523 influences glucose uptake (Rodriguez-Araujo et al., 2013) and is differentially expressed in  
524 fatty vs normal liver in extreme obese cohort (DiStefano et al., 2014). Its expression correlates  
525 with the invasiveness of pancreatic cancers (Gong et al., 2012) as well as gynecological  
526 disorders (Fujii et al., 2019; Kowalczyk-Zieba et al., 2019; Wasniewski & Woclawek-  
527 Potocka, 2018; X. Yu, Zhang, & Chen, 2016).

528 In Analyses 3 and 4, we set to determine whether genes associated with GWG also  
529 impact BMI. Basing on the summary results from the GIANT cohort we show a number of  
530 such genes. Some of their SNPs were previously associated with BMI, body fat distribution,

531 attendance to gym or sport groups, sleep duration, height, lipoprotein levels, low HDL-level  
532 cholesterol. At the same time, in a much smaller ARIC cohort we find 15 variants associated  
533 with diabetes adjusted BMI. None of them is listed in GWAS Catalog or PheWas, however  
534 other variants in these genes were already mentioned to be associated with BMI.

535 Our analysis shows association with rs12685009 in the *SLC24A2* loci, which belongs to  
536 the family of  $K^+$  dependent  $Na^+/Ca^{2+}$  exchanger (Schnetkamp, 2013). A member of the  
537 family, *SLC4* influences MC4R dependent satiety and the loss of MC4R function leads to  
538 prolonged obesity and reduced energy expenditure, while mice lacking *SLC4* display anorexia  
539 (X. F. Li & Lytton, 2014). Also, variants in *SOX6*, a transcription factor shown to be involved  
540 in the promotion of adipogenesis (Iguchi et al., 2005; Leow et al., 2016), were previously  
541 related to obesity (Correa-Rodríguez, Schmidt-RioValle, & Rueda-Medina, 2018; Y. Z. Liu et  
542 al., 2009). The rs77346003 in the *ADIPOR1* loci - a receptor for adiponectin known to  
543 regulate glucose metabolism and fatty acid oxidation associates with BMI in our study. Its  
544 variants in GWAS are related to obesity (Beckers et al., 2013; Keustermans et al., 2017;  
545 Peters et al., 2013) as well as fetal weight (Fensterseifer, Austin, Ford, & Alexander, 2018;  
546 Muñoz-Muñoz, Krause, Uauy, & Casanello, 2018). *ADIPOR1* expression correlates with  
547 improved insulin sensitivity and PGC-1A expression - a master regulator of mitochondrial  
548 gene expression (Za'don et al., 2019). Moreover, our analysis identifies eQTLs for  
549 mitochondria associated genes – *ATP5G3* (rs144768710) and *HSPB8* (rs4131847). *ATP5G3*  
550 is involved in the proton pathway and acts as an energy-driving motor (He et al., 2017; Huang  
551 et al., 2013; Spataru, Le Duc, Zagrean, & Zagrean, 2019). *HSPB8* is known to prevent  
552 oxidative tissue damage and its expression in serum is used as a biomarker for virus induced  
553 type 1 diabetes (Karthik, Ilavenil, Kaleeswaran, Sunil, & Ravikumar, 2012; X. C. Li et al.,  
554 2017; L. Yu et al., 2019). We also identified rs34427781 - an eQTL for several genes

555 (*CYP2D8P*, *NDUFA6-AS1*, *CYP2D6*, *CCDC134*, *CENPM*, *CHADL*, *FAM109B*, *NAGA*,  
556 *NDUFA6*, *OGFRP1*, *OLA1P1*, *SEPT3*, *SHISA8*, *SLC25A5P1*, *SMDT1*, *SREBF2*,  
557 *TNFRSF13C*, *WBP2NL*) to be associated with BMI. It is worth noting that a subset of these  
558 genes is involved in lipid metabolism (*NAGA*, *SREBP*) or mitochondrial signaling  
559 (*SLC25A5P1*, *NDUFA6*, *SMDT1*). *SREBP* is a master regulator of lipid and sterol  
560 biosynthesis (Düvel et al., 2010); the gene-gene interactions between variants in the *INSIG*-  
561 *SCAP-SREBP* pathway are associated with risk of obesity in Chinese children (F. H. Liu et  
562 al., 2014). The *SMDT1* gene builds the mitochondrial calcium uniporter subunit (mtCU)  
563 localized in the inner mitochondrial membrane and is responsible for the  $\text{Ca}^{2+}$  transport to the  
564 mitochondrial matrix (Pandin, Greotti, & Pozzan, 2014), while its expression was shown to  
565 associate with fetal development (Vishnyakova et al., 2019). *SLC25* genes, members of *SLC*  
566 channels mentioned above, are known to be subunits of another complex essential for proper  
567 mitochondrial dynamics and energy production – *ANC*. *ANC* is responsible for  
568 mitochondrial/cytoplasmic ADP/ATP exchange (Clémentçon, Babot, & Trézéguet, 2013) and  
569 its expression was shown to be consistently upregulated in obesity (Padilla et al., 2014).  
570 *NDUFA6* is part of the first complex of the mitochondrial respiratory chain, which expression  
571 impacts oxidative stress and energy production efficacy (Fiedorczuk & Sazanov, 2018).  
572 Lastly, the *TNFRSF13C* (*BAFF*) gene regulates insulin sensitivity (Kawasaki et al., 2013) and  
573 is associated with autoimmune diseases (Moisini & Davidson, 2009).

574 The above results are congruent with the gene-based fBAT analysis, that points us  
575 towards genes which correlate with BMI (*GATAD2*, *LHFPL6*) and immune  
576 response/autoimmunity (*HLA-DQB*, *HLA-DRB*, *HLA-B*). *GATAD2A* is involved in  
577 embryonic development (Wang et al., 2017) and associates with obesity (Saxena et al., 2012).  
578 A copy number variation of *LHFPL6* associates with average daily gain in cattle (Xu et al.,

579 2019), while its rs4073643 with systemic lupus erythematosus in the Chinese population  
580 (Zhang et al., 2016). While the association between HLA variants and T1DM is very well  
581 known (Zhao et al., 2016), HLA loci are linked to other phenotypes (Karnes et al., 2017): type  
582 2 diabetes (Ng et al., 2014), fatty liver disease (Doganay et al., 2014) as well as BMI (Shen et  
583 al., 2018) or waist-to-hip circumference (Wen et al., 2016). More importantly, these variants  
584 associate with trends in fetal birth weight (Capittini et al., 2009), level of inflammation in  
585 visceral adipose tissue in pregnant women (Eyerahi et al., 2018) and different mRNA levels in  
586 visceral omental adipose tissue of pregnant women with gestational diabetes (Deng et al.,  
587 2018). The *HLA-DQAI*, *HLADQB1* are differentially methylated in siblings born before vs  
588 after maternal bariatric surgery (Berglind et al., 2016).

589 In this study we identified several loci which contribute to the genetic correlation  
590 between BMI and GWG. Variants identified in those loci are associated with genes linked to  
591 insulin signaling, glucose homeostasis, mitochondrial metabolism, ubiquitinylation and  
592 inflammatory responses and placenta functioning, not only in the diabetic cohorts, but also in  
593 the general population. The genetic contribution to GWG is clearly connected with BMI  
594 associated loci.

595

596

## 597 REFERENCES

- 598 Andersson, E. S., Silventoinen, K., Tynelius, P., Nohr, E. A., Sørensen, T. I. A., &  
599 Rasmussen, F. (2015). Heritability of Gestational Weight Gain - A Swedish Register-  
600 Based Twin Study. *Twin Research and Human Genetics*.  
601 <https://doi.org/10.1017/thg.2015.38>  
602 Barbati, S. A., Colussi, C., Bacci, L., Aiello, A., Re, A., Stigliano, E., ... Nanni, S. (2017).

- 603 Transcription factor CREM mediates high glucose response in cardiomyocytes and in a  
604 male mouse model of prolonged hyperglycemia. *Endocrinology*.  
605 <https://doi.org/10.1210/en.2016-1960>
- 606 Bashir, M., Naem, E., Taha, F., Konje, J. C., & Abou-Samra, A. B. (2019). Outcomes of type  
607 1 diabetes mellitus in pregnancy; effect of excessive gestational weight gain and  
608 hyperglycaemia on fetal growth. *Diabetes and Metabolic Syndrome: Clinical Research  
609 and Reviews*. <https://doi.org/10.1016/j.dsx.2018.08.030>
- 610 Beckers, S., De Freitas, F., Zegers, D., Mertens, I. L., Verrijken, A., Van Camp, J. K., ... Van  
611 Hul, W. (2013). No conclusive evidence for association of polymorphisms in the  
612 adiponectin receptor 1 gene, AdipoR1, with common obesity. *Endocrine*.  
613 <https://doi.org/10.1007/s12020-012-9736-6>
- 614 Berglind, D., Müller, P., Willmer, M., Sinha, I., Tynelius, P., Näslund, E., ... Rasmussen, F.  
615 (2016). Differential methylation in inflammation and type 2 diabetes genes in siblings  
616 born before and after maternal bariatric surgery. *Obesity*.  
617 <https://doi.org/10.1002/oby.21340>
- 618 Capittini, C., Pasi, A., Bergamaschi, P., Tinelli, C., De Silvestri, A., Mercati, M. P., ...  
619 Cuccia, M. (2009). HLA haplotypes and birth weight variation: Is your future going to be  
620 light or heavy? *Tissue Antigens*. <https://doi.org/10.1111/j.1399-0039.2009.01282.x>
- 621 Celia, B., Richard, H., Chineze, O., Casey, E., Elizabeth, P., & Macauley, A. (2016). The St  
622 Vincent Declaration–25 years on: review of maternal deaths reported to the Confidential  
623 Enquiry in the UK and the lessons learnt. *European Journal of Obstetrics & Gynecology  
624 and Reproductive Biology*. <https://doi.org/10.1016/j.ejogrb.2016.07.354>
- 625 Cheng, M., Mei, B., Zhou, Q., Zhang, M., Huang, H., Han, L., & Huang, Q. (2018).  
626 Computational analyses of obesity associated loci generated by genome-wide association

- 627 studies. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0199987>
- 628 Clémentçon, B., Babot, M., & Trézéguet, V. (2013). The mitochondrial ADP/ATP carrier  
629 (SLC25 family): Pathological implications of its dysfunction. *Molecular Aspects of*  
630 *Medicine*. <https://doi.org/10.1016/j.mam.2012.05.006>
- 631 Correa-Rodríguez, M., Schmidt-RioValle, J., & Rueda-Medina, B. (2018). SOX6 rs7117858  
632 polymorphism is associated with osteoporosis and obesity-related phenotypes. *European*  
633 *Journal of Clinical Investigation*. <https://doi.org/10.1111/eci.13011>
- 634 Deng, X., Yang, Y., Sun, H., Qi, W., Duan, Y., & Qian, Y. (2018). Analysis of whole  
635 genome-wide methylation and gene expression profiles in visceral omental adipose  
636 tissue of pregnancies with gestational diabetes mellitus. *Journal of the Chinese Medical*  
637 *Association*. <https://doi.org/10.1016/j.jcma.2017.06.027>
- 638 DiStefano, J. K., Kingsley, C., Craig Wood, G., Chu, X., Argyropoulos, G., Still, C. D., ...  
639 Gerhard, G. S. (2014). Genome-wide analysis of hepatic lipid content in extreme obesity.  
640 *Acta Diabetologica*. <https://doi.org/10.1007/s00592-014-0654-3>
- 641 Doganay, L., Katrinli, S., Colak, Y., Senates, E., Zemheri, E., Ozturk, O., ... Doganay, G. D.  
642 (2014). HLA DQB1 alleles are related with nonalcoholic fatty liver disease. *Molecular*  
643 *Biology Reports*. <https://doi.org/10.1007/s11033-014-3688-2>
- 644 Dori-Dayan, N., Cukierman-Yaffe, T., Zemet, R., Cohen, O., Mazaki-Tovi, S., & Yoeli-  
645 Ullman, R. (2020). 1013: Gestational weight gain does-not affect insulin requirement  
646 during pregnancy in women with Type 1 diabetes. *American Journal of Obstetrics and*  
647 *Gynecology*. <https://doi.org/10.1016/j.ajog.2019.11.1029>
- 648 Düvel, K., Yecies, J. L., Menon, S., Raman, P., Lipovsky, A. I., Souza, A. L., ... Manning, B.  
649 D. (2010). Activation of a metabolic gene regulatory network downstream of mTOR  
650 complex 1. *Molecular Cell*. <https://doi.org/10.1016/j.molcel.2010.06.022>

- 651 Ellis, K. L., Zhou, Y., Beshansky, J. R., Ainehsazan, E., Selker, H. P., Cupples, L. A., ...  
652 Peter, I. (2015). Genetic modifiers of response to glucose-insulin-potassium (GIK)  
653 infusion in acute coronary syndromes and associations with clinical outcomes in the  
654 IMMEDIATE trial. *Pharmacogenomics Journal*. <https://doi.org/10.1038/tpj.2015.10>
- 655 Eyerahi, B. F., Mancilla-Herrera, I., Espino Y Sosa, S., Ortiz-Ramirez, M., Flores-Rueda, V.,  
656 Ibarguengoitia-Ochoa, F., ... Estrada-Gutierrez, G. (2018). Macrophage populations in  
657 visceral adipose tissue from pregnant women: Potential role of obesity in maternal  
658 inflammation. *International Journal of Molecular Sciences*.  
659 <https://doi.org/10.3390/ijms19041074>
- 660 Fang, C. Y., Yu, C. C., Liao, Y. W., Hsieh, P. L., Lu, M. Y., Lin, K. C., ... Tsai, L. L. (2019).  
661 LncRNA LINC00974 activates TGF- $\beta$ /Smad signaling to promote oral fibrogenesis.  
662 *Journal of Oral Pathology and Medicine*. <https://doi.org/10.1111/jop.12805>
- 663 Fensterseifer, S. R., Austin, K. J., Ford, S. P., & Alexander, B. M. (2018). Effects of maternal  
664 obesity on maternal and fetal plasma concentrations of adiponectin and expression of  
665 adiponectin and its receptor genes in cotyledonary and adipose tissues at mid- and late-  
666 gestation in sheep. *Animal Reproduction Science*.  
667 <https://doi.org/10.1016/j.anireprosci.2018.08.033>
- 668 Fiedorczuk, K., & Sazanov, L. A. (2018). Mammalian Mitochondrial Complex I Structure and  
669 Disease-Causing Mutations. *Trends in Cell Biology*.  
670 <https://doi.org/10.1016/j.tcb.2018.06.006>
- 671 Finlin, B. S., Zhu, B., Confides, A. L., Westgate, P. M., Harfmann, B. D., Dupont-  
672 Versteegden, E. E., & Kern, P. A. (2017). Mast cells promote seasonal white adipose  
673 beiging in humans. *Diabetes*. <https://doi.org/10.2337/db16-1057>
- 674 Forde, N., Duffy, G. B., McGettigan, P. A., Browne, J. A., Mehta, J. P., Kelly, A. K., ...

- 675 Evans, A. C. O. (2012). Evidence for an early endometrial response to pregnancy in  
676 cattle: Both dependent upon and independent of interferon tau. *Physiological Genomics*.  
677 <https://doi.org/10.1152/physiolgenomics.00067.2012>
- 678 Fujii, T., Nagamatsu, T., Schust, D. J., Ichikawa, M., Kumasawa, K., Yabe, S., ... Fujii, T.  
679 (2019). Placental expression of lysophosphatidic acid receptors in normal pregnancy and  
680 preeclampsia. *American Journal of Reproductive Immunology*.  
681 <https://doi.org/10.1111/aji.13176>
- 682 Gamazon, E. R., Wheeler, H. E., Shah, K. P., Mozaffari, S. V., Aquino-Michaels, K., Carroll,  
683 R. J., ... Im, H. K. (2015). A gene-based association method for mapping traits using  
684 reference transcriptome data. *Nature Genetics*. <https://doi.org/10.1038/ng.3367>
- 685 Gentili, A., Zaibi, M. S., Alomar, S. Y., De Vuono, S., Ricci, M. A., Alaeddin, A., ...  
686 Lupattelli, G. (2016). Circulating Levels of the Adipokines Monocyte Chemotactic  
687 Protein-4 (MCP-4), Macrophage Inflammatory Protein-1 $\beta$  (MIP-1 $\beta$ ), and Eotaxin-3 in  
688 Severe Obesity and Following Bariatric Surgery. *Hormone and Metabolic Research*.  
689 <https://doi.org/10.1055/s-0042-108731>
- 690 Gong, Y. L., Tao, C. J., Hu, M., Chen, J. F., Cao, X. F., Lv, G. M., & Li, P. (2012).  
691 Expression of lysophosphatidic acid receptors and local invasiveness and metastasis in  
692 Chinese pancreatic cancers. *Current Oncology*. <https://doi.org/10.3747/co.19.1138>
- 693 Han, K., Hassanzadeh, S., Singh, K., Menazza, S., Nguyen, T. T., Stevens, M. V., ... Sack,  
694 M. N. (2017). Parkin regulation of CHOP modulates susceptibility to cardiac  
695 endoplasmic reticulum stress. *Scientific Reports*. [https://doi.org/10.1038/s41598-017-](https://doi.org/10.1038/s41598-017-02339-2)  
696 [02339-2](https://doi.org/10.1038/s41598-017-02339-2)
- 697 He, J., Ford, H. C., Carroll, J., Ding, S., Fearnley, I. M., & Walker, J. E. (2017). Persistence of  
698 the mitochondrial permeability transition in the absence of subunit c of human ATP

699 synthase. *Proceedings of the National Academy of Sciences of the United States of*  
700 *America*. <https://doi.org/10.1073/pnas.1702357114>

701 Hebbar, P., Abu-Farha, M., Alkayal, F., Nizam, R., Elkum, N., Melhem, M., ... Thanaraj, T.  
702 A. (2020). Genome-wide association study identifies novel risk variants from RPS6KA1,  
703 CADPS, VARS, and DHX58 for fasting plasma glucose in Arab population. *Scientific*  
704 *Reports*. <https://doi.org/10.1038/s41598-019-57072-9>

705 Hormozdiari, F., van de Bunt, M., Segrè, A. V., Li, X., Joo, J. W. J., Bilow, M., ... Eskin, E.  
706 (2016). Colocalization of GWAS and eQTL Signals Detects Target Genes. *American*  
707 *Journal of Human Genetics*. <https://doi.org/10.1016/j.ajhg.2016.10.003>

708 Hsiung, C.-N., Chang, Y.-C., Lin, C.-W., Chang, C.-W., Chou, W.-C., Chu, H.-W., ... Shen,  
709 C.-Y. (2019). The Causal Relationship of Circulating Triglycerides and Glycated  
710 Hemoglobin: A Mendelian Randomization Study. *The Journal of Clinical Endocrinology*  
711 *& Metabolism*. <https://doi.org/10.1210/clinem/dgz243>

712 Hu, Y. W., Guo, F. X., Xu, Y. J., Li, P., Lu, Z. F., McVey, D. G., ... Ye, S. (2019). Long  
713 noncoding RNA NEXN-AS1 mitigates atherosclerosis by regulating the actin-binding  
714 protein NEXN. *Journal of Clinical Investigation*. <https://doi.org/10.1172/JCI98230>

715 Huang, Y., Wang, L., Bennett, B., Williams, R. W., Wang, Y. J., Gu, W. K., & Jiao, Y.  
716 (2013). Potential role of Atp5g3 in epigenetic regulation of alcohol preference or obesity  
717 from a mouse genomic perspective. *Genetics and Molecular Research*.  
718 <https://doi.org/10.4238/2013.September.18.1>

719 Huyghe, J. R., Jackson, A. U., Fogarty, M. P., Buchkovich, M. L., Stančáková, A., Stringham,  
720 H. M., ... Mohlke, K. L. (2013). Exome array analysis identifies new loci and low-  
721 frequency variants influencing insulin processing and secretion. *Nature Genetics*.  
722 <https://doi.org/10.1038/ng.2507>

- 723 Hwang, J. Y., Sim, X., Wu, Y., Liang, J., Tabara, Y., Hu, C., ... Kim, B. J. (2015). Genome-  
724 wide association meta-analysis identifies novel variants associated with fasting plasma  
725 glucose in East Asians. *Diabetes*. <https://doi.org/10.2337/db14-0563>
- 726 Ignacio, R. M. C., Gibbs, C. R., Lee, E. S., & Son, D. S. (2016). Differential chemokine  
727 signature between human preadipocytes and adipocytes. *Immune Network*.  
728 <https://doi.org/10.4110/in.2016.16.3.189>
- 729 Iguchi, H., Ikeda, Y., Okamura, M., Tanaka, T., Urashima, Y., Ohguchi, H., ... Sakai, J.  
730 (2005). SOX6 attenuates glucose-stimulated insulin secretion by repressing PDX1  
731 transcriptional activity and is down-regulated in hyperinsulinemic obese mice. *Journal of*  
732 *Biological Chemistry*. <https://doi.org/10.1074/jbc.M505392200>
- 733 *Influence of Pregnancy Weight on Maternal and Child Health*. (2007). *Influence of*  
734 *Pregnancy Weight on Maternal and Child Health*. <https://doi.org/10.17226/11817>
- 735 Kang, M., Lee, S. M., Kim, W., Lee, K. H., & Kim, D. Y. (2019). Fubp1 supports the lactate-  
736 Akt-mTOR axis through the upregulation of Hk1 and Hk2. *Biochemical and Biophysical*  
737 *Research Communications*. <https://doi.org/10.1016/j.bbrc.2019.03.005>
- 738 Karnes, J. H., Bastarache, L., Shaffer, C. M., Gaudieri, S., Xu, Y., Glazer, A. M., ... Denny, J.  
739 C. (2017). Phenome-wide scanning identifies multiple diseases and disease severity  
740 phenotypes associated with HLA variants. *Science Translational Medicine*.  
741 <https://doi.org/10.1126/scitranslmed.aai8708>
- 742 Karthik, D., Ilavenil, S., Kaleeswaran, B., Sunil, S., & Ravikumar, S. (2012). Proteomic  
743 analysis of plasma proteins in diabetic rats by 2D electrophoresis and MALDI-TOF-MS.  
744 *Applied Biochemistry and Biotechnology*. <https://doi.org/10.1007/s12010-012-9544-8>
- 745 Katoh, M. (2013). Functional proteomics, human genetics and cancer biology of GIPC family  
746 members. *Experimental and Molecular Medicine*. <https://doi.org/10.1038/emm.2013.49>

- 747 Kawai, V. K., Nwosu, S. K., Kurnik, D., Harrell, F. E., & Stein, C. M. (2019). Variants in  
748 BMI-Associated Genes and Adrenergic Genes are not Associated with Gestational  
749 Weight Trajectory. *Obesity*. <https://doi.org/10.1002/oby.22505>
- 750 Kawasaki, K., Abe, M., Tada, F., Tokumoto, Y., Chen, S., Miyake, T., ... Onji, M. (2013).  
751 Blockade of B-cell-activating factor signaling enhances hepatic steatosis induced by a  
752 high-fat diet and improves insulin sensitivity. *Laboratory Investigation*.  
753 <https://doi.org/10.1038/labinvest.2012.176>
- 754 Kershaw, E. E., Hamm, J. K., Verhagen, L. A. W., Peroni, O., Katic, M., & Flier, J. S. (2006).  
755 Adipose triglyceride lipase: Function, regulation by insulin, and comparison with  
756 adiponutrin. *Diabetes*. <https://doi.org/10.2337/diabetes.55.01.06.db05-0982>
- 757 Keustermans, G., Van Der Heijden, L. B., Boer, B., Scholman, R., Nuboer, R., Pasterkamp,  
758 G., ... Schipper, H. S. (2017). Differential adipokine receptor expression on circulating  
759 leukocyte subsets in lean and obese children. *PLoS ONE*.  
760 <https://doi.org/10.1371/journal.pone.0187068>
- 761 Kobayashi, H., Yamazaki, S., Takashima, S., Liu, W., Okuda, H., Yan, J., ... Koizumi, A.  
762 (2013). Ablation of Rnf213 retards progression of diabetes in the Akita mouse.  
763 *Biochemical and Biophysical Research Communications*.  
764 <https://doi.org/10.1016/j.bbrc.2013.02.015>
- 765 Kominiarek, M. A., & Peaceman, A. M. (2017). Gestational weight gain. *American Journal of*  
766 *Obstetrics and Gynecology*. <https://doi.org/10.1016/j.ajog.2017.05.040>
- 767 Kowalczyk-Zieba, I., Woclawek-Potocka, I., Wasniewski, T., Boruszewska, D., Grycmacher,  
768 K., Sinderewicz, E., ... Wolczynski, S. (2019). LPAR2 and LPAR4 are the Main  
769 Receptors Responsible for LPA Actions in Ovarian Endometriotic Cysts. *Reproductive*  
770 *Sciences*. <https://doi.org/10.1177/1933719118766263>

- 771 Kusić, D., Connolly, J., Kainulainen, H., Semenova, E. A., Borisov, O. V., Larin, A. K., ...  
772 Burniston, J. G. (2020). Striated muscle-specific serine/threonine-protein kinase beta  
773 segregates with high versus low responsiveness to endurance exercise training.  
774 *Physiological Genomics*. <https://doi.org/10.1152/physiolgenomics.00103.2019>
- 775 Lawlor, D. A., Fraser, A., Macdonald-Wallis, C., Nelson, S. M., Palmer, T. M., Smith, G. D.,  
776 & Tilling, K. (2011). Maternal and offspring adiposity-related genetic variants and  
777 gestational weight gain. *American Journal of Clinical Nutrition*.  
778 <https://doi.org/10.3945/ajcn.110.010751>
- 779 Leow, S. C., Poschmann, J., Too, P. G., Yin, J., Joseph, R., McFarlane, C., ... Stükel, W.  
780 (2016). The transcription factor SOX6 contributes to the developmental origins of  
781 obesity by promoting adipogenesis. *Development (Cambridge)*.  
782 <https://doi.org/10.1242/dev.131573>
- 783 Li, B., Verma, S. S., Veturi, Y. C., Verma, A., Bradford, Y., Haas, D. W., & Ritchie, M. D.  
784 (2018). Evaluation of predixcan for prioritizing GWAS associations and predicting gene  
785 expression. In *Pacific Symposium on Biocomputing*.  
786 [https://doi.org/10.1142/9789813235533\\_0041](https://doi.org/10.1142/9789813235533_0041)
- 787 Li, C., Ge, M., Yin, Y., Luo, M., & Chen, D. (2012). Silencing expression of ribosomal  
788 protein L26 and L29 by RNA interfering inhibits proliferation of human pancreatic  
789 cancer PANC-1 cells. *Molecular and Cellular Biochemistry*.  
790 <https://doi.org/10.1007/s11010-012-1404-x>
- 791 Li, X. C., Hu, Q. K., Chen, L., Liu, S. Y., Su, S., Tao, H., ... He, L. J. (2017). HSPB8  
792 promotes the fusion of autophagosome and lysosome during autophagy in diabetic  
793 neurons. *International Journal of Medical Sciences*. <https://doi.org/10.7150/ijms.20653>
- 794 Li, X. F., & Lytton, J. (2014). An essential role for the K<sup>+</sup>-dependent Na<sup>+</sup>/Ca<sup>2+</sup>-exchanger,

- 795 NCKX4, in melanocortin-4-receptor-dependent satiety. *Journal of Biological Chemistry*.  
796 <https://doi.org/10.1074/jbc.M114.564450>
- 797 Lin, H. M., Lee, J. H., Yadav, H., Kamaraju, A. K., Liu, E., Zhigang, D., ... Rane, S. G.  
798 (2009). Transforming growth factor- $\beta$ /Smad3 signaling regulates insulin gene  
799 transcription and pancreatic islet  $\beta$ -cell function. *Journal of Biological Chemistry*.  
800 <https://doi.org/10.1074/jbc.M805379200>
- 801 Liu, F. H., Song, J. Y., Shang, X. R., Meng, X. R., Ma, J., & Wang, H. J. (2014). The gene-  
802 gene interaction of INSIG-SCAP-SREBP pathway on the risk of obesity in Chinese  
803 children. *BioMed Research International*. <https://doi.org/10.1155/2014/538564>
- 804 Liu, Y. Z., Pei, Y. F., Liu, J. F., Yang, F., Guo, Y., Zhang, L., ... Deng, H. W. (2009).  
805 Powerful bivariate Genome-wide association analyses suggest the SOX6 gene  
806 influencing both obesity and osteoporosis phenotypes in males. *PLoS ONE*.  
807 <https://doi.org/10.1371/journal.pone.0006827>
- 808 Liu, Z., Li, C., Kang, N., Malhi, H., Shah, V. H., & Maiers, J. L. (2019). Transforming  
809 growth factor (TGF) cross-talk with the unfolded protein response is critical for hepatic  
810 stellate cell activation. *Journal of Biological Chemistry*.  
811 <https://doi.org/10.1074/jbc.RA118.005761>
- 812 Locke, A., Kahali, B., Berndt, S., Justice, A., & Pers, T. (2015). Genetic studies of body mass  
813 index yield new insights for obesity biology. *Nature*, *518*(7538), 197–206.  
814 <https://doi.org/10.1038/nature14177>.Genetic
- 815 Loucks, C. M., Bialas, N. J., Dekkers, M. P. J., Walker, D. S., Grundy, L. J., Li, C., ...  
816 Leroux, M. R. (2016). PACRG, a protein linked to ciliary motility, mediates cellular  
817 signaling. *Molecular Biology of the Cell*. <https://doi.org/10.1091/mbc.E15-07-0490>
- 818 Ludwig-Slomczynska, A. H., Seweryn, M. T., Kapusta, P., Pitera, E., Mantaj, U., Cyganek,

- 819 K., ... Wolkow, P. P. (2018). Mitochondrial GWAS and association of nuclear-  
820 mitochondrial epistasis with BMI in T1DM patients. *BioRxiv*.  
821 <https://doi.org/10.1101/436519>
- 822 Luecke, E., Cohen, A. K., Brillante, M., Rehkopf, D. H., Coyle, J., Hendrick, C. E., &  
823 Abrams, B. (2018). Similarities in Maternal Weight and Birth Weight Across  
824 Pregnancies and Across Sisters. *Maternal and Child Health Journal*.  
825 <https://doi.org/10.1007/s10995-018-2602-2>
- 826 Mancuso, N., Shi, H., Goddard, P., Kichaev, G., Gusev, A., & Pasaniuc, B. (2017).  
827 Integrating Gene Expression with Summary Association Statistics to Identify Genes  
828 Associated with 30 Complex Traits. *American Journal of Human Genetics*.  
829 <https://doi.org/10.1016/j.ajhg.2017.01.031>
- 830 Mastella, L. S., Weinert, L. S., Gnielka, V., Hirakata, V. N., Oppermann, M. L. R., Silveiro,  
831 S. P., & Reichelt, A. J. (2018). Influence of maternal weight gain on birth weight: A  
832 gestational diabetes cohort. *Archives of Endocrinology and Metabolism*.  
833 <https://doi.org/10.20945/2359-3997000000009>
- 834 McWhorter, K. L., Bowers, K., Dolan, L. M., Deka, R., Jackson, C. L., & Khoury, J. C.  
835 (2018). Impact of gestational weight gain and prepregnancy body mass index on the  
836 prevalence of large-for-gestational age infants in two cohorts of women with type 1  
837 insulin-dependent diabetes: A cross-sectional population study. *BMJ Open*.  
838 <https://doi.org/10.1136/bmjopen-2017-019617>
- 839 Meschede, J., Šadić, M., Furthmann, N., Miedema, T., Sehr, D. A., Müller-Rischart, A. K., ...  
840 Winklhofer, K. F. (2020). The *parkin*-coregulated gene product PACRG promotes TNF  
841 signaling by stabilizing LUBAC. *Science Signaling*.  
842 <https://doi.org/10.1126/scisignal.aav1256>

- 843 Moisini, I., & Davidson, A. (2009). BAFF: A local and systemic target in autoimmune  
844 diseases. *Clinical and Experimental Immunology*. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2249.2009.04007.x)  
845 [2249.2009.04007.x](https://doi.org/10.1111/j.1365-2249.2009.04007.x)
- 846 Muñoz-Muñoz, E., Krause, B. J., Uauy, R., & Casanello, P. (2018). LGA-newborn from  
847 patients with pregestational obesity present reduced adiponectin-mediated vascular  
848 relaxation and endothelial dysfunction in fetoplacental arteries. *Journal of Cellular*  
849 *Physiology*. <https://doi.org/10.1002/jcp.26499>
- 850 Muro, S., Miyake, Y., Kato, H., Tsutsumi, K., & Yamamoto, K. (2015). Serum anti-60s  
851 ribosomal protein L29 antibody as a novel prognostic marker for unresectable pancreatic  
852 cancer. *Digestion*. <https://doi.org/10.1159/000371545>
- 853 Ng, M. C. Y., Shriner, D., Chen, B. H., Li, J., Chen, W. M., Guo, X., ... Surdulescu, G.  
854 (2014). Meta-Analysis of Genome-Wide Association Studies in African Americans  
855 Provides Insights into the Genetic Architecture of Type 2 Diabetes. *PLoS Genetics*.  
856 <https://doi.org/10.1371/journal.pgen.1004517>
- 857 Nunnery, D., Ammerman, A., & Dharod, J. (2018). Predictors and outcomes of excess  
858 gestational weight gain among low-income pregnant women. *Health Care for Women*  
859 *International*. <https://doi.org/10.1080/07399332.2017.1391263>
- 860 Padilla, J., Jenkins, N. T., Thorne, P. K., Martin, J. S., Scott Rector, R., Wade Davis, J., &  
861 Harold Laughlin, M. (2014). Identification of genes whose expression is altered by  
862 obesity throughout the arterial tree. *Physiological Genomics*.  
863 <https://doi.org/10.1152/physiolgenomics.00091.2014>
- 864 Pendin, D., Greotti, E., & Pozzan, T. (2014). The elusive importance of being a mitochondrial  
865 Ca<sup>2+</sup> uniporter. *Cell Calcium*. <https://doi.org/10.1016/j.ceca.2014.02.008>
- 866 Peters, K. E., Beilby, J., Cadby, G., Warrington, N. M., Bruce, D. G., Davis, W. A., ... Hung,

867 J. (2013). A comprehensive investigation of variants in genes encoding adiponectin  
868 (ADIPOQ) and its receptors (ADIPOR1/R2), and their association with serum  
869 adiponectin, type 2 diabetes, insulin resistance and the metabolic syndrome. *BMC*  
870 *Medical Genetics*. <https://doi.org/10.1186/1471-2350-14-15>

871 Piccolis, M., Bond, L. M., Kampmann, M., Pulimeno, P., Chitraju, C., Jayson, C. B. K., ...  
872 Farese, R. V. (2019). Probing the Global Cellular Responses to Lipotoxicity Caused by  
873 Saturated Fatty Acids. *Molecular Cell*. <https://doi.org/10.1016/j.molcel.2019.01.036>

874 Rodriguez-Araujo, G., Nakagami, H., Hayashi, H., Mori, M., Shiuchi, T., Minokoshi, Y., ...  
875 Kaneda, Y. (2013). Alpha-synuclein elicits glucose uptake and utilization in adipocytes  
876 through the Gab1/PI3K/Akt transduction pathway. *Cellular and Molecular Life Sciences*.  
877 <https://doi.org/10.1007/s00018-012-1198-8>

878 Rudkowska, I., Pérusse, L., Bellis, C., Blangero, J., Després, J. P., Bouchard, C., & Vohl, M.  
879 C. (2015). Interaction between Common Genetic Variants and Total Fat Intake on Low-  
880 Density Lipoprotein Peak Particle Diameter: A Genome-Wide Association Study.  
881 *Journal of Nutrigenetics and Nutrigenomics*. <https://doi.org/10.1159/000431151>

882 Rys, P. M., Ludwig-Slomczynska, A. H., Cyganek, K., & Malecki, M. T. (2018). Continuous  
883 subcutaneous insulin infusion vs multiple daily injections in pregnant women with type 1  
884 diabetes mellitus: A systematic review and meta-analysis of randomised controlled trials  
885 and observational studies. *European Journal of Endocrinology*.  
886 <https://doi.org/10.1530/EJE-17-0804>

887 Saxena, R., Elbers, C. C., Guo, Y., Peter, I., Gaunt, T. R., Mega, J. L., ... Keating, B. J.  
888 (2012). Large-scale gene-centric meta-analysis across 39 studies identifies type 2  
889 diabetes loci. *American Journal of Human Genetics*.  
890 <https://doi.org/10.1016/j.ajhg.2011.12.022>

- 891 Schnetkamp, P. P. M. (2013). The SLC24 gene family of Na<sup>+</sup>/Ca<sup>2+</sup>-K<sup>+</sup> exchangers: From  
892 sight and smell to memory consolidation and skin pigmentation. *Molecular Aspects of*  
893 *Medicine*. <https://doi.org/10.1016/j.mam.2012.07.008>
- 894 Scifres, C. M., Feghali, M. N., Althouse, A. D., Caritis, S. N., & Catov, J. M. (2014). Effect  
895 of excess gestational weight gain on pregnancy outcomes in women with type 1 diabetes.  
896 *Obstetrics and Gynecology*. <https://doi.org/10.1097/AOG.0000000000000271>
- 897 Secher, A. L., Parellada, C. B., Ringholm, L., Ásbjörnsdóttir, B., Damm, P., & Mathiesen, E.  
898 R. (2014). Higher gestational weight gain is associated with increasing offspring birth  
899 weight independent of maternal glycemic control in women with type 1 diabetes.  
900 *Diabetes Care*. <https://doi.org/10.2337/dc14-0896>
- 901 Shen, J., Guo, T., Wang, T., Zhen, Y., Ma, X., Wang, Y., ... Baier, L. J. (2018). HLA-B\*07,  
902 HLA-DRB1\*07, HLA-DRB1\*12, and HLA-c\*03:02 strongly associate with BMI: Data  
903 from 1.3 million healthy Chinese adults. *Diabetes*. <https://doi.org/10.2337/db17-0852>
- 904 Siega-Riz, A. M., Bodnar, L. M., Stotland, N. E., & Stang, J. (2020). The Current  
905 Understanding of Gestational Weight Gain Among Women with Obesity and the Need  
906 for Future Research. *NAM Perspectives*. <https://doi.org/10.31478/202001a>
- 907 Song, N. J., Kim, S., Jang, B. H., Chang, S. H., Yun, U. J., Park, K. M., ... Park, K. W.  
908 (2016). Small molecule-induced complement factor D (Adipsin) promotes lipid  
909 accumulation and adipocyte differentiation. *PLoS ONE*.  
910 <https://doi.org/10.1371/journal.pone.0162228>
- 911 Spataru, A., Le Duc, D., Zagrean, L., & Zagrean, A. M. (2019). Ethanol exposed maturing rat  
912 cerebellar granule cells show impaired energy metabolism and increased cell death after  
913 oxygen-glucose deprivation. *Neural Regeneration Research*.  
914 <https://doi.org/10.4103/1673-5374.245474>

- 915 Speidel, D., Salehi, A., Obermueller, S., Lundquist, I., Brose, N., Renström, E., & Rorsman,  
916 P. (2008). CAPS1 and CAPS2 Regulate Stability and Recruitment of Insulin Granules in  
917 Mouse Pancreatic  $\beta$  Cells. *Cell Metabolism*. <https://doi.org/10.1016/j.cmet.2007.11.009>
- 918 Stephenson, S. E. M., Aumann, T. D., Taylor, J. M., Riseley, J. R., Li, R., Mann, J. R., ...  
919 Lockhart, P. J. (2018). Generation and characterisation of a parkin-Pacrg knockout  
920 mouse line and a Pacrg knockout mouse line. *Scientific Reports*.  
921 <https://doi.org/10.1038/s41598-018-25766-1>
- 922 Sugihara, M., Morito, D., Ainuki, S., Hirano, Y., Ogino, K., Kitamura, A., ... Nagata, K.  
923 (2019). The AAA+ ATPase/ubiquitin ligase mysterin stabilizes cytoplasmic lipid  
924 droplets. *Journal of Cell Biology*. <https://doi.org/10.1083/jcb.201712120>
- 925 Suzuki, T., Gao, J., Ishigaki, Y., Kondo, K., Sawada, S., Izumi, T., ... Katagiri, H. (2017). ER  
926 Stress Protein CHOP Mediates Insulin Resistance by Modulating Adipose Tissue  
927 Macrophage Polarity. *Cell Reports*. <https://doi.org/10.1016/j.celrep.2017.01.076>
- 928 Tang, J., Zhuo, H., Zhang, X., Jiang, R., Ji, J., Deng, L., ... Sun, B. (2014). A novel  
929 biomarker Linc00974 interacting with KRT19 promotes proliferation and metastasis in  
930 hepatocellular carcinoma. *Cell Death and Disease*.  
931 <https://doi.org/10.1038/cddis.2014.518>
- 932 Vishnyakova, P. A., Tarasova, N. V., Volodina, M. A., Tsvirkun, D. V., Sukhanova, I. A.,  
933 Kurchakova, T. A., ... Vysokikh, M. Y. (2019). Gestation age-associated dynamics of  
934 mitochondrial calcium uniporter subunits expression in feto-maternal complex at term  
935 and preterm delivery. *Scientific Reports*. <https://doi.org/10.1038/s41598-019-41996-3>
- 936 Voerman, E., Santos, S., Inskip, H., Amiano, P., Barros, H., Charles, M. A., ... Gaillard, R.  
937 (2019). Association of Gestational Weight Gain With Adverse Maternal and Infant  
938 Outcomes. *JAMA*. <https://doi.org/10.1001/jama.2019.3820>

- 939 Wang, Z., Kang, J., Deng, X., Guo, B., Wu, B., & Fan, Y. (2017). Knockdown of GATAD2A  
940 suppresses cell proliferation in thyroid cancer in vitro. *Oncology Reports*.  
941 <https://doi.org/10.3892/or.2017.5436>
- 942 Warrington, N. M., Richmond, R., Fenster, B., Myhre, R., Gaillard, R., Paternoster, L., ...  
943 Lawlor, D. A. (2018). Maternal and fetal genetic contribution to gestational weight gain.  
944 *International Journal of Obesity*. <https://doi.org/10.1038/ijo.2017.248>
- 945 Wasniewski, T., & Woclawek-Potocka, I. (2018). Altered expression of lysophosphatidic acid  
946 receptors, in association with the synthesis of estrogens and androgens in type 1  
947 endometrial cancer biology. *Gynecological Endocrinology*.  
948 <https://doi.org/10.1080/09513590.2017.1409707>
- 949 Wen, W., Kato, N., Hwang, J. Y., Guo, X., Tabara, Y., Li, H., ... Shu, X. O. (2016). Genome-  
950 wide association studies in East Asians identify new loci for waist-hip ratio and waist  
951 circumference. *Scientific Reports*. <https://doi.org/10.1038/srep17958>
- 952 Xu, L., Yang, L., Wang, L., Zhu, B., Chen, Y., Gao, H., ... Li, J. (2019). Probe-based  
953 association analysis identifies several deletions associated with average daily gain in beef  
954 cattle. *BMC Genomics*. <https://doi.org/10.1186/s12864-018-5403-5>
- 955 Yu, L., Liang, Q., Zhang, W., Liao, M., Wen, M., Zhan, B., ... Cheng, X. (2019). HSP22  
956 suppresses diabetes-induced endothelial injury by inhibiting mitochondrial reactive  
957 oxygen species formation. *Redox Biology*. <https://doi.org/10.1016/j.redox.2018.101095>
- 958 Yu, X., Zhang, Y., & Chen, H. (2016). LPA receptor 1 mediates LPA-induced ovarian cancer  
959 metastasis: An in vitro and in vivo study. *BMC Cancer*. [https://doi.org/10.1186/s12885-](https://doi.org/10.1186/s12885-016-2865-1)  
960 [016-2865-1](https://doi.org/10.1186/s12885-016-2865-1)
- 961 Za'don, N. H. A., Kamal, A. F. M., Ismail, F., Othman, S. I. T., Appukutty, M., Salim, N., ...  
962 Ludin, A. F. M. (2019). High-intensity interval training induced PGC-1 $\alpha$  and Adipor1

963 gene expressions and improved insulin sensitivity in obese individuals. *Medical Journal*  
964 *of Malaysia*.

965 Zhang, Y., Yang, J., Zhang, J., Sun, L., Hirankarn, N., Pan, H. F., ... Yang, W. (2016).  
966 Genome-wide search followed by replication reveals genetic interaction of CD80 and  
967 ALOX5AP associated with systemic lupus erythematosus in asian populations. *Annals of*  
968 *the Rheumatic Diseases*. <https://doi.org/10.1136/annrheumdis-2014-206367>

969 Zhao, L. P., Alshiekh, S., Zhao, M., Carlsson, A., Larsson, H. E., Forsander, G., ... Fureman,  
970 A. L. (2016). Next-generation sequencing reveals that HLA-DRB3, -DRB4, and -DRB5  
971 may be associated with islet autoantibodies and risk for childhood type 1 diabetes.  
972 *Diabetes*. <https://doi.org/10.2337/db15-1115>

973 Zhu, B., Rippe, C., Holmberg, J., Zeng, S., Perisic, L., Albinsson, S., ... Swärd, K. (2018).  
974 Nexilin/NEXN controls actin polymerization in smooth muscle and is regulated by  
975 myocardin family coactivators and YAP. *Scientific Reports*.  
976 <https://doi.org/10.1038/s41598-018-31328-2>

977

## 978 **TABLE LEGEND**

979 Table 1. The clinical characteristics of the T1DM cohort.

980 Table 2. The list of variants localized to genes associated with BMI in the GIANT cohort  
981 which associate with GWG.

982 Table 3. The list of variants localized to genes associated with BMI in T2D-GENES and  
983 ARIC cohorts which associate with GWG.

984 Table 4. The list of variants localized to genes associated with BMI (A) and GWG (B) in  
985 ARIC cohort.

986 Table 5. The list of variants localized to genes associated with BMI in ARIC cohort and  
987 associated with GWG as shown by fast-BAT.

988

## 989 **FIGURE LEGEND**

990 Figure 1. The scheme of the workflow. TWAS analyses to find genes associated with BMI in  
991 the general cohort (GIANT) and two diabetic cohorts (T2D-GENES, ARIC) as well as with  
992 GWG in T1DM cohort, followed by variant associations between the two phenotypes.

993 Figure 2. Venn diagram of the association of imputed gene expression which influence BMI  
994 in the Giant cohort, GWG in T1DM cohort and both phenotypes.

995 Figure 3. Overlap of the correlation of imputed expression of genes which influence BMI in  
996 ARIC and T2D-GENES cohorts, GWG in T1DM cohort and both phenotypes.

997 Figure 4. The enrichment analysis of variants localized to genes associated with GWG among  
998 those associated with BMI in the GIANT cohort.

999 Figure 5. The schematic representation of pathways(blue) affected by genes which variants  
1000 associate with GWG (pink).

1001 Figure 6. Genes which variants associate with GWG (pink) and that were reported to affect  
1002 glucose levels or TGF $\beta$  signaling (blue).

1003

## 1004 **SUPPLEMENTARY INFORMATION LEGEND**

1005 Table S1a. The results of PrediXcan on BMI in T1D cohort in Subcutaneous Adipose Tissue.

1006 Table S1b. The results of PrediXcan on BMI in T1D cohort in Visceral Adipose Tissue.

- 1007 Table S1c. The results of PrediXcan on GWG in T1D cohort in Subcutaneous Adipose  
1008 Tissue.
- 1009 Table S1d. The results of PrediXcan on GWG in T1D cohort in Visceral Adipose Tissue.
- 1010 Table S2. The overlap of PrediXcan results in Subcutaneous and Visceral Adipose Tissue in  
1011 the Giant cohort.
- 1012 Table 3a. The results of PrediXcan on Giant cohort on BMI only.
- 1013 Table S3b. The results of PrediXcan on Giant cohort on BMI and GWG.
- 1014 Table S3c. The results of PrediXcan on Giant cohort on GWG only.
- 1015 Table S4a. The results of PrediXcan on BMI in T2D cohort in Subcutaneous Adipose Tissue.
- 1016 Table S4b. The results of PrediXcan on BMI in T2D cohort in Visceral Adipose Tissue.
- 1017 Table S4c. The results of PrediXcan on BMI in ARIC cohort in Subcutaneous Adipose  
1018 Tissue.
- 1019 Table S4d. The results of PrediXcan on BMI in ARIC cohort in Visceral Adipose Tissue.
- 1020 Table S5a. The results of PrediXcan on T2D&ARIC cohorts on BMI only.
- 1021 Table S5b. The results of PrediXcan on T2D&ARIC cohorts on BMI and GWG.
- 1022 Table S5c. The results of PrediXcan on T2D&ARIC cohorts on GWG only.
- 1023 Table S6. The list of variants associated with GWG in BMI associated genes in T2D&ARIC  
1024 cohorts.
- 1025 Table S7. The FUMA analysis on BMI associated variants in GWG associated genes in  
1026 T1DM cohort.
- 1027 Table S8a. Go Enrichment analysis on the overlap between BMI and GWG genes in the Giant  
1028 cohort.
- 1029 Table S8b. Go Enrichment analysis on BMI only associated genes in the Giant cohort.
- 1030 Table S8c. Go Enrichment analysis on GWG only associated genes in the Giant cohort.
- 1031 Table S9a. Go Enrichment analysis on the overlap between BMI and GWG genes in T2D and  
1032 ARIC cohorts.
- 1033 Table S9b. Go Enrichment analysis on BMI only associated genes in T2D and ARIC cohorts.

1034 Table S9c. Go Enrichment analysis on GWG only associated genes in T2D and ARIC  
1035 cohorts.

1036

1037 Figure S1a. The LD analysis for GPN3 gene.

1038 Figure S1b. The LD analysis for PMS2P3 gene.

1039 Figure S1c. The LD analysis for STAG3L1 gene.

1040 Figure S2. The KEGG analysis on GWG associated variants in BMI associated genes in  
1041 T2D&ARIC cohorts.

1042 Figure S3a. Go Enrichment analysis on GWG only associated genes in the Giant cohort.

1043 Figure S3b. Go Enrichment analysis on the overlap between BMI and GWG genes in the  
1044 Giant cohort.


1045 Figure S4a. Go Enrichment analysis on GWG only associated genes in T2D and ARIC  
1046 cohorts.

1047 Figure S4b. Go Enrichment analysis on the overlap between BMI and GWG genes in T2D  
1048 and ARIC cohorts.


1049

1050


1051




*Figure 1. The scheme of the workflow. TWAS analyses to find genes associated with BMI in the general cohort (GIANT) and two diabetic cohorts (T2D-GENES, ARIC) as well as with GWG in T1DM cohort, followed by variant associations between the two phenotypes.*




***Figure 2. Venn diagram of the association of imputed gene expression which influence BMI in the Giant cohort, GWG in T1DM cohort and both phenotypes.***




***Figure 3. Overlap of the correlation of imputed expression of genes which influence BMI in ARIC and T2D-GENES cohorts, GWG in T1DM cohort and both phenotypes.***



**Figure 4. The enrichment analysis of variants localized to genes associated with GWG among those associated with BMI in the GIANT cohort.**



*Figure 5. The schematic representation of pathways(blue) affected by genes which variants associate with GWG (pink).*



*Figure 6. Genes which variants associate with GWG (pink) and that were reported to affect glucose levels or TGF $\beta$  signaling (blue).*