A Metapopulation Network Model for the Spreading of SARS-CoV-2: Case Study for Ireland

Rory Humphries,[1,*] Mary Spillane,[1,†] Kieran Mulchrone,[1,‡]
Sebastian Wieczorek,[1,§] Micheal O'Riordan[1,¶] and Philipp Hövel[1,**]

[1]*School of Mathematical Sciences, University College Cork,
Western Road, Cork T12 XF64, Ireland*

(Dated: June 26, 2020)

Abstract

We present preliminary results on an all-Ireland network modelling approach to simulate the spreading the Severe Acute Respiratory Syndrome CoronaVirus 2 (SARS-CoV-2), commonly known as the *coronavirus*. In the model, nodes correspond to locations or communities that are connected by links indicating travel and commuting between different locations. While this proposed modelling framework can be applied on all levels of spatial granularity and different countries, we consider Ireland as a case study. The network comprises 3440 *electoral divisions* (EDs) of the Republic of Ireland and 890 *superoutput areas* (SOAs) for Northern Ireland, which corresponds to local administrative units below the NUTS 3 regions. The local dynamics within each node follows a phenomenological SIRX compartmental model including classes of Susceptibles, Infected, Recovered and Quarantined (X) inspired from Science **368**, 742 (2020). For better comparison to empirical data, we extended that model by a class of Deaths. We consider various scenarios including the 5-phase roadmap for Ireland. In addition, as proof of concept, we investigate the effect of dynamic interventions that aim to keep the number of infected below a given threshold by dynamically adjusting containment measures on a national (Ireland), which could also be implemented at a regional (county) or local (ED/SOA) level.

---

[*] r.humphries@umail.ucc.ie

[†] mary.spillane@umail.ucc.ie

[‡] k.mulchrone@ucc.ie

[§] sebastian.wieczorek@ucc.ie

[¶] micheal.oriordain@ucc.ie

[**] philipp.hoevel@ucc.ie

The model proposed in this discussion paper and the numerical results arising from it are to be understood as a contribution to the scientific and public discussion of possible spreading scenarios and the impact of interventions measures. We will clearly state the model ingredients and assumptions and elaborate on conclusions that can be drawn based on these limitations; some of which are unrealistic for any practical purpose, but worth exploring, e.g., for worst-case scenarios.

After the introduction of the model equations for the local and networked dynamics in sections I and II, respectively, we continue with a case study of Ireland in section III. Besides an unrealistic, worst-case scenario, we consider different levels of increased awareness during the post lock-down period in section III B. In addition, in section III C, we explore effects of dynamic interventions to keep the prevalence below a given level.

I. LOCAL DYNAMICS

We start with the model for the local, well mixed dynamics at a single node as the basis for the network model:

$$\dot{S} = - \beta c \frac{SI}{N} \tag{1a}$$

$$\dot{I} = \beta c \frac{SI}{N} - \mu I - \alpha I - \kappa I \tag{1b}$$

$$\dot{X} = \kappa I - \mu X - \alpha X \tag{1c}$$

$$\dot{R} = \mu I + \mu X \tag{1d}$$

$$\dot{D} = \alpha I + \alpha X. \tag{1e}$$

The above model is a simple extension of the original Kermack and McKendrick SIR model [1] and inspired by Ref. [2]. There are five compartments, which individuals may belong to, and they may only belong to one at any time:

Susceptible (S): Those who have not been infected by the disease.

Infected (I): Those who are infected with the disease and are capable of transmission.

Quarantined (X): Those who are infected with the disease, but are quarantined or self-isolating and thus not transmitting.

Recovered (R): Those who have recovered from the disease and are considered immune.

Dead (D): Those who have died from the disease.

The above system of ordinary differential equations (ODEs) assume homogeneous mixing of the population, which can be a good approximation for small communities. The parameters of the ODE model (1) refer to the rates of moving from one compartment to another and population size:

β**:** probability of infection on contact.

c**:** rate of contact/mixing.

μ**:** rate of recovery.

κ**:** rate of moving from infected to quarantined.

α**:** rate of death.

N**:** total population size, $S + I + X + R + D = N$.

II. NETWORK MODEL

While the ODE model (1) can be justified for small communities, it fails at capturing the dynamics that occurs at large spatial scales and cannot account for heterogeneities in a population and interaction between metapopulations via mixing/commuting [3]. For this reason, we develop a network framework by splitting the population up into a number of smaller communities based on their geographic location. Ideally, these communities are small enough to justify a well mixed ODE model at the local level.

Every community corresponds to a node in the network. Two nodes are connected by a link, which represent the number of people travelling between the involved communities. The number of people travelling can vary with time or remain static. The resulting network model is given by

$$\dot{S}_i = - \beta c \left(S_i - \sum_j S_{ij} \right) \frac{I_i + \sum_k (I_{ki} - I_{ik})}{N_i + \sum_k (N_{ki} - N_{ik})}$$

$$- \beta c \sum_j S_{ij} \frac{I_j + \sum_k (I_{kj} - I_{jk})}{N_j + \sum_k (N_{kj} - N_{jk})} \tag{2a}$$

$$\dot{I}_i = \beta c \left(S_i - \sum_j S_{ij} \right) \frac{I_i + \sum_k (I_{ki} - I_{ik})}{N_i + \sum_k (N_{ki} - N_{ik})}$$

$$+ \beta c \sum_j S_{ij} \frac{I_j + \sum_k (I_{kj} - I_{jk})}{N_j + \sum_k (N_{kj} - N_{jk})} - \mu I - \alpha I - \kappa I \tag{2b}$$

$$\dot{X}_i = \kappa I_i - \mu X_i - \alpha X_i \tag{2c}$$

$$\dot{R}_i = \mu I_i + \mu X_i \tag{2d}$$

$$\dot{D}_i = \alpha I_i + \alpha X_i. \tag{2e}$$

For notational convenience, the compartments are now labelled by a subscript, i.e., S_i, I_i, X_i, R_i, and D_i, to denote the number of individuals belonging to the respective compartments of i-th node in the network ($S_i + I_i + X_i + R_i + D_i = N_i$). Similarly, the number of individuals travelling from node i to node j belonging to a given compartment is denoted by $S_{ij}, I_{ij}, X_{ij}, R_{ij}$, and D_{ij} resulting in $N_{ij} = S_{ij} + I_{ij} + X_{ij} + R_{ij} + D_{ij}$.

Note that the network model includes interactions between nodes only in the susceptible and infected compartments. Accordingly, we account for the number of (i) individuals that remain in their home node and (ii) those who travel to other nodes in the first and second term of Eqs. (2a) and (2b), respectively.

Given a number of travellers N_{ij} between nodes, the network model (2) assumes that the number of travellers belonging to the respective compartments is determined by a relation such as $S_{ij} = N_{ij} \frac{S_i}{N_i}$. Thus, the number of travelling susceptibles is proportional to the

number of susceptibles in the home node. The number of travelling individuals in the other compartments is computed in the same fashion.

N_{ij} may be randomly generated at each time step. The distance travelled between locations often follows a power-law or heavy-tail distribution and can be described by a gravity or radiation population model [4–9]. Alternatively, empirical movement data can inform the model. From these data on can either build a static network using, e.g., the jump-size distributions, or randomly generate a network at each time-step. Note that the proposed network model does not account for any importation from outside Ireland.

All scenarios discussed in the next sections are simulated using our library for modelling epidemics on networks, EPIGRAPH [10], which is implemented in C++ and freely available on GITHUB.

III. CASE STUDY: IRELAND

For the remainder of this discussion paper, we elaborate on the spreading of SARS-CoV-2 in Ireland as a case study. The network model given by Eqs. (2) is informed by publicly available data detailed in the next section.

A. Network Data

For the Republic of Ireland, each of the 3440 nodes in the network represents an *electoral division* as defined by the central statistics office (CSO), for which small area population statistics are published from the 2016 census. [11]. The links in the network are the number of commuters between each electoral division according to CSO data [12]. Similarly, for Northern Ireland, nodes refer to 890 *superoutput areas* according to Northern Ireland Statistics and Research Agency (NISRA) [13]. This spatial resolution corresponds to the local administrative units below the category-3 regions of the *Nomenclature of Territorial Units for Statisticss* (NUTS) [14].

B. 5-Phase Roadmap without Additional Countermeasures

In order to demonstrate the general feasibility of the proposed modelling framework, we consider the 5-phase *roadmap for reopening society & business* by reducing lock-down restrictions as outlined by the Irish Government [15].

Our model attempts to follow the historic evolution of the disease up until phase 2. Then, we allow the model to run under the predicted affects of the remaining lock-down phases. See Tab. I for implications on travel and gathering restrictions. It is important to note that at this stage, we do not consider any reintroduction or roll back to earlier phases during the course of the simulated outbreak. This corresponds to a worst-case scenario and is very much unrealistic for all practical purposes, because interventions will be reintroduced, if case numbers rise again.

The parameters are chosen such that it matches the actual number of deaths in Ireland. See Tab. II for details. It is important to note, however, that this model is in the early stages of testing and has not undergone any sensitivity training on the parameters. The effective infection probability β_{eff} is computed as $\beta_{\text{eff}} = \beta c$. The basic reproduction number R_0 of the

Phase	Mobility	Social distancing
pre lock-down	no restrictions	no restrictions
initial lock-down	travel with 2km radius	essential meetings only
1	travel with 5km radius	essential meetings only
2	travel with 20m radius	maximum of 4 people
3	travel with 20km radius	maximum of 4 people
4	no restrictions	small gatherings
5	no restrictions	large gatherings
post lock-down	no restrictions	no restrictions

TABLE I: Travel and gathering restrictions during the 5 phases of the Irish roadmap.

Phase	β	c	μ	κ	α	β_{eff}	R_0
pre lock-down	0.37	1	0.1	0.1	0.0028	0.37	1.82
initial lock-down	0.37	0.4	0.1	0.1	0.0028	0.15	0.73
1	0.37	0.3	0.1	0.1	0.0028	0.11	0.55
2	0.37	0.4	0.1	0.1	0.0028	0.15	0.73
3	0.37	0.5	0.1	0.1	0.0028	0.19	0.91
4	0.37	0.6	0.1	0.1	0.0028	0.22	1.09
5	0.37	0.7	0.1	0.1	0.0028	0.26	1.28
post lock-down	0.37	1	0.1	0.1	0.0028	0.37	1.82

TABLE II: Parameters for the 5 phases of the Irish roadmap. β_{eff} and R_0 are computed from the system parameters.

local dynamics in Eqs. (1) can be calculated as

$$R_0 = \frac{\beta c}{\mu + \kappa + \alpha}. \tag{3}$$

A value of $R_0 > 1$ ($R_0 < 1$) refers to an unstable (stable) disease-free fixed point ($S^* = N$ and $I^* = X^* = R^* = D^* = 0$) and results in a large outbreak (burn out of the disease). In other words, $R_0 > 0$ indicates that a single individual can on average infect more than one other person during the infection period. The basic reproduction number exceed the critical value of $R_0^{\mathrm{crit}} = 1$ during pre lock-down and after phase 3. Conversely, only initial lock-down, phases 1 and 2 have a values well below unity and – neglecting network-related effects – exhibit a burn out of an outbreak and self-guided decrease of the number of infected.

We introduce a small number of diseased individuals to the Dublin area on the 22nd of January, that is, 50 days before the initial lock-down, and run the model such that the disease spreads unimpeded. On the 12th of March, we introduce the initial lock-down measures, which take some time to reach full effect. Individuals' movement is limited to 2km and this is adhered to with 80% compliance. This level of compliance is supported by preliminary findings of a nation-wide series of phone interviews accounting for essential/justified and non-essential travel [16]. We account for social-distancing and self-isolation (X compartment) by reducing the rate of mixing c within communities (cf. Tab. II for the respective parameter values). During phases 1 and 2/3, the travel radius increased to 5km and 20km, respectively, and we assume the same level of compliance. Travel restrictions are removed in phases 4 and 5 (cf. Tab. I). Each of the numbered phases have a duration of 21 days.

Figures 1 depicts the time series of all five compartments – aggregated over all nodes – in semi-logarithmic scale. One can see two distinct peaks: (i) a small peak that occurs before

FIG. 1: Number of individuals (aggregated over the entire population/network) belonging to each compartment as stated in the legend over time with a log scale on the y-axis. Parameters as in Tab. II.

the lock-down and (ii) a second peak that occurs once all lock-down measures are lifted. It is clear that unless some of the measure are maintained, then the lock-down will just have the effect of delaying the epidemic and result in a second wave with a prevalence that – in this unrealistic worst-case scenario – is 100 times higher than the first one and exceeds any realistic capacity of the health-case system. The number of infected eventually reaches zero due to a depletion of the pool of susceptibles. However, the disease-free fixed point $I^* = 0$ remains inherently unstable. Since a finite number of susceptible remain, a perturbation to the networked system, e.g., introduction or importation from outside, has the potential to trigger additional outbreaks.

As stated above, the parameters are chosen to match the actual number of deaths in Ireland. This, in turn, results in a larger first peak compared to the reported numbers of confirmed cases on the 12th of March ranging from 27 according to [17] to 70 according to [18] The simulated results, however, are not meant to reflect this range of confirmed cases, but following the logic of the model, refer to all infected individuals including aymptomatic, non-diagnosed and untested.

Figure 2a and 2b show spatially resolved snapshots of the prevalence of the disease per km^2 at times $t = 50d$ (March 12) and $t = 320d$ (December 7), respectively (cf. [19] for a map based on data of reported, confirmed cases). Figure 2a corresponds to the disease prevalence at the first peak before the lock-down. The disease remains in the East of the country for the most part with the exception of big towns and cities. In contrast, at $t = 320d$ (Fig. 2b),

FIG. 2: The spatial distribution of infected individuals per km^2 around the two distinct peaks shown in Fig. 1 at (a) Day 50 (b) Day 320. Parameters as in Tab. II.

the outbreak covers the entire country and affects every electoral division. See Fig. 1 for a full time series of the aggregated variables.

FIG. 3: Evolution of the aggregated, infected compartment I for varied post lock-down conditions. The blue curve corresponds to the parameters given in Tab. II (cf. Fig. 1 for a semi-logarithmic scale). The other curves vary the parameter c in the post lock-down phase as stated in the legend: $c = 0.9$, $c = 0.8$ and $c = 0.7$.

Figure 3 shows the effect of different levels of awareness during the post lock-down phase, which is modelled by different values for the mixing parameter c. For convenience, the plots

only depict the time series of the aggregate infected compartment I in a linear scale. A reduction to 70% of the original probability of infection significantly leads to a significantly smaller and delayed second peak (red curve), while the basic reproduction number of the local dynamics remains above unity (cf. Tab. II).

FIG. 4: Evolution of the aggregated, infected I (blue), quarantined X (green) and deceased D (red) compartments for two level of compliance with lock-down restrictions: 80% (solid, cf. Fig. 1), and 70% (dashed). Other parameters as in Tab. II.

Similarly to Figure 3, the network model (2) allows considering different levels of compliance with travel restrictions. The effect is exemplarified for aggregated values of I, X and D in Fig. 4 for compliance of 80% (solid curves, cf. Fig. 1) and 70% (dashed curves). One can see that for lower compliance, the second peak occurs earlier and has a higher amplitude. This results in a slightly increased number of deaths.

In order to show how the heavy-tail distribution of travel across the network causes a lack of sensitivity in initial conditions, we run the model for a number of random initial seeds where for each run only a single individual in one electoral division or superoutput area. The resulting time series are shown in Fig. 5.

One can see that both amplitude and timing of the large, second peak are stable and it is impossible to visually differentiate between any of the curves for different initial conditions. In the presented 4 cases, the initial location is selected uniformly at random from all network nodes. We note that deliberately selecting urban or rural locations will have no qualitative impact. To further corroborate the results, initial infections of a significant, finite fraction of the population will be explored. This might account for a considerable amount of unreported cases at the start of the recording of SARS-CoV-2 cases.

FIG. 5: Time series for 4 random initial conditions. Parameters as in Tab. II with $c = 0.7$ in the post lock-down period.

C. Dynamic Interventions

Finally, we discuss the likely scenario of a flexible lock-down policy that monitors to the prevalence of the disease within the island and aims to keep it below the capacity of the health-care system. As a simple measure, we define some threshold I_{th} as the maximum number of infected that can be present before we go back into lock-down and go through earlier phases again. Such a procedure corresponds to dynamic [20–22] or active [23] interventions. Specifically, we simulate until phase 3 as before to maintain the agreement with reported cases. Afterwards, if the condition $I_{\text{th}} > \sum_i I_i$ is met, the system goes back to phase 1 and then progresses from there until the condition is met again.

In detail, we consider two scenarios with different threshold parameters: (i) $I_{\text{th}} = 10^4$, as shown in Fig. 6 and (ii) $I_{\text{th}} = 10^3$ as depicted in Fig. 7. In both figures, the worst-case scenario of Fig. 1 is added as dashed curves. Note that the curves of recovered R (orange) and deaths D (red) are only increasing at a low rate. For the case of $I_{\text{th}} = 10^4$, for instance, these values arrive at 10^5 after two years and thus, way below any effect of herd immunity.

Under the assumptions the considered network model, it would take many years for a significant depletion of the pool of susceptibles. Extrapolating from an extended run of the model similar to Fig. 6, we find that it would take roughly 27 years for the number of recovered to reach half the population, leading to a total number of approximately 92,000 deaths over the course of many waves. Even with a large threshold $I_{\text{th}} = 10^4$ for the maximum number of infected cases, the feasibility of such a strategy is questionable due to the time it would take to reach herd immunity of on the island.

The re-introduction of lock-down measures to account for rising numbers of infected does not fundamentally change the stability of the disease-free fixed point $I^* = 0$. As soon as the restrictions are relaxed, e.g., during phases 3 to 5, it will become unstable again and the outbreak pattern repeats.

A change of stability towards a stable fixed point requires other forms of interventions.

FIG. 6: The number of individuals belonging to each compartment over time (using the parameters in Tab. II) under a dynamic lock-down strategy taking $I_{\text{th}} = 10^4$. The dashed lines are the equivalent model without the dynamic lock-down strategy shown in Fig. 1.

One could consider, for example, a fishing-type extension of the model [24–26]. This will basically turn the rate κ in the local dynamics (1) into a function of the number of infected. e.g., $\kappa(I) = \frac{h}{1+I/I_c}$ with a consumption/fishing rate h (here: rate of isolation) and a reference capacity I_c for $\kappa(I) = h/2$. This I dependence will account for contact-tracing effects, which are especially effective for low numbers of infected. As a consequence, a high rate h of isolation will be able to push the basic reproduction number R_0 [cf. Eq. (3)] below the critical value of $R_0^{\text{crit}} = 1$ and change the stability of the fixed point $I^* = 0$.

IV. CONCLUSION

We have proposed a dynamical network model to study the spreading of SARS-Cov-2 that accounts for travel/commuting between nodes of the network. As a case study, we have presented results from numerical simulations for Ireland, where the model is informed from publicly available data on geographical regions, population, and mobility in both the Republic of Ireland and Northern Ireland. The network model is flexible in terms of spatial granularity and parameter selection. We have focused on the governmental 5-phase roadmap for reopening society & business and considered different initial conditions and parameters. We have observed that the roadmap will lead to a decline of case numbers. However, once the restrictions are lifted, the numbers will rise again, even if an increased level of awareness is considered. Finally, we have elaborated on a procedure of dynamic interventions that re-introduce lock-down measures, if a certain threshold of infected is reached. Following such a

FIG. 7: The number of individuals belonging to each compartment over time (using the parameters in table II) under a dynamic lock-down strategy taking $I_{\mathrm{th}} = 10^3$. The dashed lines are the equivalent model without the dynamic lock-down strategy shown in Fig. 1.

protocol, the overall number of infected can be controlled and safely kept below a prescribed threshold, which can be selected according to the capacity of the health-care system. We have considered that containment measures would be re-introduced on a country-wide level, but a more regional approach (county level) appears feasible as well. In any case, while the outbreak can be managed by dynamic interventions, it might take a period of several years until a significant fraction of the population will have lived through the disease and – under the assumption of immunity – and will have led to an effect of herd immunity.

Let us finish by stressing again that the times series and maps are preliminary findings and subject to a systematic sensitivity analysis. Nonetheless, we hope that the results presented in this discussion paper will stimulate scientific and public discussion about the purpose and effect of lock-down measures in the months ahead.

ACKNOWLEDGMENTS

The work is supported by Health Research Board Ireland in the framework of the COVID19 rapid response programme (grant number: COV19-2020-117). We are grateful to Andreas Amann, Patricia Kearny, Gerry Kileen, and Ivan Perry for helpful discussions.

[1] W. O. Kermack and A. G. McKendrick, Proc. R. Soc. A **115**, 700 (1927).

[2] B. F. Maier and D. Brockmann, Science **368**, 742 (2020).

[3] V. Belik, T. Geisel, and D. Brockmann, Phys. Rev. X **1**, 011001 (2011).

[4] D. Brockmann, L. Hufnagel, and T. Geisel, Nature **439**, 462 (2006).

[5] D. Brockmann and L. Hufnagel, Phys. Rev. Lett. **98**, 178301 (2007).

[6] C. Song, Z. Qu, N. Blumm, and A. L. Barabási, Science **327**, 1018 (2010), http://www.sciencemag.org/cgi/reprint/327/5968/1018.pdf.

[7] C. Song, T. Koren, P. Wang, and A. L. Barabási, Nat. Phys. **6**, 818 (2010).

[8] F. Simini, M. C. González, A. Maritan, and A. L. Barabási, Nature **484**, 96 (2012).

[9] S. Grauwin, M. Szell, S. Sobolevsky, P. Hövel, F. Simini, M. Vanhoof, Z. Smoreda, A. L. Barabási, and C. Ratti, Sci. Rep. **7**, 46677 (2017).

[10] R. Humphries, EpiGraph, `https://github.com/rory-humphries/EpiGraph-cpp` (2020).

[11] Census 2016 Open Data Site, `http://census2016.geohive.ie/datasets/774b6c87f0ec49c48e0e7999fc0e752b_1` (accessed: 17 June 2020).

[12] Census 2016 Profile 6, Commuting in Ireland, `https://www.cso.ie/en/csolatestnews/presspages/2017/census2016profile6-commutinginireland/` (accessed: 17 June 2020).

[13] Northern Ireland Statistics and Research Agency, `https://www.nisra.gov.uk/statistics/regional-analysis-and-trends/local-government-district` (accessed: 17 June 2020).

[14] (Eurostat, Nomenclature of Territorial Units for Statistics – NUTS, `https://ec.europa.eu/eurostat/web/nuts/background`.

[15] Department of the Taoiseach and Department of Health: Roadmap for reopening society and business, published at: 1 May 2020, last updated: 12 June 2020, `https://www.gov.ie/en/news/58bc8b-taoiseach-announces-roadmap-for-reopening-society-and-business-and-u/`, `https://assets.gov.ie/73722/ffd17d70fbb64b498fd809dde548f411.pdf` (accessed: 17 June 2020).

[16] HRB project: COVID19 rapid response programme (grant number: COV19-2020-117), unpublished data of three waves of telephone surveys.

[17] Health Protection Surveillance Centre (HPSC), `https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/casesinireland/` (accessed: June 22, 2020).

[18] Worldometer, `https://www.worldometers.info/coronavirus/country/ireland/` (accessed: June 22, 2020).

[19] Central Statistics Office (CSO): Confirmed cases of COVID-19 by Electoral Division, `https://census.cso.ie/covid19/`.

[20] R. Chowdhury, K. Heng, M. S. R. Shawon, G. Goh, D. Okonofua, C. Ochoa-Rosales, V. Gonzalez-Jaramillo, A. Bhuiya, D. Reidpath, S. Prathapan, *et al.*, European journal of epidemiology **35**, 389 (2020).

[21] G. Killeen and S. Kiware, medRxiv 10.1101/2020.04.15.20066845 (2020), https://www.medrxiv.org/content/early/2020/04/20/2020.04.15.20066845.full.pdf.

[22] S. M. Kissler, C. Tedijanto, E. Goldstein, Y. H. Grad, and M. Lipsitch, Science **368**, 860 (2020).

[23] A. D. Ames, T. G. Molnar, A. W. Singletary, and G. Orosz, medRxiv 10.1101/2020.06.17.20133264 (2020), https://www.medrxiv.org/content/early/2020/06/19/2020.06.17.20133264.full.pdf.

[24] R. J. Beverton and S. J. Holt, *On the dynamics of exploited fish populations*, Vol. 11 (Springer Science & Business Media, 1993).

[25] V. Christensen and C. J. Walters, Ecol. Modell. **172**, 109 (2004).

[26] A. Amann, How to choose the best strategy for fighting an infectious disease, private communication (2020).