A Metapopulation Network Model for the Spreading of SARS-CoV-2: Case Study for Ireland[★]

Rory Humphries[a,1], Mary Spillane[a,2], Kieran Mulchrone[a,3], Sebastian Wieczorek[a,4], Micheal O'Riordain[a,b,4] and Philipp Hövel[a,*,5]

[a]*School of Mathematical Sciences, University College Cork, Western Road, Cork T12 XF64, Ireland*
[b]*Department of Surgery, Mercy University Hospital, Grenville Place, Cork, T12 WE28, Ireland*

ARTICLE INFO

Keywords:
compartmental model
SARS-CoV-2
coronavirus
non-pharmaceutical containment strategies
network model

ABSTRACT

We present preliminary results on an all-Ireland network modelling approach to simulate the spreading the Severe Acute Respiratory Syndrome CoronaVirus 2 (SARS-CoV-2), commonly known as the *coronavirus*. In the model, nodes correspond to locations or communities that are connected by links indicating travel and commuting between different locations. While this proposed modelling framework can be applied on all levels of spatial granularity and different countries, we consider Ireland as a case study. The network comprises 3440 *electoral divisions* (EDs) of the Republic of Ireland and 890 *superoutput areas* (SOAs) for Northern Ireland, which corresponds to local administrative units below the NUTS 3 regions. The local dynamics within each node follows a phenomenological SIRX compartmental model including classes of Susceptibles, Infected, Recovered and Quarantined (X) inspired from Science **368**, 742 (2020). For better comparison to empirical data, we extended that model by a class of Deaths. We consider various scenarios including the 5-phase roadmap for Ireland. In addition, as proof of concept, we investigate the effect of dynamic interventions that aim to keep the number of infected below a given threshold. This is achieved by dynamically adjusting containment measures on a national scale, which could also be implemented at a regional (county) or local (ED/SOA) level. We find that – in principle – dynamic interventions are capable to limit the impact of future waves of outbreaks, but on the downside, such a strategy can last several years until herd immunity will be reached.

On 30th January 2020, the World Health Organisation (WHO) declared the newly emergent novel coronavirus pathogen, Severe Acute Respiratory Syndrome CoronaVirus 2 (SARS-CoV-2), to be a *public health emergency of international concern* [31, 32]. This characterisation was later updated to pandemic status on 11th March 2020 [33]. Following its initial detection in a cluster of patients experiencing acute respiratory symptoms in the city of Wuhan, in the Hubei province of China [9], the spread of SARS-CoV-2, which causes the coronavirus disease 2019 (COVID-19), to 218 countries and territories in the months since has led to unprecedented social and economic disruption, [13, 34]. Despite widespread efforts to contain the virus through a series of non-pharmaceutical interventions, global figures show reported cases standing at over 63 million, with deaths from the virus totalling 1.5 million up to late November 2020 [12, 16]. It is clear that there is an urgent and ongoing need to understand how the virus spreads in order to effectively protect populations.

As of yet there are no significantly effective pharmaceutical measures available that prevent the spread of the disease and so one of our best resources, until such measures arrive, are mathematical models [1, 6, 20–23] that can give us insight into the spreading process of the disease. These models can help decide on the best course of action to take in the form of non-pharmaceutical interventions [3, 24–26].

[★]This article is the results of the research project funded by the Health Research Board in the framework of the COVID19 rapid response programme (grant number: COV19-2020-117).

[*]Corresponding author

✉ r.humphries@umail.ucc.ie (R. Humphries); mary.spillane@umail.ucc.ie (.M. Spillane); k.mulchrone@ucc.ie (.K. Mulchrone); sebastian.wieczorek@ucc.ie (S. Wieczorek); micheal.oriordain@ucc.ie (.M. O'Riordain); philipp.hoevel@ucc.ie (.P. Hövel)

🖳 http://publish.ucc.ie/researchprofiles/D019/kmulchrone (.K. Mulchrone);
http://research.ucc.ie/profiles/D019/sebastianwieczorek (.S. Wieczorek);
https://www.muh.ie/index.php/consultants/19-mr-michael-o-riordain (.M. O'Riordain);
http://publish.ucc.ie/researchprofiles/D019/philipphoevel (.P. Hövel)

ORCID(s): 0000-0002-1370-4272 (.P. Hövel)

The model proposed in this discussion paper and the numerical results arising from it are to be understood as a contribution to the scientific and public discussion of possible spreading scenarios and the impact of interventions measures. We will clearly state the model ingredients and assumptions and elaborate on conclusions that can be drawn based on these limitations; some of which are unlikely for any practical purpose, but worth exploring, e.g., for worst-case scenarios.

We start with an introduction of the model equations for the local and networked dynamics in sections 1 and 2, respectively. This includes a discussion of the local dynamics from a geometric point of view, where we sketch the trajectories and equilibria in phase space. Then, we continue with a case study of Ireland in section 3. Besides an unlikely, worst-case scenario, we consider different levels of increased awareness during the post-lockdown period in section 3.2. In addition, in section 3.3, we explore effects of dynamic interventions to keep the prevalence below a given level. These dynamic lockdowns are considered at different levels of spatial granularity.

1. Local Dynamics

We start with the model for the local, well mixed dynamics at a single node as the basis for the network model:

$$\dot{S} = -\beta c \frac{SI}{N} \tag{1a}$$

$$\dot{I} = \beta c \frac{SI}{N} - \mu I - \alpha I - \kappa I \tag{1b}$$

$$\dot{X} = \kappa I - \mu X - \alpha X \tag{1c}$$

$$\dot{R} = \mu I + \mu X \tag{1d}$$

$$\dot{D} = \alpha I + \alpha X. \tag{1e}$$

The above model is a simple extension of the original Kermack and McKendrick SIR model [17], inspired by Ref. [22]. There are five compartments, N individuals, and each individual may belong to just one compartment at any time:

Susceptible $S(t)$ Individuals who have not been infected by the disease.

Infected $I(t)$ Individuals who are infected with the disease and are capable of transmission.

Quarantined $X(t)$ Individuals who are infected with the disease, but are quarantined or self-isolating and thus not transmitting.

Recovered $R(t)$ Individuals who have recovered from the disease and are considered immune.

Dead $D(t)$ Individuals who have died from the disease.

The above system of ordinary differential equations (ODEs) assume homogeneous mixing of the population, which can be a good approximation for small communities, and small number of deaths. The parameters of the ODE model (1) refer to the rates of moving from one compartment to another and population size:

β probability of infection on contact.

c rate of contact/mixing.

μ rate of recovery.

κ rate of moving from infected to quarantined.

α rate of death.

N total population size $N = S(t) + I(t) + X(t) + R(t) + D(t) = \text{const.}$.

Much insight into the network dynamics can be gained from geometric analysis of the local, single-node dynamics. In the (S, I, X, R, D) phase space, system (1) has a whole family of stationary solutions (equilibria). They are given by

$$I_e = X_e = 0 \quad \text{and} \quad S_e + R_e + D_e = N,$$

Figure 1: Sketch of the outbreak dynamics in the local SIXRD model (1) for $R_0 > 1$, shown as a (red) trajectory from the disease-free equilibrium $S_e = N$ to one of the stable equilibria with $0 < S_e < S_0$ in the projection of the (S, I, X, R, D) phase space onto the (S, I, R) subspace. S_0 is the herd immunity threshold separating the triangular family of equilibria into the (pink) unstable and (blue) stable parts.

correspond to disease-free conditions, and form a triangular surface shown in Fig. 1. This family includes the important pre-disease equilibrium

$$S_e = N, I_e = X_e = R_e = D_e = 0.$$

To discuss stability of equilibria, it is convenient to introduce the *basic reproduction number*:

$$R_0 = \frac{\beta\, c}{\alpha + \kappa + \mu}. \tag{2}$$

When $0 < R_0 < 1$, all equilibria are stable, meaning that no disease outbreaks can occur. When $R_0 = 1$, the pre-disease equilibrium turns unstable, and disease outbreaks become possible for $R_0 > 1$. When $R_0 > 1$, disease outbreaks become possible. To be more specific, the family of equilibria is divided into the (pink) unstable part with $S_0 < S_e \leq N$, which contains the pre-disease equilibrium, and the (blue) stable part with $0 \leq S_e < S_0$, shown in Fig. 1. The instability threshold separating the two parts is given by

$$S_0 = \frac{N}{R_0}. \tag{3}$$

Guided by these observations, we define *herd immunity* in terms of stable equilibria as follows:

Definition. *In the local SIXRD model* (1)*, we define* herd immunity *as a disease-free state of the system such that a small increase in infected I decays monotonically to zero (no outbreak):*

$$0 \leq S < S_0 \quad and \quad I = X = 0.$$

In other words, herd immunity is represented by a stable equilibrium. We call $S_0 = N / R_0$ the herd immunity threshold.

In Fig. 1, a disease outbreak is is depicted as the (red) trajectory from the pre-disease equilibrium, giving a peak in infected $I(t)$ around the herd immunity threshold S_0, to one of the (blue) herd immunity equilibria that lie below S_0. The lower the S_0, the higher the peak in $I(t)$. Thus, bringing the herd immunity threshold S_0 closer to N for a period of time and then back to its original position, e.g., via imposing temporary public-health restrictions, can modify the outbreak trajectory so that ultimate herd immunity is achieved in a controlled way, that is with a peak in $I(t)$ that remains below some desired level.

2. Network Model

While the ODE model (1) can be justified for small communities, it fails at capturing the dynamics that occurs at large spatial scales and cannot account for heterogeneities in a population and interaction between metapopulations via mixing/commuting [4]. For this reason, we develop a network framework by splitting the population up into a number of smaller communities based on their geographic location. Ideally, these communities are small enough to justify a well mixed ODE model at the local level. Every community corresponds to a node in the network. Two nodes are connected by a link, which represent the number of people travelling between the involved communities. The number of people travelling can vary with time or remain static. The resulting network model is given by

$$
\dot{S}_i = -\beta c \left(S_i - \sum_{j=1}^n S_{ij} \right) \frac{I_i + \sum_{k=1}^n (I_{ki} - I_{ik})}{N_i + \sum_{k=1}^n (N_{ki} - N_{ik})}
$$
$$
- \beta c \sum_{j=1}^n S_{ij} \frac{I_j + \sum_{k=1}^n (I_{kj} - I_{jk})}{N_j + \sum_{k=1}^n (N_{kj} - N_{jk})}
\tag{4a}
$$

$$
\dot{I}_i = \beta c \left(S_i - \sum_{j=1}^n S_{ij} \right) \frac{I_i + \sum_{k=1}^n (I_{ki} - I_{ik})}{N_i + \sum_{k=1}^n (N_{ki} - N_{ik})}
$$
$$
+ \beta c \sum_{j=1}^n S_{ij} \frac{I_j + \sum_{k=1}^n (I_{kj} - I_{jk})}{N_j + \sum_{k=1}^n (N_{kj} - N_{jk})} - \mu I - \alpha I - \kappa I
\tag{4b}
$$

$$
\dot{X}_i = \kappa I_i - \mu X_i - \alpha X_i
\tag{4c}
$$
$$
\dot{R}_i = \mu I_i + \mu X_i
\tag{4d}
$$
$$
\dot{D}_i = \alpha I_i + \alpha X_i.
\tag{4e}
$$

For notational convenience, the compartments are now labelled by a subscript, i.e., S_i, I_i, X_i, R_i, and D_i, to denote the number of individuals belonging to the respective compartments of i-th node in the network ($S_i + I_i + X_i + R_i + D_i = N_i$, $i = 1, \cdots, n$). Similarly, the number of individuals travelling from node i to node j belonging to a given compartment is denoted by $S_{ij}, I_{ij}, X_{ij}, R_{ij}$, and D_{ij} resulting in $N_{ij} = S_{ij} + I_{ij} + X_{ij} + R_{ij} + D_{ij}$. Note that the network model includes interactions between nodes only in the susceptible and infected compartments. Accordingly, we account for the number of (i) individuals that remain in their home node and (ii) those who travel to other nodes in the first and second term of Eqs. (4a) and (4b), respectively.

Given a number of travellers N_{ij} between nodes, the network model (4) assumes that the number of travellers belonging to the respective compartments is determined by a relation such as $S_{ij} = N_{ij} \frac{S_i}{N_i}$. Thus, the number of travelling susceptibles is proportional to the number of susceptibles in the home node. The number of travelling individuals in the other compartments is computed in the same fashion. N_{ij} may be randomly generated at each time step. The distance travelled between locations often follows a power-law or heavy-tail distribution and can be described by a gravity or radiation population model [7, 8, 14, 27–29]. Alternatively, empirical movement data can inform the model. From these data one can either build a static network using, e.g., the jump-size distributions, or randomly generate a network at each time-step. Note that the proposed network model does not account for any importation from outside the considered network.

For the fully networked case [cf. Eqs. (4)], the basic reproduction number R_0 can be computed from the system's next generation matrix G [30]. This takes into account the couplings between the nodes when computing the stability of the disease-free equilibria. The (i, j) entry of this matrix corresponds to the expected number of new infections in node i caused by an infected individual introduced in node j. The next generation matrix, as derived in Appendix A, is given as

$$
G_{ij} = \frac{\beta c}{\mu + \alpha + \kappa} \left[\left(S_i - \sum_{k=1}^n S_{ik} \right) \frac{\delta_{ij} \left(1 - \sum_{k=1}^n \frac{N_{ik}}{N_i} \right) + \frac{N_{ji}}{N_j}}{N_i + \sum_{k=1}^n (N_{ki} - N_{ik})} \right.
\tag{5}
$$

$$+ \sum_{k=1}^{n} S_{ik} \frac{\delta_{jk} \left(1 - \sum_{l=1}^{n} \frac{N_{jl}}{N_j} \right) + \frac{N_{jk}}{N_j}}{N_k + \sum_{l=1}^{n}(N_{lk} - N_{kl})} \Bigg].$$

Then, the reproduction number R_0^* for the networked system is given by the spectral radius ρ of the the next generation matrix,

$$R_0^* = \rho(G). \tag{6}$$

This provides a condition necessary for the stability all disease-free equilibria ($I_i = X_i = 0, S_i + R_i + D_i = N_i$ for all $i = 1, \cdots, n$). It also gives a natural definition for herd immunity in the network, as the networked reproduction number R_0 is now a function of all susceptible populations (cf. the definition of heard immunity for the local dynamics in Sec. 1). All scenarios discussed in the next sections are simulated using our library for modelling epidemics on networks, EPIGRAPH [15], which is implemented in C++ and freely available on GITHUB.

3. Case study: Ireland

For the remainder of this discussion paper, we elaborate on the spreading of SARS-CoV-2 in Ireland as a case study. The network model given by Eqs. (4) is informed by publicly available data detailed in the next section.

3.1. Network Data

For the Republic of Ireland, each of the 3440 nodes in the network represents an *electoral division* as defined by the central statistics office (CSO), for which small area population statistics are published from the 2016 census. [1]. The links in the network are the number of commuters between each electoral division according to CSO data [2]. Similarly, for Northern Ireland, nodes refer to 890 *superoutput areas* according to Northern Ireland Statistics and Research Agency (NISRA) [3]. This spatial resolution corresponds to the local administrative units below the category-3 regions of the *Nomenclature of Territorial Units for Statisticss* (NUTS) [4].

3.2. 5-Phase Roadmap without Additional Countermeasures

In order to demonstrate the general feasibility of the proposed modelling framework, we consider the 5-phase *roadmap for reopening society & business* by reducing lockdown restrictions as outlined by the Irish Government in May 2020 [5]. Our model attempts to follow the historic evolution of the disease up until phase 2. Then, we allow the model to run under the predicted affects of the remaining lockdown phases. See Tab. 1 for implications on travel and gathering restrictions. Appendix B provides a comparison to reported case numbers. It is important to note that at this stage, we do not consider any reintroduction or roll back to earlier phases during the course of the simulated outbreak. This corresponds to a worst-case scenario and is highly unlikely from a practical point of view, because interventions will be reintroduced, if case numbers rise again.

The parameters are chosen such that it matches the actual number of deaths in Ireland. See Tab. 2 for details. It is important to note, however, that this model is in the early stages of testing and has not undergone any sensitivity training on the parameters. The effective infection probability β_{eff} is computed as $\beta_{\text{eff}} = \beta c$. As defined in Eq. (2), the basic reproduction number of the local dynamics [cf. Eqs. (1)] exceeds the critical value of $R_0^{\text{crit}} = 1$ during pre lockdown and after phase 3. Conversely, only initial lockdown, phases 1 and 2 have a R_0 value well below unity and

---

[1]Census 2016 Open Data Site, `http://census2016.geohive.ie/datasets/774b6c87f0ec49c48e0e7999fc0e752b_1` (accessed: 17 June 2020).

[2]Census 2016 Profile 6, Commuting in Ireland, `https://www.cso.ie/en/csolatestnews/presspages/2017/census2016profile6-commutinginireland/` (accessed: 17 June 2020)

[3]Northern Ireland Statistics and Research Agency, `https://www.nisra.gov.uk/statistics/regional-analysis-and-trends/local-government-district` (accessed: 17 June 2020).

[4](Eurostat, Nomenclature of Territorial Units for Statistics – NUTS, `https://ec.europa.eu/eurostat/web/nuts/background`

[5]Department of the Taoiseach and Department of Health: Roadmap for reopening society and business, published at: 1 May 2020, last updated: 12 June 2020, `https://www.gov.ie/en/news/58bc8b-taoiseach-announces-roadmap-for-reopening-society-and-business-and-u/`, `https://assets.gov.ie/73722/ffd17d70fbb64b498fd809dde548f411.pdf` (accessed: 17 June 2020).

Phase	Date	Mobility	Social distancing
pre lockdown	22-01-20	no restrictions	no restrictions
initial lockdown	27-03-20	travel with 2km radius	essential meetings only
1	18-05-20	travel with 5km radius	essential meetings only
2	08-06-20	travel with 20m radius	maximum of 4 people
3	29-06-20	travel with 20km radius	maximum of 4 people
4	20-07-20	no restrictions	small gatherings
5	10-08-20	no restrictions	large gatherings
post lockdown	31-08-20	no restrictions	no restrictions

Table 1: Travel and gathering restrictions during the 5 phases of the Irish roadmap.

Phase	β	c	μ	κ	α	β_{eff}	R_0
pre lockdown	0.37	1	0.1	0.1	0.0028	0.37	1.82
school closure	0.37	0.7	0.1	0.1	0.0028	0.26	1.28
initial lockdown	0.37	0.4	0.1	0.1	0.0028	0.15	0.73
1	0.37	0.4	0.1	0.1	0.0028	0.15	0.73
2	0.37	0.5	0.1	0.1	0.0028	0.19	0.91
3	0.37	0.6	0.1	0.1	0.0028	0.22	1.09
4	0.37	0.65	0.1	0.1	0.0028	0.24	1.19
5	0.37	0.7	0.1	0.1	0.0028	0.26	1.28
post lockdown	0.37	0.75	0.1	0.1	0.0028	0.28	1.37

Table 2: Parameters for the 5 phases of the Irish roadmap. β_{eff} and R_0 are computed from the system parameters.

– neglecting network-related effects – exhibit a burn out of an outbreak and self-guided decrease of the number of infected.

We introduce a small number of diseased individuals to the Dublin area on 22nd January, that is, 50 days before the initial lockdown, and run the model such that the disease spreads unimpeded. On the 12th of March, we introduce the initial lockdown measures, which take some time to reach full effect. Individuals' movement is limited to 2km and this is adhered to with 80% compliance. This level of compliance is supported by preliminary findings of a nation-wide series of phone interviews accounting for essential/justified and non-essential travel [6]. We account for social distancing and self-isolation (X compartment) by reducing the rate of mixing c within communities (cf. Tab. 2 for the respective parameter values). During phases 1 and 2/3, the travel radius increased to 5km and 20km, respectively, and we assume the same level of compliance. Travel restrictions are removed in phases 4 and 5 (cf. Tab. 1). Each of the numbered phases have a duration of 21 days.

Figures 2 depicts the time series of all five compartments – aggregated over all nodes – in semi-logarithmic scale. One can see two distinct peaks: (i) a small peak that occurs before the lockdown and (ii) a second peak that occurs once all lockdown measures are lifted. It is clear that unless some of the measure are maintained, then the lockdown will just have the effect of delaying the epidemic and result in a second wave with a prevalence that – in this unlikely worst-case scenario – is 100 times higher than the first one and exceeds any realistic capacity of the health-case system. The number of infected eventually reaches zero due to a depletion of the pool of susceptibles. Following the geometric analysis of the local dynamics (cf. Sec. 1), the trajectory lands at one of disease-free equilibria $I_e = 0$. A change in parameters, e.g., rate of mixing c, or a rescaling of the population to the number of remaining susceptibles can render this state unstable again. Then, a perturbation to the networked system, for instance, introduction or importation from outside, has the potential to trigger an additional outbreak.

For the simulation run given in Fig. 2 we compute the next generation (5) at each time-step and find its spectral radius giving rise to the network's basic reproduction number R_0^*, which is not only a function of the model parameters but also the number of susceptibles. For details, see Appendix A. For comparison, the local/simplistic value of R_0 given by Eq. (2) is shown as (dash-dotted) red curve. The value of R_0^* shows the potential for the disease to spread at each phase in the simulation. From Fig. 3 we can see clearly that when R_0^* falls below unity, the number of infected also begins to fall. We also note the point of herd immunity is reached is in when the number of susceptibles crossed

---

[6]HRB project: COVID19 rapid response programme (grant number: COV19-2020-117), unpublished data of four waves of telephone surveys.

Figure 2: Number of individuals (aggregated over the entire population/network) belonging to each compartment as stated in the legend over time with a log scale on the y-axis. Parameters as in Tab. 2. Each successive lockdown phase is indicated by the differently colored shaded regions on the plot. After the initial lockdown (violet), phases 1 – 5 last 3 weeks each.

the point where the R_0^* goes below 1.

Figure 3: The reproduction numbers R_0 and R_0^* over time as calculated from the basic reproduction number and spectral radius of the next generation matrix, respectively, together with the number of infected in the network in the doomsday scenario as shown in Fig. 2.

As stated above, the parameters are chosen to match the actual number of deaths in Ireland. This, in turn, results in a larger first peak compared to the reported numbers of confirmed cases on the 12th of March ranging from 27 according to [7] to 70 according to [8] The simulated results, however, are not meant to reflect this range of confirmed

---

[7]Health Protection Surveillance Centre (HPSC), `https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/casesinireland/` (accessed: June 22, 2020).

[8]Worldometer, `https://www.worldometers.info/coronavirus/country/ireland/` (accessed: June 22, 2020).

cases, but following the logic of the model, refer to all infected individuals including asymptomatic, non-diagnosed and untested.

(a)

(b)

Figure 4: The spatial distribution of infected individuals per km^2 around the two distinct peaks shown in Fig. 2 at (a) Day 50 (b) Day 320. Parameters as in Tab. 2.

Figures 4a and 4b show spatially resolved snapshots of the prevalence of the disease per km^2 at times $t = 50d$ (March 12) and $t = 320d$ (December 7), respectively (cf. [9] for a map based on data of reported, confirmed cases). Figure 4a corresponds to the disease prevalence at the first peak before the lockdown. The disease remains in the East of the country for the most part with the exception of big towns and cities. In contrast, at $t = 320d$ (Fig. 4b), the outbreak covers the entire country and affects every electoral division. See Fig. 2 for a full time series of the aggregated variables.

Figure 5 shows the effect of different levels of awareness during the post-lockdown phase, which is modelled by different values for the mixing parameter c. For convenience, the plots only depict the time series of the aggregated infected compartment I in a linear scale. A reduction to 70% of the original probability of infection significantly leads to a significantly smaller and delayed second peak (red curve), while the basic reproduction number of the local dynamics remains above unity (cf. Tab. 2).

Similarly to Figure 5, the network model (4) allows considering different levels of compliance with travel restrictions. The effect is exemplified for aggregated values of I, X and D in Fig. 6 for compliance of 80% (solid curves, cf. Fig. 2) and 70% (dashed curves). One can see that for lower compliance, the second peak occurs earlier and has a higher amplitude. This results in a slightly increased number of deaths.

In order to show how the heavy-tail distribution of travel across the network causes a lack of sensitivity in initial conditions, we run the model for a number of random initial seeds where for each run only a single individual in one electoral division or superoutput area. The resulting time series are shown in Fig. 7.

One can see that both amplitude and timing of the large, second peak are stable and it is impossible to visually differentiate between any of the curves for different initial conditions. In the presented 4 cases, the initial location is selected uniformly at random from all network nodes. We note that deliberately selecting urban or rural locations will have no qualitative impact. To further corroborate the results, initial infections of a significant, finite fraction of the population will be explored. This might account for a considerable amount of unreported cases at the start of the recording of SARS-CoV-2 cases.

3.3. Dynamic Interventions

Finally, we discuss the likely scenario of a flexible lockdown policy that monitors the prevalence of the disease within the island and aims to keep it below the capacity of the health-care system. As a simple measure, we define some threshold I_{th} as the maximum number of infected that can be present before we go back into lockdown and

---

[9]Central Statistics Office (CSO): Confirmed cases of COVID-19 by Electoral Division, https://census.cso.ie/covid19/.

Figure 5: Evolution of the aggregated, infected compartment I for varied post-lockdown conditions. The blue curve corresponds to the parameters given in Tab. 2 (cf. Fig. 2 for a semi-logarithmic scale). The other curves vary the parameter c in the post-lockdown phase as stated in the legend: $c = 0.9$, $c = 0.8$ and $c = 0.75$. Other parameters are as given in Tab. 2.

Figure 6: Evolution of the aggregated, infected I (blue), quarantined X (green) and deceased D (red) compartments for two level of compliance with lockdown restrictions: 80% (solid, cf. Fig. 2), and 70% (dashed). Other parameters as in Tab. 2.

go through earlier phases again. Such a procedure corresponds to dynamic [10, 18, 19] or active [3] interventions. Specifically, we simulate until phase 3 as before to maintain the agreement with reported cases. Afterwards, if the condition $I_{\text{th}} > \sum_i I_i$ is met, the system goes back to phase 1 and then progresses from there until the condition is met again.

Figure 7: Time series for 4 random initial conditions. Parameters as in Tab. 2 with $c = 0.75$ in the post-lockdown period.

Island Interventions. In detail, we consider two scenarios with different threshold parameters, which are inspired by the reported, maximum number of hospitalized cases during the first wave in April 2020[10]: (i) $I_{th} = 10^4$, as shown in Fig. 8 and (ii) $I_{th} = 10^3$ as depicted in Fig. 9. In both figures, the worst-case scenario of Fig. 2 is added as dashed curves. Note that the curves of recovered R (orange) and deaths D (red) are only increasing at a low rate. For the case of $I_{th} = 10^4$, for instance, these values arrive at 10^5 after two years and thus, way below any effect of herd immunity.

Under the assumptions the considered network model, it would take many years for a significant depletion of the pool of susceptibles. Extrapolating from an extended run of the model similar to Fig. 8, we find that it would take roughly 27 years for the number of recovered to reach half the population, leading to a total number of approximately 92,000 deaths over the course of many waves. Even with a large threshold $I_{th} = 10^4$ for the maximum number of infected cases, the feasibility of such a strategy is questionable due to the time it would take to reach herd immunity of on the island.

The re-introduction of lockdown measures to account for rising numbers of infected can change the stability of the disease-free equilibria $I_e = 0$. However, as soon as the restrictions are relaxed, e.g., during phases 3 to 5, an unstable family of these equilibria re-emerges and the outbreak pattern repeats. Suppressing the occurrence of unstable disease-free equilibria requires other forms of interventions. One could consider, for example, a fishing-type extension of the model [2, 5, 11]. This will basically turn the rate κ in the local dynamics (1) into a function of the number of infected, e.g., $\kappa(I) = \frac{h}{1+I/I_c}$ with a consumption/fishing rate h (here: rate of isolation) and a reference capacity I_c for $\kappa(I) = h/2$. This I dependence will account for contact-tracing effects, which are especially effective for low numbers of infected. As a consequence, a high rate h of isolation will be able to push the basic reproduction number [cf. Eq. (2) or Eq. (6)] below the critical value of $R_0^{crit} = 1$ and change the stability of the equilibria $I_e = 0$.

---

[10]Health Protection Surveillance Centre (HPSC) and the Health Service Executive (HSE): Ireland's COVID-19 Data Hub, https://covid19ireland-geohive.hub.arcgis.com (accessed: 10 July 2020).

Figure 8: The number of individuals belonging to each compartment over time (using the parameters in Tab. 2) under a dynamic lockdown strategy taking $I_{th} = 10^4$. The dashed lines are the equivalent model without the dynamic lockdown strategy shown in Fig. 2.

Figure 9: The number of individuals belonging to each compartment over time (using the parameters in Tab. 2) under a dynamic lockdown strategy taking $I_{th} = 10^3$. The dashed lines are the equivalent model without the dynamic lockdown strategy shown in Fig. 2.

Regional Interventions. Next, we consider another scenario that accounts for regional lockdowns at the level of counties. In our dynamic intervention model this corresponds to each county having its own I_{th}^c. Thus, once the county surpasses this threshold, another lockdown is triggered sending the county back to phase 1. When lockdown phases differ between counties, we assume that the travel restrictions in and out are followed with the compliance of the county being travelled to. When we run the simulations using the regional interventions, the model operates at

the level of electoral divisions and superoutput areas but their travel and interaction dynamics are consistent across counties. The parameters used for each phase are the same as those given in Tabs. 2 and 1. For this run we took the maximum number of infected per county $I_{th}^c = 1400$ and as usual the model is allowed to run without intervention until phase 3 to match the historic number of deaths.

Figure 10: Number of individuals (aggregated over counties) belonging to the I compartment. Parameters as in Tab. 2. Each successive lockdown phase is indicated by the differently colored shaded regions on the plot up until the lockdown phases for each county become desynchronised. After the initial lockdown (violet), phases 1 − 5 last 3 weeks each. The four largest counties by population are explicitly denoted in the legend.

From Fig. 10 we see some interesting dynamics when we view the evolution of the number of infected in each county. After Dublin re-enters phase 1 due to dynamic lockdown after only a few days the numbers start rising again. This is not due to its own dynamics, but due to its interaction with neighbouring counties, which are not yet in lockdown. It is only once all the other counties enter phase 1 lockdown do we see the numbers in Dublin and Cork begin to drop. We conclude that treating such counties in isolation is not a good strategy due the strong coupling between each county that is not eliminated when some counties remain free to travel even only within their own borders. Figure 11 shows the geographic distribution of the disease during the largest peak of Dublin seen in Fig. 10 at the level of electoral divisions and superoutput areas in panel (a) and at the level of counties in panel (b).

4. Conclusion

We have proposed a dynamical network model to study the spreading of SARS-CoV-2 that accounts for travel or commuting between nodes of the network. As a case study, we have presented results from numerical simulations for Ireland, where the model is informed from publicly available data on geographical regions, population, and mobility in both the Republic of Ireland and Northern Ireland. The network model is flexible in terms of spatial granularity and parameter selection. We have focused on the governmental 5-phase roadmap for reopening society & business and considered different initial conditions and parameters. We have observed that the roadmap will lead to a decline of case numbers. However, once the restrictions are lifted, the numbers will rise again, even if an increased level of awareness is considered. This is in excellent agreement with reported case numbers (cf. Appendix B).

Finally, we have elaborated on a procedure of dynamic interventions that re-introduce lockdown measures, if a certain threshold of infected is reached. Following such a protocol, the overall number of infected can be controlled and safely kept below a prescribed threshold, which can be selected according to the capacity of the health-care system. We have considered that containment measures would be re-introduced on a country-wide level, but a more regional approach (county level) appears feasible as well. In any case, while the outbreak can be managed by dynamic interventions, it might take a period of several years until a significant fraction of the population will have lived through the disease and – under the assumption of immunity – and will have led to an effect of herd immunity.

(a)

Figure 11: The geographic distribution of infected individuals in the simulation shown by the time-series in Fig. 10 on the 7th December 2020. Panel (a) shows the number of infected per km^2 in each electoral division and superoutput area. Panel (b) shows the number of infected in each county.

We hope that the results presented in this discussion paper will stimulate scientific and public discussion about the purpose and effect of lockdown measures in the months ahead (cf. Appendix C) and that is inspirational beyond the island of Ireland.

Acknowledgements

The work is supported by Health Research Board Ireland in the framework of the COVID19 rapid response programme (grant number: COV19-2020-117). We are grateful to Andreas Amann, Patricia Kearny, Gerry Kileen, and Ivan Perry for helpful discussions.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

RH, KM, SW, MOR, and PH designed the study. SW and RH developed the theory. RH and MS performed the numerical simulations and analyzed the data. All authors discussed the results and contributed to the writing of the article. All authors read and approved the final version of the manuscript.

References

[1] Al-Khani, A.M., Khalifa, M.A., AlMazrou, A., Saquib, N., 2020. The SARS-CoV-2 pandemic course in Saudi Arabia: A dynamic epidemiological model. Infectious Disease Modelling , 766–771.

[2] Amann, A., 2020. How to choose the best strategy for fighting an infectious disease. private communication.

[3] Ames, A.D., Molnar, T.G., Singletary, A.W., Orosz, G., 2020. Safety-critical control of active interventions for COVID-19 mitigation. medRxiv URL: https://www.medrxiv.org/content/early/2020/06/19/2020.06.17.20133264, doi:10.1101/2020.06.17.20133264, arXiv:https://www.medrxiv.org/content/early/2020/06/19/2020.06.17.20133264.full.pdf.

[4] Belik, V., Geisel, T., Brockmann, D., 2011. Natural human mobility patterns and spatial spread of infectious diseases. Phys. Rev. X 1, 011001. doi:10.1103/physrevx.1.011001.

[5] Beverton, R.J., Holt, S.J., 1993. On the dynamics of exploited fish populations. volume 11. Springer Science & Business Media.

[6] Blyuss, K.B., Kyrychko, Y.N., 2020. Effects of latency and age structure on the dynamics and containment of COVID-19. medRxiv .

[7] Brockmann, D., Hufnagel, L., 2007. Front propagation in reaction-superdiffusion dynamics: Taming Lévy flights with fluctuations. Phys. Rev. Lett. 98, 178301.

[8] Brockmann, D., Hufnagel, L., Geisel, T., 2006. The scaling laws of human travel. Nature 439, 462–465.

[9] Chen, N., Zhou, M., Dong, X., Qu, J., Gong, F., Han, Y., Qiu, Y., Wang, J., Liu, Y., Wei, Y., et al., 2020. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. The Lancet 395, 507–513.

[10] Chowdhury, R., Heng, K., Shawon, M.S.R., Goh, G., Okonofua, D., Ochoa-Rosales, C., Gonzalez-Jaramillo, V., Bhuiya, A., Reidpath, D., Prathapan, S., et al., 2020. Dynamic interventions to control COVID-19 pandemic: a multivariate prediction modelling study comparing 16 worldwide countries. Eur. J. Epidemiol. 35, 389–399.

[11] Christensen, V., Walters, C.J., 2004. Ecopath with Ecosim: methods, capabilities and limitations. Ecol. Modell. 172, 109–139.

[12] Dong, E., Du, H., Gardner, L., 2020. An interactive web-based dashboard to track COVID-19 in real time. Lancet Infect. Dis. 20, 533–534.

[13] Douglas, M., Katikireddi, S.V., Taulbut, M., McKee, M., McCartney, G., 2020. Mitigating the wider health effects of COVID-19 pandemic response. BMJ 369.

[14] Grauwin, S., Szell, M., Sobolevsky, S., Hövel, P., Simini, F., Vanhoof, M., Smoreda, Z., Barabási, A.L., Ratti, C., 2017. Identifying and modeling the structural discontinuities of human interactions. Sci. Rep. 7, 46677. doi:10.1038/srep46677.

[15] Humphries, R., 2020. EpiGraph. https://github.com/rory-humphries/EpiGraph-cpp.

[16] Johns Hopkins University, 2020. COVID-19 dashboard by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University. https://coronavirus.jhu.edu/map.html. (Accessed on 11/30/2020).

[17] Kermack, W.O., McKendrick, A.G., 1927. A contribution to the mathematical theory of epidemics. Proc. R. Soc. A 115, 700–721.

[18] Killeen, G., Kiware, S., 2020. Why lockdown? simplified arithmetic tools for decision-makers, health professionals, journalists and the general public to explore containment options for the novel coronavirus. medRxiv URL: https://www.medrxiv.org/content/early/2020/04/20/2020.04.15.20066845, doi:10.1101/2020.04.15.20066845, arXiv:https://www.medrxiv.org/content/early/2020/04/20/2020.04.15.20066845.full.pdf.

[19] Kissler, S.M., Tedijanto, C., Goldstein, E., Grad, Y.H., Lipsitch, M., 2020. Projecting the transmission dynamics of ARS-CoV-2 through the postpandemic period. Science 368, 860–868.

[20] Kyrychko, Y.N., Blyuss, K.B., Brovchenko, I., 2020. Mathematical modelling of the dynamics and containment of COVID-19 in Ukraine. Sci. Rep. 10, 1–11.

[21] Liu, Z., Magal, P., Seydi, O., Webb, G., 2020. A covid-19 epidemic model with latency period. Infectious Disease Modelling 5, 323–337.

[22] Maier, B.F., Brockmann, D., 2020. Effective containment explains subexponential growth in recent confirmed COVID-19 cases in china. Science 368, 742–746. doi:10.1126/science.abb4557.

[23] Pinotti, F., Di Domenico, L., Ortega, E., Mancastroppa, M., Pullano, G., Valdano, E., Boëlle, P.Y., Poletto, C., Colizza, V., 2020. Tracing and analysis of 288 early sars-cov-2 infections outside china: A modeling study. PLOS Med. 17, e1003193.

[24] Prem, K., Liu, Y., Russell, T.W., Kucharski, A.J., Eggo, R.M., Davies, N., Flasche, S., Clifford, S., Pearson, C.A., Munday, J.D., et al., 2020. The effect of control strategies to reduce social mixing on outcomes of the COVID-19 epidemic in Wuhan, China: a modelling study. The Lancet Public Health 5, e261–e270.

[25] Pullano, G., Valdano, E., Scarpa, N., Rubrichi, S., Colizza, V., 2020. Evaluating the effect of demographic factors, socioeconomic factors, and risk aversion on mobility during the COVID-19 epidemic in France under lockdown: a population-based study. The Lancet Digital Health 2, e638–e649.

[26] Schlosser, F., Maier, B.F., Hinrichs, D., Zachariae, A., Brockmann, D., 2020. COVID-19 lockdown induces structural changes in mobility networks–implication for mitigating disease dynamics. arXiv preprint arXiv:2007.01583 .

[27] Simini, F., González, M.C., Maritan, A., Barabási, A.L., 2012. A universal model for mobility and migration patterns. Nature 484, 96–100. doi:10.1038/nature10856.

[28] Song, C., Koren, T., Wang, P., Barabási, A.L., 2010a. Modelling the scaling properties of human mobility. Nat. Phys. 6, 818–823.

[29] Song, C., Qu, Z., Blumm, N., Barabási, A.L., 2010b. Limits of Predictability in Human Mobility. Science 327, 1018–1021. doi:10.1126/science.1177170, arXiv:http://www.sciencemag.org/cgi/reprint/327/5968/1018.pdf.

[30] van den Driessche, P., Watmough, J., 2002. Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. Mathematical Biosciences 180, 29 – 48. URL: http://www.sciencedirect.com/science/article/pii/S0025556402001086, doi:https://doi.org/10.1016/S0025-5564(02)00108-6.

[31] World Health Organization, 2020a. Novel coronavirus (2019-ncov). Situation Report 10.

[32] World Health Organization, 2020b. Novel coronavirus (2019-ncov). https://www.who.int/groups/covid-19-ihr-emergency-committee. (Accessed on 11/27/2020).

[33] World Health Organization, 2020c. Novel coronavirus (2019-ncov). Situation Report 51.

[34] Worldometer, 2020. Coronavirus update (live): 61,177,715 cases and 1,434,894 deaths from COVID-19 virus pandemic - worldometer. https://www.worldometers.info/coronavirus/. (Accessed on 11/27/2020).

A. Next Generation Matrix

In this appendix, we provide a detailed derivation of the next generation matrix given Eq. (5). We start by splitting the networked SIXRD equations (4) into the parts that cause new infections and all other changes. We shall denote the rate of appearance of new infections at node i in compartment U by $\mathcal{F}_{U,i}(x)$. Similarly, let $\mathcal{V}_{U,i}(x)$ be the rate of change of individuals in compartment U and node i by all other means such that $\dot{U}_i = \mathcal{F}_{U,i} - \mathcal{V}_{U,i}$. This yields the following equations:

$$\mathcal{F}_{S,i} = 0 \tag{7a}$$

$$\mathcal{F}_{I,i} = \beta c \left(S_i - \frac{S_i}{N_i} \sum_{j=1}^{n} N_{ij} \right) \frac{I_i + \sum_{j=1}^{n} \frac{I_j}{N_j} N_{ji} - \frac{I_i}{N_i} \sum_{j=1}^{n} N_{ij}}{N_i + \sum_{j=1}^{n}(N_{ji} - N_{ij})} \tag{7b}$$

$$+ \beta c \frac{S_i}{N_i} \sum_{j=1}^{n} N_{ij} \frac{I_j + \sum_{k=1}^{n} \frac{I_k}{N_k} N_{kj} - \frac{I_j}{N_j} \sum_{k=1}^{n} N_{jk}}{N_j + \sum_{k=1}^{n}(N_{kj} - N_{jk})}$$

$$\mathcal{F}_{X,i} = 0 \tag{7c}$$

$$\mathcal{F}_{R,i} = 0 \tag{7d}$$

$$\mathcal{F}_{D,i} = 0. \tag{7e}$$

It is an important note that the movement of individuals form I to X does not count as new infections.

$$\mathcal{V}_{S,i} = \beta c \left(S_i - \frac{S_i}{N_i} \sum_{j=1}^{n} N_{ij} \right) \frac{I_i + \sum_{j=1}^{n} \frac{I_j}{N_j} N_{ji} - \frac{I_i}{N_i} \sum_{j=1}^{n} N_{ij}}{N_i + \sum_{j=1}^{n}(N_{ji} - N_{ij})} \tag{8a}$$

$$+ \beta c \frac{S_i}{N_i} \sum_{j=1}^{n} N_{ij} \frac{I_j + \sum_{k=1}^{n} \frac{I_k}{N_k} N_{kj} - \frac{I_j}{N_j} \sum_{k=1}^{n} N_{jk}}{N_j + \sum_{k=1}^{n}(N_{kj} - N_{jk})}$$

$$\mathcal{V}_{I,i} = \mu I + \alpha I + \kappa I \tag{8b}$$

$$\mathcal{V}_{X,i} = -\kappa I_i + \mu X_i + \alpha X_i \tag{8c}$$

$$\mathcal{V}_{R,i} = -\mu I_i - \mu X_i \tag{8d}$$

$$\mathcal{V}_{D,i} = -\alpha I_i - \alpha X_i. \tag{8e}$$

For the above equations, we replace the coupling variables with their actual values, i.e. we take $S_{ij} = \frac{S_i}{N_i} N_{ij}$, $I_{ij} = \frac{I_i}{N_i} N_{ij}, \dots$ etc. Writing these equations in vector form we obtain

$$\mathcal{F} = \begin{bmatrix} \mathcal{F}_{I,1} \\ \vdots \\ \mathcal{F}_{I,n} \\ \mathcal{F}_{X,1} \\ \vdots \\ \mathcal{F}_{X,n} \\ \mathcal{F}_{S,1} \\ \vdots \\ \mathcal{F}_{S,n} \\ \mathcal{F}_{R,1} \\ \vdots \\ \mathcal{F}_{R,n} \\ \mathcal{F}_{D,1} \\ \vdots \\ \mathcal{F}_{D,n} \end{bmatrix}, \quad \mathcal{V} = \begin{bmatrix} \mathcal{V}_{I,1} \\ \vdots \\ \mathcal{V}_{I,n} \\ \mathcal{V}_{X,1} \\ \vdots \\ \mathcal{V}_{X,n} \\ \mathcal{V}_{S,1} \\ \vdots \\ \mathcal{V}_{S,n} \\ \mathcal{V}_{R,1} \\ \vdots \\ \mathcal{V}_{R,n} \\ \mathcal{V}_{D,1} \\ \vdots \\ \mathcal{V}_{D,n} \end{bmatrix}. \tag{9}$$

The disease-free equilibria are all solutions that satisfy

$$x_0 = (I_0, \dots, I_n, X_1, \dots, X_n, S_1, \dots, S_n, R_1, \dots, R_n, D_1, \dots, D_n) \tag{10}$$

$$= (0, \dots, 0, 0, \dots, 0, S_1, \dots, S_n, R_1, \dots, R_n, D_1, \dots, D_n) \tag{11}$$

Taking the derivatives of $\mathcal{F}$ and $\mathcal{V}$ at the disease-free equilibria, we find that

$$D\mathcal{F}(x_0) = \begin{bmatrix} F & 0 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad D\mathcal{V}(x_0) = \begin{bmatrix} V & 0 \\ J_1 & J_2 \end{bmatrix} \tag{12}$$

as shown in [30]. We obtain

$$F_{ij} = \beta c \left[\left(S_i - \frac{S_i}{N_i} \sum_{k=1}^{n} N_{ik} \right) \frac{\delta_{ij}\left(1 - \sum_{k=1}^{n} \frac{N_{ik}}{N_i}\right) + \frac{N_{ji}}{N_j}}{N_i + \sum_{k=1}^{n}(N_{ki} - N_{ik})} + \frac{S_i}{N_i} \sum_{k=1}^{n} N_{ik} \frac{\delta_{jk}\left(1 - \sum_{l=1}^{n} \frac{N_{jl}}{N_j}\right) + \frac{N_{jk}}{N_j}}{N_k + \sum_{l=1}^{n}(N_{lk} - N_{kl})} \right] \tag{13}$$

$$V_{ij} = (\mu + \alpha + \kappa)\delta_{ij} \tag{14}$$

Finally, following [30], the next generation matrix is given by

$$FV_{ij}^{-1} = \frac{\beta c}{\mu + \alpha + \kappa} \left[\left(S_i - \frac{S_i}{N_i} \sum_{k=1}^{n} N_{ik} \right) \frac{\delta_{ij}\left(1 - \sum_{k=1}^{n} \frac{N_{ik}}{N_i}\right) \frac{N_{ji}}{N_j}}{N_i + \sum_{k=1}^{n}(N_{ki} - N_{ik})} \right.$$

$$\left. + \frac{S_i}{N_i} \sum_{k=1}^{n} N_{ik} \frac{\delta_{jk}\left(1 - \sum_{l=1}^{n} \frac{N_{jl}}{N_j}\right) + \frac{N_{jk}}{N_j}}{N_k + \sum_{l=1}^{n}(N_{lk} - N_{kl})} \right]. \tag{15}$$

B. Comparison to Real-World Data

Figure 12 shows a comparison of model results for a countrywide dynamic lockdown scenario with daily and cumulative figures for reported deaths and infections from the Health Surveillance Centre (HPSC). Following the initial lockdown and phased reopening as outlined in Tab. 1, the threshold of infection was chosen to be 10000 in these simulations, before a national lockdown was reintroduced. This figure is adjustable based on current public health guidelines. Panels 12a and 12b compare the model to daily and cumulative reported figures for infections, respectively. Moreover, panels 12c and 12d compare the model results to daily and cumulative figures for deaths. The blue curves correspond to the model results, and the red curves refer to the data reported by HPSC. A 14-day moving average has been applied to data from the HPSC. Note that the infected compartment in the model does not distinguish between confirmed/undetected symptomatic/asymptomatic cases. This explains the larger number of infections in the model in which we see a clear qualitative agreement in the number of infections over time. The number of deaths, however, is more reliable and the model reproduces the actual numbers with large agreement. Overall, we find that the model results compare favourably to these real-world data.

Figure 12: Comparison of the countrywide dynamic lockdown scenario with threshold of infection $I_{thresh} = 10000$ (blue) to actual figures (red) for reported daily and cumulative infections (panels 12a and 12b) and daily and cumulative deaths (panels 12c and 12d) from the Health Protection Surveillance Centre (HPSC) in Ireland. Other parameters are as given in Tab. 2.

C. Projected Post-Christmas Lockdown

As a final scenario, we finish with a projection on the a possible post-Christmas projection during which we expect to see the third wave of lockdown. Figures 13 considers the current period of the second wave and the corresponding lockdown with level-5 movement restrictions in the Republic of Ireland. After a partial lifting of these measures scheduled for 1st December 2020, we find that the case numbers will rise again. Note that the numbers remain significantly higher than during summer 2020. This is illustrated by the maps in Fig. 14, which compares the situation by the end of June and 1st December. We project that the numbers will reach a high level similar to the introduction of the current lockdown by late-January 2021 and thus will require another lockdown of similar duration.

Figure 13: Simulating the evolution of the disease such that the historic deaths are matched up until level 5 and projected form then on. There are two subsequent periods of lockdown: the current lockdown that statred mid-November and is expected to lift 1st December. The next lockdown, which occurs in January 2021, is projected based on the number of the numbers of the November lockdown. Other parameters are as given in Tab. 2.

Figure 14: The geographic distribution of infected individuals per km^2 in each electoral division and superoutput area in the simulation shown by the time-series in Fig. 13. Panel (a) shows the geographic distribution during the lowest incidence of infected during the summer and panel (b) shows the geographic distribution at the end of the lockdown projected to end the 1st December 2020. Other parameters are as given in Tab. 2.