Modeling the Coronavirus Disease 2019 Incubation Period: Impact on Quarantine Policy

Daewoo Pak[1], Klaus Langohr[2], Jing Ning[1], Jordi Cortés Martínez[2], Guadalupe Gómez Melis[2,*], Yu Shen[1,**]

Abstract

Knowledge of the incubation period of coronavirus disease 2019 (COVID-19), defined as the time from infection to symptom onset, is critical to effectively confine COVID-19. However, the incubation period of COVID-19 is not always observed exactly due to uncertain onsets of infection and disease symptom. In this paper, we demonstrate how to estimate the distribution of incubation and its association with patient demographic factors when the dates of infection and disease onset are not explicitly. We employ a sufficiently general parametric class, the generalized odds-rate class of regression models, which includes the log-logistic proportional odds model and the Weibull proportional hazards model as special cases. We base our analysis on publicly reported, clinically confirmed COVID-19 cases with potential exposure period information. The application indicates that an age-specific quarantine policy might be more efficient than a unified one in confining COVID-19.

Keywords: COVID-19, Generalized odds-rate class of models, Incubation period, Infectious disease, Interval censoring, left censoring, Quarantine policy

[*]lupe.gomez@upc.edu

[**]yshen@mdanderson.org

[1]Department of Biostatistics, The University of Texas, MD Anderson Cancer Center, Houston, Texas, USA

[2]Department of Statistics and Operations Research, Universitat Politècnica de Catalunya, Barcelona, Spain

1. Introduction

The coronavirus disease 2019 (COVID-19) was first reported in Wuhan, China, in December 2019; an outbreak rapidly spread worldwide. The novel virus infection can be asymptomatic or unapparent during a certain period and asymptomatic persons could spread the virus unknowingly [1]. Among patients who develop symptoms, the incubation period is defined as the elapsed time between infection and appearance of the first symptom. Knowledge of the incubation period is essential for disease prevention, facilitating an optimal quarantine guideline to confine the spread.

The incubation period of COVID-19 was investigated in several reports [2, 3, 4, 5]. Despite its importance, it remains unclear how the incubation distribution could vary by gender and age and whether the current 14-day quarantine period ignoring the patient demographic factors would be sufficient for the containment of COVID-19. A more accurate estimation of the incubation period using these factors is necessary to optimize the surveillance guidelines. We model uncertain dates of infection and symptom onset with the best available information, by classifying COVID-19 confirmed cases into plausible observing scenarios. This approach allows the use of incomplete observations on the COVID-19 incubation period and thereby provides more reliable results and inferences. We also revisit the impact of the current quarantine duration on COVID-19's spread by estimating the distribution of the incubation periods of COVID-19 and its association with gender and age.

2. Exposure History Reports for Confirmed COVID-19 Cases

We outline a modeling approach to enable the analysis of publicly reported, clinically confirmed cases with symptoms from two sources [6, 7], available as of March 30, 2020. To obtain the period of potential exposure, we included the following patients in the analytic data set: 1) non-residents of Wuhan who visited Wuhan since December 1, 2019 and for whom the exposure period was the time between the earliest possible arrival to and the latest possible departure from Wuhan; 2) people with travel history of visiting the coronavirus-affected nations known at the time or taking the Diamond Princess Cruises; 3) non-travel related cases with an exposure history based on contact with infected persons. Among these, 312 cases had recorded exposure periods, or at least an exposure end date, and dates of symptom onset or hospitalization, as well as gender and age; this number reflects removal of 34 potentially duplicated cases (see Figure 1).

Figure 1: Flow diagram of confirmed cases for data analysis. Two data sources were combined from DXY [6] and Xu et al [7].

There were 111 (35.6%) cases with known exposure start and end dates, illustrated in Supplementary Figure S1. For cases without an exact start date of the exposure period, the initial date was set to have a maximum of thirty days of exposure or was set to December 1, 2019, whichever one was later. In addition, the end date of exposure was set to precede the known dates of symptom onset and hospitalization. If the exact date of symptom onset was unknown, it was assumed to have occurred before hospitalization (i.e., left-censored).

3. Model and Estimation

The distribution of the COVID-19 incubation period is estimated by modelling the interval-censored exposure duration and the possibly left-censored symptom onset time. We parameterize the underlying incubation period under the generalized odds-rate class of regression models with patient's age and gender. Let T be the incubation period and $\boldsymbol{X}$ be the covariate vector including both the patient's age and gender. Within the generalized odds-rate class of regression models, the conditional probability density function of T given $\boldsymbol{X} = \boldsymbol{x}$ is defined as

$$f(t|\boldsymbol{x}) = \phi \lambda^{-\phi} t^{\phi-1} e^{\boldsymbol{x}^T \boldsymbol{\beta}} \left\{ 1 + \rho \left(t/\lambda \right)^\phi e^{\boldsymbol{x}^T \boldsymbol{\beta}} \right\}^{-(1+\rho)/\rho},$$

where $\rho > 0$, $\lambda > 0$, $\phi > 0$, and $\boldsymbol{\beta}$ is the vector of the regression parameters. The corresponding survival function is

$$S(t|\boldsymbol{x}) = \left\{ 1 + \rho \left(t/\lambda \right)^\phi e^{\boldsymbol{x}^T \boldsymbol{\beta}} \right\}^{-1/\rho}.$$

This class of models includes the log-logistic proportional odds model ($\rho = 1$) and the Weibull proportional hazards model ($\rho \to 0$) as special cases [8], which survival functions are expressed as, respectively,

$$\text{Proportional odds model:} \quad S(t|\boldsymbol{x}) = \left\{ 1 + \left(t/\lambda \right)^\phi e^{\boldsymbol{x}^T \boldsymbol{\beta}} \right\}^{-1}, \text{ and}$$

$$\text{Proportional hazard model:} \quad S(t|\boldsymbol{x}) = e^{-(t/\lambda)^\phi e^{\boldsymbol{x}^T \boldsymbol{\beta}}}.$$

To construct the likelihood function of COVID-19 data, we distinguish the four types of observation, as shown in Figure 2. Let d denote the difference between the exposure start date and end date, and r denote the difference between the exposure end date and either the symptom onset date (if available) or the date of hospitalization. If the date of infection is at best known to be within a potential exposure period (i.e., $d > 0$), let $\delta_d = 1$; otherwise $\delta_d = 0$. If a patient's symptom onset date was observed, let $\delta_s = 1$; otherwise, $\delta_s = 0$. Under the assumption of a uniform risk for infection during the potential exposure periods, the contributions of each type to the likelihood function is given as follows:

Type I: observing potential exposure period and symptom onset date (235 patients)

Type II: observing potential exposure period and hospitalization date (60 patients)

Type III: observing exact exposure date and symptom onset date (16 patients)

Type IV: observing exact exposure date and hospitalization date only (1 patient)

Figure 2: The four types of observations available for the data analysis. The total number of patients is 312.

- Type I: observing potential exposure period and symptom onset date ($\delta_d = 1$ and $\delta_s = 1$)

$$L_1 = d^{-1} \int_0^d f(a + r|\boldsymbol{x})da = d^{-1} \left\{ S(r|\boldsymbol{x}) - S(r + d|\boldsymbol{x}) \right\},$$

- Type II: observing potential exposure period and hospitalization date ($\delta_d = 1$ and $\delta_s = 0$)

$$L_2 = d^{-1} \int_0^d \left\{ 1 - S(a + r|\boldsymbol{x}) \right\} da = 1 - d^{-1} \int_0^d S(a + r|\boldsymbol{x})da,$$

- Type III: observing exact exposure date and symptom onset date ($\delta_d = 0$ and $\delta_s = 1$)

$$L_3 = f(r|\boldsymbol{x}),$$

- Type IV: observing exact exposure date and hospitalization date only ($\delta_d = 0$ and $\delta_s = 0$)

$$L_4 = 1 - S(r|\boldsymbol{x}).$$

Considering n independent observations, the overall log-likelihood is then proportional to

$$\log L = \sum_{i=1}^n \delta_{si}\delta_{di} \log L_{1i} + (1 - \delta_{si})\delta_{di} \log L_{2i} + \delta_{si}(1 - \delta_{di}) \log L_{3i} + (1 - \delta_{si})(1 - \delta_{di}) \log L_{4i}$$

$$\propto \sum_{i=1}^n \left[\delta_{si}\delta_{di} \log \left\{ S(r_i|\boldsymbol{x}_i) - S(r_i + d_i|\boldsymbol{x}_i) \right\} + (1 - \delta_{si})\delta_{di} \log \left\{ 1 - d_i^{-1} \int_0^{d_i} S(a + r_i|\boldsymbol{x}_i)da \right\} \right.$$

$$\left. + \delta_{si}(1 - \delta_{di}) \log f(r_i|\boldsymbol{x}_i) + (1 - \delta_{si})(1 - \delta_{di}) \log \left\{ 1 - S(r_i|\boldsymbol{x}_i) \right\} \right]$$

The model parameters are estimated by maximizing the log-likelihood. In our implementation, we maximized the log-likelihood with the logarithmic transformation for the positive parameters (i.e., λ, ϕ, and ρ). The delta method was used to obtain the asymptotic normal distribution for any parameter transformation, such as the median incubation time. All analyses were performed with R, version 3.6.1 [9]. The R code is provided with the supplementary material.

4. Application

4.1. Estimation results

We applied the proposed method to the COVID-19 data sets, described in Section 2. Within the total of 312 patients, the median age was 42 (interquartile range $33 - 55$) years, and 126 (40.4%) patients were women. Age was dichotomized at its median prior to the analysis. Table 1 shows the estimation results for regression models with gender and age. Patients older than 42 years of age have, on average, longer incubation periods, compared to 42-year-old or younger patients ($p = 0.039$), whereas gender has no effect on the incubation period ($p = 0.423$). We decide not to include gender in the final model (Reduced model of Table 1), based on the likelihood ratio test with $p = 0.384$. The estimated median incubation times are 5.3 (95% CI: $4.3 - 6.3$) days and 7 (95% CI: $5.6 - 8.3$) days among younger and older patients, respectively. The estimated mean incubation period is 6.6 (95% CI: $5.4 - 7.8$) days for the younger patients, and 8.8 (95% CI: $7.2 - 10.3$) days for the older patients. The estimated distribution functions in both age groups together with the 95% confidence intervals are presented in Figure 3.

Table 1: Results of the analysis of COVID-19 under the generalized odds-rate class of regression models. The p-value of the likelihood ratio test (Full model vs Reduced model) was 0.384.

Model		$log(\lambda)$	$log(\phi)$	$log(\rho)$	β_{male}	$\beta_{age \leq 42}$
	Estimate	2.300	0.305	-3.766	0.141	0.371
Full model	SE	0.155	0.119	6.569	0.176	0.180
	P-value	<0.001	0.010	0.566	0.423	0.039
	Estimate	2.115	0.457	-1.271	-	0.449
Reduced model	SE	0.126	0.128	0.685	-	0.214
	P-value	<0.001	<0.001	0.064	-	0.035

Table 2 shows the estimated differences (in days) between the quantiles of the incubation distribution in both age groups. The incubation period difference between the two age groups was not obvious in lower quantiles, but it becomes conspicuous after the $25th$ quantile, leading to about 7-day differences between the two age groups at the $97.5th$ quantile.

Figure 3: The estimated cumulative distribution of the incubation period by two age groups.

Table 2: Estimated differences of specific quantiles of the incubation period among both age groups (patients aged > 42 years – patients aged ≤ 42 years).

Percentile	2.5th	25th	50th	75th	97.5th
Difference (days, 95% CI)	0.2 (0.0, 0.4)	1.0 (0.1, 1.8)	1.7 (0.2, 3.3)	2.9 (0.3, 5.4)	6.7 (0.6, 12.8)

4.2. Impact on quarantine policy

The finding that persons older than 42 years have, on average, longer incubation periods than those who are younger may have important implications for enacting age-specific quarantine policies. This result agrees with previous studies on Severe Acute Respiratory Syndrome that showed a relationship between age and the incubation period [10]. Using the uniform 14-day quarantine policy recommended by the World Health Organization and the US Centers for Disease Control and Prevention [11, 12], our estimators imply that 8.4% (95% CI: $3.6 - 13.1\%$) of COVID-19 patients younger than 42, and 17.1% (95% CI: $9.9 - 24.2\%$) of older patients may pose a risk of infection to others before onset of their symptoms. Using a 21-day quarantine, these percentages reduce to 2.2% (95% CI: $0 - 7.0\%$) and 5.8% (95% CI: $1.3 - 13.0\%$), respectively. To ensure that at least 90% of cases' symptoms are being manifested during quarantine periods, the required durations are estimated to be 13.1 (95% CI: $10.5 - 15.7$) days for patients 42 years of age

or younger and 17.4 (95% CI: $14.0 - 20.8$) days for patients older than 42 years. As such, a unified quarantine policy could be inefficient during a viral outbreak. These estimates were derived from the conservative assumption that the quarantine periods started immediately after infection.

5. Discussion

We have proposed a likelihood-based approach to estimate the distribution of the incubation period with the patient-related covariates using publicly available COVID-19 data sets. The proposed method allows us to use a larger data cohort consisting of more confirmed COVID-19 cases than other published papers in this field; the cases may not have the exact symptom onset time (left-censored) and include an interval-censored incubation period. One of the challenges of modeling unknown incubation periods is the choice of the statistical distribution due to their large variability. Our choice of a parametric model for the incubation period has more flexibility than other commonly used distributions, which were often compared to identify the distribution that provides the best fit to the data [2, 4].

To examine how robust the estimates were to the assumption of 30 days of maximum exposure for cases with missing start date of exposure interval, we performed some sensitivity analyses considering a maximum exposure duration of 20 days or shifting the lower bound back and forth within 15 days. The variation of these assumptions had little effect on analysis results (Supplementary Table S1). Although the proposed method assumes that the infection would happen uniformly during the potential exposure period, it can be easily modified with other forms of risk for various purposes.

The application to the COVID-19 data sets has some notable limitations. Our inferences relied on publicly reported confirmed cases that might over represent more severely symptomatic patients. Moreover, the definition of COVID-19 symptoms and hospitalization criteria could differ by country, especially during the initial outbreak. We combined the data sets from two different sources, and the potential variation in source criteria for tracing infected cases may lead to different exposure records. However, we obtained similar findings when fitting the model to each data set separately, although one indicated no statistically significant differences, but the same trend in the incubation period between the age groups (Supplementary Table S2 and S3). We dichotomized age to show the difference in the incubation period time that might exist between the two groups. However, in no case was it

our goal to identify an optimal cut-off age, since we are aware of the risks involved in the dichotomization of the explanatory variables [14, 13].

Author Contributions

Conceptualization, D.P., G.G., J.N. and Y.S.; Methodology, D.P., J.N. and Y.S.; formal analysis, D.P., K.L., J.N. and J.C.; writing–original draft preparation, D.P., G.G., J.N. and Y.S. All authors have read and agreed to the published version of the manuscript.

Funding

This research was funded by the Ministerio de Economía y Competitividad (Spain) [MTM2015-64465-C2-1-R (MINECO/FEDER)] and the Departament d'Economia i Coneixement de la Generalitat de Catalunya (Spain) [2017 SGR 622 (GRBIO)]. The authors declare no conflict of interest.

Conflicts of Interest

The authors declare no conflict of interest.

References

[1] Wilson, M. E.; Chen, L. H. Travellers Give Wings to Novel Coronavirus (2019-NCoV). *Journal of Travel Medicine* **2020**, *27*, taaa015. https://doi.org/10.1093/jtm/taaa015.

[2] Backer, J. A.; Klinkenberg, D.; Wallinga, J. Incubation Period of 2019 Novel Coronavirus (2019-NCoV) Infections among Travellers from Wuhan, China, 20–28 January 2020. *Eurosurveillance* **2020**, 25. https://doi.org/10.2807/1560-7917.ES.2020.25.5.2000062.

[3] Lauer, S. A.; Grantz, K. H.; Bi, Q.; Jones, F. K.; Zheng, Q.; Meredith, H. R.; Azman, A. S.; Reich, N. G.; Lessler, J. The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Ann Intern Med* **2020**, *172*, 577-582. https://doi.org/10.7326/M20-0504.

[4] Linton, N.M.; Kobayashi, T.; Yang, Y.; Hayashi, K.; Akhmetzhanov, A.R.; Jung, S.-M.; Yuan, B.; Kinoshita, R.; Nishiura, H. Incubation Period and Other Epidemiological Characteristics of 2019 Novel Coronavirus Infections with Right Truncation: A Statistical Analysis of Publicly Available Case Data. *J. Clin. Med.* **2020**, *9*, 538. https://doi.org/10.3390/jcm9020538.

[5] Guan, W.; Ni, Z.; Hu, Y.; Liang, W.; Ou, C.; He, J.; Liu, L.; Shan, H.; Lei, C.; Hui, D. S. C.; et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* **2020**, *382*, 1708–1720. https://doi.org/10.1056/NEJMoa2002032.

[6] DXY: COVID-19 Epidemic situation in real time. Available online: https://docs.google.com/spreadsheets/d/e/2PACX-1vQU0SIALScXx8VXDX7yKNKWWPKE1YjFlWc6VTE VSN45CklWWf-uWmprQIyLtoPDA18tX9cFDr-aQ9S6/pubhtml (accessed on 20 March 2020).

[7] Xu, B.; Gutierrez, B.; Mekaru, S.; Sewalk, K.; Goodwin, L.; Loskill, A.; Cohn, E. L.; Hswen, Y.; Hill, S. C.; Cobo, M. M.; et al. Epidemiological Data from the COVID-19 Outbreak, Real-Time Case Information. *Sci Data* **2020**, *7*, 106. https://doi.org/10.1038/s41597-020-0448-0.

[8] Dabrowska, D. M.; Doksum, K. A. Estimation and Testing in a Two-Sample Generalized Odds-Rate Model. *Journal of the American Statistical Association* **1988**, *83* , 744–749. https://doi.org/10.1080/01621459.1988.10478657.

[9] R Core Team (2019). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

[10] Cowling, B. J.; Muller, M. P.; Wong, I. O. L.; Ho, L.-M.; Louie, M.; McGeer, A.; Leung, G. M. Alternative Methods of Estimating an Incubation Distribution: Examples from Severe Acute Respiratory Syndrome. *Epidemiology* **2007**, *18*, 253–259.

[11] World Health Organization (WHO). Considerations for quarantine of individuals in the context of containment for coronavirus disease (COVID-19). Available online: https://www.who.int/publications-detail/considerations-forquarantine-of-individuals-in-the-context-of-containment-forcoronavirus-disease-(covid-19) (accessed on 12 April 2020)

[12] The White House. Press Briefing by Members of the President's Coronavirus Task Force. Available online: https://www.whitehouse.gov/briefings-statements/press-briefing-members-presidents-coronavirus-task-force (accessed on 1 February 2020).

[13] Moons K. G. , Altman D. G., Reitsma J. B., et al. Transparent Reporting of a multivariable prediction model for Individual Prognosis or Diagnosis (TRIPOD): explanation and elaboration. *Annals of internal medicine* **2015**, *162(1)*, W1-73. 2015.

[14] Royston P., Altman D. G., and Sauerbrei W. Dichotomizing continuous predictors in multiple regression: a bad idea. *Stat. Med.* **2006**, *25*, 127–141. 2015.