Statistics associated with the lethality of COVID-19 by age group and gender in Mexico

Carlos Hernandez-Suarez[a,*], Efren Murillo-Zamora[b]

[a]*Facultad de Ciencias, Universidad de Colima, Bernal Diaz del Castillo 340, Colima, Colima, 28040, MEXICO*
[b]*Departamento de Epidemiología, Unidad de Medicina Familiar No. 19, IMSS, Av. Javier Mina 301, 28000, Colima, Colima. MEXICO*

Abstract

We analyzed outcomes of 34,686 SARS-CoV-2 confirmed cases of patients attending the IMSS (Mexican Institute for National Insurance) from January 2 to June 23, 2020. We calculated relative IFR by age group and gender and introduced the concept of *adjusted lethalities*, that can be used to project the burden of the disease for a population with different demographic characteristics.

Keywords: COVID-19, SARS-CoV-2, IFR, Lethality, Age, Gender, Adjusted lethality

1. Introduction

Since it was first identified in pneumonia patients in Wuhan, China, the coronavirus disease 2019 (COVID-19) by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has been characterized by its fast-spreading and high

5 related disease burden [1]. The spectrum of symptoms is wide and ranges from asymptomatic infection to severe respiratory illness [2]. By June 24, 2020, almost 10 million confirmed cases of COVID-19 had been reported globally and a fatal outcome was registered in nearly a half million of them [3]. Gender and age-related differences in disease severity and risk of death have been documented

10 [4].

---

*Corresponding author
  Email addresses: `carlosmh@mac.com` (Carlos Hernandez-Suarez), `efren.murilloza@imss.gob.mx` (Efren Murillo-Zamora)

The lethality of a disease, also known as Infection Fatality Rate (IFR) is the fraction of infected individuals that die from the disease. It's estimation is of the upmost importance for the design of and application of containment measures and to decide if lockdowns are implemented as when these should be reinforced or lifted. Unfortunately, our data does not come from a random sample and we cannot estimate the IFR, instead, we estimate the relative share of the IFR by age group and gender and introduce the concept of *adjusted lethalities* that can be used to project the burden of the disease for a population whose demographic characteristics is known.

Methodology

We used the database of the Instituto Mexicano de Seguro Social (IMSS, acronym for Mexican Institute for National Insurance) with confirmed SARS-CoV-2 cases. The IMSS provides health services to about 10% of the population in Mexico, over 12 million persons. The database on SARS-CoV-2 we used contains confirmed cases from January 2 to June 23, 2020. We included only cases with disease outcome as dead or recovered, leaving a database with 34,686 cases, from which 57 % were males and 43% females.

When a patient is diagnosed with COVID-19, it follows one of the paths depicted in Figure 1. With this data we constructed Table 1 after dividing the population in seven age categories and two genders.

Table 2 shows a processing of the information in Table 1. Vectors $\mathbf{h}$, $\mathbf{c}$, $\mathbf{d}$ and $\mathbf{d}_c$ in Table 2 are described as follows:

$\mathbf{h}$ = relative frequency of hospitalized patients.

$\mathbf{c}$ = fraction requiring ICU among those that went into hospitalization.

$\mathbf{d}$ = fraction dying among those that went into hospitalization.

$\mathbf{d}_c$ = fraction dying among those that went into ICU.

As mentioned before, it is not possible to estimate the IFR from the data

Figure 1: Transitions in our model: $N_0 + N_1 = N$, $N_2 + N_3 + N_4 = N_1$, $N_5 + N_6 = N_3$

available (Table 1) since the death rates are conditional on patients mostly symptomatic and thus they are not a representative sample of the response to an infection. Nevertheless, there is a couple of things that we can obtain from Table 2. The fist one is the *relative lethality* and the second *adjusted lethality*.

2. Relative lethality

Table 2 tells us, for instance, that the probability that a female person in age group 40-49 that is hospitalized from infection with SARS-CoV-2 dies is 0.3548, but this does not provides information on the probability that a female infected person in age group 40-49 dies from the disease. Nevertheless, observe that if f_{ij} is the fraction in the population of individuals of gender $i, i = 1, 2$ and group age $j, j = 1, 2, \ldots, 7$., and we assume that everyone is equally likely to become infected but there is a differential response to the disease, then, if p is the fraction of individuals in the population that is infected and β_{ij} is the fraction of infected individuals of gender i and age group j that become hospitalized, then:

$$f_{ij} \; p \; \beta_{ij} \propto h_{ij} \tag{1}$$

Where h_{ij} is the fraction of hospitalized that belong to age group i and gender j. Now, if α_{ij} is the fraction of individuals hospitalized that die from COVID-19, then

$$f_{ij} \; p \; \beta_{ij}\alpha_{ij} \propto h_{ij}d_{ij} \tag{2}$$

where d_{ij} is the fraction of individuals of these hospitalized that dies.

Let θ_{ij} be the *lethality* or infection fatality rate of gender i and age group j, then, the lethality of gender i and age group j can be written as:

$$\theta_{ij} = \beta_{ij}\alpha_{ij}. \tag{3}$$

Thus, the IFR by gender and group age follows this relationship:

$$\theta_{ij} \propto h_{ij}d_{ij}/f_{ij}. \tag{4}$$

The reason why we only establish that θ_{ij} is proportional to some value is because we ignore the value of p, the true proportion of infected individuals in the population. Thus, we define θ_{ij}^* as the *relative lethalitiy*:

$$\theta_{ij}^* = h_{ij}d_{ij}/f_{ij} \tag{5}$$

Even although the value of p is not known, the ratio of any two relative lethalities $\theta_{ij}^*/\theta_{kl}^*$ reflects the ratio of the true lethalities, that is:

$$\theta_{ij}^*/\theta_{kl}^* = \theta_{ij}/\theta_{kl},$$

therefore this quotient can be used to compare the lethality of any two groups. Table 3 shows in the last column the relative infection fatality rates in (5) , after normalization. In this table $\mathbf{f}_M$ and $\mathbf{f}_F$ are the relative frequencies by age group and gender in the Mexican population. In this table we can see, for instance, that the lethality of COVID-19 in age group 60+ is $0.37511/0.19795 = 1.9$ times higher in males than in females, and that the lethality in the 60+ group compared to the group with less than 20 years age is:

$$\frac{0.37511 + 0.19705}{0.00068 + 0.00024 + 0.00041 + 0.00048} = 317.7$$

times larger. Table 3 also shows that the lethality among people 50 years or older is about 6 times larger than among people less than 50 years old.

3. Adjusted lethality

It is possible to estimate the true values of $\boldsymbol{\theta}_M$ and $\boldsymbol{\theta}_F$. The lethality of the disease is the probability that a person at random that has been infected with SARS-CoV-2 dies, which is:

$$\text{IFR} = \mathbf{f}'_M \boldsymbol{\theta}_M + \mathbf{f}'_F \boldsymbol{\theta}_F \tag{6}$$

which assumes all individuals are equally likely to become infected. If we have an estimate of the lethality of the disease, θ, then we can rescale $\boldsymbol{\theta}^*_M$ and $\boldsymbol{\theta}^*_F$ multiplying by a constant c so that

$$\theta = c \ \mathbf{f}'_M \boldsymbol{\theta}^*_M + c \ \mathbf{f}'_F \boldsymbol{\theta}^*_F \tag{7}$$

Some estimates of the IFR are in the range $0.05\% - 0.25\%$ [5, 6, 7]. Here we will use the lethality reported by [5] of 86 in 100, 000 (about 0.1%) because it is a more recent study with a large sample size, thus, we use $\theta = 0.001$. The adjusted lethalities by age group and gender are shown in Table 4.

One immediate application of Table 4 is the possibility of projecting the IFR for another population whose demographics by age and gender are known. Consider for instance the demographics of the Za'atari refugee camp, in Jordan[1], which is shown in Table 5. We use $\boldsymbol{\theta}_M$ and $\boldsymbol{\theta}_F$ from Table 4 and the $\mathbf{f}_M$ and $\mathbf{f}_F$ from the camp's demographic and thus the IFR projected for this camp is:

$$\text{IFR} = \mathbf{f}'_M \boldsymbol{\theta}_M + \mathbf{f}'_F \boldsymbol{\theta}_F = 0.00038.$$

---

[1] https://data2.unhcr.org/en/situations/syria/location/53

₉₀ Observe this adjusted lethality is about 2.6 times smaller than the overall IFR, which is due to the fact that Za'atari refugee camp is a young population with 3/4 of the refugees being less than 30 years old. Although this comparison used $\theta = 0.001$ as the overall IFR of SARS-CoV-2, the observed ratio of 2.6 is independent of this value because the adjusting constant c in 7 cancels out.

₉₅ Clearly, this projection includes only demographic factors and not other health and socioeconomic factors or availability of health services. Figure 2 shows a comparison between the demographics of the Za'atari camp and the *Adjusted lethalities* from Table 4 by age group and gender.

Figure 2: Population frequency in Za'atari camp (left) and Adjusted lethalities from Table 4 (right) by age and gender. For this particular camp, the lethality affects a group of individuals that are found at a very low frequency in the camp.

The distribution of deaths across age groups and gender.

₁₀₀ Another use of the *adjusted lethalities* in Table 4 is the estimation of the distribution of deaths across group ages and genders. The question we want to answer is: what is the probability that a dead individual belongs to a particular gender and age group? This is clearly given by $\mathbf{f}_M \circ \boldsymbol{\theta}_M$ and $\mathbf{f}_F \circ \boldsymbol{\theta}_F$ after normalization. For the Mexican population, a plot of this distribution is given ₁₀₅ in Figure 3.

Conflict of interest

Authors declare no conflict of interest.

Figure 3: The projected distribution of deaths across age groups and gender for Mexico's population.

Study approval

This work is part of the program "Building the Evidence on Protracted Forced Displacement: A Multi-Stakeholder Partnership". The program is funded by UK aid from the United Kingdom's Department for International Development (DFID), it is managed by the World Bank Group (WBG) and was established in partnership with the United Nations High Commissioner for Refugees (UNHCR). The scope of the program is to expand the global knowledge on forced displacement by funding quality research and disseminating results for the use of practitioners and policy makers. This work does not necessarily reflect the views of DFID, the WBG or UNHCR. This study had approval R-2020-601-07 by the Health Research Ethics Committee (601) of the IMSS.

[1] F. Zhou, T. Yu, R. Du, G. Fan, Y. Liu, Z. Liu, J. Xiang, Y. Wang, B. Song, X. Gu, et al., Clinical course and risk factors for mortality of adult inpatients with covid-19 in wuhan, china: a retrospective cohort study, The lancet.

[2] K. Gostic, A. C. Gomez, R. O. Mummah, A. J. Kucharski, J. O. Lloyd-

Smith, Estimated effectiveness of symptom and risk screening to prevent the spread of covid-19, Elife 9 (2020) e55570. `doi:10.7554/eLife.55570`.

[3] World Health Organization, Coronavirus disease (COVID-2019) situation reports: Situation report - 150, June 18, 2020, Accessed on June 19, 2020.
URL `https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200618-covid-19-sitrep-150.pdf?sfvrsn=aa9fe9cf_2`

[4] W. Tian, W. Jiang, J. Yao, C. J. Nicholson, R. H. Li, H. H. Sigurslid, L. Wooster, J. I. Rotter, X. Guo, R. Malhotra, Predictors of mortality in hospitalized covid-19 patients: A systematic review and meta-analysis, Journal of Medical Virology`doi:10.1002/jmv.26050`.

[5] C. Erikstrup, C. E. Hother, O. B. V. Pedersen, K. Mølbak, R. L. Skov, D. K. Holm, S. Sækmose, A. C. Nilsson, P. T. Brooks, J. K. Boldsen, C. Mikkelsen, M. Gybel-Brask, E. Sørensen, K. M. Dinh, S. Mikkelsen, B. K. Møller, T. Haunstrup, L. Harritshøj, B. A. Jensen, H. Hjalgrim, S. T. Lillevang, H. Ullum, Estimation of sars-cov-2 infection fatality rate by real-time antibody screening of blood donors, medRxiv`arXiv:https://www.medrxiv.org/content/early/2020/04/28/2020.04.24.20075291.full.pdf`, `doi:10.1101/2020.04.24.20075291`.
URL `https://www.medrxiv.org/content/early/2020/04/28/2020.04.24.20075291`

[6] H. Streeck, B. Schulte, B. Kuemmerer, E. Richter, T. Hoeller, C. Fuhrmann, E. Bartok, R. Dolscheid, M. Berger, L. Wessendorf, M. Eschbach-Bludau, A. Kellings, A. Schwaiger, M. Coenen, P. Hoffmann, M. Noethen, A.-M. Eis-Huebinger, M. Exner, R. Schmithausen, M. Schmid, B. Kuemmerer, Infection fatality rate of sars-cov-2 infection in a german community with a super-spreading event, medRxiv`arXiv:https://www.medrxiv.org/content/early/2020/05/08/2020.05.04.20090076.full.pdf`, `doi:10.1101/2020.05.04.20090076`.

URL `https://www.medrxiv.org/content/early/2020/05/08/2020.05.04.20090076`

[7] J. Ioannidis, The infection fatality rate of covid-19 inferred from seroprevalence data, medRxivarXiv:`https://www.medrxiv.org/content/early/2020/06/08/2020.05.13.20101253.full.pdf`, `doi:10.1101/2020.05.13.20101253`.
URL `https://www.medrxiv.org/content/early/2020/06/08/2020.05.13.20101253`

Table 1: Number of transitions between compartments in Figure 1 by age group and gender.

Males

Age	N	N_0	N_1	N_2	N_3	N_4	N_5	N_6
0-9	77	21	56	37	4	15	2	2
10-19	72	45	27	21	1	5	0	1
20-29	1560	1150	410	328	4	78	4	0
30-39	3467	2195	1272	856	12	404	6	6
40-49	4118	1729	2389	1310	28	1051	11	17
50-59	4387	1041	3346	1407	30	1909	6	24
60 +	6120	759	5361	1322	45	3994	6	39
Total	19801	6940	12861	5281	124	7456	35	89

Females

Age	N	N_0	N_1	N_2	N_3	N_4	N_5	N_6
0-9	52	22	30	17	6	7	3	3
10-19	112	73	39	26	1	12	1	0
20-29	1564	1270	294	239	6	49	6	0
30-39	2940	2264	676	501	12	163	7	5
40-49	3303	1936	1367	876	17	474	6	11
50-59	2819	948	1871	940	15	916	5	10
60 +	4095	567	3528	1091	26	2411	3	23
Total	14885	7080	7805	3690	83	4032	31	52

Table 2: Statistics associated with the transition of COVID-19 patients between compartments in Figure 1. $\mathbf{h}$ = distribution of patients arriving to hospitals. $\mathbf{c}$ = probability of requiring ICU among those that went into hospitalization; $\mathbf{d}$ = probability of dying among those that went into hospitalization; $\mathbf{d}_c$ = probability of dying among those that went into ICU.

Males

Age	h	c	d	$\mathbf{d}_c$
0-9	0.0027	0.0714	0.3036	0.5000
10-19	0.0013	0.0370	0.2222	1.0000
20-29	0.0198	0.0098	0.1902	0.0000
30-39	0.0616	0.0094	0.3223	0.5000
40-49	0.1156	0.0117	0.4470	0.6071
50-59	0.1619	0.0090	0.5777	0.8000
60 +	0.2594	0.0084	0.7523	0.8667

Females

Age	h	c	d	$\mathbf{d}_c$
0-9	0.0015	0.2000	0.3333	0.5000
10-19	0.0019	0.0256	0.3077	0.0000
20-29	0.0142	0.0204	0.1667	0.0000
30-39	0.0327	0.0178	0.2485	0.4167
40-49	0.0661	0.0124	0.3548	0.6471
50-59	0.0905	0.0080	0.4949	0.6667
60 +	0.1707	0.0074	0.6899	0.8846

Table 3: Normalized relative lethalities. "∘" and "⊘" are the Hadamard element-by-element multiplication and division respectively.

Males

Age	$\mathbf{f}_M$	$\mathbf{h}$	$\mathbf{d}$	$\boldsymbol{\theta}_M^*$ $\mathbf{h} \circ \mathbf{d} \oslash \mathbf{f}_M$
0-9	0.09871	0.00271	0.30357	0.00068
10-19	0.09976	0.00131	0.22222	0.00024
20-29	0.08130	0.01984	0.19024	0.00376
30-39	0.07203	0.06155	0.32233	0.02233
40-49	0.05566	0.11560	0.44705	0.07528
50-59	0.03851	0.16191	0.57770	0.19692
60 +	0.04218	0.25941	0.75229	0.37511

Females

Age	$\mathbf{f}_F$	$\mathbf{h}$	$\mathbf{d}$	$\boldsymbol{\theta}_F^*$ $\mathbf{h} \circ \mathbf{d} \oslash \mathbf{f}_F$
0-9	0.09577	0.0015	0.3333	0.00041
10-19	0.09824	0.0019	0.3077	0.00048
20-29	0.08709	0.0142	0.1667	0.00221
30-39	0.07908	0.0327	0.2485	0.00833
40-49	0.06096	0.0661	0.3548	0.03121
50-59	0.04225	0.0905	0.4949	0.08599
60 +	0.04846	0.1707	0.6899	0.19705

Table 4: Adjusted lethalities by age group and gender for the Mexican population(M=male, F=female). The *adjusted lethalities* are obtained from rescaling the *unadjusted lethalities* in Table 3, so that $\mathbf{f}'_M\boldsymbol{\theta}_M + \mathbf{f}'_F\boldsymbol{\theta}_F = 0.001$.

Age	$\boldsymbol{\theta}_M$	$\boldsymbol{\theta}_F$
0-9	0.00001	0.00001
10-19	0.00001	0.00001
20-29	0.00008	0.00005
30-39	0.00049	0.00018
40-49	0.00165	0.00068
50-59	0.00432	0.00188
60 +	0.00822	0.00432

Table 5: Population structure of Syrian refugees in Jordan.

Age	$\boldsymbol{f}_M$	$\boldsymbol{f}_F$
0-9	0.1934	0.1823
10-19	0.1190	0.1116
20-29	0.0714	0.0794
30-39	0.0574	0.0611
40-49	0.0337	0.0348
50-59	0.0127	0.0171
60 +	0.0110	0.0150
Total	0.4987	0.5013