Title page

<u>**Title:**</u> **Social-distancing effectiveness tracking of the COVID-19 hotspot Stockholm**

<u>**Author:**</u> Joachim Oberhammer, PhD (ORCID: 0000-0003-3339-9137)

<u>**Affiliation:**</u> Department of Intelligent Systems, KTH Royal Institute of Technology

<u>**Address:**</u> Department of Intelligent Systems, KTH Royal Institute of Technology, Malvinas väg 10, SE-
100 44 Stockholm, Sweden

<u>**Corresponding author:**</u> *Name:* Joachim Oberhammer; *Address:* Malvinas väg 10, SE-100 44
Stockholm, Sweden; *e-mail:* joachimo@kth.se; *phone number:* +46 737 65 23 68.

<u>**Implementation and data sharing:**</u> The source code of this model is enclosed in the appendix. The
source code and the database with model parameters will be made freely available on
https://github.com/JoachimKTH/COVID19_SEIRD_model.

<u>**Ethics approval:**</u> The study is based on publicly available data. Ethics approval was not required.

<u>**Acknowledgements:**</u> Peter L E Konings, Jasmine Gardner, Nele Brusselaers, Wouter Metsola van der
Wijngaart, are acknowledged for a first feedback on the simulation model.

Summary

Background The COVID-19 outbreak in Stockholm, Sweden, is characterized by a near-absence of governmental interventions and high fatalities in the care home population. This study analyses the outbreak and the social-distancing effectiveness timeline in the general population and the care homes.

Methods A novel distributed-compartmental, time-variant epidemiological model was designed specifically for COVID-19 transmission characteristics, featuring a/pre/symptomatic transmission, a non-linear hospital model, a weakly-coupled sub-model for the care-home population, and parametrized continuous social-distancing functions. The model parameters and the social-distancing timelines are determined by randomization and Monte-Carlo simulations analysing real-world data.

Findings Despite a high initial reproduction number (3·29) and the near-absence of governmental interventions, the model quantitated that the transmission rate in the general population was suppressed by 73%, and in the care homes by 79%. The measures in the care homes took effect 4·8 days delayed; and if applied 4 or 8 days earlier, the fatalities could have been reduced by 63·2% or 89·9%. The infected population is estimated to 16·2% (June 10). An expected underestimation of population immunity by antibody studies is confirmed. The infection fatality ratio extrapolates to 0·61% (peak: 1·34%). The model indicates a seasonal effect which effectively suppressed a new rise. An analysed large-scale public event had no large influence. The asymptomatic ratio was determined to 35%.

Interpretation The proposed model and methods have proven to analyse a COVID-19 outbreak and to re-construct the social-distancing behaviour with unprecedented accuracy, confirming even minor details indicated by mobility-data analysis, and are applicable to other regions and other emerging infectious diseases of similar transmission characteristics. The self-regulation of the population in Stockholm, influenced by advices by the authorities, was able to suppress a COVID-19 outbreak to a level far beyond that the stringency index of governmental interventions suggests. Proper timing of effective measures in the care homes is important to reduce fatalities.

Joachim Oberhammer, KTH Royal Institute of Technology, 2020-06-25

Manuscript

Introduction

Sweden, has never imposed a lockdown and only issued very few governmental restrictions. Instead,
the population has been briefed daily with advices by the Public Health Agency of Sweden
(Folkhälsomyndigheten, FHM). As of June 10, 4795 confirmed COVID-19 deaths have been reported
in Sweden (population: 10·3 million), by that time the fifth highest death rate per capita in the
world. Sweden has a 80·3% share of the deaths in North Europe, despite only having 38·2% of the
Nordic population.[1] The region Stockholm (23% of the population of Sweden) accounts for 45·4% of
all deaths in Sweden, and is characterized by a large proportion of deaths (43·3%) having occurred in
the care-homes population (only 0·6% of the total population).

Epidemiological modelling of the COVID-19 outbreak has been used for forecasting the healthcare
demand[2] and to develop containment strategies using non-pharmaceutical interventions (NPI).[3,4,5]
Modelling is also used to determine key parameters such as population immunity and infection
fatality rate (IFR), and to analyse the impact of imposing and revoking social-distancing
measures.[6,7,8,9,10] A well-established class of models is the Susceptible(S)-Exposed(E)-Infected(I)-
Recovered(R)-Deceased(D) compartmental model.[11,12,13] Due to short simulation times, in particular
deterministic SEIRD models are used for parameter estimation by randomization.[14] The impact of
interventions is usually modelled by discrete steps of the infectivity parameter.[2,6,8,15] It was found
that multiple inferring change points are more suitable for tracking the effect of specific
interventions.[7] A continuous modelling of the transmission-rate reduction is assumed to be more
suitable for countries like Sweden, without disruptive governmental interventions and adjusting to
the self-regulation behaviour of the population.

Methods

Epidemic simulation model

The deterministic, time-variant SEIRD model (Figure 1) features distributed compartments for more
realistic modelling of the different phases of the infection by time-distributed parameters
(infectivity, hospitalization and death rates, probabilities for PCR or antibody positive testing) as
shown in Figure 2:A. The model comprises two weakly coupled sub-models, for modelling the
different dynamics of the general population (GP) and the care homes (CH). The GP model features
an asymptomatic (individuals never developing clearly noticeable symptoms) and a symptomatic
branch, with the latter structured into pre-symptomatic and symptomatic compartments. Only
individuals in compartments I_1—I_4 are considered infectious in both branches, and only
compartments I_3-I_6 of the symptomatic branch have associated hospitalization and death rates.
Environmental transmission is not modelled due to its minor contribution for COVID-19.[16,17,18] Age-
groups, work/household environment, or individual susceptibility is not modelled.[19,20]

For the GP model, the force of infection λ_{GP} (in $\Delta S_{GP} = -\lambda_{GP} \cdot S_{GP}$) is as follows:

$$\lambda_{GP} = \beta_0(t)\left(\sum \beta_{a,i} \cdot I_{a,i} + \sum \beta_{s,i} \cdot I_{s,i}\right) \cdot \frac{1}{N_{GP}} \qquad (1)$$

(S_{GP} susceptible population; $I_{a,i}$ and $I_{s,i}$ individuals and $\beta_{a,i}$ and $\beta_{s,i}$ transmission rates in
asymptomatic/symptomatic branch compartments; $\beta_0(t)$ time-variant transmission-rate matrix
scaling factor: $\boldsymbol{\beta}_{GP} = \beta_0(t) \cdot [\beta_{a,i}; \beta_{s,i}]$; N_{GP} host population, Stockholm: 2374550).

The CH model is simplified (3-day latency; 3-day infectious phase I_1 including two presymptomatic
and a symptomatic day; 7-day sickness phase I_2 without transmission, i.e. patient isolation assumed),
without distinguishing asymptomatic/symptomatic branches, since CH population is negligible (0·6%
of GP) for a transmission event analysis. Besides the internal factor α_{INT}, the force of infection λ_{CH} s
weakly coupled to the GP model by an external factor α_{EXT} modelling imported transmissions by
health care workers and visitors:

$$\lambda_{CH} = \beta_{CH}(t) \left(\alpha_{EXT} \cdot \lambda_{GP} + \alpha_{INT} \cdot \frac{I_{CH}}{N_{CH}} \right) \qquad (2)$$

(I_{CH} infectious individuals in I_1; N_{CH} CH population size, 15000 estimated,[21] time-variant factor
$\beta_{CH}(t)$). Transmission-events feedback from the CH to the GP model is assumed neligible.

The hospital model comprises a primary ward, an intensive care, and a secondary general-ward
compartment for severely-sick, critically-sick, and patients recovering from intensive care,
respectively.

Time-variant and non-linear model parameters

The GP and CH transmission rates are scaled by the time-variant factors β_0 and β_{CH}, in the following
referred to as social-distancing effectiveness functions (SDEF), which track the social-distancing
behaviour and seasonal effects. All elements of the transmission-rate matrix are independently
controllable by time-dependent reduction factors, which allows to investigate the effect of different
advices to the population, from staying-home with symptoms to whole-household quarantine
(Figure 2:B).

Since the reported number of patients in ICU in Stockholm follows not exactly the shape of the
general ward, it was assumed that with time a growing proportion of patients can be kept in regular
ward, which was implemented by a three-parameter logistics function (details in Appendix). The
hospital death rate, in relation to the hospital occupancy, was found to decrease with time, which
might reflect increasing treatment experience or a reduced number of CH patients transferred to
hospital, which is also modelled by a parametrized logistics function. A non-linear hospital and ICU
death-rate scaling factor, depending on the hospital occupancy, is also implemented by a logistics
function.

SARS-CoV-2 infection parameters

The onset of symptoms after infection is set to 5 days (discretization of the 5·2 days incubation
time).[22,23] The distribution of the transmission rates in the two presymptomatic (I_1, I_2) and the two
symptomatic (I_3, I_4) infectious compartments (Figure 2:A) is following the distribution of
transmission-pair analysis so that the proportion of the presymptomatic transmission events in the
symptomatic branch is 44%.[24] The total contagious period (I_1-I_4) is 7 day.[22] Since literature reports no
significant difference in viral spread between asymptomatic and symptomatic individuals,[25,26] the
initial infectivity in the two branches is the same, followed by a 75% reduction after the first
symptomatic day in the symptomatic branch when symptomatic individuals are assumed to reduce
social interaction.

The proportion of cases moving into the asymptomatic branch is determined by analysing the
proportion of non-symptomatic (asymptomatic and presymptomatic) individuals testing PCR-

positive, taking the dynamics of the outbreak into account, and comparing to an average reported in
the literature (pre- and post-lockdown study in Vo, Italy;[25] Diamond Princess;[27] Heinsberg study;[28]
repatriating individuals to the UK;[2] average: 41·5%; Figure 3). The best matching asymptomatic-
branch proportion is 35%, assuming that R_{eff} in the studies is around 1·0.

Antibody development in the model is assumed to happens within 1 to 3 weeks after infection[29]
using a normal distribution ($\overline{\Delta t}$=14 days, 95%CI[4·5, 23·5]). The mapped probability distribution of
individuals to test PCR positive in different phases[30] is shown in Figure 2:A.

For the care-home model (CH), the outcome is decided in average seven days after onset of
symptoms (FHM: two thirds of all CH fatalities in the first week). The CH death rate is 39·75%
(reported CFR for CH, Stockholm, May 25). The hospital transfer rate is set to 5% (assumption based
on: only 10·7% of all care-home inhabitants in Sweden are younger than 75 years;[31] and FHM: only a
small fraction of care-home patients benefits from critical and intensive care), which is also reflected
by the high proportion of CH deaths and the low ICU death rate.

The ICU is modelled after a report by the Swedish Intensive-Care Register (SIR) of April 30
(median/mean/max time: 6·75/7·25/31·8 days; 29·75% death rate; time on ward before transfer to
ICU: 2·1 days). [32] The average times on regular (H_1) and on post-ICU (H_2) ward are 8 and 5 days,
respectively.[33] It is assumed that most people surviving ICU recover, therefore the H_2 death rate is
arbitrarily set to 5% of the H_1 death rate. The hospitalization-rate distribution, i.e. the transfer from
the GP symptomatic-branch compartments, was set so that the average time from onset of
symptoms to hospitalization is 8·16 days.[32]

Model fitting to real-world data: social-distancing tracking

The starting values and randomization confidence intervals of the unknown model parameters (SDEF
for GP and CH; hospitalization rate; ward-to-ICU transfer, hospital death-rate, and the hospital non-
linearity parameters) were determined by sensitivity analysis using multi-dimensional parameter
variation, with the GP SDEF initially modelled by a three-parameter logistics function. The
parameters were then refined by randomization with Monte-Carlo simulations. Minimum mean
square error functions were used to benchmark the model outcome to real-world data: the number
of patients in critical care and in intensive care, the number of date-adjusted GP and CH deaths
(daily situation reports, Region Stockholm), and the results of two PCR studies by FHM in April.[34,35]
The results of four weekly antibody studies by FHM (April to June),[36] where non-COVID-19 patients
seeking ambulatory care in Stockholm were sampled, was not included for model fitting since FHM
reported on May 26 a potential bias towards persons with a lower risk of infection for this type of
sampling, and since in particular mild/asymptomatic people do not have a robust antibody
response.[26,37]

The model is started by injecting infected cases to the latency compartment so that the daily new
infections are matching the reported confirmed cases from contact tracing, assuming a 5-day
reporting delay and a 5—10 times larger dark number range (Figure 3:B). The injections happen
primarily in Stockholm's winter school vacation week, reported by FHM to have been the major
source of imported cases.

For the determination of the SDEF function of the GP by Monte-Carlo simulations, 15 randomized
points with shape-preserving piecewise cubic interpolation were mapped on the initial logistics
function. The points were optimized by three subsequent Monte-Carlo batches with 1 million
simulation runs each, where the 95%CI calculated from the corrected sample standard deviation of
the 100 best functions of a previous batch were used as the 95%CI for the randomization in a

subsequent batch (convergence shown in Figure 4:A). The optimization of the hospital parameters
was done by a subsequent 1-million-run batch (Figure 4:C). For the CH model, the SDEF was
modelled as a three-parameter logistic function, determined with the internal and external
transmission-rate factors by a 1-million simulation-run batch. All parameters were fine-adjusted with
a combined 1-million run batch. More details on the model and data fitting in the Appendix.

Results

Mapping the outbreak in Stockholm

The SDEF determined for the GP with the advices to the population by FHM, governmental
measures, and key events of the outbreak mapped on the timeline, with comparison to Google's
mobility data analysis[38] and the Oxford Governmental Stringency Index,[15] is shown in Figure 5. From
an initial 1·0 (February 1), the function falls to a minimum of 0·29 (April 5). The major reduction
happens between March 6 and 20, when the majority of the governmental restrictions and advices
were issued. The slope and the increase after April 4 are in good agreement with the mobility data,
which recorded for Stockholm a major slope between March 7 and March 19 (-31% retail/recreation
mobility; -45% public-transport; -42% workplace-mobility) and shows a minimum for the Easter
weekend, for which media reported a traffic reduction by 90% as compared to previous years. A
more detailed mapping of the Easter week (April 5-15) was carried out by three additional
randomization points, while randomizing also all surrounding and following points in two 1-million
simulation-run batches (Figure 4:B), revealing a 3·9% improved error-function, a deeper SDEF
minima to 0·27, and a rise thereafter to the previous level (comparison of the two SDEF in Figure 5).
After a local peak (May 13; 0·34) the social distancing function falls again, contrary to the mobility
data. A time-shifting of the SDEF by ±2 days results in a large variation in peak hospital beds (-763 (-
29%) to 1517 (+39%); Figure 4:D), which confirms the previously reported sensitivity to the timing of
NPIs.[7] Verification tests of the SDEF and its independency on the starting point choice were done by
shifting the starting points ±1—±3 days (500000 Monte-Carlo runs each). The SDEF function was
confirmed by all verification functions, with the local minima (April 5) and the local maxima (May 13)
of the original SDEF matching the 95%CI (estimated from the 100 best functions each) of 5 out of 6
verification functions. The average deviation of the 50% transition was 0·32 days (95%CI[0·28, 0·36])
(details in Appendix).

The delay between the SDEFs of the CH to the GP was determined to 4·8 days (95%CI[4·67, 4·92]),
and the final level is lower for the CH (0·2048, 95%CI[0·2038, 0·2058]) than for the GP, indicating
that measures in the CH were taken later but were more effective. Figure 4:E shows that the CH
deaths could have been reduced by 63·2% (606 people, 95%CI[604, 608]) if the transmission-rate
reduction in the CHs would have been simultaneously with the GP, or, if additional four days earlier,
even by 89·9% (95%CI[89·6, 90·2]). If the imported infections by CH workers and visitors would have
been halved early on, the CH deaths could have been reduced by 44·6% (95%CI[44·4, 44·6]; Figure
4:F). With 43·3% of Stockholm's total deaths attributed to the care homes, these reductions would
have decreased the total fatalities by 15·9%, 43·7%, and 24%, respectively. The hospital death-rate
nonlinearity for peak day (April 6) was determined to +6·6% (95%CI[-5·5, 18·8]). By June 10, 16·2%
(95%CI[15·9, 16·48]) of the hospital patients had needed intensive care.

Figure 6 summarizes the model results, showing an excellent fit to the real-world data
($\bar{R}^2$=0·9890/0·9886/0·9995/0·9986; internally-studentized residuals shown). The basic reproduction
number R_0 was determined to 3·29, by replacing the starting sequence with an equivalent single

injection (Figure 6:F). R_{eff}, after falling below 1·0 on March 27 and reaching a minimum of 0·84 on
April 1, increases to 0·99 on May 7, after which it falls again. The confidence intervals given in the
following, for data referring to the total number of infected, are dominated not by the data-fitting
but by the PCR-study reference-points uncertainty. By June 10, 16·2% (95%CI[10·9, 22·8]) of the
population in Stockholm had been infected, increasing by 0·8% (95%CI[0·54, 1·12]) per day (Figure
6:D). The population immunity predicted by the model matches the results of FHM's antibody
studies at the upper quartile of their confidence-intervals (Figure 6:D), confirming the expected
underestimation (average difference: -1·57%, 95%CI[-1·06, -2·22]). The number of infectious people
has been reduced from its peak of 44140 (95%CI[29680, 62269]; April 1) to 15120 (95%CI[10167,
21330]), and the daily new infections are down from a peak of 6719 (95%CI=[4518, 9478], March 23)
to 1590 (95%CI[1069, 2243]), same level as of March 8 (Figure 6:E). The hospitalization rate was
2·27% (95%CI[1·52, 3·20]). The peak infection fatality rate (IFR) was 1·34% (95%CI[0·90, 1·89]), and
would extrapolate to 0·61% (95%CI[0·41, 0·86]; Figure 6:F). The asymptomatic ratio is determined to
35% by analysing the literature with this model.

Europe's last pre-corona large-scale public event: the Eurovision Song Contest final in Stockholm

The Eurovision Song Contest (ESC) Swedish Final (Stockholm, March 7, 27000 attendees), took place
at a time when Sweden was the only country in Europe which had not banned large-scale public
gatherings. Several COVID-19 super-spreader are known,[28,39] and can have severe impact an
outbreak, as in South Korea where 61·3% of all cases by March 15 were associated with a church
service on February 9.[40] The model determines the epidemiological parameters in Stockholm on
March 7 to: R_{eff}=2·76, 6674 infected, 2011 contagious, 1449 daily new infections, 23·3 contagious
people present at the event. The potential impact of the event was analysed by injecting an average
number *N* of transmissions per contagious person, scaling the model starting sequence to maintain
same outcome, and then removing the injected ESC incidences to determine the outcome without
the event. For *N*=5/10/20, the ESC contributes by 115/226/439 additional transmissions, and, if
cancelled or conducted without public, the peak hospital occupancy would have been reduced from
1084 to 1069/1055/1028 patients (-1·4/-2·7/-5·4%). A local raise in the number of hospital patients,
even for *N*=50 determined to only 4·6 patients, is not visible in the real-world data with a localized
(17 points) standard-deviation of 11·4 patients. If happened on February 2 or 3 weeks earlier, and a
single contagious individual present would have triggered 20 or 50 transmission-chains, the event
would have added 12·9% or 30·8% to the epidemic, respectively.

Discussion

The proposed distributed-compartmental epidemiological model, with COVID-19 specific infection
characteristics including a sub-model for the care homes, was able to track the Stockholm outbreak
and to quantitate the social-distancing effectiveness function with an unprecedented accuracy so
that even small details in the timeline, indicated by mobility data analysis, were resolved. The model
is constructed as a trade-off between model complexity, accuracy, and speed for suitability for large-
scale data fitting methods. This study has quantitated that by primarily relying on the advices by
authorities and on the self-regulation of the population, as in Sweden, it is possible to at least
temporarily suppress COVID-19 by 73%, far beyond the government stringency-index indication. The
determined social-distancing behaviour of the general population follows the mobility data analysis,
whereas the determined care homes transmission-rate reduction, delayed by several days, follows
the delayed government stringency index timeline. This indicates that the general-population self-
regulation effect was ahead of time of the effect of governmental measures. The decrease of the

social-distancing function after a local peak in May, contradicted by mobility data, indicates the
onset of a summer seasonal effect and has prevented R_{eff} from climbing again above 1·0. The model
outcome, in agreement with previous studies, is very sensitive to the timing of the social-distancing
function. An anticipated underestimation of the population immunity by antibody studies is
confirmed. This study has quantitated the importance of applying early care-home protection
measures. The analysis of Europe's last large-scale public event in pre-corona times has found that at
an already wide-spread community transmission such an event has a significantly lower impact as
compared to happening in a more sensitive earlier phase. The proposed model and methods are
applicable to other regions and other emerging infectious diseases with similar transmission
characteristics as COVID-19.

REFERENCES

Johns Hopkins Coronavirus Resource Center, data as of June 8. https://coronavirus.jhu.edu/
Ferguson N M et al. Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality and healthcare demand. https://doi.org/10.25561/77482. Published March 13, 2020.
Maier B F, Brockmann D. Effective containment explains subexponential growth in recent confirmed COVID-19 cases in China. *Science* 2020; **368/6492**: 742—746. DOI: 10.1126/science.abb4557.
Adamik B et al. Mitigation and herd immunity strategy for COVID-19 is likely to fail. *medRXiv* 2020; https://doi.org/10.1101/2020.03.25.20043109, published May 5, 2020.
Bittihn P, Golestanian R. Containment strategy for an epidemic based on fluctuations in the SIR model. *arXiv* 2003.08784, published April 13, 2020.
Salje H et al. Estimating the burden of SARS-CoV-2 in France. *Science* 2020; DOI: 10.1126/science.abc3517.
Dehning J et al. Inferring change points in the spread of COVID-19 reveals the effectiveness of interventions. *Science* 2020; DOI: 10.1126/science.abb9789.
Lai S. et al. Effect of non-pharmaceutical interventions to contain COVID-19 in China. *Nature* 2020; https://doi.org/10.1038/s41586-020-2293-x.
Flaxman S et al. Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe. *Nature* 2020; https://doi.org/10.1038/s41586-020-2405-7.
Hsiang S et al. The effect of large-scale anti-contagion policies on the COVID-19 pandemic. *Nature* 2020; https://doi.org/10.1038/s41586-020-2404-8.
Diekmann O, Heesterbeek H, Britton T. Mathematical Tools for Understanding Infectious Disease Dynamics, Princeton University Press, 2013.
Hethcote H. The mathematics of infectious diseases. *SIAM Rev.* 2000; **42**: 599—653. doi/10.1137/S0036144500371907.
Grassly N C, Fraser C. Mathematical models of infectious disease transmission. *Nat Rev Microbio* 2008; **6**: 477—87. doi:10.1038/nrmicro1845.
Manly B. Randomization, Bootstrap and Monte Carlo Methods in Biology. Chapman & Hall, London, 1997.
Oxford COVID-19 Government Response Tracker (OxCGRT) stringency index, https://github.com/OxCGRT/covid-policy-tracker (accessed June 15, 2020).
Döhla M et al. SARS-CoV-2 in environmental samples of quarantine households. *medRxiv* 2020; doi: https://doi.org/10.1101/2020.05.28.20114041. (posted June 2, 2020).
Ferretti L et al. Quantifying SARS-CoV-2 transmission suggests epidemic control with digital contact tracing. *Science* 2020; **368**: eabb6936 DOI: 10.1126/science.abb6936.
Van Doremalen M et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *N Engl J Med* 2020; **382**:1564-1567 DOI: 10.1056/NEJMc2004973.
Endo A et al. Estimating the overdispersion in COVID-19 transmission using outbreak sizes outside China. *Wellcome Open Research* 2020; **5**:67; last update April 28, 2020.
Gomes G M et al. Individual variation in susceptibility or exposure to SARS-CoV-2 lowers the herd immunity threshold. *medRxiv*, 2020; doi: https://doi.org/10.1101/2020.04.27.20081893 (posted May 21, 2020).

Joachim Oberhammer, KTH Royal Institute of Technology, 2020-06-25

Covid-19 på särskilda boenden för äldre samt hemtjänst. Stockholm Region County, Stockholm, 2020-04-22, updated 2020-06-02.

Lauer S A. The incubation period of 2019-nCoV from publicly reported confirmed cases: estimation and application," *Ann Intern Med* 2020; M20-0504. doi: 10.7326/M20-0504.

Li Q et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia. *N Engl J Med* 2020; **382**:1199-1207, DOI: 10.1056/NEJMoa2001316.

He X et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med* 2020; **26**: 672–675, https://doi.org/10.1038/s41591-020-0869-5.

Lavezzo E et al. Suppression of COVID-19 outbreak in the municipality of Vo, Italy. *medRxiv* 2020; https://doi.org/10.1101/2020.04.17.20053157 (posted April 18, 2020).

Long Q et al. Clinical and immunological assessment of asymptomatic SARS-CoV-2 infections. *Nat Med* 2020; https://doi.org/10.1038/s41591-020-0965-6

Mizumoto K, Kagaya K, Zarebski A, Chowell G, Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. *Euro Surveill.* 2020; **25(10)**: pii=2000180. https://doi.org/10.2807/1560-7917.

Streeck H et al. Infection fatality rate of SARS-CoV-2 infection in a German community with a super-spreading event," *medRxiv* 2020; doi: https://doi.org/10.1101/2020.05.04.20090076 (posted June 2, 2020).

Hang AT et al. A systematic review of antibody mediated immunity to coronaviruses: antibody kinetics, correlates of protection, and association of antibody responses with severity of disease. *medRxiv* 2020; https://doi.org/10.1101/2020.04.14.20065771 (posted April 17, 2020)

Kucirka L M, Lauer S A, Laeyendecker O, Boon D, Lessler J. Variation in False-Negative Rate of Reverse Transcriptase Polymerase Chain Reaction–Based SARS-CoV-2 Tests by Time Since Exposure. *Ann Intern Med* 2020; doi:10.7326/M20-1495.

Appendix to report: Statistik om socialinsatser till äldre 2019" by Socialstyrelsen. National Board of Health and Welfare Sweden, Stockholm, 2020-04-28.

Report: Covid-19 på Svenska intensivvårdsavdelningar. Svenska Intensivvårdsregistret (SIR, Swedish Intensive-care Register), Stockholm, 2020-04-30.

Report: Skattning av behov av slutenvårdplatser covid-19. Folkhälsomyndigheten (Health Agency of Sweden), Stockholm, 2020-05-13.

Report: Förekomsten av covid-19 i region Stockholm, 26 mars–3 april 2020. Folkhälsomyndigheten, Stockholm, 2020-04-17.

Press release: Nya resultat från undersökning av förekomsten av covid-19 i Sverige. Folkhälsomyndigheten, Stockholm, 2020-05-12.

Report: Påvisning av antikroppar efter genomgången covid-19 i blodprov från öppenvården (Delrapport 1). Folkhälsomyndigheten, Stockholm, 2020-06-18.

Liu T et al. Prevalence of IgG antibodies to SARS-CoV-2 in Wuhan - implications for the ability to produce long-lasting protective antibodies against SARS-CoV-2. *medRxiv* 2020; https://doi.org/10.1101/2020.06.13.20130252 (posted June 16, 2020).

Google COVID-19 Community Mobility Reports for Stockholm, combined raw data Feb. 15 to June 7 (accessed June 11, 2020).

Hamner L et al. High SARS-CoV-2 Attack Rate Following Exposure at a Choir Practice — Skagit County, Washington, March 2020. *Morbidity and Mortality Weekly Report* 2020; **69/19**, US Department of Health and Human Services/Centers for Disease Control and Prevention, May 15, 2020.

Kim S et al. Evaluation of COVID-19 epidemic outbreak caused by temporal contact-increase in South Korea," *Int J Inf Dis* 2020; **96**: 454—457.

Joachim Oberhammer, KTH Royal Institute of Technology, 2020-06-25

Figures

Figure 1. Flow chart of the deterministic, time-variant, distributed-compartmental model with a
detailed sub-model for the general population including a symptomatic and an asymptomatic
branch, a weakly-coupled sub-model for the care-home population, and a common hospital model.

Figure 2. Compartments with time-distributed infection parameters of the general population
model: A: asymptomatic and symptomatic branches with associated key parameters (delay times;
transmission rates; probability for positive RT PCR tests based on Mizumoto et al.[30]). B: Modelling of
the effect of advices by Swedish authorities, by reducing the infectivity of different compartments
(the last advice, whole household quarantine, was never issued in Sweden).

Joachim Oberhammer, KTH Royal Institute of Technology, 2020-06-25

Figure 3. A: Estimation of the asymptomatic branch ratio, by analysing the proportion of identified
non-symptomatic individuals to the literature (Vo,1 and Vo,2: pre- and post-lockdown study in the
Italian city of Vo;[25] DP: study from cruising ship *Diamond Princess*;[27] Heinsberg study, range as given
in the study for no symptoms and 1 out of 15 symptoms;[28] ICL: repatriating individuals to the UK[2]). B:
Starting sequence of the model, mapping the reported contact-tracing cases, multiplied by a dark-
figure factor of 5—10, and assuming a 5-day reporting delay, to the daily new infections in the
model.

Figure 4. Real-world data fitting results by Monte-Carlo simulations (A-C), and outcome analysis by parameter variation (D-F). A: social-distancing effectiveness function (SDEF) determined by three simulation batches with 1 million runs each (convergence of standard-deviation of selected points shown in the insert); B: results of 2 million runs with additional optimization points for refining the Easter-week minimum; C: optimization of ICU transfer rate; D: effect of time shift of SDEF of GP by ±2 days; E: time shift of SDEF of CH; F: effect of reduction of the infections by externals into the CH. (GP=general population; CH=care homes).

Joachim Oberhammer, KTH Royal Institute of Technology, 2020-06-25

Figure 5. Social distancing effectiveness function (SDEF) determined by the proposed model and Monte-Carlo simulations data-fitting, with advices and governmental restrictions mapped on the timeline (A=advice by Public Health Agency; GR=governmental restriction; CR=care-home restriction), including a SDEF variation with additional randomization points for resolving the Easter week minima indicated by the mobility data; middle: Google mobility data for Stockholm;[38] bottom: Oxford government stringency index for Sweden.[15]

Figure 6. Simulation results of the model mapped to the real-world data of the COVID-19 outbreak in Stockholm until June 10, 2020. (GP=general population, CH=care home population)