

A model of population dynamics with complex household structure and mobility: implications for transmission and control of communicable diseases

Rebecca H. Chisholm^{1,2}, Bradley Crammond², Yue Wu³, Asha C. Bowen^{3,4}, Patricia T. Campbell^{2,5}, Steven Y. C. Tong^{6,7,8}, Jodie McVernon^{2,5}, and Nicholas Geard^{5,9}

¹**Department of Mathematics and Statistics, La Trobe University, Bundoora VIC, Australia**

²**Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne VIC, Australia**

³**Wesfarmers Centre for Vaccines and Infectious Diseases, Telethon Kids Institute, Perth, WA, Australia**

⁴**Infectious Diseases Department, Perth Children's Hospital, Perth, WA, Australia**

⁵**Victorian Infectious Diseases Reference Laboratory Epidemiology, University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne VIC, Australia**

⁶**Doherty Department, University of Melbourne at the Peter Doherty Institute for Infection and Immunity, Melbourne VIC, Australia**

⁷**Victorian Infectious Diseases Service, The Royal Melbourne Hospital, Melbourne VIC, Australia**

⁸**Global and Tropical Health Division, Menzies School of Health Research, Charles Darwin University, Darwin NT, Australia**

⁹**School of Computing and Information Systems, Melbourne School of Engineering, The University of Melbourne, Melbourne VIC, Australia**

Corresponding author:
Rebecca H. Chisholm^{1,2}

Email address: r.chisholm@latrobe.edu.au

ABSTRACT

Households are known to be high-risk locations for the transmission of communicable diseases. Numerous modelling studies have demonstrated the important role of households in sustaining both communicable diseases outbreaks and endemic transmission, and as the focus for control efforts. However, these studies typically assume that households are associated with a single dwelling and have static membership. This assumption does not appropriately reflect households in some populations, such as those in remote Australian Indigenous communities, which can be distributed across more than one physical dwelling, leading to the occupancy of individual dwellings changing rapidly over time. In this study, we developed an individual-based model of an infectious disease outbreak in communities with demographic and household structure reflective of a remote Australian Indigenous community. We used the model to compare the dynamics of unmitigated outbreaks, and outbreaks constrained by a household-focused prophylaxis intervention, in communities exhibiting fluid versus stable dwelling occupancy. Our findings suggest that fluid dwelling occupancy can lead to larger and faster outbreaks, interfere with the effectiveness of household-focused interventions, and may contribute to the considerable burden of communicable diseases in communities exhibiting this type of structure.

44 INTRODUCTION

45 For many infectious diseases, it is assumed that the risk of transmission within households exceeds that
46 in the wider community due to the increased opportunity they provide for repeated and prolonged close
47 contact between the people who live in them (Goeyvaerts et al. (2018); Endo et al. (2019)). Due to
48 this increased risk, households are often the focus of infectious disease control strategies. For example,
49 household contacts of invasive Group A *Streptococcus* cases are estimated to have a 2000-fold increased
50 risk of developing the disease themselves (Oliver et al. (2019)). For Meningococcal disease, the equivalent
51 increase in risk is estimated to be between 500–800-times (De Wals et al. (1981)). As such, prophylaxis
52 of household contacts of cases for both of these infectious diseases is recommended to prevent further
53 spread (Oliver et al. (2019); De Wals et al. (1981)).

54 Much of our understanding of household structure, and hence its representation in mathematical
55 models of disease transmission comes from descriptions of census data. However, these descriptions
56 frequently rely on the notion of a stable ‘nuclear household’ (*i.e.*, comprising two parents and their
57 children). This notion may fail to capture the complexities and nuances of populations with very different
58 household structure and dynamics. In many settings, households differ in their composition—the people
59 they contain and their relationships to each other. Households may contain extended family members,
60 multiple family units, and unrelated people. For example, in Thailand, the proportion of households *not*
61 considered to be ‘nuclear’ is estimated at close to 50% (Dommaraju and Tan (2014)). In Vietnam, this
62 proportion is estimated to be one third, the majority of which are so-called ‘stem households’ which
63 include adults, their parents, and possibly their children (Dommaraju and Tan (2014)). The proportion of
64 households where there is co-residence of children under 15 years of age with older people over 60 differs
65 greatly throughout the world—in Senegal it is 37% , but just 0.2% in the Netherlands (United Nations,
66 Department of Economic and Social Affairs, Population Division (2017)).

67 Patterns of membership of households may also vary over time. People may spend time in multiple
68 housing units, blurring the relationship between the household as a unit of social organisation and the
69 physical dwelling (Smith (1992)). For example, a study in Northern Malawi found that households were
70 distributed across between one and twelve dwellings (mean of 1.7 dwellings per household), with between
71 one and nineteen persons occupying each dwelling per night (mean 3.0) (Fine et al. (1997)). Australian
72 Indigenous households can also be distributed across more than one physical dwelling. One study of the
73 occupancy of a single dwelling in an Australian Indigenous community over time revealed that in addition
74 to core residents, the dwelling was also regularly occupied (although less frequently) by an extended
75 household comprising other relatives and close associates (Musharbash (2008)). Over the course of
76 just over a year, more than 100 unique people were observed to stay at the dwelling for at least one night.
77 The flux in occupancy of individual dwellings potentially results in an increased risk of introduction
78 into dwellings, and a continually changing population at risk of household-level infection transmission,
79 particularly if there is also high rates of overcrowding (Morphy (2007)). The implications of this type of
80 fluid dwelling occupancy on infectious disease transmission and control are unknown.

81 In this study, we introduce an individual-based model incorporating a more flexible representation of
82 household membership distributed across multiple dwellings. We calibrate our model to an Australian
83 Indigenous community to capture observed demographic, household and mobility characteristics of the
84 population. We then use the model to simulate unmitigated and mitigated (through a household-focused
85 prophylaxis intervention) outbreaks of an influenza-like illness where the risk of infection transmission
86 between contacts residing in the same dwelling is greater than those in the wider community. Model
87 outputs are compared to those from a more traditional household model assuming stable dwelling
88 occupancy, to quantify the impact of distributed households and fluid dwelling occupancy on the dynamics
89 and control of communicable diseases outbreaks.

90 MATERIALS AND METHODS

91 Individual-based model of population and infection dynamics

92 *Population structure.*

93 Our individual-based model tracks the age and current residence of individuals in a community over time.
94 The community is comprised of N individuals and H physical dwellings. An individual’s age is updated
95 each day, and individuals are lost due to natural death at an age-dependent rate. When a death occurs, a
96 new individual aged zero is born into the population so that the population size N is constant.

97 **Population mobility.**

98 The mobility model is based on the Australian Indigenous mobility framework proposed in (Musharbash
99 (2008)). This study tracked the number of people that stayed at least one night in a particular dwelling
100 in the remote Australian Aboriginal community, Yuendumu during the 221 nights for which this data
101 was recorded (these 221 nights were not continuous, but occurred during the 467 day study period). The
102 cumulative number of nights stayed by each person was reported. The authors identified four types of
103 residents, based on the amount of time spent in the dwelling: so called core residents, who were present
104 60-100% of the time, regular residents, who were present 20-34% of the time, other residents who stayed
105 less frequently on an on-and-off basis and were present 4-16% of the time, and many sporadic short-term
106 visitors who stayed for between 1–6 nights.

107 In line with this framework, individuals in our model are assigned (uniformly at random) to a fixed set
108 of three dwellings within their community, which we assume remain fixed over the time frames we are
109 considering in this study (less than one year), and which we refer to as their *dwelling set*. These dwellings
110 represent their core residence, where they spend most nights, a regularly-visited residence, and an on-off
111 residence, where they stay less frequently on an on-and-off basis (see Figure 1). We refer to individuals
112 who have the same core residence as a *core household*, while an individual's *extended household* consists
113 of all core, regular and on-off residents of their dwelling set.

114 An individual's *current residence* can change due to population mobility. We have two types of
115 mobility in the model: between-dwelling mobility (intra-community); and between-community mobility
116 (inter-community).

117 Within the community, each day, an individual's current residence is chosen to be either their core,
118 regular or on/off residence with respective probabilities p_c , p_r and p_o , where $p_c > p_r > p_o$. There is also
119 a small probability p_s , where $p_s < p_o$ and $\sum_i p_i = 1$, that their current residence will be a dwelling chosen
120 uniformly at random.

121 To capture inter-community mobility, each day, A individuals (where A is a Poisson distributed random
122 variable with mean αN , and α is the mean per capita migration rate) are chosen uniformly at random to be
123 replaced by immigrants (thus ensuring that community size remains constant). Immigrants are assumed to
124 have a similar age to individuals in the population. This is implemented by specifying that an immigrant
125 will have the same age as an individual selected uniformly at random from the population. Immigrants are
126 assigned a current residence chosen uniformly at random from all dwellings in the community, as are the
127 dwellings which make up their dwelling set (their core, regular and on/off residences). All immigrants are
128 assumed to be susceptible to infection.

129 In an extended version of the model (described in the supplementary material) we consider an
130 additional type of mobility – the regular influx of temporary visitors into the community due to two types
131 of events: funerals (which take place after the death of a community resident) and reoccurring events,
132 such as sporting matches or festivals. This type of mobility leads to temporary changes in the community
133 size.

134 **Infection dynamics.**

135 We use an *SEIR* (Susceptible-Exposed-Infectious-Recovered) transmission model to simulate an outbreak
136 of an influenza-like illness in the community (see Figure 1c). Individuals in the community are classified
137 according to their infection status: they are either susceptible to infection (i.e., they can acquire the
138 infection from an infectious contact), exposed (i.e., they have a latent infection and are not infectious),
139 infectious (i.e., they have an active infection and can infect susceptible contacts), or recovered (i.e., they
140 have recovered from the infection and are protected from re-infection). This infection status can change
141 over time due to a transmission event, the progression to infectiousness, or due to the clearance of an
142 infection (detailed below).

143 Each day, individuals with the same current residence make contact with each other (we refer to
144 these contacts as household contacts), and we simulate daily contacts that occur between individuals in
145 the wider community (i.e., between individuals with different current residences, which we refer to as
146 community contacts). These community contacts occur at age-dependent rates $c_{u,v}$, where $c_{u,v}$ is the daily
147 rate of contact of an individual in age-category u with individuals in age category v . Community contacts
148 are chosen uniformly at random from the pool of individuals in the relevant age category.

149 If a susceptible person makes contact with an infected individual with a different current residence, the
150 susceptible person becomes infected (entering the exposed class) with probability q . Household contacts
151 (between individuals with the same current residence) are assumed to be more intense than community




Figure 1. Population, mobility and infection model. (a) Intra- and inter-community mobility results in the movement of infectious (I) and non-infectious individuals (S, E, R) within and between communities. (b) Individuals identify with three dwellings in their community: their core residence, where they spend most nights, a regularly-visited residence, and an on-off residence, where they stay less frequently on an on-and-off basis. Individuals may also sporadically stay in a dwelling chosen uniformly at random from all dwellings in the community. (c) $SEIR$ outbreak model. Individuals are born susceptible to infection, can become exposed to infection through contact with an infectious person, before progressing to infectiousness, and then become immune to re-infection following recovery from infection.

152 contacts. We translate this increased intensity into a probability of transmission per contact that is higher
 153 by a factor of $\hat{q} \geq 1$ for these household contacts, compared to community contacts. The duration of latent
 154 and active infection are assumed to be exponentially distributed with respective mean duration of $1/\sigma$
 155 and $1/\gamma$. Once an individual clears an infection (and enters the recovered class), they can no longer be
 156 infected.

157 Simulated outbreaks were seeded with one infectious individual (chosen uniformly at random), and
 158 with the rest of the population in the susceptible class, and were run until the end of the outbreak (when
 159 there were zero infected individuals left in the population).

160 **Dynamics of a core household-focused prophylaxis intervention.**

161 Finally, we also consider outbreaks where a prophylaxis intervention is administered to the core household
 162 members of an infectious person.

163 We assume that this prophylaxis intervention is administered once an infected person enters the
 164 infectious state. We do not explicitly model the onset of symptoms in the model. However, if symptom
 165 onset corresponds to the onset of infectiousness, then the timing of this intervention corresponds to there
 166 being a 0-day delay in the core household receiving prophylaxis from symptom onset.

167 We consider outbreak scenarios where the intervention is 100% and 50% effective at protecting the
 168 core household from contracting the infection, if they hadn't been previously infected and/or recovered.

169 **Model parameterisation and description of outbreak simulation scenarios**

170 We parameterised the model to be consistent with demography and mobility in a remote Indigenous
 171 Australian community.

172 For all outbreak scenarios, we considered two community sizes $N \in \{2500, 500\}$ with respective
 173 numbers of houses $H \in \{358, 80\}$, that are reflective of a large and small-medium community in the
 174 NT (Australian Bureau of Statistics (2018a)). With these values, the mean number of core residents per
 175 house was 7 and 6.3, respectively. We also explored scenarios in populations with lower numbers of core
 176 residents per house (i.e., with the number of houses set to $H \in \{833, 160\}$, so that the mean number of
 177 core residents per house was 3 and 3.1, respectively), to explore the impact of fluid dwelling occupancy in
 178 less-crowded communities.

179 Mortality rates and the initial age distribution were taken from the most recent census data of
 180 Aboriginal and Torres Strait Islander Australians in the Northern Territory (NT), Australia (Australian
 181 Bureau of Statistics (2018b)) (see Figure 2(a)). We set the intra-community mobility probabilities to be

182 $\{p_c, p_r, p_o, p_s\} = \{0.66, 0.23, 0.09, 0.02\}$ based on data (summarised above) of house occupancy over
183 time in a single household from the remote Australian Aboriginal mobility study in Yuendumu (Mushar-
184 bash (2008)). Inter-community mobility patterns are not described in this setting although, anecdotally,
185 Aboriginal Australians are described as having a higher than average rate of mobility compared to non-
186 Aboriginal Australians (Morphy (2007)). We set the per capita expected migration rate α to be between
187 $[0.002, 0.004]$ per day, which corresponds to, on average, between $[5, 10]$ migration events per week when
188 the population size $N = 2500$.

189 To date, there have been no studies measuring contact patterns outside of households in remote
190 Indigenous Australian communities. Age-dependent contact data that differentiates between household
191 and non-household contacts is available for rural populations in Kenya (Kiti et al. (2014)), and we used
192 this to specify the age-dependent community contact rates $c_{u,v}$ in our model.

193 Infection parameters were chosen to be consistent with influenza-like illness: the mean duration of
194 latency $1/\sigma$ was set to be between $[1, 3]$ days, as was the duration of infectiousness $1/\gamma$. We do not have data
195 to inform the within-house transmission factor \hat{q} . Therefore, we considered two different scenarios: a high
196 household-infection risk scenario where \hat{q} is set between $[3, 5]$, and a medium household-infection risk
197 scenario where \hat{q} is set between $[1, 3]$. We set the community transmission probability q to be between
198 $[0.002, 0.004]$ which, in the high household-infection risk scenario, led to outbreaks where greater than
199 50% of the population became infected, when the outbreaks took off. Results are also provided in the
200 supplementary material where we assumed a higher transmissibility of the infection with q set to be
201 between $[0.004, 0.006]$.

202 To account for uncertainty in the model parameters, for each population and infection scenario consid-
203 ered, we generated 1000 samples from the parameter space using Latin Hypercube Sampling (Blower and
204 Dowlatabadi (1994)). The parameters α , $1/\sigma$, $1/\gamma$, \hat{q} , and q were sampled from uniform distributions
205 with upper and lower bounds as described above. All other parameters were held constant.

206 All outbreak scenarios were re-run in a population assuming stable dwelling occupancy (i.e., with
207 the intra-community mobility probabilities set to $\{p_c, p_r, p_o, p_s\} = \{1, 0, 0, 0\}$), and again in populations
208 where the core household-focused prophylaxis intervention, described above, was implemented, to
209 understand the implications of fluid dwelling occupancy on outbreak dynamics and control.

210 The model is implemented in MATLAB and the code needed to regenerate all figures and tables is avail-
211 able at <https://github.com/rhchisholm/transmission-complex-households>.

212 RESULTS

213 Population mobility model leads to dwelling occupancy distributions consistent with 214 observations in an Indigenous Australian community

215 To determine whether our model leads to dwelling occupancy distributions that are consistent with
216 that observed in Yuendumu, we first set up our model population to have similar characteristics to this
217 community. According to the most recent census data, Yuendumu has a population size $N = 759$, an
218 average household size of 4.3 (which we used to estimate the number of dwellings $H = 176$), and people
219 have a median age of 28 (Australian Bureau of Statistics (2016)). We then simulated population and
220 mobility dynamics using our model, collecting occupancy data from all dwellings over 221 days, and
221 compared this to the occupancy distribution from the Yuendumu study (Musharbash (2008)). A sample
222 of these outputs shown in Figure 2 reveals clear steps corresponding to different resident types, as was
223 observed in the original study, with the width of these steps reflecting the number of residents of each
224 type (core, regular, on/off and sporadic visitors) associated with the dwellings, which differed between
225 dwellings. Equivalent model outputs from the extended model with event migration were also consistent
226 with the observed data (Figure S1), indicating that both models capture, at least in a qualitative way, the
227 fluid dwelling occupancy observed in an Indigenous Australian community.

228 Fluid dwelling occupancy leads to faster, and more-intense outbreaks

229 We then used our model to simulate outbreaks of an influenza-like illness in communities with a population
230 size and core dwelling size distributions reflective of large and small-medium Indigenous communities in
231 the NT, Australia (the population sizes were 2500 and 500, and the mean number of core residents per
232 dwelling was 7 and 6.3, respectively) (Australian Bureau of Statistics (2018a)). Key model outputs are
233 shown in the main manuscript for large communities, and the analogous outputs for the small-medium
234 communities are provided in the electronic supplementary material. All outbreaks were seeded with a




Figure 2. Population and mobility model outputs for a community with similar characteristics to Yuendumu (NT, Australia). (a) Age distribution of the population in years; (b) Core, regular and on/off dwelling size distributions; and (c) dwelling occupancy over 221 days for 12 dwellings in the model. Each bar represents a unique individual who stayed at the dwelling for at least one night. The height of the bar represents the cumulative number of nights the individual was present. Individuals are shown in order of decreasing occupancy. The title of each plot shows the number of {core, regular, on/off} residents for that dwelling at the end of the simulation.

235 single infectious person, and model outputs (summarised in Tables 1 and S1) were compared to those
 236 from equivalent simulations in communities assuming stable dwelling occupancy.

237 We found that infection introductions were just as likely to lead to outbreaks in communities with
 238 fluid dwelling occupancy as they were in communities with stable dwelling occupancy. However, for
 239 outbreaks which did take off, those which occurred in communities with fluid dwelling occupancy were
 240 consistently more intense than those in communities with stable dwelling occupancy (Figure 3(a)). That
 241 is, in communities with fluid dwelling occupancy, outbreaks were typically larger in overall size (the total
 242 number of people infected during outbreaks), had a higher, and earlier peak (the time in the outbreak
 243 when the number of infectious people was highest), and had a shorter duration than those in communities
 244 with stable dwelling occupancy.

245 These differences in outbreak intensity were less noticeable when we considered outbreak scenarios
 246 (i) with a lower increased risk of infection transmission between contacts residing in the same dwelling
 247 compared to those in the wider community (Figure 3(b)); (ii) in communities with a lower extent of
 248 household overcrowding (Figure 3(c)); and/or (iii) with a more transmissible pathogen (Figure S2(a),(c)).
 249 These results were robust to the sizes of the communities considered (Figure S3), and to the inclusion of
 250 event-based mobility in the fluid dwelling-occupancy model (Figures S4 and S5).

Table 1. Statistics from model scenarios of unmitigated outbreaks in communities of size 2500, including percentage of simulations that led to an outbreak (take off %), and the median (50% CIs) of the outbreak duration and final size.

Scenario	Dwelling occupancy	Take off (%)	Duration (days)	Final size
Baseline	fluid	52.8	112 (91,136)	1760 (1304,2040)
Baseline	stable	52.3	129 (105,160)	1494 (1073,1789)
Lower \hat{q}	fluid	37.7	139 (104,179)	1051 (507,1414)
Lower \hat{q}	stable	34.3	141 (112,183)	995 (473,1319)
Less crowded	fluid	33.8	147 (117,184)	910 (446,1276)
Less crowded	stable	33	140 (88,181)	725 (118,1070)




Figure 3. The impact of fluid dwelling occupancy on influenza-like outbreaks in a population of size $N = 2500$ assuming (a) a high-level; (b) a medium-level, of increased risk of transmission from household contacts compared to community contacts; and (c) less crowding in dwellings. The lines and shading show the median and interquartile ranges of the population prevalence of infection over time when there is fluid dwelling occupancy (red solid line, red shading); compared to when there is stable dwelling occupancy (black dashed line and grey shading).

251 **Higher outbreak intensity is driven by an increased number of unique, and higher-risk,**
 252 **household contacts.**

253 To understand why communities with fluid dwelling occupancy experienced more intense outbreaks, we
 254 inspected the number and types of contacts of infectious people over the course of outbreaks (Figure 4
 255 and S6). We found that the greatest relative difference between the contact patterns of infectious people
 256 between the fluid occupancy model (with and without event-based migration) and stable occupancy
 257 model was in relation to the number of unique individuals they contacted within dwellings, which was
 258 much greater in communities assuming fluid dwelling occupancy compared to stable dwelling occupancy,
 259 independent of the community size considered. Neither the number of unique community contacts, nor
 260 the total number of contacts of infectious people within or outside of dwellings were as affected by the
 261 type of dwelling occupancy model assumed, which suggests that the higher outbreak intensity observed
 262 in model communities with fluid versus stable dwelling occupancy was driven by the increased number of
 263 unique, and higher-risk, household contacts.

264 **Fluid dwelling occupancy decreases the impact of a core household-focused prophylaxis**
 265 **intervention**

266 Finally, we explored the effect of fluid dwelling occupancy on the impact of a core household-focused
 267 prophylaxis intervention that could be implemented during outbreaks. This intervention was administered
 268 to an infected person's core household at the time of infectiousness onset, which protected the core
 269 household from contracting the infection, if they hadn't previously been infected and/or recovered. We
 270 found that, while this intervention did reduce outbreak size (Table 2), it had less impact in communities




Figure 4. The impact of fluid dwelling occupancy on the distribution of the number of contacts of infectious people during outbreaks in a population of size $N = 2500$ assuming (a) a high-level; (b) a medium-level, of increased risk of transmission from household contacts compared to community contacts; and (c) less crowding in dwellings. Each disk with error bars shows the mean of means \pm one pooled standard deviation of either the total number of contacts, or the total number of unique individuals contacted (as indicated in the plots) during the infectious period of infected individuals, when there is fluid dwelling occupancy (red, solid lines); fluid dwelling occupancy with event-based migration (blue, dotted lines); compared to when there is stable dwelling occupancy (black, dash-dot lines).

271 with fluid dwelling occupancy, compared to those with stable dwelling occupancy (Figures 5(a),(b),(d),(e)
 272 and S7(a),(d)). It also had less impact when implemented in communities with less crowding in dwellings,
 273 likely because the average population coverage of the intervention per treated core household was reduced
 274 (Figures 5(c),(f) and S7(b),(c),(e),(f)). These differences were more pronounced when the intervention
 275 was assumed to be 100% effective at protected the core household of a case from contracting the infection,
 276 compared to when we assumed it was 50% effective. Again, these results were robust to the sizes
 277 of the communities considered (Figure S8), and to the inclusion of event-based mobility in the fluid-
 278 household membership model (Figure S9–S10). In some scenarios where we assumed the intervention
 279 was 50% effective, the duration of the outbreak was increased by the intervention, although the total size
 280 was reduced (Figures 5(d), S7(d),(f), S8(j),(l), S9(d),(j) and S10(j)). This occurred more frequently in
 281 communities of size 2500, and when we assumed a higher relative contact intensity for household versus
 282 community contacts.

Table 2. Statistics from model scenarios of mitigated outbreaks in communities of size 2500, including the percentage reduction in the median value of the outbreak duration and final size, compared to the equivalent unmitigated scenarios, for 100% and 50% effective interventions.

Scenario	Dwelling occupancy	Median duration reduction (%) with intervention effect		Median final size reduction (%) with intervention effect	
		100%	50%	100%	50%
Baseline	fluid	29	-4	87	48
Baseline	stable	55	-2	97	49
Lower \hat{q}	fluid	45	11	87	61
Lower \hat{q}	stable	59	10	94	61
Less crowded	fluid	35	10	77	39
Less crowded	stable	37	13	85	54




Figure 5. The impact of fluid dwelling occupancy on the effect of a household-focused prophylaxis intervention that is (a–c) 100% effective; and (d–f) 50% effective in a population of size $N = 2500$ assuming (a,d) a high-level; and (b,e) a medium-level, of increased risk of transmission from household contacts compared to community contacts; and (c,f) less crowding in dwellings. The lines and shading show the median and interquartile ranges of the population prevalence of infection over time when there is fluid dwelling occupancy (unmitigated outbreak: red solid line and shading; with intervention: magenta dash-dot line and shading); compared to when there is stable dwelling occupancy (unmitigated outbreak: black dashed line and shading; with intervention: blue dotted line and shading).

283 DISCUSSION

284 It is generally assumed that households are associated with a single physical dwelling which is considered
 285 to be a high-risk location for the transmission of many infectious diseases. However, the assumption of a
 286 one-to-one correspondence between households and dwellings does not appropriately reflect households
 287 in some populations, such as the Australian Indigenous communities considered here, where households
 288 can extend across multiple physical dwellings leading to fluid groups of people occupying individual
 289 dwellings. In this study, we showed that communities made up of such extended households have the
 290 potential to experience larger and more intense outbreaks of infectious diseases spread by close contact,
 291 particularly when there are high levels of household crowding.

292 **Outbreaks spread rapidly in communities characterised by fluid dwelling occupancy due** 293 **to close mixing in and between interconnected households.**

294 In our model with fluid dwelling occupancy, the extended household of an individual does not, in general,
 295 overlap with that of others in their extended household. Thus, multiple extended households can be
 296 connected via shared members, leading to large pools of individuals at greater risk of quickly contracting
 297 an infection and spreading it to other extended households, and to faster and larger outbreaks. For
 298 pathogens where there is even greater relative risk of infection transmission between household contacts
 299 compared to between community contacts, the risk of onward transmission beyond an extended household
 300 is amplified further, leading to even larger discrepancies in outbreak intensity between model communities
 301 characterised by fluid versus stable dwelling occupancy.

302 These reflections also help to explain why smaller discrepancies in outbreak intensity were observed
 303 between communities with different dwelling occupancy models when either household crowding was
 304 reduced or a more-transmissible pathogen was considered. In both of these scenarios, the lower-risk
 305 community contacts contributed much more to widespread transmission because, in the first scenario,
 306 the number of household contacts was significantly reduced, and in the latter scenario, the overall risk of

307 infection transmission from the more-frequent community contacts had increased.

308 **Implications for infectious disease control**

309 Our findings contribute to the evidence base that supports reducing household overcrowding as an effective
310 strategy to decrease the risk of severe outbreaks in populations with fluid dwelling occupancy (World
311 Health Organization (2018)). They also highlight the limitations of household-focused interventions in
312 these settings, which suggests that such interventions should be scaled up to reflect the interconnectedness
313 of households. Our finding also suggest that an intervention that reduces the number of unique household
314 contacts during an outbreak by, for example, limiting the amount of movement between dwellings, may
315 reduce outbreak intensity for certain pathogens. Further work could explore the effectiveness of such
316 interventions.

317 **Model limitations and further work**

318 The mechanistic model of intra-community mobility proposed in this study was based on data describing
319 the cumulative occupancy over a period of time of a single dwelling in one remote Australian Indigenous
320 community (Musharbash (2008)). While the occupancy distributions generated from our model do
321 resemble this data, it remains an open question whether our model is an accurate reflection of the
322 mechanisms which led to these cumulative patterns. It is also an open question how generalisable this
323 model is to other dwellings in the same community in which the data was collected, to other remote
324 Australian Indigenous communities, and to other population settings where households are distributed
325 across multiple dwellings. Longitudinal data of intra-community mobility from multiple dwellings, in
326 multiple communities, and from different populations could help to inform these open questions.

327 Nevertheless, our study highlights why accounting for correct household structure and dynamics in
328 models of infectious diseases that spread through close contacts can be important in outbreak scenario
329 analysis. Pathogens which spread via other means, for example, those which spread via vectors or which
330 are sexually transmitted, may not necessarily have different outbreak dynamics in communities with fluid
331 versus stable dwelling occupancy. Further work could explore the implications of the dwelling occupancy
332 model for such pathogens, as well as those which are endemic in populations.

333 **ACKNOWLEDGMENTS**

334 We thank all participants of the public health consultation workshop: Community-guided and evidence-
335 based strategies to reduce the burden of *Strep A* in Australian Indigenous communities, held in Darwin
336 Australia, February 2020.

337 **REFERENCES**

- 338 Australian Bureau of Statistics (2016). Census of population and housing: General Community Profile,
339 Yuendumu (SSC70301), 'Table G02: Selected medians and averages', cat. no 2001.0.
- 340 Australian Bureau of Statistics (2018a). Census of Population and Housing: Characteristics of Aboriginal
341 and Torres Strait Islander Australians, 2016, 'East Arnhem 2016 census story'. Technical Report cat
342 no. 2076.0, ABS, Canberra.
- 343 Australian Bureau of Statistics (2018b). Life Tables for Aboriginal and Torres Strait Islander Australians,
344 2015-2017, 'Table 1.6: Life tables for Aboriginal and Torres Strait Islander Australians, Northern
345 Territory-2015-2017', data cube: Excel spreadsheet, cat. no. 3302.0.55.003.
- 346 Blower, S. M. and Dowlatabadi, H. (1994). Sensitivity and uncertainty analysis of complex models
347 of disease transmission: an HIV model, as an example. *International Statistical Review/Revue
348 Internationale de Statistique*, pages 229–243.
- 349 De Wals, P., Hertoghe, L., Borlee-Grimee, I., De Maeyer-Cleempoel, S., Reginster-Haneuse, G., Dachy,
350 A., Bouckaert, A., and Lechat, M. (1981). Meningococcal disease in Belgium. Secondary attack rate
351 among household, day-care nursery and pre-elementary school contacts. *Journal of Infection*, 3:53–61.
- 352 Dommaraju, P. and Tan, J. (2014). Households in Contemporary Southeast Asia. *Journal of Comparative
353 Family Studies*, 45(4):559–580.
- 354 Endo, A., Uchida, M., Kucharski, A. J., and Funk, S. (2019). Fine-scale family structure shapes influenza
355 transmission risk in households: Insights from primary schools in Matsumoto city, 2014/15. *PLoS
356 Computational Biology*, 15(12).

- 357 Fine, P. E., Steme, J., Pönnighaus, J. a., Bliss, L., Saul, J., Chihana, A., Munthali, M., and Wamdorff,
358 D. (1997). Household and dwelling contact as risk factors for leprosy in northern Malawi. *American*
359 *Journal of Epidemiology*, 146(1):91–102.
- 360 Goeyvaerts, N., Santermans, E., Potter, G., Torneri, A., Van Kerckhove, K., Willem, L., Aerts, M., Beutels,
361 P., and Hens, N. (2018). Household members do not contact each other at random: implications for
362 infectious disease modelling. *Proceedings of the Royal Society B*, 285(1893):20182201.
- 363 Kiti, M. C., Kinyanjui, T. M., Koech, D. C., Munywoki, P. K., Medley, G. F., and Nokes, D. J. (2014).
364 Quantifying age-related rates of social contact using diaries in a rural coastal population of Kenya.
365 *PloS One*, 9(8):e104786.
- 366 Morphy, F. (2007). Uncontained subjects: Population and household in remote Aboriginal Australia.
367 *Journal of Population Research*, 24(2):163–184.
- 368 Musharbash, Y. (2008). *Yuendumu everyday: contemporary life in remote Aboriginal Australia*. Aboriginal
369 Studies Press.
- 370 Oliver, J., Thielemans, E., McMinn, A., Baker, C., Britton, P. N., Clark, J. E., Marshall, H. S., Blyth, C. C.,
371 Francis, J., Buttery, J., et al. (2019). Invasive group A Streptococcus disease in Australian children:
372 2016 to 2018—a descriptive cohort study. *BMC public health*, 19(1):1–10.
- 373 Smith, D. E. (1992). *A national survey of indigenous Australians: options and implications*, volume 3 of
374 *CAEPR Research Monograph*, chapter The cultural appropriateness of existing survey questions and
375 concepts. Centre for Applied Economic Policy Research.
- 376 United Nations, Department of Economic and Social Affairs, Population Division (2017). Household
377 Size and Composition around the World 2017 – Data Booklet (ST/ESA/SER. A/405).
- 378 World Health Organization (2018). *WHO housing and health guidelines*, chapter 3: Household crowding.
379 Geneva.