

Covid-19 Risk Among Airline Passengers: Should the Middle Seat Stay Empty?

Abstract

We use recent data and research results to approximate the probability that an air traveler in coach will contract Covid-19 on a US domestic flight two hours long, both when all coach seats are full and when all but middle seats are full. The point estimates we reach based on data from late June 2020 are 1 in 4,300 for full flights and 1 in 7,700 when middle seats are kept empty. These estimates are subject to both quantifiable and nonquantifiable sources of uncertainty, and sustain known margins of error of a factor about 2.5. However, because uncertainties in key parameters affect both risk estimates the same way, they leave the relative risk ratio for “fill all seats” compared to “middle seat open” close to 1.8 (i.e., close to $1/4,300 / (1/7,700)$). We estimate the mortality risks caused by Covid-19 infections contracted on airplanes, taking into account that infected passengers can in turn infect others. The point estimates—which use 2019 data about the percentage of seats actually occupied on US flights—range from one death per 400,000 passengers to one death per 600,000. These death-risk levels are considerably higher than those associated with plane crashes but comparable to those arising from two hours of everyday activities during the pandemic.

Arnold Barnett
Sloan School of Management
MIT
Cambridge, MA 02142

abarnett@mit.edu

617 686-1485

E62-568, MIT

(George Eastman Professor of Management Science, Professor of Statistics)

Introduction

1 Among the many ways the Covid-19 crisis is unprecedented is a public disagreement
2 among US airlines on a safety question. As of July 2020, Alaska, Delta, jetBlue, and Southwest
3 Airlines are keeping middle seats open on their flights to limit infection risk, while Allegiant,
4 American, Spirit, and United Airlines are selling all seats when demand warrants. United
5 Airlines has vigorously defended its policy, describing “middle seats only” as a “PR strategy and
6 not a safety strategy.” Its chief communications officer depicted social distancing as impossible
7 on an airplane, saying:

8 “When you're on board the aircraft, if you're sitting in the aisle, and the middle seat is
9 empty, the person across the aisle from you is within six feet of you. The person at the
10 window is within six feet of you. The people in the row in front of you are within six feet
11 of you. The people in the row behind you are within six feet of you.”

12 Yet prominent experts have expressed dismay at the “fill all seats” policy. When
13 American Airlines announced that it would sell as many seats as it could, Dr. Anthony Fauci, the
14 top infectious diseases official at the US National Institutes for Health, told a Senate hearing
15 that “obviously, that's something that is of concern.” Dr. Robert Redfield, the director of the US
16 Centers for Disease Control and Prevention, agreed, declaring that “I can tell you that when
17 they announced that the other day, obviously there was substantial disappointment with
18 American Airlines,”

19 Motivated by this disagreement, this paper is an attempt to estimate the level of risk to
20 US airline passengers under both “middle seats empty” and “fill all seats” policies. Making
21 those estimates entails major complications and uncertainties, which can easily lead one to
22 throw up one’s hands. But even a rough approximation of the risks at issue seems preferable
23 to clashes of unsubstantiated conjectures. This paper strives for such an approximation.

24 As we will discuss, a first-order estimate is that coach passengers on full flights two
25 hours long on popular US jets suffer a 1 in 4300 risk of contracting Covid-19 from a nearby
26 passenger. Under “middle seat empty,” the risk is approximately 1 in 7700, a factor of 1.8
27 lower. Both these estimates are subject to sizable quantifiable and unquantifiable
28 uncertainties, though the factor of 1.8 is considerably less so. Given these estimates and some
29 others, one could expect approximately one death from Covid-19 per 400,000 passengers on
30 flights by airlines that would sell all seats if they could. Under “middle seat empty,” the
31 corresponding figure is about one death per 600,000 passengers.

32 *Materials and Methods*

33 To estimate the risk to an uninfected passenger from a passenger experiencing Covid-19, it
34 is necessary to consider three questions:

- 35 • What is the probability that a given passenger on board is contagious with Covid-19?
- 36 • What is the probability that universal masking can prevent a contagious passenger from
37 spreading the disease?
- 38 • How does the risk of infection depend on the locations on the aircraft of both the
39 contagious and uninfected passenger?

40 The general formula for combining the answers to these questions is:

41
$$P = Q * Q_M * Q_L \quad (I)$$

42 where P= the probability that a particular uninfected passenger contracts Covid-19 during the
43 flight

44 Q = the probability that a given passenger on the flight has covid-19

45 (It is assumed the Q is small enough that having two or more contagious passengers
46 near the uninfected one is a remote risk.)

47 Q_M = the probability that universal mask-wearing on aircraft *fails to prevent*
48 transmission of Covid-19

49 Q_L = the conditional probability that a contagious passenger transmits Covid-19 to the
50 uninfected one *if the mask fails*

51 Q_L and thus P can depend on whether the operating policy is “fill all seats” or “middle seat
52 empty”

53 In the forthcoming analysis, we make estimates of these three quantities. Among the
54 primary assumptions underlying these estimates are:

- 55 1. The number of actual cases of Covid-19 in the US is a large multiple of the number of
56 confirmed cases. However, asymptomatic carriers of the disease are considerably less
57 contagious than pre-symptomatic and symptomatic ones, and air travelers are
58 considerably less likely to be contagious than the citizenry as a whole.
- 59 2. All passengers are wearing masks, and masks are highly effective at preventing
60 transmission of Covid-19.

61 3. An uninfected passenger is only threatened with Covid-19 by a contagious passenger
62 sitting in the same row, the row ahead, or the row behind. Because of strong air-
63 purification mechanisms on aircraft, the risks posed by other passengers are of
64 secondary importance.

65 Specifically:

66 *The Estimation of Q*

67 For a given passenger from a particular American state, the risk of contagiousness is
68 estimated in several steps:

- 69 • First, one finds N_7 , the number of confirmed new Covid-19 infections in that state over
70 the last seven days [1]. Seven days is chosen because that is the approximate length of the
71 contagiousness period for someone experiencing Covid-19. (The average such period is a
72 bit below seven days in asymptomatic cases and higher than seven in symptomatic ones;
73 see [2,3].)
- 74 • One then divides N_7 by N_{POP} , the state's estimated population in 2020, to obtain
75 N_7/N_{POP} as the state's per capita rate of new *confirmed cases* over the last week.
- 76 • Then, in accordance with recent estimates from the US Centers for Disease Control [4] ,
77 one multiplies N_7 by *ten* to approximate the actual number of new infections in the state
78 over the previous week.
- 79 • Then one recognizes that people with Covid-19 who board airplanes are presumably
80 either asymptomatic, pre-symptomatic, or mildly symptomatic. (Those with severe
81 symptoms are unlikely to be flying.) Because of evidence that asymptomatic Covid-19
82 carriers constitute about 40% of all carriers and are only about 40% as contagious as the

83 others [5], one multiplies the prior product by a factor of $\frac{3}{4}$. (This factor of $\frac{3}{4}$ arises because
84 the number of contagious passengers with Covid-19 is approximately $.4*.4 + .6*1 = .76$ of
85 the number of passengers with the disease.)

- 86 • Then one multiplies by a factor of $\frac{1}{2}$ approximately to reflect of the premise that
87 passengers who fly are generally more affluent (and less likely to encounter Covid-19 risks)
88 than the citizenry at large. (This factor treats the half of the population less vulnerable to
89 the disease as $\frac{1}{3}$ as likely to suffer it as the half that is more vulnerable. It further treats
90 air travelers as members of the less-vulnerable half.)
- 91 • Finally, one divides by N_{POP} , the state's estimated population in 2020, to obtain N_7/N_{POP} as
92 the state's per capita rate of new *confirmed* cases over the last week.

93 The estimate of Q consistent with these specifications is:

$$94 \quad Q \approx (N_7 * (10) * \left(\frac{3}{4}\right) * \left(\frac{1}{2}\right)) / N_{POP} = 3.75N_7 / N_{POP} \quad (I)$$

95 *The Estimation of Q_M*

96

97 Here we assume that all passengers wear masks. For Q_M , a meta-analysis in *The Lancet*
98 by Chu et al [6] estimated that mask wearing cuts transmission risk given contagiousness from
99 17.4% to 3.1%, a reduction of 82%. Ignoring the possibility that the masks under study were
100 more effective than those worn by airline passengers, we estimate Q_M as $1 - .82 = .18$.

101 *The Estimation of Q_L*

102 In this analysis, we focus on the coach section of a Boeing 737 or Airbus 320 jet and a
103 flight of two hours (which is about average for a US domestic flight). In a given row, there are
104 six seats: with A, B, and C (the window, middle, and aisle seats, respectively to the left of the
105 aisle,) and D, E, and F (respectively the aisle, middle and window sets right of the aisle).

106 We focus on a particular passenger who is traveling alone, and assume that the primary
107 infection risk for this passenger arises from other passengers in the same row. We further
108 assume that additional risk arises from passengers in the row ahead and row behind. For two
109 reasons, we treat the risk posed by other passengers as negligible:

- 110 • We posit that the airlines are correct when they contend that the powerful
111 air-purification systems on jet aircraft largely negate the risk of aerosol
112 transmission of Covid-19. Thus, when a contagious passenger is in her seat in
113 row 22, she poses little risk to another traveler seated in row 14.
- 114 • A study by Hertzberg et. al. in the *Proceedings of the National Academy of*
115 *Sciences* [7] suggested that infection risk depends on the duration of exposure to
116 contagious person (as did Brundage [8]). We accept this premise, which implies
117 there is limited risk posed by (say) a contagious passenger who passes one's row
118 en route to the lavatory. Moreover, we treat the risks associated with boarding
119 the aircraft, leaving the aircraft, visiting the lavatory, and touching surfaces in
120 the passenger cabin, as second-order effects. If this assumption understates

121 such risks, then the infection estimates presented here would be construed as
122 lower-bounds on Covid-19 risk.

123 A given passenger can get infected, however, by droplets from a contagious passenger
124 in the same row. But here the risk depends on the distance between the two passengers. The
125 meta-analysis of more than 100 studies in *The Lancet* (Chu et al, 2020; [6]) yields the
126 approximation that infection risk is about 13% given physical contact with the contagious
127 person, and that it falls by essentially a factor of two as the distance from that person increases
128 by one meter. The equation reflecting this pattern of exponential decay is:

129
$$R_T \approx 0.13 * e^{-.69d} \quad (II)$$

130 where d = distance in meters between contagious and uninfected person

131 This formula assumes no barriers between the contagious and uninfected persons. If
132 there were (say) a layer of plexiglass between the two, then transmission risk would essentially
133 drop to zero.

134 In each coach row in a Boeing 737 or an Airbus 320, the individual seats are
135 approximately 18 inches wide, while the aisle width is about 30 inches. Under the “fill all seats”
136 policy on a full flight, all six of the ABCDEF seats will be occupied. Under “no middle seats,” A/C
137 and D/F will be occupied on a full flight but not B/E. Assuming that (II) refers to passengers
138 without masks, one can use it to estimate the transmission risk posed by others in the same
139 row to an A-seat passenger, given that the contagious passenger’s mask fails (as happens with
140 probability Q_M):

$$141 \quad Q_L(A \text{ same row}) \approx \begin{cases} R_T(A, C) + R_T(A, D) + R_T(A, F) \text{ under middle seat open} \\ R_T(A, B) + R_T(A, C) + R_T(A, D) + R_T(A, E) + R_T(A, F) \text{ under "fill all seats"} \end{cases}$$

142 (III)

143 where $R_T(A, X)$ = transmission probability *absent masks* given a contagious passenger in seat X
144 of a given row and an uninfected passenger in seat A of that row

145 Equation (III) taken literally treats infections caused by passengers in different seats as
146 mutually exclusive events. But they are not mutually exclusive: it is possible that
147 contagious persons are seated in both seats 16C and 16F. The actual assumption –
148 consistent with data--is that Q is small enough that having several contagious people
149 close to one another is a second-order effect, with probabilities involving Q^2 or higher
150 powers of Q. In practical terms, therefore, the events of interest are mutually
151 exclusive.

152 We therefore make the approximation that:

$$153 \quad R_T(A, X) \approx .13e^{-.69d(A,X)} \quad (IV)$$

154 where $d(A, X)$ = distance from a person's head in the *middle* of seat A to another
155 person's head in the middle of seat X.

156 For the jets under consideration, the quantity $d(A, B)$ is about 18 inches, while
157 $d(A, C)$ is $18+18= 36$ inches, $d(A, D) = 36 + 9 + 30 + 9 = 84$ inches, $d(A, E) = 84+18= 102$
158 inches, and $d(A, F) = 102 + 18= 120$ inches. Because a meter is 39.37 inches, $d(A, B)$ in
159 meters is $18/39.37 = .457$, etc.

160 Analogous expressions arise when the uninfected passenger is in the B, C, ..F seat.

161 One can use (III) and (IV) to obtain:

162
$$Q_L(A \text{ same row}) \approx \begin{cases} 0.115 & \text{under middle seat empty} \\ 0.232 & \text{under "fill all seats"} \end{cases}$$

163 Using similar reasoning, one can likewise determine that:

164
$$Q_L(B \text{ same row}) \approx .282 \text{ under "fill all seats"}$$

165
$$Q_L(C \text{ same row}) \approx \begin{cases} 0.155 & \text{under middle seat empty} \\ 0.291 & \text{under "fill all seats"} \end{cases}$$

166
$$Q_L(D \text{ same row}) = Q_L(C \text{ same row});$$

167
$$Q_L(E \text{ same row}) = Q_L(B \text{ same row});$$

168
$$Q_L(F \text{ same row}) = Q_L(A \text{ same row});$$

169 Averaging across all the passengers in a given row yields:

170
$$Q_L(\text{same row}) = \begin{cases} .268 & \text{under "fill all seats"} \\ .135 & \text{under middle seat empty} \end{cases}$$

171 Hertzberg et al [7] concluded from their computer simulations that, for droplet-
172 mediated respiratory diseases, contagious passengers pose appreciable transmission risk to
173 uninfected travelers within one meter. They therefore concluded that, beyond the same row,
174 transmissions can occur from passengers in the row ahead of an uninfected passenger and in
175 the row behind. Here we process distances using (III) rather than a one-meter yes/no

176 threshold, but we concur that passengers in the two neighboring rows pose first-order
177 transmission risk.

178 In Hertzberg et. al. all contagious passengers within one meter pose equal levels of
179 transmission risk, regardless of whether they are in the same row as the uninfected passenger.
180 However, the authors noted that they did not consider the possibility that seatbacks would
181 impede transmissions between rows. Here we do not ignore that possibility.

182 While seatbacks can somewhat block droplets from a contagious passenger, they are
183 presumably less effective than plexiglass, which all but eliminates transmission. Lacking
184 available studies about the benefit conferred by seatbacks, we make the estimate that they are
185 about $\frac{3}{4}$ as effective as plexiglass. More specifically, we assume that:

- 186 • When the flight is full, the six passengers one row ahead of the uninfected passenger
187 collectively pose $\frac{1}{4}$ the transmission risk of the five passengers in the same row.
- 188 • When the flight is full, the six passengers one row behind the uninfected passenger
189 collectively pose $\frac{1}{4}$ the transmission risk of the five passengers in the same row.
- 190 • When the flight follows “middle seats empty” but is otherwise full, the four passengers
191 one row ahead of the uninfected passenger collectively pose $\frac{2}{3}$ the transmission risk of
192 the six passengers in that row had the flight been full.

- 193 • When the flight follows “middle seats empty” but is otherwise full, the four passengers
194 one row behind the uninfected passenger collectively pose 2/3 the transmission risk of
195 the six passengers in that row had the flight been full.

196 If this factor-of-four reduction overstates the effectiveness of the seatbacks against contagion,
197 then our risk estimates tied to neighboring rows could well be too low.

198 Under these approximations:

$$Q_L(\text{full flight}) = 1.5Q_L(\text{full flight, same row})$$

199 $Q_L(\text{middle seat empty}) = Q_L(\text{middle seat empty, same row}) + \left(\frac{2}{3}\right) * \left(\frac{1}{2}\right) Q_L(\text{full flight same row})$

200 Thus:

201
$$Q_L = \begin{cases} .402 & \text{for full flight} \\ .224 & \text{when middle seat empty} \end{cases}$$

202 We should stress that Equation (II) summarizes the experiences reported in many
203 disparate studies, probably none of which reflects the exact conditions in a US jet flight two
204 hours long. The risk of infection presumably relates to duration of exposure, but the durations
205 in the summarized studies are often unknown. Moreover, US airlines have argued convincingly
206 that ventilation during their flights suppresses disease spread more effectively than that in a
207 typical indoor setting. Yet the mix of environments in the studies that contributed to (II) is not
208 clear. It is also understood that disease spread is greater when the contagious person is
209 speaking than when he is silent. But the proportional breakdown of exposure time between
210 speaking and silence is unavailable, both for airplanes and in the summarized studies.

211 A further complication is that, while we are assuming that Equation (II) arises when the
212 contagious person wears no mask, there were apparently some masked individuals in the
213 studies that generated the equation (private communication from Dr. Chu). As noted, we
214 apply an 82% reduction in infection risk because of masks, but (II) already reflects the benefit of
215 masks to some extent.

216 These circumstances need not compromise the exponential-decay factor in (ii), under
217 which transmission risk drops by a factor of two per meter of distance. But the 13% factor
218 could well be affected, with some considerations suggesting the factor is too high and others
219 that it is too low. This analysis offers a baseline risk estimate using the Chu et al. results at face
220 value, which seems reasonable absent further information about the combinations of
221 conditions that underlie those results.

222 When the estimates of Q , Q_M , and Q_L are at hand, they are multiplied under (I) to obtain
223 a point estimate of P .

224 *Confidence Intervals for the Point Estimates*

225 The point estimates of Covid-19 risk are subject to various sources of uncertainty, some
226 of them subject to quantification. The basic relationship:

$$227 \quad P \approx Q * Q_M * Q_L \quad (I)$$

228 can be rewritten as:

$$229 \quad \ln P \approx \ln Q + \ln Q_M + \ln Q_L$$

230 The confidence intervals in the literature for individual parameters typically suggest that the
231 variables in (I) are approximately lognormal. (e.g. the 95% confidence interval extends from

232 roughly half the point estimate to double that estimate), and the estimation errors for different
233 variables can be considered independent. Thus, $\ln P$ would be essentially normal, and a 95%
234 confidence interval for P can be obtained through exponentiating the 2.5%ile and 97.5%ile of
235 $\ln P$. The calculated confidence intervals pertain to known uncertainty in the risk estimates,
236 and not to the uncertainties in Equation (II) discussed above.

237 *Results*

238 *Point Estimates*

239 New confirmed covid-19 cases were sharply increasing in many American states during
240 the last week of June 2020, but continuing to decline in others. Exemplifying the states
241 experiencing spikes was Texas, with 42,254 new cases over 6/24/20 to 6/30/20, while typifying
242 states long past their peaks of new infections was New York, with 5,200 cases over that period.
243 Because the population of Texas in 2020 was estimated at 29.1 million in mid-2020, its per
244 capita rate of new infections that week (i.e., N_7/N_{POP}) was 1/689. For New York, with a
245 population of 19.5 million, the corresponding rate was 1/3750.

246 Under (I), the probability a passenger from a particular state has contagious covid-19 is
247 approximated by:

$$248 \quad Q \approx 3.75(N_7/N_{POP})$$

249 Meaning that:

$$250 \quad Q \approx \begin{cases} \frac{1}{184} & \text{for Texas} \\ \frac{1}{1000} & \text{for New York} \end{cases}$$

251 For US domestic jet flights as a group, one might approximate Q by taking the average of
252 these estimates for higher-infection-rate Texas and lower-rate New York, which yields $Q \approx \frac{1}{310}$
253 Even on a flight from Dallas to New York City, there will be Texas natives, New York natives, and
254 transfer passengers who originated elsewhere, so a mid-range estimate seems suitable. Had
255 we instead used new infections over a week in mid-July 2020 (when this is written), the
256 estimate of Q would have increased by a factor of 1.6. We use the late June estimate in the
257 hope that the mid-July upsurge is temporary.

258 As noted, the probability Q_M for mask failure is estimated as 0.18, while .402 and .224 are
259 treated as, respectively, the transmission probabilities absent masks (Q_L) under “fill every seat”
260 and “middle seat empty.” In consequence, Equation (I) generates the following estimates for
261 dates around 6/30/20:

$$262 \quad P(\text{infection}) \approx \begin{cases} \frac{1}{4,300} & \text{on full US flights} & \text{under "fill all seats)} \\ \frac{1}{7,700} & \text{on full US flights under "middle seat empty"} \end{cases}$$

263 The first of these risk estimates is the average for the passengers in the six filled seats in each
264 row. The second is the average for passengers in the four seats occupied under “middle seat
265 empty.”

266 *Confidence Intervals*

267 As noted earlier, there is nonquantifiable uncertainty in our use of Equation (II), related to
268 ventilation, speech/silence, duration of exposure, and mask usage. But the literature offers
269 margins of error in addition to point estimates for three parameters in the risk calculations:

- 270 • The 95% confidence interval for the rate of exponential decay with distance extends
271 from a factor of 1.08 per meter to a factor of 3.76. (We have used a point estimate of
272 two, corresponding to -0.69 in (II).)
- 273 • The 95% confidence interval for the failure probability of masks extends from .07 to
274 .34 (point estimate .18)
- 275 • The 95% confidence interval for the ratio of actual cases of Covid-19 to confirmed
276 cases extends from 8.7 to 12.7. (point estimate 10).

277 These confidence intervals suggest that the distributions for the parameter estimates can
278 be approximated as lognormal. For example, the interval for exponential decay goes from
279 about half the point estimate of two to roughly double that estimate. Treating
280 uncertainties in the three parameters as independent, we can estimate the overall degree
281 of imprecision that they (alone) cause in the estimate of P.

282 In the relationship based on (I):

$$283 \quad \ln P = \ln Q + \ln Q_M + \ln Q_L,$$

284 the independence and normality of the three random variables on the right implies that $\ln P$
285 itself is normally distributed. First we find the mean and standard deviations of $\ln Q$, $\ln Q_M$, and
286 $\ln Q_L$, and then we use them to find the mean, standard deviation, and 95% confidence interval
287 for $\ln P$.

288 For $\ln Q_M$, we note that its 2.5th percentile is $\ln(.07)$ and its 97.5th percentile is $\ln(.34)$.
289 From this information and the assumption of lognormality, we readily deduce that the mean of
290 $\ln Q_M$ is the average of $\ln(.07)$ and $\ln(.34)$, while its standard deviation is $(\ln(.34) - \ln(.07))/3.92$.
291 Because $\ln(.34) = -1.08$ and $\ln(.07) = -2.66$, $\ln(Q_M)$ is approximately normal with mean -1.87
292 and standard deviation 0.40 . The calculations for $\ln Q$ and $\ln Q_L$ proceed similarly, with the
293 distribution for $\ln Q_L$ dependent on whether the policy under study is “fill all seats” or “middle
294 seats empty.”
295 The 95% confidence intervals for P are:

296
$$\left\{ \begin{array}{l} \frac{1}{10,500} \text{ to } \frac{1}{1,700} \text{ for full flights under fill all seats.} \\ \frac{1}{19,300} \text{ to } \frac{1}{3,000} \text{ for full flights under middle seat empty} \end{array} \right.$$

297 In both instances, there is factor-of-2.5 uncertainty in the point estimates offered earlier (i.e.
298 $\frac{1}{4,300}$ for fill all seats, $\frac{1}{7,700}$ for middle seats empty). It important to note, however, that the
299 two estimates of P arise from the same parameters, and that the uncertainty in those
300 parameters affects both estimates in essentially the same way. The upshot is that the ratio of
301 $P(\text{fill all seats})$ to $P(\text{middle seat empty})$, which is 1.8 based on the point estimates. i.e.,
302 $(1/4,300)/(1/7,700)$, would stay stable despite the considerable uncertainty that affects the
303 numerator and denominator of the ratio.

304 *Discussion*

305 The reason that Covid-19 is so fearsome is that it entails a risk of death. For a coach
 306 passenger with a 1% chance of dying from the virus—which is slightly above the chance of 0.7%
 307 now assumed to apply to the full population [9]-- the estimated mortality risk on a full flight
 308 two hours long would be about 1 in 430,000 and about 1 in 770,000 million when all but middle
 309 seats are full. But even an airline willing to fill every seat does not expect to do so: it
 310 presumably aspires to a passenger load factor (passenger miles divided by seat miles) around
 311 85.1%, which prevailed in 2019 on US flights. Having 85.1% of seats taken is consistent with
 312 roughly 55% of middle seats full and 45% empty. With that load factor, a 1% chance of dying
 313 from Covid-19 given infection would yield a death risk under “fill all seats” of about 1 in 540,000
 314 rather than 1 in 430,000. Airlines that keep middle seats empty could well try to fill nearly all of
 315 them, so their mortality risk would remain about 1 in 770,000.

316 Actually, covid-19 infections on planes can cause deaths to some people who were not
 317 passengers (e.g., a 22-year traveler gets infected, and passes the virus on to his elderly
 318 grandparents). A quantity familiar in this pandemic is R_0 , the average number of new
 319 infections generated directly by an infected person. The quantity $E(\text{Further Inf})$, the mean total
 320 number of further infections that person causes, follows:

321
$$E(\text{Further Inf}) = \frac{R_0}{1 - R_0} \quad \text{for } R_0 < 1 \quad (V)$$

322 Using the conservative estimate that $R_0 = 0.5$, $E(\text{Further Inf}) = 1$ under (V). Assuming a
 323 0.7% death risk for both the person infected on a flight and the people further infected, the
 324 expected total number of deaths would be $0.007 + 1 \cdot 0.007 = .014$. Thus, if a given passenger on
 325 a full flight has a 1 in 4,300 chance of getting infected, the resulting number of deaths would on

326 average be about $(1/4300) \cdot .014 = 1$ in 310,000. If the flight is 85.1% full, the estimate would
327 decline to 1 in 390,000. If the flight is 66.7% full under middle seat empty, the estimate would
328 decline further to 1 in 550,000.

329 All these death-risk estimates are considerably higher than the risk of perishing in a US
330 air crash unrelated to Covid-19, which is about 1 in 34 million [10]). It is not clear, however,
331 that two hours spent on an airplane entails higher infection risk (or mortality risk) than two
332 hours of everyday activities. In late June 2020, approximately 45,000 Americans were
333 confirmed to have contracted Covid-19 each day. Given that actual infections are estimated to
334 be about ten times confirmed ones [4], roughly 450,000 new infections arose per day among
335 the 330 million Americans. That works out to a daily infection probability of
336 $450,000/330$ million, which is $1/733$. Assuming 16 waking hours, the chance of infection over a
337 two-hour period would be approximately $(2/16) \cdot (1/733) = 1/5900$, which is quite close to our
338 infection risk estimates for a two-hour flight. (If air travelers have lower Covid-19 risk than
339 average citizens—as we have assumed—that could be because they engage in relatively few
340 everyday activities (e.g. fewer rides on buses or visits to supermarkets)). But that circumstance
341 does not affect the risk estimate for such activities.)

342 *Final Remarks*

343 Calculations like the ones here are highly approximate and, as has been evident during
344 this pandemic, projections about it often fall far from the mark. It would therefore be
345 desirable to use actual passenger outcomes to determine what fraction of travelers contracted
346 Covid-19 on their flights. If, averaged over US carriers, the risk level per passenger is estimated
347 as (say) 1 in 6,500, then approximately 90 cases of Covid-19 should emerge each day at a time

348 (like early July 2020) when 600,000 US passengers are flying daily. In reality, load factors fell
349 below 50% in July 2020 even on “fill all seats” airlines, meaning that the number of cases that
350 actually arose would be closer to 50 per day.

351 But it would not be easy to determine how many Covid-19 cases arose on US flights on a
352 given day. If only 10% of infections are confirmed, 50 infections would yield five confirmed
353 ones. Is it plausible that five of the 45,000 confirmed US infections per day arose during air
354 journeys over the previous week? Because follow-ups of known Covid-19 infections are now
355 weak in the US, it is not clear that the circumstances of those five infections would be
356 identified. Further complicating the situation is the fact that air passengers who subsequently
357 get Covid-19 may have been infected elsewhere besides the airplane. Despite these difficulties,
358 it would be worth some effort to substantiate or refute projections that are tied to strong
359 assumptions.

360 The calculations here, however imperfect, do suggest a measurable reduction
361 in Covid-19 risk when middle seats on aircraft are deliberately kept open. The question is
362 whether relinquishing 1/3 of seating capacity is too high a price to pay for the added
363 precaution.

Acknowledgements

To be supplied

References

- (1) Worldometer.com “Age, Sex, Existing Conditions of COVID-19 Cases and Deaths, data for confirmed cases by day and American state, available at <https://www.worldometers.info/coronavirus/coronavirus-age-sex-demographics/>
- (2) A. Byrne et. al.(2020), “Inferred Durations of Infectious Period of SARS-Cov-2: Rapid Scoping Review and Analysis of Available Evidence for Asymptomatic and Symptomatic Covid-19 Cases,” available at <https://www.medrxiv.org/content/10.1101/2020.04.25.20079889v1.full.pdf>
- (3) Y. Ling et. al.(2020), “Persistence and Clearance of Viral RNA in 2019 Novel Coronavirus Disease Rehabilitation Patients,” *Chinese Medical Journal* 133(9), pp. 1039-1043, May 5, 2020, available at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7147278/>
- (4) F. Havers et al, “Sero-prevalence of Antibodies to Sars-Cov-2 in 10 Sites in the United States, March 23-May 12, 2020, *JAMA Internal Medicine*, July 21, 2020, available at <https://jamanetwork.com/journals/jamainternalmedicine/fullarticle/2768834>
- (5) CSPEC (2020), “Defining High-Value Information for Covid-19 Decision Making,” available at <https://www.medrxiv.org/content/10.1101/2020.04.06.20052506v1.full.pdf>
- (6) D. Chu et. al.(2020) “Physical Distancing, Face Masks, and Eye Protection, To Prevent Person-to-Person Transmission of SARS-Cov-2 and COVID-19: A Systematic Review and Meta-Analysis, *The Lancet*, 395(10242), pp. 1973-1987, June 27, 2020
- (7). V.S. Hertzberg, H. Weiss, et. al. (2018) “Behaviors, Movements, and Transmission of Droplet-Mediated Respiratory Diseases During Transcontinental Airline Flights, *Proceedings of the National Academy of Sciences*, 115(14), pp. 3623-3627, April 3, 2018. This paper supersedes an earlier paper by Hertzberg and Weiss, which suggested that infection could spread beyond one row away from the contagious person. (VS. Hertzberg H. Weiss .(2016), On the 2-row rule for Infectious Disease Transmission on Aircraft. *Ann Glob Health*; 82: pp 819–823.)

(8) E. Brundage (2020), “The Risks: Know Them, Avoid Them,” available at <https://www.erinbromage.com/post/the-risks-know-them-avoid-them>

(9) G. Meyerowitz-Katz and L. Merone “A Systematic Review and Meta-analysis of Published Research Data on Covid-19 Infection-Fatality Rates,” July 7, 2020, available at <https://www.medrxiv.org/content/10.1101/2020.05.03.20089854v4>

(10) A. Barnett (2020), “Aviation Safety: A Whole New World?” *Transportation Science* 54(1), pp. 84-96, January-February 2020,

