Manual ability in hand surgery patients: validation of the ABILHAND scale in four diagnostic groups

Short title: Validation of the ABILHAND scale for hand surgery

Ghady El Khoury[1,2*], Olivier Barbier[1], Xavier Libouton[1], Jean-Louis Thonnard[2], Philippe Lefèvre[2], Massimo Penta[2,3]

[1] Service d'orthopédie et traumatologie, Cliniques universitaires Saint-Luc, Bruxelles, Belgique

[2] Institute of Neuroscience, Université catholique de Louvain, Bruxelles, Belgique

[3] Arsalis SRL, Glabais, Belgium

* Corresponding author

Email: ghady.elkhoury@uclouvain.be (GEK)

Abstract

Background: Patients treated in hand surgery (HS) belong to different demographic groups and have varying impairments related to different pathologies. HS outcomes are measured to assess treatment results, complication risks and intervention reliability. A one-dimensional and linear measure would allow for unbiased comparisons of manual ability between patients and different treatment effects.

Objective: To adapt the ABILHAND questionnaire through Rasch analysis for specific use in HS patients and to examine its validity.

Methods: A preliminary 90-item questionnaire was presented to 216 patients representing the diagnoses most frequently encountered in HS, including distal radius fracture (n=74), basal thumb arthritis (n=66), carpal tunnel syndrome (n=53), and heavy wrist surgery (n=23). Patients were assessed during the early recovery and in the late follow-up period (0-3 months, 3-6 months and >6 months), leading to a total of 305 assessments. They rated their perceived difficulty with queried activities as impossible, difficult, or easy. Responses were analyzed using the RUMM2030 software. Items were refined based on item-patient targeting, fit statistics, differential item functioning, local independence and item redundancy. Patients also completed the QuickDASH, 12-item Short Form Survey (SF-12) and a numerical pain scale.

Results: The rating scale Rasch model was used to select 23 mostly bimanual items on a 3-level scale, which constitute a unidimensional, linear measure of manual ability with good reliability across all included diagnostic groups (Person-Separation Index = 0.90). The resulting scale was found to be invariant across demographic and clinical subgroups and over time. ABILHAND-HS

36   patient measures correlated significantly (p<0.001) with the QuickDASH (r=-0.77), SF-12 Physical

37   Component Summary (r=0.56), SF-12 Mental Component Summary (r=0.31), and pain scale (r=-

38   0.49).

39   **Conclusion:** ABILHAND-HS is a robust person-centered measure of manual ability in HS patients.

Introduction

In hand surgery (HS), as in other medical specialties, outcome evaluations are needed to assess the effectiveness and reliability of the intervention, as well as to reinforce patient education regarding the risks and outcomes of the procedure and, potentially, to justify therapeutic practices to payers [1]. Physician-documented reports of HS outcomes based on clinical examination and imaging should be complemented with patient reported outcomes assessed by questionnaires designed to capture patients' perspectives with respect to the impact of their conditions and interventions on their daily lives [2–4].

Current views of health and disability have been shaped by the World Health Organization's International Classification of Functioning, Disability, and Health [5], which parses disease consequences into three domains: impairment of anatomical structures (e.g. bones, muscles, ligaments) or body functions (e.g. motor skills, sensitivity), activity limitations (e.g. manual activities), and participation restrictions (e.g. in hobbies and work). The impact of a pathology or a surgical intervention in these three domains is also conditioned by personal factors (motivation, capacity to develop compensatory strategies) and environmental factors (social or professional context). Although impairment measurements such as imaging can provide clues regarding functional prognosis, it does not provide good information about performance in everyday life, especially of the hands, which are important for a great variety of activities [6–8]. For example, demonstration of a bone fracture union is insufficient to determine whether a patient is capable of resuming their usual activities or returning to work [9–12].

The patient-reported questionnaires that have been most commonly used in HS [13] are the Disability of the Arm, Shoulder and Hand questionnaire (DASH) [14], the Patient Rated Wrist

Evaluation (PRWE) [15], and the Carpal Tunnel Questionnaire (CTQ) [16]. Each of these

questionnaires has been reported to have good psychometric properties, but each has a

particular focus on its own area(s) of disablement. The DASH assesses body functions, activities,

and participation [17] and can be divided into 3 subscales based on dimensionality [18].

Meanwhile, the PRWE is specific to the wrist joint and the CTQ is specific to carpal tunnel

syndrome (CTS). The Michigan Hand Outcomes Questionnaire (MHQ) [19] is a multidimensional

hand-specific outcomes instrument consisting of six subscales, measuring overall hand function,

activities of daily living, pain, work performance, aesthetics, and satisfaction. It measures

impairment by hand (left and right separately), rather than overall disability. Interpretation of

total scores on multi-dimensional instruments can be less than straightforward given that

patients can show simultaneous improvement in one domain with deterioration in another

[20,21]. Assessment of functional recovery on a unidimensional [22] and linear [21] scale would

allow for quantitative comparisons of ability among different patients and treatments. Such a

scale can be developed with state-of-the-art psychometric methods, such as the Rasch model

[23,24].

The ABILHAND questionnaire is a Rasch-model built measure of manual ability [25] that provides

an invariant linear scale and allows for quantitative comparisons of manual ability between

patients and over time. The scale has been validated in populations with rheumatoid arthritis

[26], chronic stroke [8], pediatric cerebral palsy [27], systemic sclerosis [28] and neuromuscular

diseases [29]. These previous validations have shown that the difficulty of most manual activities

was diagnosis-dependent [30]. Therefore, the objective of this work was to adapt the ABILHAND

scale to the most frequent diagnoses treated in HS.

Methods

Questionnaire adaptation to HS patients

The ABILHAND is a measure of manual ability that assesses one's ability to manage daily activities requiring upper limb use, regardless of strategy [25]. The necessary permissions were obtained from the developer of the original questionnaire to modify it. To develop a HS-adapted ABILHAND, a preliminary item list was compiled from previous versions of the ABILHAND questionnaire, the DASH, PRWE, CTQ, and MHQ items together with some new items. This pool of items was submitted to nine HS experts (hand surgeons, physical medicine and rehabilitation physicians, physical therapists, and occupational therapists), who were asked to assess each item's relevance to hand surgery patients on a yes/no basis and propose additional items that might be affected by the relevant pathologies (e.g. sensation for CTS and wrist loading for distal radius fractures (DRF)). A final list of 90 items constituted the experimental ABILHAND-HS questionnaire (S4 Appendix).

Patients

A convenience sample of 216 patients was recruited from February 2018 to February 2019 at the HS consultation center at Cliniques Universitaires Saint-Luc, Belgium representing the following four diagnostic categories: CTS, DRF, basal thumb arthritis (BTA), and heavy wrist surgery (HWS, including 1st row carpectomy and partial or total wrist arthrodesis). The inclusion criteria for patients were being >18 years old and being able to read and understand French. The exclusion criteria included comorbidities that may impede manual ability substantially (i.e. tremor, paralysis and active rheumatologic disease) and any mental or cognitive dysfunction (i.e. dementia and

105   mental retardation). The patient characteristics are summarized in Table 1. Patients provided

106   written informed consent to participate. This study was approved by the ethical committee of

107   Cliniques Universitaires Saint-Luc-Université catholique de Louvain (N° B403201523492).

109 **Table 1. Sample characteristics (n = 216).**

Characteristic	N (%)[a]
Gender	
Women	145 (67%)
Men	71 (33%)
Mean age (range), years	60.3 (19–93)
Education level	
Basic	109 (51%)
Postsecondary	107 (49%)
Work status	
Student	2 (1%)
Unemployed	22 (10%)
(Self-)Employed	83 (38%)
Retired	109 (51%)
Hand dominance	
Right	194 (90%)
Left	15 (7%)
Ambidextrous	7 (3%)
Involved dominant hand	
Yes	136 (63%)
No	80 (37%)
Diagnostic group	
Distal radius fracture (DRF)	74 (34%)
Basal thumb arthritis (BTA)	66 (31%)
Carpal tunnel syndrome (CTS)	53 (24%)
Heavy wrist surgery (HWS)	23 (11%)
Follow-up assessments (n = 305)	
0–3 months (57 DRF, 52 CTS, 22 BTA, 1 HWS)	132 (43%)
3–6 months (38 DRF, 16 CTS, 3 BTA, 1 HWS)	58 (19%)
>6 months (30 DRF, 18 CTS, 46 BTA, 21 HWS)	115 (38%)

[a]**Except where otherwise indicated**

Procedures

The French-language experimental ABILHAND-HS items were presented in five random orders to avoid a systematic item sequence bias. Patients were asked to indicate their perceived difficulty associated with completing the activities without technical or human assistance, independent of the hand used to perform the activity on a three-level scale: impossible (0), difficult (1), or easy (2) [8]. Activities not attempted during the last week were treated as missing responses. Patients also completed the QuickDASH [31], 12-item Short Form Survey (SF-12) questionnaire [32] and a 10-level numerical pain scale, for external validation purposes.

Patients were first assessed as soon as they presented to their hand surgery consultation appointments and had experienced manual activities in their own environment: after hand surgery and cast removal for DRF, BTA and HWS and at the first consultation for non-operated CTS and BTA. For the first assessment, patients were interviewed by the principal investigator in order to ensure clarity, obtain feedback from participants, and make sure instructions are properly followed. Patients were also asked to suggest additional items they felt the questionnaire was missing. However, these were either gender related (e.g. fastening a bra) or very specific and were thus not retained. Follow-up assessments were completed in our consulting office or returned by mail, leading to a total of 305 completed assessments, which provides sufficient power to support the planned Rasch analysis [33].

Rasch analysis

The 90-item experimental ABILHAND-HS questionnaire responses were analyzed using the Rasch model in RUMM2030 software (RUMM Laboratory Pty Ltd., Perth, Australia). The Rasch model

[24], a prescriptive model, requires that specified response probabilities depend on only item

difficulty and patient ability. Polytomous datasets with thresholds between successive response

categories can be analyzed with either a rating scale model that constrains all threshold locations

to be equal across items [34] or a partial credit model that allows threshold locations to vary

across items [35]. Patient abilities and item difficulties are located along a common linear,

unidimensional continuum that defines the latent variable of interest (i.e. manual ability). The

locations are expressed in logits, calculated as the logarithm of the pass/fail probability ratio of

an item or threshold. The logit locations were converted into centiles to facilitate clinical

interpretation on a linear scale ranging from 0% (smallest ability) to 100% (largest ability) [36].

Expected responses, determined based on the patient and item locations, were compared to the

responses actually reported to compute residual and fit statistics, which were then used to assess

the scale's unidimensionality [37]. A good fit of the data with the model affirms invariant locations

along the continuum and indicates that the measure can be used to compare manual ability

across patients and diagnoses.

Item selection

From the experimental version of the questionnaire, the ABILHAND-HS was refined through

successive analyses of 305 assessments with the goal of selecting items that define a

unidimensional and clinically relevant scale of manual ability. P values < 0.05 were considered

significant for each of the following analysis steps:

1) *Item-patient targeting.* Based on examination of patient distributions and item locations,

items that showed a floor effect (too easy) or did not target the patients sample ability were

removed.

2) *Rating scale*. Items with disordered thresholds and items with thresholds that were too

narrow (<1.4 logits) or too wide (>5 logits) were removed before applying the rating scale

model [38].

3) *Unidimensionality*. Only items that delineated a common manual ability construct according

to the following four criteria were retained: (1) standardized residuals obtained over three

class intervals had to be within ±2.5 with a non-significant χ^2 [37]; (2) no observable major

differential item functioning (DIF) [39], uniform or non-uniform, shown by a 2-way analysis of

variance of the residuals with Bonferroni correction [40], according to gender (male vs.

female), age (above vs. below the median age of 63 years), pathology (CTS vs. DRF vs. BTA vs.

HWS), involved hand (dominant vs. non-dominant), level of education (basic vs. superior), and

follow-up (0-3 months vs. 3–6 months vs. >6 months); (3) overall fit of the response set based

on a non-significant item-trait interaction χ^2 [37]; and (4) statistically similar patient locations,

according to paired t-tests, calculated with items that loaded either positively or negatively

on the first residuals principal component [41–43].

4) *Local independence.* When items were found to be querying redundant content [44],

demonstrated by a residual correlation > 0.3, the item with the poorer fit statistic was deleted

[45].

5) *Item redundancy*. To shorten the scale, when two or more items had similar locations on the

continuum, the one with the best fit was retained.

Scale reliability

The Person-Separation Index (PSI), i.e. the proportion of total variance (including error) that is attributed to patient location variance, was used to determine the ABILHAND-HS scale's reliability and its degree of precision with the dataset, and thus how many statistically different ability strata can be distinguished along the scale [46].

Construct validity

The construct validity of the ABILHAND-HS was examined with a comparison of means for associations with gender, involved hand, and diagnosis. The relationships of the ABILHAND-HS with age, the QuickDASH scale, the numerical pain scale, the SF-12 Physical Component Summary (PCS), and the SF-12 Mental Component Summary (MCS) were assessed with a correlation analysis.

Patient perceptions were compared between ABILHAND-HS and QuickDASH items by adding the six QuickDASH activity items to the anchored data matrix. The locations of similar items were then compared between the scales.

Statistical analyses

Statistical analyses were completed in IBM SPSS Statistics for Windows, version 25 (IBM Corp., Armonk, N.Y., USA). Data normality was verified for statistical tests using the Shapiro-Wilk test and Q-Q plots. Parametric tests were used for normal data and continuous variables, non-parametric tests for non-normal data and ordinal variables. A Mann-Whitney u-test (two-tailed) was used for gender differences, an independent-samples t-test (two-tailed) for association with

the involved hand, and an analysis of variance for diagnosis. Pearson correlation coefficient was

calculated for association with age, while relationships with the QuickDASH scale, the numerical

pain scale, the SF-12 Physical Component Summary (PCS), and the SF-12 Mental Component

Summary (MCS) were assessed with Spearman correlation coefficients. P values < 0.05 were

considered significant. Mean values are reported with standard deviations (SD). Chi-square and t

values are reported with degrees of freedom (df).

Results

Item selection for the ABILHAND-HS scale

201 Successive analyses led to the selection of 23 items defining a unidimensional manual ability scale

202 in HS. Of the 90 experimental items, 34 were removed because they were too easy (e.g. 'Drinking

203 a glass of water'), 3 items had too-narrow thresholds (e.g. 'Using a touch screen'), 4 items were

204 misfitting (e.g. 'Carrying a shopping bag'), and 26 items had a location redundant with another

205 better fitting item (e.g. 'Peeling onions' was deleted in favor of 'Peeling potatoes with a knife').

Metric properties

The calibration obtained for the 23 mostly bimanual activities retained for ABILHAND-HS is

reported in Table 2 in descending difficulty order. The standardized residuals obtained matched

the expected standard normal distribution for items [mean (SD), -0.30 (0.99)] and for patients

[0.31 (0.97)], indicating that the ABILHAND-HS scale is globally unidimensional. An invariant item

location was obtained for more- and less- able patients as shown by a nonsignificant item-trait

interaction (χ^2 = 57.76, 46 df, p = 0.11). An invariant patient ability was obtained with items with

different content as shown by a non significant t-test when using items that loaded positively or

negatively on the first principal residual component (t = 1.24, 304 df, p = 0.22).

Table 2. Calibration of the 23 items of the ABILHAND-HS.

Item	Bi-manual	Difficulty logits (centiles)	SE logits	Residual z	Fit χ²	P
a. Doing push-ups	x	3.54 (78)	0.21	0.61	0.46	0.79
b. Playing a racket sport	x	2.30 (68)	0.25	0.04	1.66	0.44
c. Cutting a hedge	x	2.00 (65)	0.18	0.05	3.83	0.15
d. Opening a screw-topped jar	x	1.30 (59)	0.12	0.48	6.19	0.05
e. Applauding vigorously	x	1.11 (58)	0.16	2.09	2.00	0.37
f. Lifting a full pan	x	0.96 (57)	0.12	-0.53	5.69	0.06
g. Wringing a towel	x	0.86 (56)	0.12	-2.00	0.14	0.93
h. Opening a can with a can opener	x	0.76 (55)	0.12	-1.83	0.49	0.78
i. Hammering a nail	x	0.45 (52)	0.16	-0.70	5.31	0.07
j. Shaking bed sheets	x	0.13 (50)	0.17	-1.95	3.49	0.17
k. Using a screwdriver	x	-0.05 (48)	0.14	-0.14	0.54	0.77
l. Peeling potatoes with a knife	x	-0.16 (47)	0.13	-0.75	5.75	0.06
m. Ironing	x	-0.38 (45)	0.15	-0.44	3.12	0.21
n. Taking the cap off a bottle	x	-0.52 (44)	0.13	-0.32	2.75	0.25
o. Cutting one's nails	x	-0.55 (44)	0.13	-0.05	0.33	0.85
p. Shuffling and dealing cards	x	-0.77 (42)	0.16	-1.21	6.18	0.05
q. Wiping windows		-0.77 (42)	0.15	0.01	1.29	0.52
r. Tying shoelaces	x	-0.96 (41)	0.14	-0.61	2.02	0.36
s. Tearing open a pack of chips	x	-1.16 (39)	0.15	1.70	1.31	0.52
t. Fastening the zipper of a jacket	x	-1.45 (37)	0.14	0.32	1.58	0.45
u. Turning a car steering wheel	x	-1.69 (35)	0.18	-0.26	0.78	0.68
v. Putting on gloves	x	-2.24 (30)	0.19	-0.80	1.99	0.37
w. Spreading butter on a slice of bread	x	-2.68 (26)	0.18	-0.60	0.86	0.65

Analysis of DIF of the ABILHAND-HS with six criteria yielded only four instances of uniform DIF among the 23 items (Table 3). A small magnitude DIF was revealed among diagnoses (Fig 1) with no substantial impact on scale invariance, as evidenced by a good overall fit. Note that items were not specifically calibrated to the HWS group because of a limited sample size (n=23) [47]. No DIF was observed between the first and last assessments, showing satisfactory invariance to support the scale follow-up stability. Likewise, an intraclass correlation coefficient across the first

223 and last assessments was 0.94, indicating excellent item-difficulty-hierarchy consistency and

224 providing confidence for data pooling over different time points [48]. The PSI in this sample was

225 equal to 0.90, indicating the distinguishability of four strata of manual ability [46].

226 **Table 3. Differential item functioning (DIF) summary.**

Label	Person factor	Magnitude (logits)	Type	Difficulty
Taking the cap off a bottle	Gender	1.36	Uniform	Women > Men
Opening a screw-topped jar	Gender	0.93	Uniform	Women > Men
Opening a screw-topped jar	Diagnosis	0.61	Uniform	BTA > CTS > DRF
Using a screwdriver	Involved hand	1.26	Uniform	Non-dominant > Dominant hand involved

Fig 1. Differential item functioning (DIF) plots comparing the item difficulty hierarchy between

subgroups. In each plot, the lines represent the 95% confidence interval of an ideal invariance

between subgroups; the items are represented by the dots or by their letter if they display

significant DIF. The most difficult items (dots) are plotted in the top right part of each plot.

When comparing the item difficulty hierarchy between each diagnostic group relative to the

whole sample, most of the ABILHAND-HS items lie within 95% confidence interval of the ideal

invariance, indicating an invariant difficulty across diagnostic groups. When comparing the item

difficulty hierarchy between the first and last assessment, all items fall within the 95%

confidence interval of an ideal invariance, affirming invariance of item difficulties between the

assessments.

Scale description

The ABILHAND-HS structure and targeting of HS patients are illustrated in Fig 2, showing an

average patients' manual ability of 1.17 logits (SD = 1.85 logits; i.e. 58 (15) centiles). Twenty-four

patients (7.9%) were able to perform all 23 activities easily, and were thus identified as extreme

patients. Extreme patients tended to be younger men evaluated more than 6 months after

treatment, and were more likely to have a CTS rather than a HWS. The three response categories

were well distinguished in HS patients, with an inter-threshold distance of 2.93 logits (24 centiles),

indicating that, regardless of patient ability, rating an item as 'easy' is about 20 (i.e. $e^{2.93}$=18.7)

times more difficult than rating it as 'impossible'. Although the threshold distribution (range, -

4.15 to 5 logits) was well targeted to the range of patient abilities, the patients' ability levels

skewed high, indicating that the scale could measure patients that are more severely disabled

than in this sample.

Fig 2. Structure of the ABILHAND-HS scale. Top: distribution of manual ability measures for the

whole sample expressed in logits (log of the pass/fail probability ratio) and centiles (fraction of

the measurement range). Twenty-four patients (7.9%) were able to perform all 23 activities

easily, and were thus identified as extreme patients. None of the participants reported that they

could not perform any of the 23 activities. Middle: most probable patient response to each item

based on the patient manual ability and on the difficulty of the item's response category. The

average item difficulty was set to 0 logits and the items are ordered from most (top) to least

(bottom) difficult. The distance between thresholds (middle bar) is constant for all items (2.93

logits or 24 centiles). A patient with a manual ability measure of 0 logits would be expected to

perform the first 3 activities easily, to have some difficulty with the following 17 activities, and

to be unable to perform the 3 most difficult activities. A patient with a measure of 2.1 logits

should be able to perform all activities easily or with some difficulty. Bottom: conversion of

ordinal raw scores into a linear continuum of manual ability for complete response sets. The raw

scores ranged from 0 to 46 (sum of scores of 0–2 for 23 items). This curve is linear in its central

(30^{th}~70^{th} percentile) range, with sigmoid flattening outside the central range, highlighting a

non-linear relationship, especially at the extremities of the score range.

Construct validity

ABILHAND-HS measures were normally distributed across the whole sample and subgroups,

except for men (W = 0.97; 100 df; p = 0.038). An effect of gender on ABILHAND-HS manual ability

measures was observed, with men [1.88 (2.39) logits; median 1.74 logits] reporting a significantly

higher mean manual ability than women [1.32 (1.93) logits; median 1.4 logits; U = 8642; p =

0.026]. Manual ability was not found to be significantly associated with age (R = -0.04; p = 0.47),

the hand involved (t = 0.96; 303 df; p = 0.37), or the patient's diagnosis (F = 1.92; 3 df; p = 0.12).

Although variance across diagnosis groups was not significant, we did observe a broad spectrum

of manual ability. Patients with CTS reported the highest manual ability [1.9 (2.0) logits], followed

by patients with BTA [1.5 (2.3) logits], DRF [1.4 (2.1) logits], and HWS [0.9 (1.8) logits].

The relationships between ABILHAND-HS measures and scores obtained with other instruments

are shown in Fig 3. Briefly, ABILHAND-HS correlated strongly with QuickDASH scores, moderately

with SF-12 PCS scores and pain scale scores, and weakly with SF-12 MCS scores. We observed

substantial similarity with respect to manual ability scale locations between the ABLHAND-HS and

QuickDASH activity items (Fig 4).

Fig 3. Correlations of ABILHAND-HS scores with QuickDASH, PCS, MCS, and numerical pain

scale scores. Spearman correlation coefficients are indicated in the top right of each graph. All

correlations were statistically significant (p < 0.001).

Fig 4. Comparison of difficulty levels (vertical axis) between similar items of the ABILHAND-HS

(left) and QuickDASH (right) scales. QuickDASH item responses were added to the anchored data

matrix of ABILHAND-HS responses to equate both measures.

Discussion

Here, we report the adaptation and validation of an ABILHAND-HS questionnaire for use with HS

patients. Impairments present in our study cohort included weakness (e.g. following DRF), loss of

sensation (e.g. in CTS), and stiffness (e.g. in BTA), with some patients presenting with a

combination of these impairments. The ABILHAND-HS was constructed to measure manual ability

on a common, linear, and unidimensional scale wherein the 23 activities retained delineate an

invariant item difficulty hierarchy independent of patient diagnosis. All ABILHAND-HS activities

with the exception of one involve both hands and, consistent with our clinical experience, the

most difficult ones require high levels of force (e.g. 'doing push-ups' loads the wrist in extension).

Of the experimental 90 items, those that could be interpreted in different ways, for instance using

the injured or uninjured hand, were misfitting and thus omitted (e.g. 'carrying a shopping bag').

The sample size was adequate for the statistical interpretation of fit statistics [33], and was within

the same range of studies dealing with the development of outcome measures [14,15,19,31]. The

fit statistics for the 23 retained items support the item hierarchy invariance across the latent trait

[49]. A few instances of minor DIF were retained to maintain the scale's construct validity [50].

The resulting scale is well targeted to the studied HS population, despite a small persistent ceiling

effect, most likely due to missing responses for the most difficult activities. This observation of

apparent ceiling effect involves 7.9% (24/305) of the records, which is well below the maximum

recommended allowance of 15% [51].

Although reliability indices should be compared with caution across potentially different study

conditions, it is noteworthy that the PSI obtained for the ABILHAND-HS (0.90) was higher than

prior values obtained for the activities subscales of the PRWE [52] (0.78 and 0.81 for the usual

and specific activities subscales, respectively, in DRF patients), for the Patient-Rated Wrist and

Hand Evaluation [53] (0.83 in HS patients), for the QuickDASH scale [54] (0.84 in patients with

various upper limb dysfunctions) and for the Manual Ability Measure [55] (MAM-16; 0.83 for HS

patients), while being equal to that for the DASH manual functioning subscale [18]. PSI values

reflect sensitivity to clinical evolution over time, with greater values indicating a greater number

of distinguishable ability strata. We obtained person separation among patients using three

response levels (impossible, difficult, and easy), consistent with previous studies showing patients

unable to discriminate more than three levels of difficulty (ABILHAND [8], DASH activity items

[18] and QuickDASH activity items [54]).

Accurate communication of scale administration instructions is critical for targeting patient

manual ability as defined by the ABILHAND-HS. Generally, patients focus on their ability to

perform the queried activities with their injured hand; likewise, the PRWE explores use of the

affected hand explicitly [15]. The ABILHAND-HS, like the QuickDASH, is oriented towards real daily

life behaviors and is intended to be independent of the limb(s) or strategy used and unbiased by

activities that are never performed with the affected hand or avoided during recovery [8]. Our

findings of stable item calibrations and lack of DIF across the assessments indicate that the

ABILHAND-HS can be used confidently to assess the patient recovery at different time points

during follow-up. Moreover, the stability of items hierarchy between the first and last evaluation

indicate that the results were not influenced by the method of administration (interview with the

investigator versus self-reported).

ABILHAND-HS construct validation results fit well with our clinical observations. The patients with

the highest manual ability scores on the ABILHAND-HS also had the highest SF-12 PCS and SF-12

MCS scores as well as the lowest QuickDASH and numerical pain scale scores, which was also

observed in other validation studies [15,19]. Correlations of ABILHAND-HS with other

instruments, including the QuickDASH, SF-12 and a pain scale are also consistent with prior

findings suggesting that generic instruments are less sensitive than specific ones [56]. The present

ABILHAND-HS manual ability scores were not related significantly to age, consistent with other

versions of the ABILHAND [8,26,29]. Our findings of a small, but significant gender effect, with

men tending to report a higher manual ability than women (mean difference, 0.56 logits), varied

across HS diagnoses but, generally, were consistent with previous reports in patients with DRFs

and wrist arthrodesis [57–59]. A possible explanation is that manual ability is related to grip

strength, which is more important in men compared to women [8,60]. The construct validity of

the ABILHAND-HS was further supported by our confirmation of a similar item difficulty hierarchy

for QuickDASH items in our patients sample. Notably, those ABILHAND-HS activities that require

a great amount of force (e.g. 'Opening a screw-topped jar') have been reported to likewise be

among the most difficult items in the DASH and QuickDASH [18,54] and in the MAM-16 [55].

The ABILHAND-HS, developed using Rasch methodology, has several advantages over

questionnaires developed using classical test theory. These are summarized in Table 4. Firstly, the

ABILHAND-HS can tolerate missing responses, which enables it to remain valid even in patients

who scarcely perform some of the queried activities. Secondly, the ability to analyze response

patterns can identify those patients whose responses do not fit the model due to random or

careless answers, a particular injury or comorbidities. Finally, the high precision of the ABILHAND-

HS items minimizes the need for interpretation, thereby allowing more reliable comparisons

between patients (e.g. recreational activities involving force or impact are broken down into the

items 'doing push-ups', 'practicing a racket sport').

Table 4. Pros and cons of the ABILHAND-HS.

	Feature	Benefit
Pros	Unidimensional and linear scale	Quantitative comparisons of manual ability can be made between patients, between treatments and along follow up
	Invariant scale calibration validated in four diagnostic groups	Unbiased comparisons of manual ability can be made within and between clinical subgroups (different HS diagnostics, stage of recovery)
	Precise item definition	Inaccurate responses and guessing are avoided
	Amenable to incomplete responses	The test is specific to activities really performed by the patient
	Capacity to analyze response patterns	The test can identify unexpected patient responses linked to patient specific behaviors, random or careless answers or comorbidities
	Precision of the measure	The standard error of measurement is specific to each patient measure allowing statistical assessment during follow-up

	Feature	Compensation
Cons	Yet another test for hand surgery outcomes evaluation	Takes five minutes to complete
	Complex statistical background	Necessary for the analyst, but not for routine clinical use
	Use of dedicated computer programs	A free web service (www.rehab-scales.org) can be used to interpret patient responses
	Ceiling effect of 7.9% of records	Well below the maximum recommended allowance of 15%

Limitations of this research include a sample of patients with hand and wrist disabilities from one hand surgery outpatient clinic. The unbalanced diagnostic groups and genders might have influenced the item calibrations. However, the gender distribution is similar in studies involving DRF and CTS [15,16,31] and the DIF analysis allowed us to select items where the diagnosis and gender effects were absent. Future studies with a larger sample size should confirm or refine our

findings. Furthermore, the way participants responded to the 90 items in the questionnaire may

not be the same as responses to the final 23-item instrument. One limitation to the availability of

Rasch model-based questionnaires is that they have a complex statistical background and require

the use of dedicated computer programs that are not easy to learn and implement. To facilitate

and spread the use of the ABILHAND-HS, a website (www.rehab-scales.org) developed by

Université catholique de Louvain and Arsalis, a spin-off of the ABILHAND authors' laboratory, can

be used to convert the questionnaire raw scores into manual ability measures. The web service is

free-to-use for daily practice in clinical and research applications although a license is required

for commercial applications and for clinical trials.

Conclusion

ABILHAND-HS was demonstrated to be a successful adaptation for application in HS patients. The

resulting scale was shown to be a valid, patient-oriented, clinically meaningful and precise

instrument. It targets commonly performed manual activities and allows stable and linear

measurement of manual ability over multiple time points in patients treated for DRF, BTA, CTS,

or HWS. The scale reveals unexpected responses that may provide clues regarding the patient's

clinical state, as summarized at www.rehab-scales.org. The questionnaire is available online, and

the web service is free-to-use for daily practice in clinical and research applications although a

license is required for commercial applications and for clinical trials. Future research should

include more patients with HWS, as well as other diagnoses such as tendinopathies, ligamentous

injuries and complex hand injuries, and an assessment of scale responsiveness.

References

1. Dubert T. Outcome measurements in hand and upper limb surgery. Chirurgie de la Main. 2014;33:
235–246.

2. Berkanovic E, Hurwicz M-L, Lachenbruch PA. Concordant and discrepant views of patients' physical
functioning. Arthritis & Rheumatism. 1995;8: 94–101.

3. Hewlett SA. Patients and Clinicians Have Different Perspectives on Outcomes in Arthritis. J
Rheumatol. 2003;30: 877–9.

4. Ziebland S, Fitzpatrick R, Jenkinson C. Tacit models of disability underlying health status
instruments. Social science & medicine. 1993;37: 69–75.

5. World Health Organization, editor. International classification of functioning, disability and health:
ICF. Geneva: World Health Organization; 2001.

6. Barbier O, Penta M, Thonnard J-L. Outcome evaluation of the hand and wrist according to the
International Classification of Functioning, Disability, and Health. Hand Clinics. 2003;19: 371–378.

7. Bobos P, Lalone EA, Grewal R, MacDermid JC. Do Impairments Predict Hand Dexterity After Distal
Radius Fractures? A 6-Month Prospective Cohort Study. HAND. 2018;13: 441–447.

8. Penta M, Tesio L, Arnould C, Zancan A, Thonnard J-L. The ABILHAND Questionnaire as a Measure of
Manual Ability in Chronic Stroke Patients: Rasch-Based Validation and Relationship to Upper Limb
Impairment. Stroke. 2001;32: 1627–1634.

9. Giladi AM, Chung KC. Measuring Outcomes in Hand Surgery. Clinics in Plastic Surgery. 2013;40:
313–322.

10. Jaremko JL, Lambert RGW, Rowe BH, Johnson JA, Majumdar SR. Do radiographic indices of distal
radius fracture reduction predict outcomes in older adults receiving conservative treatment?
Clinical Radiology. 2007;62: 65–72.

11. Synn AJ, Makhni EC, Makhni MC, Rozental TD, Day CS. Distal Radius Fractures in Older Patients: Is
Anatomic Reduction Necessary? Clin Orthop Relat Res. 2009;467: 1612–1620.

12. Young BT, Rayan GM. Outcome following nonoperative treatment of displaced distal radius
fractures in low-demand patients older than 60 years. The Journal of Hand Surgery. 2000;25: 19–
28.

13. Changulani M, Okonkwo U, Keswani T, Kalairajah Y. Outcome evaluation measures for wrist and
hand – which one to choose? International Orthopaedics. 2008;32: 1–6. doi:10.1007/s00264-007-
0368-z

14. Hudak PL, Amadio PC, Bombardier C. Development of an upper extremity outcome measure: the
DASH (disabilities of the arm, shoulder and hand) [corrected]. The Upper Extremity Collaborative
Group (UECG). Am J Ind Med. 1996;29: 602–608.

15.   MacDermid JC, Turgeon T, Richards RS, Beadle M, Roth JH. Patient rating of wrist pain and
disability: a reliable and valid measurement tool. J Orthop Trauma. 1998;12: 577–586.

16.   Levine DW, Simmons BP, Koris MJ, Daltroy LH, Hohl GG, Fossel AH, et al. A self-administered
questionnaire for the assessment of severity of symptoms and functional status in carpal tunnel
syndrome. JBJS. 1993;75: 1585–1592.

17.   Coenen M, Kus S, Rudolf K-D, Müller G, Berno S, Dereskewitz C, et al. Do patient-reported outcome
measures capture functioning aspects and environmental factors important to individuals with
injuries or disorders of the hand? Journal of Hand Therapy. 2013;26: 332–342.

18.   Franchignoni F, Giordano A, Sartorio F, Vercelli S, Pascariello B, Ferriero G. Suggestions for
Refinement of the Disabilities of the Arm, Shoulder and Hand Outcome Measure (DASH): A Factor
Analysis and Rasch Validation Study. Archives of Physical Medicine and Rehabilitation. 2010;91:
1370–1377.

19.   Chung KC, Pillsbury MS, Walters MR, Hayward RA. Reliability and validity testing of the Michigan
Hand Outcomes Questionnaire. The Journal of Hand Surgery. 1998;23: 575–587.

20.   Merbitz C, Morris J, Grip JC. Ordinal scales and foundations of misinference. Archives of physical
medicine and rehabilitation. 1989;70: 308–312.

21.   Wright BD, Linacre JM. Observations are always ordinal; measurements, however, must be interval.
Arch Phys Med Rehabil. 1989;70: 857–860.

22.   Thurstone L. The measurement of values. Chicago: The University of Chicago Press; 1959.

23.   Grimby G, Tennant A, Tesio L. The use of raw scores from ordinal scales: Time to end malpractice?
Journal of Rehabilitation Medicine. 2012;44: 97–98.

24.   Rasch G. Probabilistic models for some intelligence and attainment tests. Chicago, IL: The
University of Chicago Press; 1980.

25.   Penta M, Thonnard J-L, Tesio L. ABILHAND: a Rasch-built measure of manual ability. Archives of
physical medicine and rehabilitation. 1998;79: 1038–1042.

26.   Durez P, Fraselle V, Houssiau F, Thonnard J, Nielens H, Penta M. Validation of the ABILHAND
questionnaire as a measure of manual ability in patients with rheumatoid arthritis. Ann Rheum Dis.
2007;66: 1098–1105.

27.   Arnould C, Penta M, Renders A, Thonnard J-L. ABILHAND-Kids: a measure of manual ability in
children with cerebral palsy. Neurology. 2004;63: 1045–1052.

28.   Vanthuyne M, Smith V, Arat S, Westhovens R, de Keyser F, Houssiau FA, et al. Validation of a
manual ability questionnaire in patients with systemic sclerosis. Arthritis & Rheumatism. 2009;61:
695–703.

29. Vandervelde L, Van den Bergh PYK, Penta M, Thonnard JL. Validation of the ABILHAND
questionnaire to measure manual ability in children and adults with neuromuscular disorders.
Journal of Neurology, Neurosurgery & Psychiatry. 2010;81: 506–512.

30. Arnould C, Vandervelde L, Batcho CS, Penta M, Thonnard J-L. Can manual ability be measured with
a generic ABILHAND scale? A cross-sectional study conducted on six diagnostic groups. BMJ open.
2012;2: e001807.

31. Beaton DE, Wright JG, Katz JN. Development of the QuickDASH: Comparison of Three Item-
Reduction Approaches. Journal of Bone and Joint Surgery (American). 2005;87A: 9.

32. Ware J, Kosinski M, Keller SD. A 12-Item Short-Form Health Survey: construction of scales and
preliminary tests of reliability and validity. Med Care. 1996;34: 220–233.

33. Hagell P, Westergren A. Sample Size and Statistical Conclusions from Tests of Fit to the Rasch
Model According to the Rasch Unidimensional Measurement Model (Rumm) Program in Health
Outcome Measurement. J Appl Meas. 2016;17: 416–431.

34. Andrich D. A rating formulation for ordered response categories. Psychometrika. 1978;43: 561–
573.

35. Masters GN. A rasch model for partial credit scoring. Psychometrika. 1982;47: 149–174.

36. van Nes SI, Vanhoutte EK, van Doorn PA, Hermans M, Bakkers M, Kuitwaard K, et al. Rasch-built
Overall Disability Scale (R-ODS) for immune-mediated peripheral neuropathies. Neurology.
2011;76: 337–345.

37. Andrich D, Sheridan B, Luo G. Polytomous data. In: Interpreting RUMM2030: part II. Western
Australia: Perth; 2013.

38. Linacre JM. Optimizing rating scale category effectiveness. J Appl Meas. 2002;3: 85–106.

39. Holland PW, Wainer H. Differential item functioning. Hillsdale, NJ, US: Lawrence Erlbaum
Associates, Inc; 1993.

40. Armstrong RA. When to use the Bonferroni correction. Ophthalmic Physiol Opt. 2014;34: 502–508.

41. Linacre JM. Detecting multidimensionality: which residual data-type works best? J Outcome Meas.
1998;2: 266–283.

42. Pallant JF, Tennant A. An introduction to the Rasch measurement model: an example using the
Hospital Anxiety and Depression Scale (HADS). Br J Clin Psychol. 2007;46: 1–18.

43. Smith EV. Detecting and evaluating the impact of multidimensionality using item fit statistics and
principal component analysis of residuals. J Appl Meas. 2002;3: 205–231.

44. Wright BD. Local dependency, correlations and principal components. Rasch Meas Trans
1996;10:509–11.

45. Ramp M, Khan F, Misajon RA, Pallant JF. Rasch analysis of the Multiple Sclerosis Impact Scale (MSIS-
29). Health Qual Life Outcomes. 2009;7: 58.

46. Fisher W. Reliability, Separation, Strata Statistics. Rasch Meas Trans. 1992;6: 238.

47. Chen W-H, Lenderking W, Jin Y, Wyrwich KW, Gelhorn H, Revicki DA. Is Rasch model analysis
applicable in small sample size pilot studies for assessing item characteristics? An example using
PROMIS pain behavior item bank data. Qual Life Res. 2014;23: 485–493.

48. Chang W-C, Chan C. Rasch analysis for outcomes measures: some methodological considerations.
Archives of Physical Medicine and Rehabilitation. 1995;76: 934–939.

49. Tennant A, Conaghan PG. The Rasch measurement model in rheumatology: What is it and why use
it? When should it be applied, and what should one look for in a Rasch paper? Arthritis &
Rheumatism. 2007;57: 1358–1362.

50. Hagquist C, Andrich D. Recent advances in analysis of differential item functioning in health
research using the Rasch model. Health and Quality of Life Outcomes. 2017;15: 181.

51. McHorney CA, Tarlov AR. Individual-patient monitoring in clinical practice: are available health
status surveys adequate? Quality of Life Research. 1995;4: 293–307.

52. Esakki S, MacDermid JC, Vincent JI, Packham TL, Walton D, Grewal R. Rasch analysis of the patient-
rated wrist evaluation questionnaire. Arch Physiother. 2018;8:5.

53. Packham T, MacDermid JC. Measurement properties of the Patient-Rated Wrist and Hand
Evaluation: Rasch analysis of responses from a traumatic hand injury population. Journal of Hand
Therapy. 2013;26: 216–224.

54. Franchignoni F, Ferriero G, Giordano A, Sartorio F, Vercelli S, Brigatti E. Psychometric properties of
QuickDASH – A classical test theory and Rasch analysis study. Manual Therapy. 2011;16: 177–182.

55. Chen CC, Granger CV, Peimer CA, Moy OJ, Wald S. Manual Ability Measure (MAM-16): a
preliminary report on a new patient-centred and task-oriented outcome measure of hand function.
Journal of Hand Surgery. 2005;30: 207–216.

56. Aktekin LA, Eser F, Başkan BM, Sivas F, Malhan S, Öksüz E, et al. Disability of Arm Shoulder and
Hand Questionnaire in rheumatoid arthritis patients: relationship with disease activity, HAQ, SF-36.
Rheumatol Int. 2011;31: 823–826.

57. Amorosa LF, Vitale MA, Brown S, Kaufmann RA. A Functional Outcomes Survey of Elderly Patients
who Sustained Distal Radius Fractures. Hand (N Y). 2011;6: 260–267.

58. Cowie J, Anakwe R, McQueen M. Factors Associated with One-Year Outcome after Distal Radial
Fracture Treatment. Journal of Orthopaedic Surgery. 2015;23: 5.

59. Owen DH, Agius PA, Nair A, Perriman DM, Smith PN, Roberts CJ. Factors predictive of patient
outcome following total wrist arthrodesis. The Bone & Joint Journal. 2016;98-B: 647–653.

60.   Arnould C, Penta M, Thonnard J. Hand impairments and their relationship with manual ability in
517   children with cerebral palsy. Journal of Rehabilitation Medicine. 2007;39: 708–714.

Supporting information

S1 Text. Common versus specific scales.

S2 Text. The ABILHAND-HS questionnaire (English version) instructions and scoring sheets. The scoring sheets are available in 10 different randomizations, as found on www.rehab-scales.org.

S3 Text. The ABILHAND-HS questionnaire (French version) instructions and scoring sheets. The scoring sheets are available in 10 different randomizations, as found on www.rehab-scales.org.

S4 Appendix. The experimental 90-item ABILHAND questionnaire.