

1 **Impact of anxiety associated with COVID-19 on tinnitus**

2

3 Li Xia<sup>1</sup>, Gang He<sup>1</sup>, Yong Feng<sup>1</sup>, Xiaoxu Yu<sup>1</sup>, Xiaolong Zhao<sup>1</sup>, Zhengnong Chen<sup>2</sup>, Shankai  
4 Yin<sup>2</sup>, Jian Wang<sup>3\*</sup>, Jiangang Fan<sup>1\*</sup>, Chuan Dong<sup>1\*</sup>

5

6 1 Department of Otolaryngology-Head and Neck Surgery, Sichuan Provincial  
7 People's Hospital and Sichuan Academy of Medical Sciences, Chengdu, Sichuan, 610072,  
8 China.

9 2 Department of Otolaryngology-Head and Neck Surgery, Shanghai Jiao Tong  
10 University Affiliated Sixth People's Hospital, 200233, Shanghai.

11 3 School of Communication Science and Disorders, Dalhousie University, Halifax,  
12 Nova Scotia B3H 4R2, Canada.

13 \*Corresponding authors: Chuan Dong (dc1220@sina.com) and Jian Wang  
14 (jian.wang@dal.ca) and Jiangang Fan (entscfjg@163.com)

15

16

17

18

19

20

21

22

23 **Abstract**

24 **Background**

25 We investigated how the anxiety associated with COVID-19 impacts the severity of  
26 tinnitus and the outcomes of tinnitus therapy.

27 **Methods and Findings**

28 A retrospective research design was used to compare the clinical characteristics of  
29 tinnitus between patients from March to April 2020 under pandemic pressure and those  
30 from the matching period in 2019. Tinnitus severity was evaluated using the Tinnitus  
31 Handicap Inventory (THI) questionnaire and the minimum masking level (MML)  
32 measure while anxiety was quantified using the Zung's Self-rating Anxiety Scale (SAS).  
33 The assessments were repeated after a 2-month interval, in which sound therapy was  
34 applied to a subgroup of patients. In all, 94 and 70 cases were reported in the 2020 and  
35 2019 groups, respectively. The effects of the pandemic on emotional status were evident  
36 from a higher incidence of anxiety and much higher SAS scores in the 2020 group. There  
37 was also an increase in the THI scores in the 2020 group, but the between-year difference  
38 was smaller than that of the SAS score. Moreover, there was no between-year difference  
39 in MML or the treatment effect, as measured via both THI and MML. Furthermore, the  
40 reduction in SAS score in the second assessment was significantly smaller in the 2020  
41 group. However, a positive correlation between the initial SAS score and the  
42 improvement was seen within the 2020 group.

43 **Conclusions**

44 Anxiety increased greatly in tinnitus subjects due to the COVID-19 pandemic. However,  
45 this was not associated with an increase in tinnitus severity in 2020. Instead, there was no

46 between-year difference in the THI score or MML or in the improvement of either  
47 measure after treatment. The smaller improvement in SAS score and the positive  
48 correlation with the initial SAS score in the 2020 group suggests that the SAS score  
49 change in this group might largely be due to the natural relief of pressure as the pandemic  
50 decelerated in China. Therefore, the anxiety change induced by the COVID-19 pandemic  
51 is not likely to have a strong impact on tinnitus.

## 52 **Introduction**

53       The spread of coronavirus disease 2019 (COVID-19) has already reached pandemic  
54 proportions, affecting the majority of countries, areas, and territories across the world [1] .  
55 By the end of June 2020, over nine million people had tested positive for COVID-19 with  
56 the death toll increasing to more than 484,000 globally [2] . Decisive containment  
57 measures in China have reduced new cases and the spread of infection [3]. However,  
58 worries about the spread of the disease, living difficulties, and financial burden related to  
59 the pandemic are likely to have had negative psychosocial impacts on residents, as  
60 reported by many recent studies [4-6]. It would be reasonable, therefore, to expect an  
61 increase in the incidence of disorders that are associated with psychological issues.

62       Tinnitus is typically referred to as the perception of sound in the absence of an  
63 acoustic stimulus or that is only generated by structures in the ear, commonly described  
64 as ringing in one or both ears [7]. While the exact mechanisms of tinnitus remain unclear,  
65 many risk or promoting factors have been identified, including sensorineural hearing loss,  
66 vestibular schwannoma, ototoxic medications, and emotional stress [8]. Tinnitus has been  
67 linked to stress and related disorders in many previous studies. This link has been  
68 thoroughly reviewed, repeatedly, by different authors (e.g., [9-15]). The direction and  
69 causality of this link remain unclear, as pointed out in many previous studies, although  
70 individuals' emotional states appear to be an important factor mediating the effects of  
71 tinnitus loudness on tinnitus-related distress [16-18]; anxiety, somatization, and in  
72 particular depression have also been identified as possible mediators of tinnitus-related  
73 distress [19-22].

74       The clinicians in our department noticed that the tinnitus patients seen since the

75 hospital was reopened after COVID-19 had more emotional complaints than before. We  
76 thought that this might be related to the various pressures experienced by the patients  
77 during the pandemic event and the lockdown. Therefore, the COVID-19 pandemic and  
78 lockdown might provide a good opportunity to investigate whether anxiety impacts  
79 tinnitus as a promoting or enhancing factor. The present study explored whether anxiety  
80 was increased by the COVID-19 pandemic in subjects with tinnitus, and if so whether  
81 this increase in anxiety affected the severity of tinnitus and the outcomes of tinnitus  
82 therapies.

83


## 84 **Methods**

### 85 **Study Design**

86 In this retrospective study, clinical data from outpatients visiting our department (the  
87 Hearing Center of Otolaryngology Department of the Sichuan Provincial People's  
88 Hospital and Sichuan Academy of Medical Sciences, Chengdu, Sichuan, People's  
89 Republic of China) were collected over the same periods, from March 1 to April 14, in  
90 both 2020 and 2019. This period in 2020 was the first 6 weeks of the reopening of our  
91 department to non-emergency visits after the nationwide lockdown for COVID-19 in  
92 China (from January 23 to February 29, 2020) that coincided with the deceleration phase  
93 of the pandemic and the resumption of economic activities. In this period, there were  
94 concerns about a resurgence of COVID-19 [23].

95 The same protocol was followed for the treatment of patients during both years. On  
96 the initial visit, after collecting their history, every patient received a comprehensive  
97 audiological and psychological assessment. After the assessment, they were treated with

98 one of three methods based on reported efficacy, financial cost, and the patient's  
99 preference: sound therapy with educational counseling and relaxation therapy, sound  
100 amplification with educational counseling and relaxation therapy, or educational  
101 counseling and relaxation therapy only. Two months after the initial appointment, every  
102 participant was examined in a second assessment. Fig 1 shows a flowchart of the major  
103 procedures of this study. Although no procedure was experimental, we sought and  
104 received approval for the study from the Review Board of the Sichuan Provincial  
105 People's Hospital and Sichuan Academy of Medical Sciences (permit number: 2020–355).  
106 This study was conducted according to the principles expressed in the Declaration of  
107 Helsinki [24]



108

109 **Fig 1. Flowchart of the major procedures in this study.** \*numbers in parentheses are those of cases that  
110 were lost to the study. THI: tinnitus handicap inventory, SAS: Zung's Self-Rating Anxiety Sale, ST: sound  
111 therapy, HA: hearing aid, EC: educational counseling.

112

### 113 **Audiological Tests and Tinnitus Evaluation**

114 The procedures for all tests were explained to the patients before they were

115 conducted. All patients were examined using monocular otoscopy to identify any sign of  
116 blockage or inflammation in ear canals or perforation in the tympanic membrane.  
117 Tympanometry was tested at the most common 226 Hz probe tone, using an AT235  
118 impedance meter (Interacoustics, Assens, Denmark); the type of tympanogram was  
119 determined for each ear (with type A as normal). Those who were abnormal in those tests  
120 were not included in this study.

121 The hearing status was tested with pure-tone audiometry (AC40, Interacoustics) in a  
122 soundproofed room. The air conduction threshold was examined for frequencies ranging  
123 from 250 Hz to 8 kHz using TDH 39 headphones (Telephonics, NY, USA) and bone  
124 conduction hearing was examined from 500 Hz to 4 kHz using a B-72 bone-conduction  
125 vibrator (Radioear, PA, USA), each in octave steps. The hearing thresholds were  
126 determined at each frequency using the standard Hughson–Westlake up–down procedure.  
127 Thresholds of 20 dB HL or lower were considered normal. The minimum masking level  
128 (MML) was tested in each ear with tinnitus, this test evaluates the maskability of tinnitus  
129 by external sounds. Broadband noise with a flat power spectrum was used for this  
130 evaluation, which was generated by a table-top sound generator (BTD01, BetterLife  
131 Medical Technology Co., Ltd., Jiangsu, China). To measure the MML, the level of the  
132 noise was gradually increased by the tester in 1 dB steps until the patient stated that the  
133 tinnitus had become nearly inaudible: then this level was recorded as the MML.

134

### 135 **Educational Counseling and Relaxation Therapy**

136 The counseling was performed by the audiologists for each patient with tinnitus to  
137 acknowledge the patient’s suffering, and to help the patient understand tinnitus,



138 demystify the condition, and correct any false preconceptions (duration 0.5–1 h) [25].  
139 Relaxation therapy consisted of home-based exercises, such as listening to music,  
140 avoiding unnecessary tension, and tai chi [26, 27]. Patients were advised to execute this  
141 for two sessions of 30 min per day over a period of 8 weeks.

142

### 143 **Sound Therapy**

144 The first step of the sound therapy was to identify the nature of the tinnitus in pitch  
145 and loudness. Pitch matching was conducted using the same sound generator (BTD01) as  
146 in the MML test to produce pure tones for tonal tinnitus or narrow-band noise for non-  
147 tonal tinnitus. The match was established by adjusting the central frequency and  
148 bandwidth, which could be changed from 100 Hz to 1 kHz, around the center frequency.  
149 In loudness matching, the matched tone or noise was presented continuously, and the  
150 level of the matching signal was adjusted from low to high until the tinnitus could hardly  
151 be heard. In this report, loudness matching results are presented in dB SL. Using the pitch  
152 and loudness matching data, a sound file was generated for each individual to produce a  
153 sound matching their tinnitus in frequency and level. This sound file was the uploaded to  
154 an ear level sound generator (BTM-N6, BetterLife Medical Technology Co., Ltd.) that  
155 was dispensed to the patient. The patients were instructed to listen to the sound file for  
156 30 min each time, and to gradually increase from once to 3–6 times per day, every day,  
157 during the whole course of home-based therapy, which lasted for 2 months.

158

### 159 **Questionnaires**

160 The tinnitus patients recruited in this study all completed two questionnaires at the

161 initial visit and again during the follow-up, two months later. The Chinese version of the  
162 Tinnitus Handicap Inventory (THI) questionnaire was used in this study [28], consisting  
163 of 25 questions to assess the difficulty caused by tinnitus with respect to its functional,  
164 emotional, and catastrophic aspects [29, 30].

165 A Chinese version of Zung's Self-rating Anxiety Scale (SAS) questionnaire was used,  
166 which was adapted from a previous report [31, 32]. The raw scores were multiplied by  
167 1.25 to generate the index scores [32]. We used a value of 45 as the cut-off for anxiety,  
168 instead of 50, as reported in the most recent publication [33].

169

## 170 **Statistical Analyses**

171 All parametric data are presented as mean  $\pm$  standard deviation unless otherwise  
172 specified. When the parameters of participants were compared between two groups, the t-  
173 test was used or, if among multiple groups, analysis of variance (ANOVA) was used for  
174 continuous variables and the chi-square test for categorical variables, including sex, age,  
175 and site of tinnitus, and for risk factors among groups. Treatment outcomes were  
176 evaluated by comparing the scores of THI and SAS before and after the treatments, using  
177 a paired t-test or ANOVA. All analyses were performed using the SPSS 19.0.0 software at  
178 a significance level of 0.05.

179

## 180 **Results**

181 A total of 99 cases were collected between March 1 and April 14, 2020 (this year),  
182 and 76 in the same period in 2019 (Fig 1). Table 1 compares the demographics and  
183 tinnitus characteristics between the 2 years' groups. The case load for tinnitus appeared to

184 be higher in 2020 than in the same period in 2019 (99 vs. 76, or an increase of 30.3%).  
 185 Such an increase could be fully attributed to a normal fluctuation due to the temporary  
 186 pause on non-emergency patients in our hospital between January and February 2020.  
 187 The two groups were matched by age and sex despite their different sizes.

188 Table 1. Comparison of initial clinical characteristics of patients between 2020 and 2019  
 189

|  | March-April 2020 | March-April 2019 | <i>p</i> -value |
|--|------------------|------------------|-----------------|
| Sex (M:F)                              | 43:56            | 39:37            | .3              |
| Age (years, mean ± standard deviation) | 50.8 ± 15.1      | 51.5 ± 14.8      | .834            |
| Educational background                 |                  |                  | .658            |
| Bachelor's degree and above            | 54               | 44               |                 |
| Less than bachelor's degree            | 45               | 32               |                 |
| Duration (month)                       | 25 ± 53.6        | 31.8 ± 51.0      | .064            |
| Site                                   |                  |                  | .02             |
| Bilateral                              | 36               | 41               |                 |
| Unilateral                             | 63               | 35               |                 |
| Anxiety involved/total #               | 74/99 (74%)      | 41/76 (53%)      | .004            |
| Risk factors                           |                  |                  |                 |
| Sensorineural hearing loss             | 69               | 58               | .331            |
| Noise exposure                         | 1                | 0                | 1               |
| Hypertension                           | 3                | 6                | .179            |
| Hyperthyroidism                        | 1                | 0                | 1               |
| Head/neck trauma                       | 1                | 0                | 1               |


190  
 191 The chi-square test was used for the between-group comparison of sex, educational background, site,  
 192 anxiety, and the risk factor of sensorineural hearing loss; the t-test was used for age; a Mann–Whitney rank-  
 193 sum test was used for duration; and a Fisher's exact test was used for the risk factors of noise exposure,  
 194 hypertension, hyperthyroidism, and head or neck trauma.

195

### 196 **Impact of Anxiety on THI and MML**

197 In the 2020 group, 74 out of 99 subjects had an SAS score higher than 45 (the  
 198 criterion for anxiety), thus the incidence of anxiety was 74.7% in this group, significantly  
 199 higher than that in the 2019 group (41/76, 53.9%,  $\chi^2 = 8.256$ ,  $p = 0.004$ ). To further  
 200 investigate the difference in the involvement of anxiety in tinnitus, a two-way ANOVA

201 was performed on the SAS scores against the factors of year group (2020 vs. 2019) and  
202 anxiety (with anxiety vs. without). Overall, the group had a significant effect, with the  
203 2020 group having higher SAS scores ( $61.9 \pm 11.9$ ) than the 2019 group ( $48.2 \pm 8.2$ ;  
204  $F_{1, 171} = 74.684$ ,  $p < 0.001$ , Fig 2A). The anxiety effect also significantly interacted with  
205 the group effect ( $F_{1, 171} = 44.486$ ,  $p < 0.001$ ). The average SAS score was significantly  
206 higher in 2020 patients than in 2019 patients, according to a *post hoc* pairwise test using  
207 the Tukey method ( $68.0 \pm 6.4$  in 2020 vs.  $53.5 \pm 7.8$  in 2019;  $q = 18.146$ ,  $p < 0.001$ ,  
208 Fig 2B); this method was used throughout for all such tests. This result suggests that the  
209 COVID-19 outbreak not only increased the incidence of anxiety but also the degree of  
210 anxiety in those who suffered from it. No significant differences in the averaged SAS  
211 score was seen between the patients without anxiety in 2020 and 2019 ( $43.9 \pm 1.5$  vs.  
212  $42.0 \pm 2.2$ ;  $q = 1.738$ ,  $p = 0.219$ ; Figs 2A and 2B).



213

214 Fig 2. Initial comparisons of SAS scores, THI scores and MML against year group and anxiety.

215 (A) The SAS score changes with the two main factors (year group and anxiety) and significance was found  
216 for both. (B) The year difference in SAS score in subgroups with and without anxiety. The SAS scores were  
217 higher in the 2020 subgroup with anxiety compared to those in 2019; there was no between-year difference  
218 in the subgroups without anxiety. (C) The THI score changes with the two main factors. (D) The year  
219 difference in THI in subgroups with and without anxiety. The THI increase in 2020 was seen only in the  
220 subgroup with anxiety. (E) The MML score changes with the two main factors. Overall, MML was  
221 significantly higher in the 2020 group, and there were no differences in MML between subjects with and  
222 without anxiety. (F) *Post hoc* comparison of MML. Within 2020, subjects with anxiety appeared to have a  
223 significantly higher MML, while the opposite was seen in 2019. Moreover, the subjects without anxiety in  
224 2020 showed a lower MML than the 2019 subjects without anxiety. Significance: \* $p < 0.05$ , \*\*  $p < 0.01$ ,  
225 \*\*\*  $p < 0.001$ . THI: tinnitus handicap inventory, SAS: Zung's Self-Rating Anxiety Sale, MML: minimum  
226 masking levels.

227 The THI score in the 2020 group was  $40.1 \pm 6.9$ , significantly higher than that in the  
228 2019 group ( $33.6 \pm 8.7$ ) as shown by the group effect in a two-way ANOVA  
229 ( $F_{1,171} = 15.223$ ,  $p < 0.001$ ), which also demonstrated a significant effect for anxiety:  
230  $39.1 \pm 8.5$  for subjects with anxiety and  $33.9 \pm 6.9$  for those without ( $F_{1,171} = 8.697$ ,  $p =$   
231  $0.004$ ). However, there was not a significant interaction between two factors  
232 ( $F_{1,171} = 2.867$ ,  $p = 0.092$ ). From a *post hoc* pairwise comparison in the subjects with  
233 anxiety, the THI score in 2020 was  $41.7 \pm 7.7$ , which was significantly higher than in  
234 those with anxiety in 2019 ( $34.3 \pm 8.3$ ;  $q = 6.631$ ,  $p < 0.001$ ) and those without anxiety in  
235 2020 ( $35.6 \pm 5.0$ ;  $q = 4.63$ ,  $p < 0.05$ ). However, there were no significant differences in  
236 THI score between the years in subjects without anxiety (Fig 2C and 2D).

237 The between-year difference in THI was further analyzed using a breakdown of the  
238 scores in the emotional, functional, and catastrophic questionnaire sections. A significant  
239 difference was seen between years in the emotional score using the Mann–Whitney rank-

240 sum test ( $14.636 \pm 3.7$  in 2020 and  $12.211 \pm 3.5$  in 2019,  $U = 5130$ ,  $p < 0.001$ ) and in the  
241 functional score ( $18.515 \pm 3.6$  in 2020 and  $15.342 \pm 4.5$  in 2019,  $U = 5353.5$ ,  $p < 0.001$ ),  
242 but not in the catastrophic scores ( $7.0 \pm 2.5$  vs.  $6.0 \pm 2.5$ ,  $U = 4462$ ,  $p = 0.031$ ). This  
243 result suggests that the increase in THI in 2020 could be partially related to the increase  
244 in anxiety.

245 A two-way ANOVA was performed on MMLs against both year group and anxiety.  
246 The subjects in 2020 had significantly lower MMLs ( $8.3 \pm 3.5$  dB SL) compared to those  
247 in the 2019 group ( $10.0 \pm 4.2$  dB SL;  $F_{1,171} = 15.281$ ,  $p < 0.001$ ), while the effects of  
248 psychological status were not significant ( $F_{1,171} = 0.569$ ,  $p = 0.452$ ; Fig 2E). However,  
249 the interaction between anxiety and year was significant ( $F_{1,171} = 19.184$ ,  $p < 0.001$ ).  
250 *Post hoc* pairwise tests showed that 2019 patients without anxiety had a higher MML  
251 ( $11.6 \pm 4.3$  dB SL) than the patients with anxiety in 2020 ( $8.6 \pm 3.6$  dB SL,  $q = 5.148$ ,  $p <$   
252  $0.001$ ). However, the opposite was seen in 2020, in that patients with anxiety had an  
253 MML of  $8.9 \pm 3.7$  dB SL, which was slightly but significantly higher than those without  
254 anxiety in 2020 ( $6.7 \pm 2.0$  dB SL,  $q = 3.616$ ,  $p \leq 0.01$ ; Fig 2F). Furthermore, the 2019  
255 subgroup without anxiety showed a higher MML than the 2019 subgroup with anxiety  
256 and the 2020 subgroup without (Fig 2F). Considering all of these analyses together, no  
257 clear indication can be seen as to whether anxiety played a role in the loudness of tinnitus.  
258

## 259 **Treatment Outcomes**

260 The 94 patients in the 2020 group completed their face-to-face follow-up 2 months  
261 after the first assessment, while this number was 70 in the 2019 group (Fig 1). The  
262 numbers of patients who received sound therapy, a hearing aid, or counseling alone were

263 38, 14, and 42, respectively, while the respective numbers were 58, 10, and 2 in the 2019  
 264 group. Due to the large difference in the number of patients requesting counseling alone  
 265 between the years and the small sample sizes in patients receiving hearing aids, we  
 266 analyzed only the treatment outcomes for sound therapy.

267

268 Table 2. Between-year comparison of the characteristics of tinnitus patients receiving sound therapy  
 269

|   | May-June 2020 | May-June 2019 | p-value |
|---|---------------|---------------|---------|
| Sex (M:F)                                 | 16:22         | 30:28         | .356    |
| Age (year old, mean ± standard deviation) | 48.9 ± 16.5   | 50.2 ± 14.1   | .871    |
| Educational background                    |               |               | .658    |
| Bachelor and superior                     | 21            | 36            |         |
| Inferior to bachelor                      | 17            | 22            |         |
| Duration (month)                          | 31.8 ± 54.3   | 25.5 ± 43.7   | .428    |
| Site                                      |               |               | .312    |
| Bilateral                                 | 18            | 32            |         |
| Unilateral                                | 20            | 26            |         |
| Anxiety involved/total #                  | 29/38 (76%)   | 32/58 (55%)   | .035    |
| Risk factors                              |               |               |         |
| Sensorineural hearing loss                | 24            | 40            | .555    |
| Noise exposure                            | 0             | 0             | \       |
| Hypertension                              | 1             | 6             | .396    |
| Hyperthyroidism                           | 0             | 0             | \       |
| Head/neck trauma                          | 0             | 0             | \       |

270

271 The chi-square test was used for between-group comparisons of sex, educational background, site, anxiety,  
 272 and the risk factor of sensorineural hearing loss; the t-test was used for age; and the Fisher's exact test was  
 273 used for the risk factor of hypertension.

274

275 As shown in Table 3, out of the 38 patients treated in 2020, 27 (71%) reported a  
 276 significant reduction of tinnitus loudness from the treatment, which was significantly  
 277 lower than the rate in 2019 in which 51 out of 58 (87.9%) reported a mitigation of  
 278 tinnitus ( $\chi^2 = 4.293$ ,  $p = 0.038$ ). This rate was slightly lower in the subjects with anxiety  
 279 in 2020, in which only 20 out of 29 subjects (68.9%) reported an improvement, while it



280 was 90.6% (29/32) in 2019 subjects with anxiety ( $\chi^2 = 4.516$ ,  $p = 0.034$ ). There was no  
 281 significant between-year difference in this rate in subjects without anxiety (7 out of 9 or  
 282 80.3% in 2020 and 22 out of 26 or 82.8% in 2019;  $\chi^2 = 0.93$ ,  $p = 0.76$ ). These results  
 283 suggest that the increased anxiety probably reduced the effectiveness of the treatment.

284

285 Table 3. Self-reported improvement during follow-up from tinnitus treatment by sound therapy

286


|              | <b>Overall</b> | <b>Anxiety</b> | <b>No Anxiety</b> | <b>Anxiety effect within year</b> |
|--------------|----------------|----------------|-------------------|-----------------------------------|
| 2020         | 27/38          | 20/29          | 7/9               | $p = 1^*$                         |
| 2019         | 51/58          | 29/32          | 22/26             | $p = 0.689^*$                     |
| Group effect | $p = 0.038$    | $p = 0.034$    | $p = 0.635^*$     |                                   |

287

288 \*comparison via Fisher's exact test; all other comparisons were conducted using the chi-square test

289

290 To fully evaluate the effects of the sound treatment, the changes in SAS, THI, and  
 291 MML scores were compared between years, as shown in Fig 3. First, the SAS score was  
 292 significantly reduced in both years after treatment, as indicated by the significant  
 293 treatment effect in two-way repeated measure (RM) ANOVA ( $F_{1,94} = 71.614$ ,  $p < 0.001$ ).  
 294 However, the SAS score was generally higher in 2020 than in 2019, as indicated by the  
 295 significant year effect in the ANOVA ( $F_{1,94} = 61.19$ ,  $p < 0.001$ ; Fig 3A). There was no  
 296 significant interaction between the factors of year and treatment. The SAS score was  
 297 reduced in both years in *post hoc* tests for the effects of the treatment ( $q = 6.911$ ,  $p <$   
 298  $0.001$  in 2020 and  $q = 10.48$ ,  $p < 0.001$  in 2019). Interestingly, however, the SAS score of  
 299 the 2020 group after being reduced by the treatment ( $58.0 \pm 10.6$ ) was significantly  
 300 higher than the before-treatment SAS score in the 2019 group ( $48.3 \pm 8.5$ ) according to a  
 301 Mann–Whitney rank-sum test ( $U = 534$ ,  $p < 0.001$ ). These results suggest that the anxiety  
 302 associated with COVID-19 had not been counteracted by the treatment.



303

304 **Fig 3. Treatment-related changes in SAS, THI and MML scores.** Two-way repeated measure analysis of  
 305 variance (ANOVA) was performed on these measures against the treatment and the year group (A–C), and  
 306 on the SAS score improvements against the anxiety and the year group (D). A significant effect for  
 307 treatment was seen in all three measures tested, and the improvements of the patients without anxiety in  
 308 2020 were significantly lower. Significance: \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  in ANOVA; ### $p < 0.001$   
 309 in a Mann–Whitney rank-sum test. SAS: Zung’s Self-rating Anxiety Scale (SAS), THI: Tinnitus Handicap  
 310 Inventory, MML: minimum masking level.

311

312 To further identify the between-year difference and the effects of anxiety on the SAS

313 score improvements, another two-way ANOVA was performed (Fig 3D). The  
314 improvement in the SAS score appeared to be slightly but significantly smaller in 2020  
315 ( $5.0 \pm 8.6$ ) than in 2019 ( $6.1 \pm 3.8$ ,  $F_{1,92} = 6.046$ ,  $p = 0.016$ ). A significant effect for  
316 anxiety was revealed, in that the tinnitus patients with anxiety received more benefit from  
317 the sound therapy ( $6.6 \pm 6.2$  in anxiety group vs.  $4.0 \pm 5.9$  in no-anxiety group;  
318  $F_{1,92} = 10.447$ ,  $p = 0.002$ ).


319 A significant interaction between the two factors was also seen ( $F_{1,92} = 9.037$ ,  $p =$   
320  $0.002$ ). Correspondingly, the amount of the SAS score reduction from the treatment in the  
321 patients without anxiety was significantly smaller ( $-1.5 \pm 7.2$ ) than those with anxiety in  
322 2020 ( $7.0 \pm 8.0$ ,  $q = 5.364$ ,  $p < 0.001$ ) and those without anxiety in 2019 ( $6.0 \pm 3.5$ ,  
323  $q = 4.636$ ,  $p = 0.002$ ; Fig 3D). That is, the patients without anxiety in 2020 had less  
324 improvement in SAS score. However, no difference was seen in SAS score improvement  
325 in 2019 between the subjects with and without anxiety.

326 The sound therapy appeared to reduce the THI scores in both years, as indicated by  
327 the significant treatment effect in the two-way RM ANOVA ( $F_{1,94} = 58.405$ ,  $p < 0.001$ ). A  
328 significant effect of the year was evidenced by an overall higher THI in the subjects,  
329 overall, in the 2020 group ( $40.7 \pm 6.7$ ) than for those in the 2019 group ( $32.7 \pm 8.3$ ,  
330  $F_{1,94} = 27.073$  and  $p < 0.001$ ), although a significant interaction between year and  
331 treatment was not seen ( $F_{1,94} = 1.054$ ,  $p = 0.307$ ). The THI scores in the 2020 group  
332 improved from  $40.7 \pm 6.7$  to  $37.7 \pm 8.0$  according to the Tukey method ( $p < 0.001$ ), while  
333 those in 2019 improved from  $32.7 \pm 8.3$  to  $28.7 \pm 7.6$  ( $p < 0.001$ ; Fig 3B). It is noticeable  
334 that the post-treatment THI in the 2020 group was even higher than pre-treatment in 2019.  
335 Furthermore, the improvement (the pre-treatment score minus the post-treatment) was

336 slightly larger in the 2019 group ( $4.0 \pm 3.0$ ) than in 2020 ( $3.0 \pm 5.5$ ), but the difference  
337 was not significant ( $U = 1211$ ,  $p = 0.413$ ).

338 A significant effect of the treatment on MML was also shown by a two-way RM  
339 ANOVA ( $F_{1,94} = 69.105$ ,  $p < 0.001$ ) against the treatment and year group. Significant  
340 improvements were observed in both the 2020 and 2019 groups (from  $9 \pm 4.4$  dB SL to  
341  $7.3 \pm 4.2$  dB SL and from  $10.0 \pm 3.8$  dB SL to  $7.9 \pm 3.8$  dB SL respectively, Fig 3C).  
342 There were no significant differences between years in the overall MML ( $F_{1,94} = 0.916$ ,  
343  $p = 0.341$ ) and no significant interaction between year and treatment ( $F_{1,94} = 1.247$ ,  $p =$   
344  $0.267$ ). The improvement of MML in 2019 ( $2.1 \pm 1.7$  dB) was a little higher than that in  
345 2020 ( $1.6 \pm 2.7$  dB), but this was not statistically significant ( $U = 1247$ ,  $p = 0.255$ ).

346 To further evaluate whether the initial anxiety was a factor that impacted on the  
347 treatment, the Pearson product moment correlation was calculated between the initial  
348 SAS score and the changes in the SAS, THI, and MML scores. There was a weak,  
349 positive, linear relationship between the initial SAS score and the SAS score  
350 improvement across all of the subjects pooled from both years ( $r = 0.253$ ,  $p = 0.01$ ,  
351 Fig 4A). When separated into the two years, a moderate positive correlation was seen in  
352 the 2020 group ( $r = 0.511$ ,  $p = 0.001$ ) and no significant correlation was found in the  
353 2019 group (Fig 4B). These results suggest that the treatment was more effective for  
354 mitigating anxiety in subjects with higher SAS scores, which was associated with the  
355 COVID-19 pandemic in 2020. However, no significant correlation was seen between the  
356 initial SAS score and reductions in MML or THI score, either in the total sample or  
357 separated into different year groups ( $p > 0.05$ ). This suggests that the initial anxiety status  
358 was not related to the effectiveness of the treatment of sound therapy for tinnitus.



359

360 **Fig 4. Correlations between the initial SAS score and the improvement in SAS score.** (A) The  
361 correlation in all subjects pooled across both years. (B) The correlations by year. A significant, moderate  
362 correlation was seen only in the 2020 group, in which the average initial SAS score was much higher. SAS:  
363 Zung's Self-rating Anxiety Scale.

364

365

## 366 Discussion

367 In this retrospective study, we compared the severity of our tinnitus patients over a 6-  
368 week period between 2020, which was under the influence of the COVID-19 pandemic,  
369 and 2019. A significant increase was seen in 2020 in the incidence of subjects who were  
370 judged as having anxiety by the current criterion of their SAS score (Fig 2A and 2B).  
371 Furthermore, the average SAS score was much higher in 2020 than in 2019 in the  
372 subgroups with anxiety (Fig 2C and 2D). This demonstrates that the psychological stress  
373 in dealing with various pressures related to the COVID-19 pandemic not only increased  
374 the prevalence of anxiety in the tinnitus patients, but also greatly elevated the level of  
375 anxiety in the subjects who suffered from it. We attributed the changes to increased stress  
376 under the COVID-19 pandemic because the subjects in both years were matched in many  
377 major factors (such as age and sex; Table 1) that might have affected tinnitus. However,  
378 the increased anxiety was not clearly linked to the maskability of the tinnitus itself, as  
379 evaluated by MML (Fig 2E and 2F). The proportion of subjects who reported a mitigation

380 in tinnitus after sound therapy was slightly but significantly lower in 2020 than in 2019.  
381 This reduction was likely linked to the increased anxiety in 2020. While the treatment  
382 reduced SAS score more effectively in 2019, the improvement in SAS score was  
383 positively correlated with initial SAS score only in the 2020 group, in which patients  
384 without anxiety showed no improvement at all (in fact, with a slightly increased SAS  
385 score, Fig 3D). Furthermore, the initial state of anxiety appeared to be a factor only in the  
386 improvement of the SAS score, not in the THI score or MML. Although the increased  
387 anxiety was one of the reasons for the increased THI score, there was no clear link  
388 between the changes in anxiety and the maskability of tinnitus, as evaluated by MML.

389 There is no doubt that a significant increase in psychological stress developed as a  
390 result of the COVID-19 pandemic. Many recently published articles have revealed the  
391 high prevalence of anxiety across China during the COVID-19 pandemic, from 28.8% to  
392 35.1% [34, 35], compared to the previously reported prevalences of 5.6% and 7.6% for  
393 the years of 2009 and 2019, respectively [36, 37]. A cross-sectional survey, using the  
394 same anxiety questionnaire as adapted in the present study, reported an average SAS  
395 score of  $45.89 \pm 1.1$  among front-line clinical staff during the pandemic [38]. This value  
396 was located between the scores for our subjects with and without anxiety ( $68.0 \pm 6.0$  vs.  
397  $43.9 \pm 1.5$ ), and lower than the average for all of our subjects in the 2020 group  
398 ( $61.9 \pm 10.9$ ). This implies that our tinnitus patients seen in 2020 have experienced  
399 extremely high psychological pressure, even higher than those medical doctors who were  
400 in the most challenging job during the pandemic. The number of tinnitus subjects seen in  
401 the 6-week period in 2020 was higher than that last year. However, this increase may be  
402 largely attributable to the accumulation of patients during the hospital closure in the

403 national lockdown.

404       The COVID-19 pandemic provides a good opportunity to investigate whether stress  
405 or anxiety could impact tinnitus as a causative or promotive factor. The association  
406 between tinnitus and anxiety has been investigated in many previous studies and has been  
407 well reviewed [9-15] . However, no information is available on the direction and  
408 causality between the two ends of the link [13, 15, 39-41], although many studies have  
409 implied that psychological states, such as those related to common stressors, influence  
410 perception of or coping with tinnitus [42, 43]. In this regard, two related systems are  
411 involved in tinnitus: (1) the brain regions along the hypothalamic–pituitary–adrenal axis  
412 (see reviews [9, 13]), which is the main neuroendocrine system involved in stress  
413 response, and (2) the limbic system including the hippocampus and amygdala, which  
414 regulates the perception of tinnitus and the adaptation (thereby, the ability to cope with  
415 stress) [44-49]. While the data from the previous studies have indicated the possible role  
416 of emotional factors in tinnitus via those systems, the relationship was mostly  
417 investigated in cross-sectional comparisons across subjects with different levels of  
418 tinnitus and those without.

419       Furthermore, several other limitations exist for the generalization of the findings in  
420 relevant research. First, different populations have been targeted by different studies. For  
421 example, one study was completed amongst male veterans [50], one study reported more  
422 recently was on subjects over 75 years old [40], while other studies were conducted on a  
423 more general population [51]. Second, various methods have been used to measure  
424 anxiety, including ICD-9 codes [50] , the Hospital Anxiety and Depression Scale (HADS)  
425 [40, 52] , the Anxiety Sensitivity Index [51] , the State Trait Anxiety Inventory [53], as

426 well as the SAS [54]. Third, the cause of anxiety also varied, or was not clear at all, in  
427 different studies. In the studies above, one showed a clear goal to examine how the  
428 anxiety related to sleep disorders affected tinnitus [54] , while in most other studies the  
429 cause of anxiety was not specified [40, 50, 51, 53, 54]. This variation makes it difficult, if  
430 not impossible, to compare the outcomes across different studies. For example, in one  
431 study amongst male veterans, the incidence of tinnitus subjects with anxiety (quantified  
432 via the ICD-9 code) was reported to be very high (79.1%) [50] . This high incidence of  
433 anxiety probably related to the special population investigated in this study, because the  
434 incidence has been found to be much lower in other studies of the general population. For  
435 example, in a recent study, most subjects with tinnitus (68%) were free of anxiety [53]. In  
436 that study, the anxiety was assessed using the State Trait Anxiety Inventory and the result  
437 was consistent with another study in which only 23.4% of the tinnitus participants were  
438 diagnosed as having anxiety via the Anxiety Sensitivity Index [51] .

439 It is also noticeable that, in most of the previous studies, the goal was to evaluate  
440 how tinnitus worked as a promoting factor for anxiety [40] , or to simply established  
441 whether there was a link between anxiety and tinnitus [40, 50, 51, 53, 54] . In the present  
442 study, we evaluated whether anxiety plays a causative role in promoting the development  
443 or the enhancement of tinnitus. The increased stress under COVID-19 resulted in an  
444 increase in anxiety with a clear cause and was encountered, inevitably, by most of the  
445 population who lived in the same area. Although large variation existed across different  
446 individuals in relationship to their jobs and financial situations, as well as their closeness  
447 to COVID-19 patients, the stress factor associated with this study was much more  
448 homogeneous than those that had been examined in previous studies. We therefore think



449 that the between-year differences in the tinnitus clinic afforded a good chance to verify  
450 whether anxiety plays a causative or promotive role for tinnitus.

451 Consistency exists across different studies in that an increase in anxiety is positively  
452 associated with the severity of tinnitus whether it is examined with the THI score [50] ,  
453 MML [55], or not quantitatively but with the answers to two simple questions [56].

454 Unfortunately, the level of anxiety was also evaluated with different methods in these  
455 studies (ICD-9 code, HADS, and symptoms of depression and anxiety, respectively). In  
456 the present study, SAS was used as the evaluation tool to quantify the degree of anxiety,  
457 because the SAS has good psychometric credentials and continues to be widely used in  
458 clinical screening [33]. The severity of tinnitus was quantified by both THI and MML in  
459 the present study. Therefore, we will compare our results with those previous studies with  
460 similar methodologies. In one study reporting on 543 tinnitus cases in China, the average  
461 SAS score was  $37.26 \pm 8.99$  (mean  $\pm$  standard deviation) [54]. If this study focused on  
462 the association between psychiatric disorders and tinnitus, the SAS scores were most  
463 likely to be the raw value, which should be converted by multiplying by 1.25 before  
464 comparing with the results in the present study. After this conversion, the mean SAS  
465 score becomes 46.575, which is much lower than the initial SAS score of 61.9 in our  
466 2020 group and comparable with the value in our 2019 group (48.2, or 22.1% lower than  
467 2020 group; Fig 2A). The average THI score in this study was  $40.87 \pm 22.68$ , which was  
468 comparable to our 2020 group ( $40.1 \pm 6.9$ ) and the results from another previous study  
469 ( $40.1 \pm 24.1$ , [53] ). A higher THI ( $>51$  on average) was also reported in tinnitus subjects  
470 with anxiety [51] . In addition, our 2019 group showed a lower THI score overall  
471 ( $33.6 \pm 8.7$ ), yielding a between-year difference of 16.2% on average (Fig 2D). Taken

472 together, the much higher anxiety in SAS score in the 2020 group in the present study  
473 was largely, but not fully, matched by the between-year difference in the THI score.  
474 Furthermore, the between-year difference in THI scores were seen in the functional and  
475 emotional sections, but not in the catastrophic section.

476 Discrepancies were also noted between the present study and previous ones,  
477 particularly in the evaluation of tinnitus loudness. In the present report, the MML was  
478 lower in the 2020 group, which had much higher SAS and THI scores than the 2019  
479 group (Fig 3F). This conflicts with one previous study in which tinnitus patients with  
480 anxiety had a larger MML than those without [51] . However, results that were  
481 qualitatively similar to ours were also reported in a study in which the group with a  
482 higher anxiety score (by HADS) exhibited a significantly lower MML  
483 ( $49.3 \pm 22.5$  dB HL) than the group with a lower anxiety score ( $57.9 \pm 18.9$  dB HL),  
484 although the difference in the HADS scores between the groups was not statistically  
485 significant [57]. More importantly, in the present study, the MML was only slightly  
486 higher in subjects with anxiety in 2020 than those without, while the SAS score had a  
487 large difference between the two subgroups in 2020. Taken together, we believe that the  
488 anxiety induced by the pandemic has had an extremely limited, if any, impact on tinnitus  
489 loudness.

490 Therefore, the anxiety increase caused by the COVID-19 pandemic seems to have  
491 had a limited impact on tinnitus itself. This idea is further supported by the outcomes of  
492 the sound therapy in the present study, compared to the results from previous reports. The  
493 effects of the treatment were evaluated using three measurements (SAS, THI, and MML)  
494 in the present study. The SAS score improvement was slightly but significantly less in the

495 2020 group in which the initial SAS score was very high. However, the improvement in  
496 SAS score was well correlated with the initial SAS score in 2020 but not in 2019. This is  
497 superficially paradoxical but explainable if the improvement of the SAS score in the 2020  
498 group was not because of the treatment, but instead because of the natural release of the  
499 underlying stress in association with the deceleration of the pandemic during the two-  
500 month interval between the two evaluations in the present study. With this idea, it  
501 becomes understandable that those who had a higher SAS score might achieve more  
502 recovery during this period when the pandemic situation decelerated, and it may not have  
503 been a result from the therapy. Otherwise, the relationship should have be similar to that  
504 seen in the 2019 group.

505 This idea is further supported by the fact that there was no significant between-year  
506 difference in the improvements in the THI score and MML in the present study. Our  
507 results on the relationship between anxiety and the treatment effects on both THI and  
508 MML also contradicted those reported in previous studies. For example, THI score  
509 reductions were reported to be larger in subjects with anxiety (e.g., by higher HADS  
510 scores in [52]) and furthermore, the reduction in anxiety was found to parallel the  
511 improvement in THI by sound therapy [58, 59]. In the present study, however, the  
512 improvements in both THI score and MML were not larger in the 2020 group, in which  
513 anxiety was much more severe. This conflict can also be explained by an assumption that  
514 the anxiety induced by the COVID-19 pandemic has had little or no impact on tinnitus  
515 severity, and therefore no impact on the treatment outcome.

## 516 **Limitations**

517 There were several limitations to our study. This was a retrospective study in which

518 only the SAS was used to evaluate anxiety. This makes it difficult to compare our study  
519 with previous ones. In addition, only sound therapy with counseling produced reliable  
520 data that allowed a comparison between years and the number of subjects using this  
521 therapy decreased in 2020. Many tinnitus patients in 2020 selected educational  
522 counseling alone for their treatment, over the other two, probably due to financial  
523 constraints. The combination of counseling with another therapy has been suggested by  
524 others to achieve better outcomes [60-63]. In addition, the overall sample size in the  
525 present study was small as the data were collected only from one hospital within a limited  
526 period. Although the data and conclusion are solid in the present study, further  
527 investigation would be helpful to verify the conclusion with a larger sample.

## 528 **Conclusion**

529 There was a substantial increase in anxiety of tinnitus subjects in 2020 under the  
530 pressures produced by the COVID-19 pandemic. However, this increase was not matched  
531 by the between-year difference in tinnitus loudness but matched better the change in THI  
532 score. The efficacy of sound therapy on tinnitus itself showed no significant difference  
533 between years. While the post-treatment SAS score was still much higher in the 2020  
534 group, there was a positive correlation between the SAS score improvement and the  
535 initial SAS score in that group. This might have resulted from the natural reduction in  
536 stress from the pandemic reaching another stage in China. Therefore, the anxiety caused  
537 by the pandemic is unlikely to be a promoting or enhancing factor for tinnitus.

538

## 539 **Acknowledgement**

540 The authors acknowledge the colleagues for participating in this study. We are deeply

541 indebted to the families who participated in the study.

## 542 **Financial Disclosure**

543 This study was supported by the research grants from Health and Family Planning  
544 Commission of Sichuan Province of China [18PJ078], the National Natural Science  
545 Foundation of China [81770998], Natural Science & Engineering Research Council of  
546 Canada [RGPIN 2017-04493].

## 547 **Competing interests**

548 The authors have declared that no competing interests exist.

## 549 **References**

550

- 551 1. Remuzzi A, Remuzzi G. COVID-19 and Italy: what next? *Lancet*. 2020;395(10231):1225-8.
- 552 2. World Health Organization. Coronavirus disease 2019 (COVID-19) Situation Report – 158 2020  
553 [2020 June 26]. Available from: [https://www.who.int/docs/default-source/coronaviruse/situation-](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200626-covid-19-sitrep-158.pdf?sfvrsn=1d1aae8a_2)  
554 [reports/20200626-covid-19-sitrep-158.pdf?sfvrsn=1d1aae8a\\_2](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200626-covid-19-sitrep-158.pdf?sfvrsn=1d1aae8a_2).
- 555 3. Liu NN, Tan JC, Li J, Li S, Cai Y, Wang H. COVID-19 Pandemic: Experiences in China and  
556 implications for its prevention and treatment worldwide. *Curr Cancer Drug Targets* [Internet]. 2020 Apr 14.  
557 Available from: <http://www.eurekaselect.com/180883/article>.
- 558 4. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The  
559 psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet*.  
560 2020;395(10227):912-20.
- 561 5. Lu W, Wang H, Lin Y, Li L. Psychological status of medical workforce during the COVID-19  
562 pandemic: A cross-sectional study. *Psychiatry Res*. 2020;288:112936.
- 563 6. Wang G, Zhang Y, Zhao J, Zhang J, Jiang F. Mitigate the effects of home confinement on  
564 children during the COVID-19 outbreak. *Lancet*. 2020;395(10228):945-7.
- 565 7. Bauer CA. Tinnitus. *N Engl J Med*. 2018;378(13):1224-31.
- 566 8. Baguley D, McFerran D, Hall D. Tinnitus. *Lancet*. 2013;382(9904):1600-7.

- 567 9. Ziai K, Moshtaghi O, Mahboubi H, Djalilian HR. Tinnitus Patients Suffering from Anxiety and  
568 Depression: A Review. *Int Tinnitus J.* 2017;21(1):68-73.
- 569 10. Pattyn T, Van Den Eede F, Vanneste S, Cassiers L, Veltman DJ, Van De Heyning P, et al. Tinnitus  
570 and anxiety disorders: A review. *Hear Res.* 2016;333:255-65.
- 571 11. Durai M, Searchfield G. Anxiety and depression, personality traits relevant to tinnitus: A scoping  
572 review. *Int J Audiol.* 2016;55(11):605-15.
- 573 12. Malouff JM, Schutte NS, Zucker LA. Tinnitus-related distress: A review of recent findings. *Curr*  
574 *Psychiatry Rep.* 2011;13(1):31-6.
- 575 13. Mazurek B, Boecking B, Brueggemann P. Association Between Stress and Tinnitus-New  
576 Aspects. *Otol Neurotol.* 2019;40(4):e467-e73.
- 577 14. Zirke N, Seydel C, Szczepek AJ, Olze H, Haupt H, Mazurek B. Psychological comorbidity in  
578 patients with chronic tinnitus: analysis and comparison with chronic pain, asthma or atopic dermatitis  
579 patients. *Qual Life Res.* 2013;22(2):263-72.
- 580 15. Wallhauser-Franke E, Brade J, Balkenhol T, D'Amelio R, Seegmuller A, Delb W. Tinnitus:  
581 distinguishing between subjectively perceived loudness and tinnitus-related distress. *PLoS One.*  
582 2012;7(4):e34583.
- 583 16. Schlee W, Pryss RC, Probst T, Schobel J, Bachmeier A, Reichert M, et al. Measuring the  
584 Moment-to-Moment Variability of Tinnitus: The TrackYourTinnitus Smart Phone App. *Front Aging*  
585 *Neurosci.* 2016;8:294.
- 586 17. Probst T, Pryss R, Langguth B, Schlee W. Emotional states as mediators between tinnitus  
587 loudness and tinnitus distress in daily life: Results from the "TrackYourTinnitus" application. *Sci Rep.*  
588 2016;6:20382.
- 589 18. Probst T, Pryss R, Langguth B, Schlee W. Emotion dynamics and tinnitus: Daily life data from  
590 the "TrackYourTinnitus" application. *Sci Rep.* 2016;6:31166.
- 591 19. Bartels H, Middel B, Pedersen SS, Staal MJ, Albers FW. The distressed (Type D) personality is  
592 independently associated with tinnitus: a case-control study. *Psychosomatics.* 2010;51(1):29-38.
- 593 20. Bartels H, Pedersen SS, van der Laan BF, Staal MJ, Albers FW, Middel B. The impact of Type D  
594 personality on health-related quality of life in tinnitus patients is mainly mediated by anxiety and

- 595 depression. *Otol Neurotol*. 2010;31(1):11-8.
- 596 21. Trevis KJ, McLachlan NM, Wilson SJ. Cognitive Mechanisms in Chronic Tinnitus:  
597 Psychological Markers of a Failure to Switch Attention. *Front Psychol*. 2016;7:1262.
- 598 22. Trevis KJ, McLachlan NM, Wilson SJ. Psychological mediators of chronic tinnitus: The critical  
599 role of depression. *J Affect Disord*. 2016;204:234-40.
- 600 23. Bedford J, Enria D, Giesecke J, Heymann DL, Ihekweazu C, Kobinger G, et al. COVID-19:  
601 towards controlling of a pandemic. *Lancet*. 2020;395(10229):1015-8.
- 602 24. World Medical Association. WMA declaration of Helsinki: Ethical principles for medical  
603 research involving human subjects 2018 [2020 June 26 ]. Available from: [https://www.wma.net/policies-](https://www.wma.net/policies-post/wma-declaration-of-helsinki-ethical-principles-for-medical-research-involving-human-subjects/)  
604 [post/wma-declaration-of-helsinki-ethical-principles-for-medical-research-involving-human-subjects/](https://www.wma.net/policies-post/wma-declaration-of-helsinki-ethical-principles-for-medical-research-involving-human-subjects/).
- 605 25. Langguth B. Treatment of tinnitus. *Curr Opin Otolaryngol Head Neck Surg*. 2015;23(5):361-8.
- 606 26. Tyler RS. Mindfulness-Based Therapy: New Study Underscores Tinnitus Treatment Options. *The*  
607 *Hearing Journal*. 2014;67(1).
- 608 27. Arif M, Sadlier M, Rajenderkumar D, James J, Tahir T. A randomised controlled study of  
609 mindfulness meditation versus relaxation therapy in the management of tinnitus. *J Laryngol Otol*.  
610 2017;131(6):501-7.
- 611 28. Kam AC, Cheung AP, Chan PY, Leung EK, Wong TK, van Hasselt CA, et al. Psychometric  
612 properties of the Chinese (Cantonese) Tinnitus Handicap Inventory. *Clin Otolaryngol*. 2009;34(4):309-15.
- 613 29. Meng Z, Zheng Y, Liu S, Wang K, Kong X, Tao Y, et al. Reliability and validity of the chinese  
614 (mandarin) tinnitus handicap inventory. *Clin Exp Otorhinolaryngol*. 2012;5(1):10-6.
- 615 30. Newman CW, Jacobson GP, Spitzer JB. Development of the Tinnitus Handicap Inventory. *Arch*  
616 *Otolaryngol Head Neck Surg*. 1996;122(2):143-8.
- 617 31. Gao LL, Ip WY, Sun K. Validation of the short form of the Chinese Childbirth Self-Efficacy  
618 Inventory in Mainland China. *Res Nurs Health*. 2011;34(1):49-59.
- 619 32. Zung WW. A Self-Rating Depression Scale. *Arch Gen Psychiatry*. 1965;12:63-70.
- 620 33. Dunstan DA, Scott N. Norms for Zung's Self-rating Anxiety Scale. *BMC Psychiatry*.  
621 2020;20(1):90.
- 622 34. Huang Y, Zhao N. Generalized anxiety disorder, depressive symptoms and sleep quality during

- 623 COVID-19 outbreak in China: a web-based cross-sectional survey. *Psychiatry Res.* 2020;288:112954.
- 624 35. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate Psychological Responses and  
625 Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among  
626 the General Population in China. *International journal of environmental research and public health.*  
627 2020;17(5):1729.
- 628 36. Huang Y, Wang Y, Wang H, Liu Z, Yu X, Yan J, et al. Prevalence of mental disorders in China: a  
629 cross-sectional epidemiological study. *The Lancet Psychiatry.* 2019;6(3):211-24.
- 630 37. Phillips MR, Zhang J, Shi Q, Song Z, Ding Z, Pang S, et al. Prevalence, treatment, and  
631 associated disability of mental disorders in four provinces in China during 2001-05: an epidemiological  
632 survey. *Lancet.* 2009;373(9680):2041-53.
- 633 38. Wu K, Wei X. Analysis of Psychological and Sleep Status and Exercise Rehabilitation of Front-  
634 Line Clinical Staff in the Fight Against COVID-19 in China. *Medical science monitor basic research.*  
635 2020;26:e924085-e.
- 636 39. Lugo A, Edvall NK, Lazar A, Mehraei G, Lopez-Escamez JA, Bulla J, et al. Relationship  
637 between headaches and tinnitus in a Swedish study. *Sci Rep.* 2020;10(1):8494.
- 638 40. Daniioth L, Brotschi G, Croy I, Friedrich H, Caversaccio MD, Negoias S. Multisensory  
639 environmental sensitivity in patients with chronic tinnitus. *J Psychosom Res.* 2020;135:110155.
- 640 41. Park E, Kim H, Choi IH, Han HM, Han K, Jung HH, et al. Psychiatric Distress as a Common  
641 Risk Factor for Tinnitus and Joint Pain: A National Population-Based Survey. *Clin Exp Otorhinolaryngol*  
642 [Internet]. 2019 Dec 18. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31842535>  
643 <https://www.e-ceo.org/upload/pdf/ceo-2019-00563.pdf>.
- 644 42. Lazarus RS. Toward better research on stress and coping. *Am Psychol.* 2000;55(6):665-73.
- 645 43. Lazarus RS. Coping theory and research: past, present, and future. *Psychosom Med.*  
646 1993;55(3):234-47.
- 647 44. Lockwood AH, Salvi RJ, Coad ML, Towsley ML, Wack DS, Murphy BW. The functional  
648 neuroanatomy of tinnitus: evidence for limbic system links and neural plasticity. *Neurology.*  
649 1998;50(1):114-20.
- 650 45. Kapolowicz MR, Thompson LT. Plasticity in Limbic Regions at Early Time Points in



- 651 Experimental Models of Tinnitus. *Front Syst Neurosci.* 2019;13:88.
- 652 46. Raghavan P, Steven A, Rath T, Gandhi D. Advanced Neuroimaging of Tinnitus. *Neuroimaging*  
653 *Clin N Am.* 2016;26(2):301-12.
- 654 47. Leaver AM, Seydell-Greenwald A, Rauschecker JP. Auditory-limbic interactions in chronic  
655 tinnitus: Challenges for neuroimaging research. *Hear Res.* 2016;334:49-57.
- 656 48. Chen YC, Xia W, Chen H, Feng Y, Xu JJ, Gu JP, et al. Tinnitus distress is linked to enhanced  
657 resting-state functional connectivity from the limbic system to the auditory cortex. *Hum Brain Mapp.*  
658 2017;38(5):2384-97.
- 659 49. Zhang J, Chen YC, Feng X, Yang M, Liu B, Qian C, et al. Impairments of thalamic resting-state  
660 functional connectivity in patients with chronic tinnitus. *Eur J Radiol.* 2015;84(7):1277-84.
- 661 50. Hu J, Xu J, Streelman M, Xu H, Guthrie O. The Correlation of the Tinnitus Handicap Inventory  
662 with Depression and Anxiety in Veterans with Tinnitus. *Int J Otolaryngol.* 2015;2015:689375.
- 663 51. Moon KR, Park S, Jung Y, Lee A, Lee JH. Effects of Anxiety Sensitivity and Hearing Loss on  
664 Tinnitus Symptom Severity. *Psychiatry Investig.* 2018;15(1):34-40.
- 665 52. Takahashi M, Kabaya K, Sekiya K, Matsuda F, Sekiya Y, Esaki S, et al. An improved system for  
666 grading and treating tinnitus. *Auris Nasus Larynx.* 2018;45(4):711-7.
- 667 53. Cho CG, Chi JH, Song JJ, Lee EK, Kim BH. Evaluation of anxiety and depressive levels in  
668 tinnitus patients. *Korean J Audiol.* 2013;17(2):83-9.
- 669 54. Xu Y, Yao J, Zhang Z, Wang W. Association between sleep quality and psychiatric disorders in  
670 patients with subjective tinnitus in China. *Eur Arch Otorhinolaryngol.* 2016;273(10):3063-72.
- 671 55. Barros Suzuki FA, Suzuki FA, Yonamine FK, Onishi ET, Penido NO. Effectiveness of sound  
672 therapy in patients with tinnitus resistant to previous treatments: importance of adjustments. *Braz J*  
673 *Otorhinolaryngol.* 2016;82(3):297-303.
- 674 56. McCormack A, Edmondson-Jones M, Fortnum H, Dawes PD, Middleton H, Munro KJ, et al.  
675 Investigating the association between tinnitus severity and symptoms of depression and anxiety, while  
676 controlling for neuroticism, in a large middle-aged UK population. *Int J Audiol.* 2015;54(9):599-604.
- 677 57. Petridou AI, Zagora ET, Petridis P, Korres GS, Gazouli M, Xenelis I, et al. The Effect of  
678 Antioxidant Supplementation in Patients with Tinnitus and Normal Hearing or Hearing Loss: A

- 679 Randomized, Double-Blind, Placebo Controlled Trial. *Nutrients*. 2019;11(12).
- 680 58. Mahboubi H, Haidar YM, Kiumehr S, Ziai K, Djalilian HR. Customized Versus Noncustomized  
681 Sound Therapy for Treatment of Tinnitus: A Randomized Crossover Clinical Trial. *Ann Otol Rhinol*  
682 *Laryngol*. 2017;126(10):681-7.
- 683 59. Westin VZ, Schulin M, Hesser H, Karlsson M, Noe RZ, Olofsson U, et al. Acceptance and  
684 commitment therapy versus tinnitus retraining therapy in the treatment of tinnitus: a randomised controlled  
685 trial. *Behav Res Ther*. 2011;49(11):737-47.
- 686 60. Jastreboff PJ, Gray WC, Gold SL. Neurophysiological approach to tinnitus patients. *Am J Otol*.  
687 1996;17(2):236-40.
- 688 61. Sereda M, Xia J, El Refaie A, Hall DA, Hoare DJ. Sound therapy (using amplification devices  
689 and/or sound generators) for tinnitus. *The Cochrane database of systematic reviews*.  
690 2018;12(12):CD013094-CD.
- 691 62. Cima RF, Maes IH, Joore MA, Scheyen DJ, El Refaie A, Baguley DM, et al. Specialised  
692 treatment based on cognitive behaviour therapy versus usual care for tinnitus: a randomised controlled trial.  
693 *Lancet*. 2012;379(9830):1951-9.
- 694 63. Jastreboff PJ, Jastreboff MM. Tinnitus Retraining Therapy (TRT) as a method for treatment of  
695 tinnitus and hyperacusis patients. *J Am Acad Audiol*. 2000;11(3):162-77.
- 696