

Development and Validation of a Web-based Prediction Model for Acute Kidney Injury after surgery

Sang H. Woo^{1*}, Jillian Zavodnick¹, Lily Ackermann¹, Omar Maarouf², Jingjing Zhang², Scott W. Cowan³

¹ Department of Medicine- Division of Hospital Medicine, Thomas Jefferson University, Philadelphia, PA 19107

² Department of Medicine- Division of Nephrology, Thomas Jefferson University, Philadelphia, PA 19107

³ Department of Surgery, Thomas Jefferson University

Running title: preoperative renal risk assessment, acute kidney injury

Correspondence:

Sang Hoon Woo, MD ; Department of Medicine, Thomas Jefferson University

email: sang.woo@jefferson.edu

Abstract:

Background and objectives Acute kidney injury after surgery is associated with high mortality and morbidity. The purpose of this study is to develop and validate a risk prediction tool for the occurrence of postoperative acute kidney injury requiring renal replacement therapy.

Design, setting, participants, measurements This retrospective cohort study had 2,299,502 surgical patients over 2015-2017 from the American College of Surgeons National Surgical Quality Improvement Program Database (ACS-NSQIP). Eleven predictors were selected for the predictive model: age, history of congestive heart failure, diabetes, ascites, emergency surgery, preoperative serum creatinine, hematocrit, sodium, preoperative sepsis, preoperative acute renal failure and surgery type. The predictive model was trained using 2015-2016 data (n=1,487,724) and further tested using 2017 data (n=811,778). A risk model was developed using multivariate logistic regression and machine learning methods.

Main outcomes The primary outcome was postoperative 30-day acute kidney injury requiring renal replacement therapy(AKI-D)

Results The unadjusted 30-day postoperative mortality rate associated with AKI-D was 37.5%. The renal risk prediction model had high AUC (area under the receiver operating characteristic curve, training cohort: 0.89, test cohort: 0.90) for postoperative AKI-D.

Conclusions This model provides a clinically useful bedside predictive tool for postoperative acute kidney injury requiring dialysis.

INTRODUCTION

Postoperative renal failure is a serious medical complication of surgery, with a rising incidence and significantly associated morbidity and mortality. Postoperative acute kidney injury (AKI) increases five-year mortality regardless of level of injury and extent of kidney recovery.¹ For patients who develop acute kidney injury requiring dialysis (AKI-D) after elective surgery, ninety-day mortality rate was reported to be 42%; among patients undergoing cardiac surgery, dialysis conferred a thirty-day mortality rate of 63.7% as compared to 4.3% without AKI.^{2,3} Of patients who survived the episode of AKI-D, 27.2% required chronic dialysis, with its attendant morbidity and cost.³ Hospital costs are significantly higher for patients who develop any postoperative AKI.⁴

Preoperative assessment of surgical risk is valuable for determining appropriateness of surgery, facilitating multidisciplinary surgical planning, and engaging patients in shared decision-making.⁵ Tools for cardiac and pulmonary risk stratification have been widely adopted, but the same is not true for renal risk assessment.^{6,7,8,9} Existing postoperative AKI models are based on small datasets or are specific to patients undergoing cardiac surgery.^{10,11,12,13,14} The Universal American College of Surgeons (ACS) National Surgical Quality Improvement Program (NSQIP) Surgical Risk Calculator includes postoperative renal failure as one of its 8 outcomes.¹⁵ The need to input 21 clinical factors into the calculator limits the adoption of the NSQIP calculator, especially for clinicians who are mainly interested in renal outcome.¹⁵

Establishing a renal risk prediction model for surgical patients including the risk for renal replacement therapy is a clinical unmet need. The objective of this study is to develop and validate an AKI-D risk calculator using data from the NSQIP database.

METHODS

Study participants

Data were obtained from the ACS NSQIP database, which collects clinical data from more than 500 United States hospitals including patient demographics, medical conditions, laboratory data, and 30-day postoperative outcomes collected by trained surgical clinical reviewers.^{16 17} Patients undergoing surgery at participating hospitals from 2015 to 2017 were included, with a total population of 2,299,502. The dataset from 2015 to 2016 was used to develop the model (n=1,487,724), while the dataset from 2017 was held for validation. The study was approved by the Thomas Jefferson University Institutional Review Board and informed consents were waived.

Outcome

The outcome of interest was AKI-D within thirty days of surgery, a time frame chosen based on the ACS-NSQIP definition. ACS-NSQIP defines AKI-D as “worsening of renal dysfunction postoperatively requiring hemodialysis, peritoneal dialysis, hemofiltration, hemodiafiltration, or ultrafiltration in a patient who did not require dialysis preoperatively.”¹⁸ Instructions from the organization state that if a patient refuses a recommendation for dialysis, they should still be coded as developing AKI-D as dialysis was deemed necessary.¹⁸

Data Analysis

We analyzed previously-described risk factors for AKI including demographic factors, medical history, current medical condition (e.g. presence of sepsis, American Society of

Anesthesiologists (ASA) class), and preoperative functional status. Multiple preoperative laboratory tests were analyzed including serum sodium, creatinine, platelets, and hematocrit. Participant risk factors were compared using Chi-square (χ^2) for categorical variables and Wilcoxon for continuous variables. Characteristics of patients who developed AKI-D were compared by adjusted odds ratio. Multivariate logistic regression was performed with risk factors to determine the significance of their associations with the outcome.

Separate categories were made for missing values of laboratory concentrations of serum hematocrit. Backwards elimination and clinical importance were used to select predictor variables of the model. The model was trained on 2015-2016 cohort data and tested on 2017 cohort.

GridSearchCV was used to perform 5 fold cross validation to select the best model parameters. Data set from year 2017 (n=811,778) was used to further test and validate the predictive model. Area under the receiver operating characteristic curve (AUC) was used for model evaluation.

Statistical analysis was performed with Python (version 3.65), Statsmodels (version 0.8.0), and R programming (RStudio version 1.1.463). Scikit-learn was used for machine learning modeling.

RESULTS

Study population

The study included 2,299,502 patients from 2015-2017. 56.4 % were women. **Table 1** shows the characteristics of study patients from the years 2015-2016. There were 0.3% of patients who

developed AKI-D. Patients who required dialysis were older (66.2 years versus 58.5, $p<0.001$), had higher preoperative serum creatinine level (1.96 mg/dl vs. 0.92 mg/dl, $p<0.001$), and were more likely to have congestive heart failure (8.83 % vs. 0.93%, $p<0.001$), diabetes (34.2% vs 17.19%, $p<0.001$), or other comorbidities. They were more likely to have undergone emergency surgery (37.8% versus 10.4%, $p<0.001$). **eFigure 1** in the **Supplement** shows a flow chart of study participants.

Mortality, readmission, length of stay associated with acute kidney injury

Figure 1 shows a high unadjusted 30-day postoperative mortality rate (37.5%) from 2015-2017 for patients who developed AKI-D compared to those who did not. The rate of 30-day hospital readmission was also higher in these patients (19.7% vs 5.0%). The patients who developed AKI-D also had increased length of hospital stay (18.7 days vs 2.9 days).

Figure 2 illustrates unadjusted 30-day mortality rates according to surgery types from 2015-2017. These varied widely, ranging from 16.2%-53.3%. Although patients undergoing ENT surgery had a high mortality rate, the incidence of AKI-D was low at 0.03%. 1.86% of cardiac surgery patients developed AKI-D with a mortality rate of 37.9%, lower than the rate of 63.7% reported in a previous study from the years 1987-1994.²

eFigure 2 in the **Supplement** shows unadjusted rates of postoperative AKI requiring dialysis according to surgery types. Aorta surgery has the highest risk of AKI-D at 6.05% with a mortality rate of 41.4%. Other surgery types that have higher renal risk include liver, pancreas, spleen surgery (0.82%), intestinal surgery (0.8%), vascular surgery (0.59%) and thoracic surgery

(0.48%). Surgery types in lower risk categories include orthopedic upper extremity surgery (0.05%) and spine surgery (0.09%).

Effect of preoperative renal function, serum sodium and hematocrit

As shown in **Figure 3**, preoperative renal dysfunction was associated with increased unadjusted risk of AKI-D. The risk of acute kidney injury was higher with higher preoperative serum creatinine levels. Anemia also increased the risk of AKI-D, with higher unadjusted rates among patients with hematocrit less than 30% (hematocrit $\leq 24\%$: 2.3%, 24.1-30%: 1.4%, $>30\%$: 0.2%). Patients with hyponatremia and hypernatremia had higher incidence of AKI-D (≤ 130 mEq/L: 1.5%, 130.1-135 meq/L: 0.6%, 135.1-145 mEq/L: 0.2%, >145 mEq/L: 1.0%)

Prediction of acute kidney injury requiring dialysis

Table 2 lists clinical factors which were strongly associated with the occurrence of postoperative 30-day AKI-D, and were ultimately incorporated into the prediction model. History of congestive heart failure was a significant predictor with an increased adjusted odds of the outcome, OR 2.24 [CI 1.99-2.53, $p<0.001$]. Diabetes was also associated with an increased risk, OR 2.09 [CI 1.92-2.27, $p<0.001$] if insulin dependent, OR 1.35 [CI 1.24-1.48, $p<0.001$] if non-insulin dependent, a finding consistent with a previous study.¹⁹ Preoperative condition such as SIRS(systemic inflammatory response syndrome), OR 2.49[CI 2.23-2.78, $p<0.001$], sepsis, OR 3.26 [CI 2.92-3.64, $p<0.001$], and septic shock, OR 6.86 [CI 6.04-7.78, $p<0.001$] were strongly associated with postoperative AKI-D. Increasing ASA class was also significantly associated with AKI-D. Laboratory findings associated with increased risk included preoperative

serum sodium and hematocrit. Serum sodium abnormalities in either direction had an increased adjusted risk of the outcome: for serum sodium ≤ 130 mEq/L, OR of AKI-D was 1.35 (CI: 1.17-1.55, $p < 0.001$) and for serum sodium 130.1-135 mEq/L, OR was 1.21 (CI: 1.12-1.31, $p < 0.001$); for serum sodium > 145 mEq/L, OR was 1.47 (CI: 1.23-1.75, $p < 0.001$). Surgery type was also a predictive factor, with the highest rates of AKI-D in patients undergoing aortic surgery, followed by cardiac surgery and surgery of the liver, pancreas, or spleen. Intestinal and other vascular procedures were also strongly associated.

These predictors were used to develop a model for AKI-D as described in the methods section. Brier score for the training cohort was 0.003. The model has a training AUC of 0.89. **eFigure 3** in the **Supplement** shows the calibration plot of the prediction model which is well matched with a 45-degree line.

Validation of the renal risk model

The trained model was tested by applying the model to a validation cohort of 811,778 from 2017. The prediction model has excellent predictive power with AUC of 0.90. 5-fold cross validation from the training cohort also showed high average AUC (0.89). **Figure 4** shows AUC of model validation.

Examples of Renal Risk Calculator

A calculator for AKI-D risk was programmed from coefficients and intercept from multivariate logistic regression and Scikit-learn machine learning library.²⁰ . The following examples demonstrate the bedside applicability of this model as well as the impact of various clinical factors on the risk of the outcome. The mobile web model is available at <https://renalrisk.herokuapp.com/>. The mobile-web model is shown in **eFigure 4** in the **Supplement**.

Hypothetical case 1

67 year old male with a history of congestive heart failure and insulin dependent diabetes mellitus undergoing emergent hip replacement surgery. No ascites. Patient had preoperative acute kidney injury and SIRS (systemic inflammatory response syndrome). Preoperative laboratory values: serum creatinine 2.5 mg/dL, hematocrit 24%, sodium 128 mEq/L. **Probability of AKI requiring dialysis (AKI-D) within 30 days after surgery: 28.36%**

Hypothetical case 2

67 year old male with a history of congestive heart failure and insulin dependent diabetes mellitus undergoing liver (hepatectomy) surgery. Patient had no preoperative acute kidney injury and SIRS. Preoperative laboratory values: serum creatinine 2.5 mg/dL, hematocrit 32%, sodium 133 mEq/L. **Probability of AKI-D within 30 days after surgery: 5.9%**

Discussion

We have developed a risk model for postoperative AKI-D in patients undergoing broad types of surgery. This model uses simple, easily available clinical data to deliver excellent predictive ability for the outcome, with a validation AUC of 0.90.

Our study has strengths compared to prior models to predict postoperative renal failure. A prior machine learning algorithm to predict postoperative renal failure has been described with lower predictive ability (AUC ranging from 0.797 and 0.858) and requiring a complex set of EHR-extracted data.²¹ There are also previously described machine learning prediction models for acute renal failure in hospitalized patients, not specific to the postoperative population and often relying on complex data extracted from the EHR.^{22 23 24 25}

Other strengths of this study include the very large (2,299,502 patients) and recent (2015-2017) cohort used to build our model. The study population includes various surgery types including cardiac surgery, and the risk model was validated across surgery types.

Another strength of this model is the inclusion of potentially modifiable variables such as preoperative serum sodium and hematocrit, which are potential targets for preoperative optimization. Identifying modifiable pre-op risk factors for renal failure is an unmet clinical need. Whether the changes of these modifiable factors improve the renal outcome will require further investigation.

A limitation of our model is an inability to account for multiple factors as is possible in an EHR-integrated model, for example zip code as a predictor of socioeconomic factors. The simplicity of our model was an intentional choice, making it useful in a variety of settings without requiring institutional investment in EHR integration. Additionally, the large dataset

used to develop and validate the model portends a high degree of external validity. This could be further tested by using this model at institutions that do not participate in NSQIP and evaluating its performance in that population, which will assure the generalizability of the model.

Another limitation pertains to the clinical use of this model. Though certain factors investigated are modifiable (for example, preoperative hematocrit and sodium), this analysis cannot determine causality, or the impact of modifying these factors on ultimate risk of AKI-D. An individual with a sodium <130 mEq/L can be counseled on an elevated postoperative risk of AKI-D, but it is not known if correcting the sodium prior to surgery will be protective.

This analysis shows that preoperative renal function, older age, congestive heart failure, diabetes and anemia are all significantly associated with the development of postoperative AKI-D. A clinical model incorporating 11 readily-available patient factors can be easily used by the clinician at bedside and has high predictive ability for this outcome, making it a useful and individualized tool for shared surgical decision-making. Further study will be needed to see if optimizing modifiable predictors can improve renal outcomes.

Figures and tables:

Table 1. Patient characteristics (2015- 2016)

	No AKI with Dialysis (n=1,483,169)	AKI with Dialysis (n=4,555)	p-value
Age(Mean±SD)	58.5(±16.4)	66.2(±13.6)	*
Sex (Male)	43.60	61.58	*
ASA Class 1 (%)	5.97	0.26	*
2	42.56	6.78	*
3	44.98	45.42	*
4	6.32	41.12	*
5	0.17	6.41	*
Diabetes (%)	17.19	34.23	*
Disseminated Cancer (%)	2.71	5.95	*
Smoker within one year(%)	18.04	24.13	*
COPD(%)	5.05	14.97	*
Ascites(%)	0.36	3.86	*
Congestive Heart Failure(%)	0.93	8.83	*

Hypertension on medication(%)	49.41	73.83	*
Steroid(%)	4.01	9.77	*
Weight Loss(%)	1.39	4.59	*
Bleeding disorder(%)	4.60	17.85	*
Emergency surgery(%)	10.39	37.78	*
Serum Sodium(mEq/L)	139.1	138.0	*
Serum creatinine(mg/dL)	0.92	1.96	*
White blood cell ($\times 10^9/L$)	8.2	11.4	*
Hematocrit(%)	39.8	35.3	*
Platelet($\times 10^9/L$)	249.2	233.1	*

* p<0.001

Figure 1. 30-day postoperative unadjusted mortality, 30-day hospital readmission rate and length of hospital stay associated with acute kidney injury requiring dialysis.




Figure 2. 30-day postoperative unadjusted mortality rates of patients who developed AKI requiring dialysis according to surgery types




Figure 3. Unadjusted risk of postoperative AKI requiring dialysis according to preoperative serum creatinine, sodium, and hematocrit




Figure 4. Predictive model performance with validation cohort for postoperative AKI requiring dialysis




Table 2. Predictors of postoperative renal failure requiring dialysis

	Coefficient	SE	P value	Adjusted odds ratio
Intercept	-9.2277	0.122		
Age	0.0171	0.001	<0.001	1.02 (1.015-1.02)
Ascites	0.9321	0.09	<0.001	2.54 (2.13-3.03)
Congestive heart failure	0.8077	0.061	<0.001	2.24 (1.99-2.53)
Emergency surgery	0.7446	0.042	<0.001	2.11 (1.94-2.29)
Preoperative acute kidney injury	1.1624	0.064	<0.001	3.2 (2.82-3.63)
Diabetes				
Insulin dependent	0.7346	0.042	<0.001	2.09 (1.92-2.27)
Non-insulin dependent	0.3003	0.045	<0.001	1.35 (1.24-1.48)
Serum Sodium (reference="135.1-145")(mEq/L)				
0-130	0.2974	0.073	<0.001	1.35 (1.17-1.55)
130.1-135	0.1913	0.041	<0.001	1.21 (1.12-1.31)
>145	0.3826	0.091	<0.001	1.47 (1.23-1.75)
Serum Hematocrit (%)				
0-24	0.7588	0.074	<0.001	2.14 (1.85-2.47)
24.1-30	0.6566	0.042	<0.001	1.93(1.78-2.09)
Preoperative Sepsis (reference = No)				
SIRS	0.9105	0.056	<0.001	2.49 (2.23-2.78)

Sepsis	1.182	0.057	<0.001	3.26 (2.92-3.64)
Septic Shock	1.925	0.064	<0.001	6.86 (6.04-7.78)
Surgery type (reference = anorectal, appendix)				
ENT	0.2652	0.371	0.475	1.30 (0.63-2.7)
Bariatric, stomach, esophagus	1.4485	0.127	<0.001	4.26 (3.32-5.46)
brain	0.8844	0.21	<0.001	2.42 (1.61-3.65)
cardiac	2.9198	0.141	<0.001	18.54(14.06-24.44)
Endocrine, thyroid, parathyroid, adrenal, breast	-0.4172	0.203	0.040	0.66 (0.44-0.98)
Gallbladder, biliary tract	0.5608	0.14	<0.001	1.75 (1.33-2.31)
Hernia, peritoneum, omentum, diverticulum	0.934	0.121	<0.001	2.55 (2.01-3.23)
Intestine (not rectum)	1.6948	0.111	<0.001	5.45 (4.38-6.77)
Liver, pancreas, spleen	2.577	0.126	<0.001	13.16 (10.28-16.84)
OBGYN	0.1946	0.173	0.262	1.22(0.87-1.71)
Orthopedic lower extremity, pelvis	0.7366	0.118	<0.001	2.09 (1.66-2.63)
Orthopedic upper extremity, shoulder	0.0239	0.261	0.927	1.02 (0.61-1.71)
Skin, subcutaneous, other musculoskeletal	0.9193	0.13	<0.001	2.51 (1.95-3.23)
Spine	0.6338	0.148	<0.001	1.89 (1.41-2.52)
Thoracic (non-esophageal)	1.9322	0.149	<0.001	6.91 (5.16-9.25)
Urology	1.5805	0.124	<0.001	4.86 (3.81-6.19)
Aorta	4.0768	0.128	<0.001	58.96 (45.91-75.72)

Vascular	1.8613	0.117	<0.001	6.43 (5.11-8.1)
Serum Creatinine(mg/dL)	0.4055	0.009	<0.001	1.5 (1.47-1.53)

Acknowledgments

The American College of Surgeons National Surgical Quality Improvement Program and the hospitals participating in the ACS NSQIP are the source of the data used herein; they have not verified and are not responsible for the statistical validity of the data analysis or the conclusions derived by the authors.

Funding/Support: no funding

Bibliography

1. Bihorac A, Yavas S, Subbiah S, Hobson CE, Schold JD, Gabrielli A, Layon AJ, Segal MS: Long-term risk of mortality and acute kidney injury during hospitalization after major surgery. *Ann. Surg.* 249: 851–858, 2009
2. Chertow GM, Lazarus JM, Christiansen CL, Cook EF, Hammermeister KE, Grover F, Daley J: Preoperative renal risk stratification. *Circulation* 95: 878–884, 1997
3. Siddiqui NF, Coca SG, Devereaux PJ, Jain AK, Li L, Luo J, Parikh CR, Paterson M, Philbrook HT, Wald R, Walsh M, Whitlock R, Garg AX: Secular trends in acute dialysis after elective major surgery--1995 to 2009. *CMAJ* 184: 1237–1245, 2012
4. Hobson C, Ozrazgat-Baslanti T, Kuxhausen A, Thottakkara P, Efron PA, Moore FA, Moldawer LL, Segal MS, Bihorac A: Cost and mortality associated with postoperative acute kidney injury. *Ann. Surg.* 261: 1207–1214, 2015
5. Ajitsaria P, Eissa SZ, Kerridge RK: Risk Assessment. *Curr. Anesthesiol. Rep.* 8: 1–8, 2018
6. Fleisher LA, Fleischmann KE, Auerbach AD, Barnason SA, Beckman JA, Bozkurt B, Davila-Roman VG, Gerhard-Herman MD, Holly TA, Kane GC, Marine JE, Nelson MT, Spencer CC, Thompson A, Ting HH, Uretsky BF, Wijeyesundera DN, American College of Cardiology, American Heart Association: 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. *J. Am. Coll. Cardiol.* 64: e77-137, 2014
7. Kristensen SD, Knuuti J, Saraste A, Anker S, Bøtker HE, Hert SD, Ford I, Gonzalez-Juanatey JR, Gorenek B, Heyndrickx GR, Hoes A, Huber K, Iung B, Kjeldsen KP, Longrois D, Lüscher TF, Pierard L, Pocock S, Price S, Roffi M, Authors/Task Force Members: 2014 ESC/ESA Guidelines on non-cardiac surgery: cardiovascular assessment and management: The Joint Task Force on non-cardiac surgery: cardiovascular assessment and management of the European Society of Cardiology (ESC) and the European Society of Anaesthesiology (ESA). *Eur. Heart J.* 35: 2383–2431, 2014
8. Gupta PK, Gupta H, Sundaram A, Kaushik M, Fang X, Miller WJ, Esterbrooks DJ, Hunter CB, Pipinos II, Johanning JM, Lynch TG, Forse RA, Mohiuddin SM, Mooss AN: Development and validation of a risk calculator for prediction of cardiac risk after surgery. *Circulation* 124: 381–387, 2011
9. Smetana GW, Lawrence VA, Cornell JE, American College of Physicians: Preoperative pulmonary risk stratification for noncardiothoracic surgery: systematic review for the American College of Physicians. *Ann. Intern. Med.* 144: 581–595, 2006
10. Pannu N, Graham M, Klarenbach S, Meyer S, Kieser T, Hemmelgarn B, Ye F, James M, APPROACH Investigators and the Alberta Kidney Disease Network: A new model to predict acute

- kidney injury requiring renal replacement therapy after cardiac surgery. *CMAJ* 188: 1076–1083, 2016
11. Jiang W, Teng J, Xu J, Shen B, Wang Y, Fang Y, Zou Z, Jin J, Zhuang Y, Liu L, Luo Z, Wang C, Ding X: Dynamic Predictive Scores for Cardiac Surgery-Associated Acute Kidney Injury. *J. Am. Heart Assoc.* 5: 2016
 12. Berg KS, Stenseth R, Wahba A, Pleym H, Videm V: How can we best predict acute kidney injury following cardiac surgery?: a prospective observational study. *Eur. J. Anaesthesiol.* 30: 704–712, 2013
 13. Thakar CV, Arrigain S, Worley S, Yared J-P, Paganini EP: A clinical score to predict acute renal failure after cardiac surgery. *J. Am. Soc. Nephrol.* 16: 162–168, 2005
 14. Mehta RH, Grab JD, O'Brien SM, Bridges CR, Gammie JS, Haan CK, Ferguson TB, Peterson ED, Society of Thoracic Surgeons National Cardiac Surgery Database Investigators: Bedside tool for predicting the risk of postoperative dialysis in patients undergoing cardiac surgery. *Circulation* 114: 2208–16; quiz 2208, 2006
 15. Bilimoria KY, Liu Y, Paruch JL, Zhou L, Kmieciak TE, Ko CY, Cohen ME: Development and evaluation of the universal ACS NSQIP surgical risk calculator: a decision aid and informed consent tool for patients and surgeons. *J. Am. Coll. Surg.* 217: 833–42.e1, 2013
 16. Cohen ME, Ko CY, Bilimoria KY, Zhou L, Huffman K, Wang X, Liu Y, Kraemer K, Meng X, Merkow R, Chow W, Matel B, Richards K, Hart AJ, Dimick JB, Hall BL: Optimizing ACS NSQIP modeling for evaluation of surgical quality and risk: patient risk adjustment, procedure mix adjustment, shrinkage adjustment, and surgical focus. *J. Am. Coll. Surg.* 217: 336–46.e1, 2013
 17. Hall BL, Hamilton BH, Richards K, Bilimoria KY, Cohen ME, Ko CY: Does surgical quality improve in the American College of Surgeons National Surgical Quality Improvement Program: an evaluation of all participating hospitals. *Ann. Surg.* 250: 363–376, 2009
 18. NSQIP 2105 User Guide.
 19. Suen WS, Mok CK, Chiu SW, Cheung KL, Lee WT, Cheung D, Das SR, He GW: Risk factors for development of acute renal failure (ARF) requiring dialysis in patients undergoing cardiac surgery. *Angiology* 49: 789–800, 1998
 20. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Prettenhofer P, Weiss R, Dubourg V, Vanderplas J, Passos A, Cournapeau D, Brucher M, Perrot M, Duchesnay É: Scikit-learn: Machine Learning in Python. *Journal of Machine Learning Research* 2011
 21. Thottakkara P, Ozrazgat-Baslanti T, Hupf BB, Rashidi P, Pardalos P, Momcilovic P, Bihorac A: Application of Machine Learning Techniques to High-Dimensional Clinical Data to Forecast Postoperative Complications. *PLoS ONE* 11: e0155705, 2016
 22. Kate RJ, Perez RM, Mazumdar D, Pasupathy KS, Nilakantan V: Prediction and detection models for acute kidney injury in hospitalized older adults. *BMC Med. Inform. Decis. Mak.* 16: 39, 2016
 23. Davis SE, Lasko TA, Chen G, Siew ED, Matheny ME: Calibration drift in regression and machine

- learning models for acute kidney injury. *J. Am. Med. Inform. Assoc.* 24: 1052–1061, 2017
24. Forni LG, Dawes T, Sinclair H, Cheek E, Bewick V, Dennis M, Venn R: Identifying the patient at risk of acute kidney injury: a predictive scoring system for the development of acute kidney injury in acute medical patients. *Nephron Clin. Pract.* 123: 143–150, 2013
 25. Koyner JL, Carey KA, Edelson DP, Churpek MM: The development of a machine learning inpatient acute kidney injury prediction model. *Crit. Care Med.* 46: 1070–1077, 2018