

## Ultraviolet A Radiation and COVID-19 Deaths: A Multi Country Study

Mark Cherrie, postdoctoral researcher<sup>1</sup>, Tom Clemens, lecturer<sup>1</sup>, Claudio Colandrea, researcher<sup>1</sup>, Zhiqiang Feng, senior lecturer<sup>1</sup>, David J Webb, professor<sup>2</sup>, Chris Dibben\*, professor<sup>1</sup>, Richard B Weller\*, reader (0000-0003-2550-9586)<sup>3</sup>

### Author affiliations

<sup>1</sup>School of Geosciences, University of Edinburgh, Edinburgh, UK

<sup>2</sup>Centre for Cardiovascular Science, University of Edinburgh, Edinburgh, UK

<sup>3</sup>Centre for Inflammation Research, University of Edinburgh, Edinburgh, UK

\*These authors contributed equally

Correspondence to: Professor Chris Dibben, [chris.dibben@ed.ac.uk](mailto:chris.dibben@ed.ac.uk) School of Geosciences, University of Edinburgh EH8 9XP or

Dr Richard Weller, [r.weller@ed.ac.uk](mailto:r.weller@ed.ac.uk), Department of Dermatology, Lauriston Building, Lauriston Place, Edinburgh, EH3 9HA.

## **Abstract**

### **Objectives**

To determine whether UVA exposure might be associated with COVID-19 deaths

### **Design**

Ecological model and meta-analysis

### **Setting**

2,474 counties of the contiguous USA, 6,755 municipalities in Italy, 6,274 areas in England.

Data were collected during their 'Vitamin D winter' (monthly mean  $UV_{vitd}$  of under  $165 \text{ kJ/m}^2$ ) from Jan to April 2020.

### **Participants (instead of patients or subjects)**

The 'at-risk' population is the total county population, with measures to incorporate spatial infection into the model. The model is adjusted for: age, ethnicity, socioeconomic deprivation, long term PM2.5, and UV weighted by the vitamin D action spectrum.

### **Main outcome measures**

We derive UVA measures for each area and estimate their relationship with COVID-19 mortality with a random effect for States, in a multilevel zero-inflated negative binomial model.

In the USA and England death certificates had to record COVID-19. In Italy excess deaths in 2020 over expected from 2015-19.

### **Data sources**

Satellite derived mean daily UVA dataset from Japan Aerospace Exploration Agency. Data on deaths compiled by Center for Disease Control (USA), Office for National Statistics (England) and Italian Institute of Statistics.

### **Results**

Daily mean UVA (January-April 2020) varied between 450 to 1,000 KJ/m<sup>2</sup> across the three countries. Our fully adjusted model showed an inverse correlation between UVA and COVID-19 mortality with a Mortality Risk Ratio (MRR) of 0.73 (0.62 to 0.87) per 100KJ/m<sup>2</sup> increase UVA in the USA, 0.81 (0.71 to 0.93) in Italy and 0.51 (0.39 to 0.66) in England. Pooled MRR was 0.68 (0.53 to 0.66).

## Conclusions

Our analysis, replicated in 3 independent national datasets, suggests ambient UVA exposure is associated with lower COVID-19 specific mortality. This effect is independent of vitamin D, as it occurred at irradiances below that likely to induce significant cutaneous vitamin D3 synthesis.

Causal interpretations must be made cautiously in observational studies. Nonetheless this study suggests strategies for reduction of COVID-19 mortality

## Introduction

Seasonality<sup>1</sup> and variation in temperature<sup>2</sup>, humidity<sup>3</sup> and UV radiation<sup>4</sup> are related to the incidence of several infectious diseases. COVID-19 arose less than 6 months ago, and it is thus not possible to describe seasonal variation. Nonetheless, spatial variation in levels of environmental UV in the early pandemic allows its relationship with COVID-19 mortality to be modelled. We have previously described a novel nitric oxide driven, vitamin D independent, mechanism<sup>5</sup> by which sunlight can lower blood pressure, and at the population level we have shown that UV is associated with lower blood pressure<sup>6</sup> and a reduced incidence of myocardial infarctions<sup>7</sup>. The same UV driven mechanism may also cause seasonal variation in development of diabetes and metabolic syndrome<sup>8</sup>. Given the apparent greater severity of illness and risk of death amongst those with these conditions<sup>9 10</sup> and the importance of season, we explore whether ambient UVA exposure is associated with COVID-19 deaths across the USA and replicate this analysis in independent data sets from England and Italy.

## Methods

### *Study Setting and Participants*

We use an ecological model of COVID-19 deaths across USA counties (N=2,472), English Middle Layer Super Output Areas (MSOA's) (N=6,724) and excess deaths in Italian municipalities (N=6,775). We only include counties that were experiencing levels of UV too low (monthly mean UVvitd of under 165 KJ/m<sup>2</sup>) to be inducing significant levels of cutaneous vitamin D3 synthesis during the study period ('UV vitamin D winter'). USA COVID-19 deaths were drawn from data compiled by the Center for Systems Science and Engineering at Johns Hopkins University. Data were from the period up to the 30<sup>th</sup> of April. These data are derived from death certificates, with information on cause and circumstances of death recorded, collected by the Center for Disease Control and Prevention (CDC). A COVID-19 death represents a case where the practitioner suspected that COVID-19 played a role in the death, even if it was not directly attributable to the death. English COVID-19 deaths were drawn from data compiled by the

Office for National Statistics

(<https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/deathsinvolvedcovid19bylocalareasanddeprivation/deathsoccurringbetween1marchand17april>). Data were from the period from 1<sup>st</sup> of March to the 17<sup>th</sup> of April. To be included in this dataset COVID-19 had to be mentioned on the death certificate, with a delay of usually five days between occurrence and registration. Italian excess deaths were drawn from ISTAT (Italian Institute of Statistics) (<https://www.istat.it/it/archivio/240401>) for a sample of municipalities (7,270/7,904; 92%). Data were for the period 1<sup>st</sup> of March to 30<sup>th</sup> of April for 2015-2019 and 2020. These data included all deaths registered in an integrated database, consisting of the registry source (Anagrafe Nazionale della Popolazione Residente; ANPR, and municipalities) with the Tax Register. We classified excess deaths as the positive difference in deaths between 2020 and 2015-19, with negatives recorded as zero.

#### *Ambient UV data*

We derive mean daily UVA for each study – USA (1<sup>st</sup> of Jan – 30<sup>th</sup> of April); England (1<sup>st</sup> of Jan – 17<sup>th</sup> of April) and Italy (1<sup>st</sup> of Jan – 30<sup>th</sup> of April). The UVA dataset was produced by JAXA (Japan Aerospace Exploration Agency) using the MODerate resolution Imaging Spectroradiometer (MODIS) instrument on board NASA's Aqua and Terra satellites<sup>11</sup>. Atmospheric absorption due to the ozone and water vapour (cloudiness) were accounted for by using a simplified planetary atmosphere (clear atmosphere positioned above a cloud layer). Downward irradiance values (i.e. combined direct and diffuse radiation on a horizontal plane) for UVA (315nm-400nm) were converted to daily values by using the diurnal cycle of solar zenith angle with instantaneous atmospheric conditions. These data are available to download at a 5km by 5km spatial resolution. UVA data were aggregated for USA counties, English MSOA's and Italian municipalities and expressed as mean daily KJ/m<sup>2</sup>.

UV<sub>vitd</sub> data were aggregated for US counties and expressed in mean monthly KJ/m<sup>2</sup>. We used the highest quintile Figure 1 as the cut off for year round vitamin D synthesis, which corresponds to a monthly mean of over 165 KJ/m<sup>2</sup>. It also corresponds approximately to the

37°N parallel; Holick suggests that people living North of this latitude will not receive sufficient UV for vitamin D synthesis in the winter<sup>12</sup>.

### *Covariates*

A number of demographic, socioeconomic deprivation, long term environment and infection susceptibility variables were used in the analysis.

### Demographic

We used data on county percentage of older residents ( $\geq$  age 65), percentage of Hispanic residents and percentage of Black residents for the USA. We used data from ONS mid-year population estimates of 2018 on percentage of residents who were: aged 80 or over, aged 65-79, the 2011 census on living in care homes, Black, Indian, Pakistani/Bangladeshi and Chinese in England. We used data from ISTAT on number of foreign born residents and the percentage of residents: aged 65 and above and aged 85 and above, for Italy. Older age and BAME ethnicity are associated with higher risk of COVID-19 death, possibly due to higher prevalence of comorbidities, including hypertension, heart disease and respiratory diseases<sup>10</sup>.

### Socioeconomic Deprivation

Socioeconomic deprivation was defined in the USA by the first principal component score from A Principle Component Analysis of the following county variables: percentage in poverty, median house value, median house income, percentage owner occupied and percent of population with less than a high school education. We reversed the direction of percentage in poverty and percent of population with less than a high school education so that higher score represented higher affluence. To capture socioeconomic deprivation in England we used percentage of residents who were under 21 did not enter higher education and the income deprivation score indicating of percentage of people who received benefit due to low income (<https://www.gov.uk/government/statistics/english-indices-of-deprivation-2019>). For our Italian study, we used the Italian Deprivation index, calculated by ISFOL (research Italian institute on work and social policies), which considers income, education, living conditions, unemployment

and household composition. It is hypothesised that whilst affluent citizens were more likely to get COVID-19 in the early stages of the pandemic, poorer citizens are increasingly at higher risk of infection due to essential working and death due to pre-existing health conditions<sup>13</sup>.

## Long term environment

Long term PM<sub>2.5</sub> (2000-2016) data at a 0.01° by 0.01° resolution were modelled using satellite and monitored PM<sub>2.5</sub> station data<sup>14</sup>. We used these data for both the USA and Italy. In England long term 2014-2018 PM<sub>2.5</sub> at a 1km by 1km resolution was modelled using monitored PM<sub>2.5</sub> station data (<https://uk-air.defra.gov.uk/data/pcm-data>). Higher PM<sub>2.5</sub> is linked with a range of respiratory and cardiovascular disease and shown to increase COVID mortality rate in other analyses<sup>15</sup>. Long term mean monthly winter temperature (Dec-Feb) at a 4km by 4km resolution for 2000-2016, was modelled using satellite data<sup>16</sup> for the USA. Long term mean monthly winter temperature (Dec-Feb) at a 1km by 1km resolution for 1981-2010 was modelled using interpolation of Met Office weather stations for England (<http://data.ceda.ac.uk/badc/ukmo-hadobs/data/insitu/MOHC/HadOBS/HadUK-Grid/v1.0.1.0/1km/tas/mon-30y/v20190808>). Long term median land surface temperature (Dec-Feb) daytime monthly median value at a 1km by 1km resolution for 2000-2017, was modelled using satellite data for Italy ([https://developers.google.com/earth-engine/datasets/catalog/OpenLandMap\\_CLM\\_CLM\\_LST\\_MOD11A2-DAY\\_M\\_v01](https://developers.google.com/earth-engine/datasets/catalog/OpenLandMap_CLM_CLM_LST_MOD11A2-DAY_M_v01)). Variation in temperature was recently associated with COVID-19<sup>17</sup> mortality.

## Infection Susceptibility

Infection susceptibility was defined by county population density, urban/rural status and state percentage of positive tests in the USA. We used population density from the 2018 mid-year population estimates of ONS, percentage of residents using different forms of transport (bus, train, tube) from the 2011 census and Upper Tier Local Authority (UTLA) number of days since a local authority had 10 confirmed cases. We used ISTAT population density from 2019, the municipality area, and total cases in province in Italy up to the 30<sup>th</sup> of April (<https://github.com/pcm-dpc/COVID-19/tree/master/dati-province>). In densely populated, urban or

peri-urban areas, with high use of public transport COVID-19 transmission is faster and the prevalence of cases higher. Therefore the infections susceptibility measures adjust the models for the population ‘at-risk’.

### *Statistical Analysis*

We estimate, in a multilevel zero-inflated negative binomial model (using the glmmTMB package), the relationship between ambient UVA and COVID-19 deaths (21<sup>st</sup> January-April 30<sup>th</sup>), with a state level (N=46) level random effect. The ‘at-risk’ population is the total county population, with the [1] state level random effect, [2] a measure of the proportion of population tested positive for COVID-19 at the state level and [3] measures of infection susceptibility (county population density and urban-rural status), used to incorporate viral spread into the model. The model is adjusted at the county level for: percentage of older residents ( $\geq$  age 65), Hispanic and Black residents; socioeconomic deprivation and long term modelled 2000-2016 PM<sub>2.5</sub>, long term mean winter temperature (Dec-Feb).

Our first replication of the USA model was to investigate the relationship between ambient UVA and COVID-19 deaths in England (1<sup>st</sup> of March to 17<sup>th</sup> of April), with UTLA (N=150) level random effect. The ‘at risk’ population is the total MSOA population, with the [1] UTLA level random effect, [2] number of days since a local authority had 10 confirmed cases UTLA level and [3] measures of infection susceptibility (MSOA population density, population using public transport – bus, train and tube). The model is adjusted at the MSOA level for: Long term PM<sub>2.5</sub>, long term winter temperature, Percentage of residents: aged 80+, aged 65-79, Black, Indian, Pakistani/Bangladeshi, Chinese, in care homes, in higher education and income deprivation score (Table 1).

Our second replication of the USA model was to investigate the relationship between ambient UVA and excess deaths in Italy (1<sup>st</sup> of March to 30<sup>th</sup> of April), with Province (N=104) level random effect. The ‘at risk’ population is the total municipality population, with the [1] province level random effect, [2] number of tests for COVID-19 cases at the Province level and [3] measures of infection susceptibility (municipality population density and area). The model is



adjusted at the municipality level for: Long term PM<sub>2.5</sub>, long term winter temperature, Number of foreign born, Percentage of residents: aged 65+, aged 85+ and deprivation (Table 1).

Effect estimates were presented as mortality rate ratios with 95% confidence intervals. We predicted the number of deaths per million population at suitable levels of UVA by calculating the marginal means in the 'emmeans' package. To understand the average true effect of UVA on COVID-19 related death in all similar countries to those included, we used a random effects model as part of the 'metafor' package.

All analyses were undertaken in R 3.6.1.

### Data Availability

The UVA estimates were developed by JAXA, with MODIS data provided by NASA GSFC, and available from [ftp://apollo.eorc.jaxa.jp/pub/JASMES/Global\\_05km/](ftp://apollo.eorc.jaxa.jp/pub/JASMES/Global_05km/). The long term UV<sub>vitd</sub> estimates were developed by NCAR and available from <https://www2.acom.ucar.edu/modeling/tuv-download>. The USA COVID-19 death and county level indicators of population, socioeconomic and environmental data (PM<sub>2.5</sub> and temperature) were developed previously by Xiao Wu et al., and are available here: [https://github.com/wxwx1993/PM\\_COVID](https://github.com/wxwx1993/PM_COVID).

### Code Availability

The code and data to reproduce the current study is held at [https://github.com/markocherrie/COVID19\\_UVA](https://github.com/markocherrie/COVID19_UVA).

### Results

Daily mean UVA (January-April 2020) varied between 450-1,000 KJ/m<sup>2</sup> across the three countries, with lower average levels experienced across England (Figure 2 a,b,c). Our fully adjusted model shows reductions in Mortality Risk Ratios (MRR) of 0.73 in the USA per 100 increase in UVA (KJ/m<sup>2</sup>) (Table). We found a similar size of effect in our two replication studies; an MRR in Italy of 0.81 and in England 0.49 with a pooled estimate of 0.68 (figure 3d). This

represents a halving of the average risk of death across the lower and narrower range of UVA experienced across England and across the higher and wider range across Italy and the USA (Figure 3 a,b,c)

## Discussion

Our analysis suggests that higher ambient UVA exposure is associated with lower COVID-19 specific mortality. This effect was independent of temperature and UV within the vitamin D action spectrum. Relatively higher UVA, across a country experiencing lower average levels of UVA during and immediately preceding the pandemic (England), is associated with greater falls in mortality risk compared to countries (Italy and the USA) with a higher average level suggesting an exposure tolerance effect.

There has been considerable debate around the possible role of Vitamin D supplementation for prevention of COVID-19, but to date the data do not support this intervention<sup>18</sup>. We have been able to show a possible benefit from UV at wavelengths that do not cause synthesis of vitamin D. This suggests that explanations for seasonal and geographic variations in COVID-19 prevalence<sup>19</sup> need to consider mechanisms other than just vitamin D.

Seasonal variations in disease incidence can be caused by environmental, behavioural, and immunological factors with the relative importance of these varying by location and disease. UV may have a direct effect on the viability of SARS-CoV2 virus in airborne droplets and on fomites, thus reducing both infection rates, and also the size of inoculum in those becoming infected, with correspondingly reduced disease severity<sup>20 21</sup>. UVA radiation photo-releases nitric oxide (NO) from stores in the skin whence it is mobilised to the systemic circulation, causing vasodilation and reduction in blood pressure<sup>5</sup>, offering cardiovascular and metabolic benefits from UV exposure<sup>5 8</sup>. As cardio-metabolic disease and possibly hypertension<sup>22</sup> increase the risk of death from COVID-19, any UV driven improvements in these risk factors would be expected to reduce mortality<sup>23</sup>. Nitric oxide may also have a specific effect on COVID-19. It inhibits replication of SARS CoV<sup>24</sup>, probably by S-nitrosating the spike protein, thus preventing the post translational palmitoylation of the spike protein, required for fusion with its cognate angiotensin converting enzyme 2 receptor(ACE2R).<sup>25</sup> The spike protein of SARS CoV is highly

homologous to that of SARS CoV2<sup>26 27</sup> suggesting that NO may similarly limit binding to ACE2R by SARS CoV2 with reduction in disease transmission and severity. Endothelial damage, with reduced homeostatic endothelial NO synthase activity may underlie widespread organ involvement<sup>28</sup>. This would be mitigated by photochemical NO production.

There are several weaknesses in this study. UVA and covariates are measured at the small area level not as individual level exposures, there could be misclassification of deaths and infection was only estimated within the model and with indirect measures. However any resulting measurement errors seem unlikely to be correlated with spatial variation in UVA and therefore biasing. Importantly we replicate the finding across three countries with very different health systems and economic and social structures, pandemic situations and measure COVID-19 related deaths in two different ways.

The study was observational and therefore any causal interpretation needs to be taken with caution. However, if the relationship identified proves to be causal, it suggests that optimising sun exposure may be a possible public health intervention. Given that the effect appears independent of a vitamin D pathway, it suggests possible new COVID-19 therapies and in particular the importance of exploring the role of circulating NO.

### **Acknowledgement:**

This work was supported by Health Data Research UK which receives its funding from **HDR UK Ltd (HDR-5012)** funded by the UK Medical Research Council, Engineering and Physical Sciences Research Council, Economic and Social Research Council, Department of Health and Social Care (England), Chief Scientist Office of the Scottish Government Health and Social Care Directorates, Health and Social Care Research and Development Division (Welsh Government), Public Health Agency (Northern Ireland), British Heart Foundation (BHF) and the Wellcome Trust

### **Ethics declaration**

## Competing interests

RW is a director of Relaxsol (company making sunscreen).

## References

1. Fisman D. Seasonality of viral infections: mechanisms and unknowns. *Clinical Microbiology and Infection* 2012;18(10):946-54. doi: 10.1111/j.1469-0691.2012.03968.x
2. Earn DJ, He D, Loeb MB, et al. Effects of school closure on incidence of pandemic influenza in Alberta, Canada. *Annals of internal medicine* 2012;156(3):173-81.
3. Shaman J, Pitzer VE, Viboud C, et al. Absolute Humidity and the Seasonal Onset of Influenza in the Continental United States. *PLOS Biology* 2010;8(2):e1000316. doi: 10.1371/journal.pbio.1000316
4. Charland KM, Buckeridge DL, Sturtevant JL, et al. Effect of environmental factors on the spatio-temporal patterns of influenza spread. *Epidemiol Infect* 2009;137(10):1377-87. doi: 10.1017/S0950268809002283 [published Online First: 2009/03/20]
5. Liu D, Fernandez BO, Hamilton A, et al. UVA irradiation of human skin vasodilates arterial vasculature and lowers blood pressure independently of nitric oxide synthase. *J Invest Dermatol* 2014;134(7):1839-46. doi: 10.1038/jid.2014.27
6. Weller RB, Wang Y, He J, et al. Does Incident Solar Ultraviolet Radiation Lower Blood Pressure? *J Am Heart Assoc* 2020;9(5):e013837. doi: 10.1161/JAHA.119.013837 [published Online First: 2020/02/29]
7. Mackay DF, Clemens TL, Hastie CE, et al. UVA and Seasonal Patterning of 56 370 Myocardial Infarctions Across Scotland, 2000-2011. *J Am Heart Assoc* 2019;8(23):e012551. doi: 10.1161/JAHA.119.012551 [published Online First: 2019/11/22]
8. Geldenhuys S, Hart PH, Endersby R, et al. Ultraviolet radiation suppresses obesity and symptoms of metabolic syndrome independently of vitamin D in mice fed a high-fat diet. *Diabetes* 2014;63(11):3759-69. doi: 10.2337/db13-1675
9. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA* 2020 doi: 10.1001/jama.2020.2648 [published Online First: 2020/02/25]
10. Guan W-j, Liang W-h, Zhao Y, et al. Comorbidity and its impact on 1590 patients with Covid-19 in China: A Nationwide Analysis. *European Respiratory Journal* 2020:2000547. doi: 10.1183/13993003.00547-2020
11. Saigusa N, Ichii K, Murakami H, et al. Impact of meteorological anomalies in the 2003 summer on Gross Primary Productivity in East Asia. *Biogeosciences* 2010;7(2):641-55. doi: DOI 10.5194/bg-7-641-2010
12. Holick MF. High Prevalence of Vitamin D Inadequacy and Implications for Health. *Mayo Clinic Proceedings* 2006;81(3):353-73. doi: <https://doi.org/10.4065/81.3.353>
13. Han Y, Li VOK, Lam JCK, et al. Who is more susceptible to Covid-19 infection and mortality in the States? *medRxiv* 2020:2020.05.01.20087403. doi: 10.1101/2020.05.01.20087403

14. Atmospheric Composition Analysis Group [Available from: [http://fizz.phys.dal.ca/~atmos/martin/?page\\_id=140](http://fizz.phys.dal.ca/~atmos/martin/?page_id=140) accessed 15/05/2020 2020.
15. Wu X, Nethery RC, Sabath BM, et al. Exposure to air pollution and COVID-19 mortality in the United States: A nationwide cross-sectional study. *medRxiv* 2020:2020.04.05.20054502. doi: 10.1101/2020.04.05.20054502
16. Idaho Uo. GRIDMET: University of Idaho Gridded Surface Meteorological Dataset 2020 [Available from: [https://developers.google.com/earth-engine/datasets/catalog/IDAHO\\_EPSCOR\\_GRIDMET](https://developers.google.com/earth-engine/datasets/catalog/IDAHO_EPSCOR_GRIDMET) accessed 15/05/2020 2020.
17. Ma Y, Zhao Y, Liu J, et al. Effects of temperature variation and humidity on the death of COVID-19 in Wuhan, China. *Science of The Total Environment* 2020;724:138226. doi: <https://doi.org/10.1016/j.scitotenv.2020.138226>
18. Torjesen I. Evidence does not support vitamin D for reducing respiratory infections, reviews conclude. *BMJ* 2020;369:m2629. doi: 10.1136/bmj.m2629
19. Rhodes JM, Subramanian S, Laird E, et al. Editorial: low population mortality from COVID-19 in countries south of latitude 35 degrees North supports vitamin D as a factor determining severity. *Alimentary Pharmacology & Therapeutics* 2020;51(12):1434-37. doi: 10.1111/apt.15777
20. Paulo AC, Correia-Neves M, Domingos T, et al. Influenza Infectious Dose May Explain the High Mortality of the Second and Third Wave of 1918–1919 Influenza Pandemic. *PLOS ONE* 2010;5(7):e11655. doi: 10.1371/journal.pone.0011655
21. Yu X, Sun S, Shi Y, et al. SARS-CoV-2 viral load in sputum correlates with risk of COVID-19 progression. *Critical Care* 2020;24(1):170. doi: 10.1186/s13054-020-02893-8
22. Gao C, Cai Y, Zhang K, et al. Association of hypertension and antihypertensive treatment with COVID-19 mortality: a retrospective observational study. *European Heart Journal* 2020 doi: 10.1093/eurheartj/ehaa433
23. Lindqvist PG. The Winding Path Towards an Inverse Relationship Between Sun Exposure and All-cause Mortality. *Anticancer Res* 2018;38(2):1173-78. doi: 10.21873/anticancer.12337
24. Åkerström S, Mousavi-Jazi M, Klingström J, et al. Nitric Oxide Inhibits the Replication Cycle of Severe Acute Respiratory Syndrome Coronavirus. *Journal of Virology* 2005;79(3):1966-69. doi: 10.1128/jvi.79.3.1966-1969.2005
25. Åkerström S, Gunalan V, Keng CT, et al. Dual effect of nitric oxide on SARS-CoV replication: Viral RNA production and palmitoylation of the S protein are affected. *Virology* 2009;395(1):1-9. doi: <https://doi.org/10.1016/j.virol.2009.09.007>
26. Grifoni A, Sidney J, Zhang Y, et al. A Sequence Homology and Bioinformatic Approach Can Predict Candidate Targets for Immune Responses to SARS-CoV-2. *Cell Host Microbe* 2020;27(4):671-80 e2. doi: 10.1016/j.chom.2020.03.002 [published Online First: 2020/03/19]
27. Wrapp D, Wang N, Corbett KS, et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* 2020;367(6483):1260-63. doi: 10.1126/science.abb2507 [published Online First: 2020/02/23]
28. Teuwen L-A, Geldhof V, Pasut A, et al. COVID-19: the vasculature unleashed. *Nature Reviews Immunology* 2020 doi: 10.1038/s41577-020-0343-0

Figure 1. Quintiles of  $UV_{vitD}$




Figure 2a: Daily mean UVA (KJ/m<sup>2</sup>) in the contiguous USA between Jan-Apr 30<sup>th</sup> 2020.




Figure 2b: Daily mean UVA (KJ/m<sup>2</sup>) in England between Jan 1<sup>st</sup> -Apr 17<sup>th</sup> 2020






Figure 2c: Daily mean UVA (KJ/m<sup>2</sup>) in the Italy between Jan 1<sup>st</sup> - Apr 30<sup>th</sup> 2020




Figure 3a: Predicted COVID-19 deaths by experienced ambient UVA exposure in the contiguous USA




Figure 3b: Predicted COVID-19 deaths by experienced ambient UVA exposure in England




Figure 3c: Predicted excess deaths by experienced ambient UVA exposure in the Italy




Figure 3d: Forrest plot of estimates for Mortality Rate Ratio per 100 KJ/m<sup>2</sup> increase in mean daily UVA




Table : Model estimates for the relationship between UVA and covid-19 death

Sample	Negative Binomial Model	Zero-Inflated Model	Estimate (mortality rate ratio; 95% CI)	St.Dev of Group
USA (N=2,474)	<p>Average UVA Jan-Apr 30<sup>th</sup> (696 KJ/m<sup>2</sup> ± 83; per 100KJ/m<sup>2</sup>)</p> <p>Long term: PM<sub>2.5</sub>; UV<sub>vitd</sub>, winter temperature; Percentage of residents: 65+; Black, Hispanic; deprivation score; urban/rural; state number of positive COVID-19 cases</p>	<p>Percentage of residents: 65+; Black, Hispanic; deprivation score; urban/rural; state number of positive COVID-19 cases</p>	0.73 (0.62-0.87)	0.61  State (n=46)
England (N=6,724)	<p>Average UVA Jan-Apr 17<sup>th</sup> (412 KJ/m<sup>2</sup> ± 18; per 100 KJ/m<sup>2</sup>)</p> <p>Long term: PM<sub>2.5</sub>, long term winter temperature; Percentage of residents: aged 80+, aged 65-79, Black, Indian, Pakistani/Bangladeshi, Chinese, in care homes, in</p>	<p>Percentage of residents: aged 80+, aged 65-79, Black, Indian, Pakistani/Bangladeshi, Chinese, in care homes, in higher education, using public transport</p>	0.49 (0.38-0.64)	0.38  Local Authorities (n=150)

	higher education; income deprivation score; Upper Tier Local Authority (UTLA) number of days since a local authority had 10 confirmed cases.	(bus, train, tube); income deprivation score; population density; UTLA number of days since a local authority had 10 confirmed cases		
Italy (N=6,775)	Average UVA Jan-Apr 30 <sup>th</sup> (717 KJ/m <sup>2</sup> ± 52; per 100KJ/m <sup>2</sup> )  Long term: PM <sub>2.5</sub> , long term winter temperature; Number of foreign born; Percentage of residents: aged 65+, aged 85+; population density; municipality area, deprivation score; total cases in province	Number of foreign born; Percentage of residents: aged 65+, aged 85+; population density; municipality area, deprivation score; total cases in province	0.81 (0.71-0.93)	0.49  Province (n=104)