

1 Ultrametric model for covid-19 dynamics: an
2 attempt to explain slow approaching herd
3 immunity in Sweden

4 Andrei Khrennikov
Linnaeus University, International Center for Mathematical Modeling
in Physics and Cognitive Sciences Växjö, SE-351 95, Sweden

5 July 3, 2020

6 **Abstract**

7 We present a mathematical model of infection dynamics that might
8 explain slower approaching the herd immunity during the covid-19 epi-
9 demy in Sweden than it was predicted by a variety of other models;
10 see graphs Fig. 2. The new model takes into account the hierar-
11 chic structure of social clusters in the human society. We apply the
12 well developed theory of random walk on the energy landscapes rep-
13 resented mathematically with ultrametric spaces. This theory was
14 created for applications to spin glasses and protein dynamics. To
15 move from one social cluster (valley) to another, the virus (its car-
16 rier) should cross a social barrier between them. The magnitude of a
17 barrier depends on the number of social hierarchy's levels composing
18 this barrier. As the most appropriate for the recent situation in Swe-
19 den, we consider linearly increasing (with respect to hierarchy's levels)
20 barriers. This structure of barriers matches with a rather soft regula-
21 tions imposed in Sweden in March 2020. In this model, the infection
22 spreads rather easily inside a social cluster (say working collective),
23 but jumps to other clusters are constrained by social barriers. This
24 model's feature matches with the real situation during the covid-19
25 epidemy, with its cluster spreading structure. Clusters need not be
26 determined solely geographically, they are based on a number of hi-
27 erarchically ordered social coordinates. The model differs crucially
28 from the standard mathematical models of spread of disease, such as
29 the SIR-model. In particular, our model describes such a specialty of
30 spread of covid-19 virus as the presence of "super-spreaders" who by

31 performing a kind of random walk on a hierarchic landscape of social
32 clusters spreads infection. In future, this model will be completed by
33 adding the SIR-type counterpart. But, the latter is not a specialty of
34 covid-19 spreading.

35 **keywords: covid-19, epidemic, infection spreading, herd**
36 **immunity, hierarchy of social clusters, energy landscapes,**
37 **random walk, trees, ultrametric spaces, social barriers, linear**
38 **growing barriers**

39 1 Introduction

40 The dynamics of propagation of the covid-19 pandemy has many un-
41 usual features that were not reflected by classical mathematical mod-
42 eling of epidemic dynamics (see, e.g., [1]-[4]). In spite the tremendous
43 efforts [5]-[11], we still do not have a mathematical model describing
44 adequately this dynamics. In this complex situation, one can consider
45 models reflecting only some *specialties of propagation of covid-19 in-*
46 *fection* throughout the human society. This paper is a step in this
47 direction. We want to *explain essentially slower than expected ap-*
48 *proaching the herd immunity¹ in Swedish population* (see, e.g., [12]-[14]
49 for reports from Public Health Institute of Sweden, [7]-[9] for attempts
50 of mathematical modeling and [32]-[36] for reports from massmedia).²
51 Our model is based on the advanced mathematics, theory of random
52 walks on hierarchic structures. We use the results of the well known
53 paper [19] on random walk on p -adic trees, the simplest hierarchic
54 structures endowed with ultrametric.

55 The main output that can be interesting say for epidemiologists are
56 the graphs presented in section 4 showing t^{-a} dynamics of probability
57 to become infected in clustered society in long run of an epidemic; so
58 to say *at the phase of approaching of some immunity level*. The model
59 differs crucially from the standard models of spread of disease, such
60 as the SIR-model [4]. Besides explaining slow growth of immunity,
61 our model describes such a specialty of spread of covid-19 virus as
62 the presence of “super-spreaders” who by performing a kind of ran-
63 dom walk on a hierarchic landscape of social clusters spreads infection
64 (section 5). In future, the model will be completed by adding the
65 SIR-type counterpart. But, the latter is not a specialty of covid-19

¹In particular, the mathematical model of covid-19 epidemic dynamics of Tom Britton [7, 8] that was used by Swedish State Health Authority predicted that the herd immunity will be approached already in May.

²As is well known and widely debated in mass-media, the Swedish government chosen its own way to handle covid-19 epidemic, namely, without any kind of lock-down. The main aim of such a policy is approaching the herd immunity.

66 spreading. We hope that this paper may attract attention of experts
67 in epidemiology and sociology (the model explores the structure of
68 social connections in the society) who may stimulate its further devel-
69 opment.

70 Our basic assumption (that, in fact, led to consideration of the ul-
71 trametric mathematical model) is that infection's distribution in the
72 society has the structure of disjoint social clusters.³ Such a cluster can
73 be a collective of some enterprise or a state department, say clerks of
74 community office of some town or the personal of some hospital. In-
75 side such a cluster people still have relatively high degree of social
76 connections (see appendix 2). However, even the mild isolation poli-
77 cy of Swedish authorities erected sufficiently high *barriers between*
78 *clusters*, because people terminated many sorts of social contacts. In
79 particular, the government did not recommend travels inside the coun-
80 try or abroad. (They were not strictly forbidden, but the majority of
81 citizens followed these recommendations.) Another characteristic fea-
82 ture of the model is that connections between social clusters that can
83 be used by the virus to spread have *the hierarchic structure*. For ex-
84 ample, the community office has higher level in the social hierarchy
85 than the working place. Any worker can have some things to do with
86 community even during the period of epidemy.

87 Thus the basic assumption of our model is the following one:

88 **Assumption:** *Infection distribution in population has the hierar-*
89 *chic social cluster structure.*

90 Starting with this assumption, we design a mathematical model
91 implying

92 **Consequence 1.** *Slower than predicted approaching the herd im-*
93 *munity.*

94 **Consequence 2.** *Slow decay of epidemy.*

95 The problem of approaching the herd immunity is especially impor-
96 tant in the light of Swedish experience and its consequences for states'
97 policies in preparation to the second wave of covid-19 or other simi-
98 lar viruses. The herd immunity is mathematically formalized through
99 the probability for a person to become infected at time t , see Fig. ??
100 (section 4).

101 To model cluster dynamics, we use random walk on ultrametric
102 spaces [19]. It was widely used in studied in physics and microbiology,

³“Disjoint” has the meaning of separation in social space. In physical space, clusters can essentially overlap. Social separation does not complete, say of the form of walls. There are social barriers, but a person (virus-carrier) can hope over barriers, with some probability.

103 see [19], [20]-[25] and references herein. Geometrically ultrametric
104 spaces have the treelike structure. The simplest trees are homogeneous
105 trees with the fixed number of branches p leaving each vertex, p -adic
106 trees.

107 Random walks on trees describe dynamics on *energy landscapes*.
108 There are given energy barriers Δ_m separating valleys, movement from
109 one valley to another valley is constrained by necessity to jump over
110 a barrier between them.

111 In our model of the covid-19 epidemic, the virus (or its carrier)
112 randomly walks in socially clustered society. In our model, it starts
113 just from a single social cluster (this assumption is used for mathe-
114 matical simplicity); it spreads relatively easily inside any cluster, but
115 to approach other social clusters it should “jump over social barriers”.
116 Dynamics depends heavily on two parameters: on the magnitudes of
117 the social barriers (the type of their growth between different levels of
118 hierarchy) and on the social analog of temperature. The magnitude
119 of a social barrier depends only on the number of hierarchy levels
120 composing it, but not on a social cluster. This is the very important
121 constraint on the population, it should be sufficiently homogeneous
122 with respect to the magnitudes of social barriers.

123 The configuration space of dynamics is the tree of social connec-
124 tions between people. In epidemic modeling, it is natural to assume
125 the presence of the hierarchic structure in social clustering of people,
126 by ranging basic social parameters coupled to infection. This rep-
127 resentation of individuals, as *vectors of hierarchically ordered social*
128 *coordinates* have been already used by the author and his collabo-
129 rators in a series of studies in cognition, psychology, and sociology
130 [26]-[31].

131 Just before submission of this preprint, I discovered the recent
132 paper of Britton et al. [15]. One of its authors, Britton, initiated in-
133 tensive mathematical modeling of spreading covid-19 in Sweden and,
134 in particular, he is famous in Sweden for his prognoses of approaching
135 the herd immunity in Swedish population [7]-[9]. However, his prog-
136 noses did not match with the real dynamics of epidemic in Sweden.
137 In the new modeling [15], Britton et al. started to take into account
138 heterogeneity of population (cf. [16]-[18]). This is a step towards cou-
139 pling with our model. Before to discuss similarities and differences of
140 two models in more detail, I should study article [15] more carefully.

141 2 Social trees

142 We represent the human society as a system of hierarchically coupled
143 disjoint clusters. There are many ways of creation of such representa-

144 tions. We present one of these possibilities that was used in my works
145 of ultrametric modeling in cognition and sociology (see, e.g., [26]-[30]).
146 The treelike representation of *social types* is based on selection of hier-
147 archically ordered social factors enumerated as $m = 0, 1, 2, 3, \dots$; factor
148 $m = 0$ is the most important, $m = 1$ is less important and so on. A
149 social type is represented by a vector

$$x = (x_0, x_1, \dots, x_{n-1}), \quad (1)$$

150 where its coordinates x_m take (typically) discrete values quantifying
151 the m -th factor. In the simplest case, x_m takes two values, “yes”/“no”,
152 or 1/0. We call numbers (x_m) *social coordinates*. The vector represen-
153 tation of social types and individuals is widely used in sociology and
154 psychology. The main distinguishing feature of our model is endowing
155 the space of vectors with the special metric reflecting the hierarchic
156 structure corresponding the order of social factors (see, e.g., [26]-[30]
157 on application to cognition and psychology). The space of all vectors
158 of the form (4) is called the hierarchic social space.

159 Since in the covid-19 situation the state plays the crucial role as
160 the policy determining organ, it is natural to select $m = 0$ as the
161 state that population is under consideration. However, since the ma-
162 jority of states selected the lock down policy that was not oriented
163 towards approaching the herd immunity, we restrict consideration to
164 the Swedish population and use $m = 0$ for the next basic social factor,
165 namely, for individual’s age, then say $m = 1$ for the presence of one
166 chronic diseases, $m = 2$ for gender, $m = 3$ for race, $m = 4$ for the
167 town of location, $m = 6$ for profession, $m = 7$ for the level of social
168 activity, $m = 5$ the district in the town, $m = 8$ for the number of chil-
169 dren, and so on. We understand that such ranking of the basic social
170 factors related to the covid-19 epidemy may be naive and incomplete.
171 The contribution of sociologists, psychologists, and epidemiologists
172 can improve the present model essentially.

173 For mathematical simplicity, we consider p -adic coordinates, $x_m =$
174 $0, 1, \dots, p - 1$, where $p > 1$ is a prime number. The space of all such
175 vectors denote by the symbol $Z_{p;n}$ (p is fixed). For analytical compu-
176 tations, it is convenient to represent elements of this space by natural
177 numbers

$$x = x_0 + x_1p + x_1p^2 + x_{n-1}p^{n-1}, x_m \in \{0, 1, \dots, p - 1\}. \quad (2)$$

178 Thus the points of the hierarchic social space can be represented by
179 natural numbers, $Z_{p;n} = \{0, \dots, p^n - 1\}$. The number of its points grows
180 exponentially with the number of social coordinates n .

181 We now turn to the definition of a metric on $Z_{p;n}$. Consider two
182 social vectors $x = (x_1, \dots, x_{n-1})$ and $y = (y_1, \dots, y_{n-1})$. Let their firsts

183 k coordinates are equal, $x_0 = y_0, \dots, x_{k-1} = y_{k-1}$, but the k th co-
 184 ordinates are different, $x_k \neq y_k$. Then the hierarchic social distance
 185 between these social types, $d(x, y) = n - k$. The first social coordinates
 186 are the most important: longer the common initial segment of vectors
 187 corresponds to closer social type, increase of k implies decrease of dis-
 188 tance between two social types. For example, let $k = n - 1$, i.e., two
 189 points differ only in the last coordinate, then $d(x, y) = 1$. This is the
 190 minimal possible distance in $Z_{p;n}$. (The coordinate x_n has the minimal
 191 degree of importance.) If the vectors differ already by the first coor-
 192 dinate, i.e., $x_0 \neq y_0$, then $d(x, y) = n$. This is the maximal possible
 193 distance between points in space $Z_{p;n}$. Distance d is ultrametric, it
 194 satisfies *the strong triangle inequality*:

$$d(x, y) \leq \max\{d(x, z), d(y, z)\}, \quad (3)$$

195 for any triple of points $x, y, z \in Z_{p;n}$. Here in each triangle, the third
 196 side is less or equal not only to the sum of two other sides (as usual),
 197 but even to their maximum.

198 As usual in a metric space we can introduce balls, $B_N(a) = \{x \in$
 199 $Z_{p;n} : d(a, x) \leq N\}$, where $N = 1, \dots, n$, and $a = (a_0, \dots, a_{n-1})$ is some
 200 point in $Z_{p;n}$, ball's center. In an ultrametric space, *any two balls are*
 201 *either disjoint or one is contained in another and any point of a ball*
 202 *can be selected as its center.*

For our modeling, it is important that the space $Z_{p;n}$ can be split
 into disjoint social clusters. (As we shall see soon, these clusters are,
 in fact, balls.) Each cluster is determined by fixing a few first (the
 most important) social coordinates,

$$Z_{p;n} = \cup_{j=0}^{p-1} C_j,$$

where $C_j = \{x : x_0 = j\}$. This cluster representation corresponds to
 the first level of social hierarchy, we distinguish points by their most
 important coordinate. Each of clusters C_j can be represented similarly
 as

$$C_j = \cup_{i=0}^{p-1} C_{ji},$$

203 where $C_{ji} = \{x : x_0 = j, x_1 = i\}$ are clusters of the deeper hierarchic
 204 level and so on, up to the single-point clusters corresponding to fixing
 205 all social coordinates.

Clusters are, in fact, ultrametric balls:

$$C_{i_0 \dots i_{k-1}} = \{x : x_0 = i_0, \dots, x_{k-1} = i_{k-1}\} = B_{n-k}(a),$$

206 where a is any point of the form $a_0 = i_0, \dots, a_{k-1} = i_{k-1}$ and arbitrary
 207 coordinates $a_j, j = k, \dots, n - 1$.

208 Geometrically space $Z_{p;n}$ is represented as a homogeneous tree with
209 p branches leaving each vertex. Cluster is a bunch of branches having
210 the common root. By extending this root we split the cluster into
211 subclusters.

212 Now we consider the procedure of extension of a social tree by
213 adding new social coordinates, so from tree $Z_{p;n}$ to tree $Z_{p;N}$, where
214 $N > n$. As the result of such an extension, each point of social space
215 $Z_{p;n}$ becomes a social cluster in social space $Z_{p;N}$. In principle, it is
216 impossible to determine a social type by fixing any finite number of
217 social coordinates. Hence, We have to consider infinite sequences of
218 coordinates:

$$x = (x_0, x_1, \dots, x_{n-1}, \dots), \quad x_j = 0, 1, \dots, p - 1. \quad (4)$$

219 Denote the space of such sequences by the symbol Z_p . This is the
220 complete hierarchic social space. Points of finite trees represent social
221 clusters.

222 3 Dynamics

223 Consider a tree with n levels of social hierarchy, $Z_{p;n}$. Suppose that
224 initially infected people are concentrated just in one social cluster
225 represented by a single point $x \in Z_{p;n}$. Consider the initial probability
226 for a person in this cluster to become infected, $P(y, 0) = \delta(y - x)$.

227 The virus plays the role of a system moving through barriers in
228 models of dynamics on energy landscapes (see [19], [20]-[25] and refer-
229 ences herein). In our case, these are social barriers between social
230 clusters of population. The virus performs a complex random walk
231 motion inside each social cluster moving in its sub-clusters, goes out
232 of it and spreads through the whole population, sometimes the virus
233 comes back to the original cluster from other social clusters that have
234 been infected from this initial source of infection, and so on. During
235 this motion the virus should cross numerous social barriers.

236 Instead of virus walking through the social tree, we can consider
237 a person. A person of the social type $x = (x_1, \dots, x_n)$ can interact
238 with persons of the social type, $y = (x_1, \dots, x_k - 1, y_k, \dots, y_n)$, where
239 $x_k \neq y_k$ by crossing the social barrier for $n - k$ levels of hierarchy. The
240 temporal sequence of social contacts of some persons can have a very
241 complicated trajectory, visiting numerous clusters (but the probability
242 of approaching a cluster depends crucially on social barriers).

243 We are interested in *the relaxation process* that is understood as
244 the evolution of the probability to become infected for a person from
245 the social cluster x , i.e., in the time dependence of the quantity $P(x, t)$.

246 We do not want to go into details, since the problem of relaxation
247 was well studied in physics and microbiology, see [19], [20]-[25] and
248 references herein. The relaxation regime depends crucially on the
249 barriers' magnitude, how rapidly they grow up on the way from one
250 cluster to another.

251 4 Random walk on on infinite trees: 252 temporal behavior of probability

253 As in the previous section, we a random walk on a finite tree. Here
254 we follow the paper of Ogielski [19]. Let us consider a finite tree with
255 n levels. Thus there are 2^n points at the last level. They enumerate
256 the total population: $i = 0, \dots, 2^n - 1$.

257 Let virus encounters a barrier of size Δ_m , in hopping a distance m
258 (crossing m levels of hierarchy), where $\Delta_1 < \Delta_2 < \dots < \Delta_m < \dots$. It
259 is supposed that barriers Δ_m are the same for all social clusters, i.e.,
260 they depend only on distance, but not on clusters.

261 The probability to jump over the barrier Δ_m has the form (up to
262 the normalization constant):

$$\mathbf{P}(m) = e^{-\frac{\Delta_m}{T}}, \quad (5)$$

263 where T is model's parameter; in physics it has the meaning of *tem-*
264 *perature* of the environment; we also interpret it similarly, as a kind of
265 *social temperature*, cf. [29]. Probability $\mathbf{P}(m)$ grows with growth of
266 social temperature T . For high social temperatures, jumps are easier.

267 Consider the energy landscape with a uniform barrier Δ , at every
268 branch point; that is, a jump of distance 1 involves surmounting a
269 barrier Δ , of distance 2, a barrier 2Δ , and so on. Hence, barriers
270 linearly grow with distance m ,

$$\Delta_m = m\Delta, m = 1, 2, \dots \quad (6)$$

271 It seems that this type of behavior is the most natural from the
272 viewpoint of social connections during the covid-19 epidemic in Swe-
273 den. Barriers are sufficiently high, but they still are not walls as during
274 the rigid quarantine (as say in Italy, France, or Russia). The probabil-
275 ity to jump over the barrier Δ_m has the form (up to the normalization
276 constant):

$$\mathbf{P}(m) = e^{-\frac{m\Delta}{T}}. \quad (7)$$

277 The probability depends on the height Δ of the minimal social barrier,
278 but, for large m , its contribution to is not so important.

279 By using random walk on the tree with n levels of hierarchy and
 280 approaching $n \rightarrow \infty$ one can derive the following asymptotic behavior
 281 of the relaxation probability:

$$P(x, t) = t^{-T \ln p / \Delta}, t \rightarrow \infty. \quad (8)$$

282 Set $\kappa = T \ln p / \Delta$. If $\kappa \ll 1$, e.g., the social temperature is low and
 283 the primary social barrier Δ is relatively large, then the probability for
 284 a person in the original social cluster x to become infected decreases
 285 rather slowly, see Fig. 1.




Figure 1: Asymptotic behavior of probability to become infected. (For fixed social temperature T , the upper graphs correspond to one-step barrier growth 10 and 100 times, respectively.)

286 Hence immunity against covid-19 increases also slowly, see Fig. 2,
 287 as function

$$1 - P_I(x, t) \sim 1 - t^{-T \ln p / \Delta}, t \rightarrow \infty. \quad (9)$$




Figure 2: Asymptotic behavior of probability to become immune; increasing of the herd immunity (for fixed social temperature T , the upper graphs correspond to one-step barrier growth 10 and 100 times, respectively).

5 Average social distance traveled by infection spreader

In our mathematical model, any infection spreader travels through the social tree, he-she visits a few social clusters and infects people in these clusters. Theory of random walks in ultrametric spaces predicts the average social distance for spreader's travel through clusters, starting at some fixed cluster x and jumping to other cluster y ,

$$\langle d(x, t) \rangle = \sum_{y \in Z_{n,p}} d(x, y) P(y, t) \quad (10)$$

For linearly growing social barriers, and $n \rightarrow \infty$, the asymptotic behavior has the following form:

$$\langle d(x, t) \rangle \sim (T/\Delta) \log t, \quad t \rightarrow \infty. \quad (11)$$

This average distance goes to infinity. (As can be expected, higher social temperature T and lower one-step social barrier Δ induce more rapid growth.) Although the log-growth is relatively slow, it, nevertheless, implies very extended spread of the infection. Unsoundness of $\langle d(x, t) \rangle$ can be associated with the presence of super-spreaders who jump even over high social barriers and spread the virus to social clusters that are far from the original source of infection. We repeat once

304 again that the distance under consideration is in social and not in
305 physical space.

306 6 Concluding remarks

307 The presented ultrametric model with random walk dynamics on energy
308 landscapes describes well the social cluster structure of spreading
309 of covid-19. The model was designed to explain mathematically
310 the slow approaching the herd immunity in Swedish population during
311 March-June 2020, essentially slower than it was predicted by all
312 mathematical models.⁴ The model elevates the role of the social dimension
313 of infection spreading comparing with its purely bio-medical
314 dimension. The bio-medical restrictions in Sweden were very soft; in
315 particular, no face-masks, 1,5 meter social distance, public transportation
316 worked more or less as usual (see appendix 2 for details). The
317 most typical infecting took place in working collectives in that social
318 contacts were not occasional, as say in metro, or in a bus, or even in
319 a restaurant during lunch, but close and permanent, on the everyday
320 basis.

321 We applied to the new area of research, to epidemiology, mathematical
322 theory developed for applications in statistical physics (spin glasses)
323 and microbiology (protein folding): ultrametric random walk
324 describing dynamics on complex energy landscapes with the hierarchic
325 structure of barriers between valleys. The presence of social barriers
326 growing with hierarchy's levels makes the evolution of epidemy
327 essentially slower than in models which do not take into account the
328 cluster-character of infection spreading. In particular, our model is
329 purely diffusional, cf. with standard SIR-model [4]. We motivate
330 consideration of linearly growing social barriers in Sweden with its
331 soft epidemiological restrictions.

332 Although the model is very simplified it reflects some features of
333 the covid-19 epidemic, especially its the social cluster character. We
334 hope that our model will stimulate further development of ultrametric
335 epidemiological models that take into account the hierarchic social
336 clustering of population.

⁴See appendix 1; finally, the state epidemiologist Anders Tegnell pointed that it seems meaningless to appeal to mathematical models at all, since they do not reflect even approximately the real situation.

Appendix 1: Growth of the herd immunity in Sweden

Our model can be interesting only for Sweden, where the “experiment” for approaching the herd immunity was performed in its clean form, without overshadowing by quarantine. The main concern of Swedish authorities, including Anders Tegnell, is that the dynamics of getting antibodies for covid-19 in Swedish population is essentially slower than it was expected. Since the end of April 2020, Anders Tegnell, the chief state epidemiologist of Sweden, started to announce (see, e.g., [32]) that very soon we, Swedish citizens, will become immune against the virus. This announcement was repeated a few times during May-June 2020, but it still not clear whether it happened or not (see, e.g., [34]-[36]). Moreover, Stockholm university made recently (June 24 []) the update that, in fact, already the level 60% can be considered as the herd immunity state. So, it seems that there is no hope to approach 70% level of immunity.

On June 24, 2020, it was announced [35] that: “A total of 7.3 percent of the blood samples collected from people in Stockholm were positive, 4.2 percent in Skane and 3.7 percent in Västra Götaland. The numbers reflect the situation in early April, as it takes time for the antibodies to develop. The authority’s modeling has previously shown that 26 percent of the population in Stockholm would have been infected on May 1. Tom Britton, professor of mathematical statistics at Stockholm University, comments on these and says that the figures are “clearly lower”, than he thought. “Firstly, I am thinking about the modeling, if there is any mistake that I and the Public Health Authority have made. Then you also wonder if some have been infected but do not have detectable antibodies, for example people who are mildly infected.”

The level of the herd immunity depends on the age group. On June 2, 2020, it was pointed out that for the youngsters up to 19 years old around 8% Andelen med antikroppar mot covid-19 bland yngre upp till 19 år var närmare åtta procent, för åldersgruppen 20-64 låg motsvarande siffra på 6-7 procent. För personer över 65 år låg siffran på cirka tre procent. Det kan finnas flera skäl till att det är lägre nivåer för vuxna och äldre, enligt Karin Tegmark Wisell.

References

- [1] R. M. Anderson, R. M. May, *Infectious Diseases of Humans: Dynamics and Control* (Oxford Univ. Press, 1991).

- 376 [2] H. Andersson, T. Britton, Stochastic Epidemic Models and Their
377 Statistical Analysis (Springer, 2000).
- 378 [3] O. Diekmann, H. Heesterbeek, T. Britton, Mathematical Tools
379 for Understanding Infectious Disease Dynamics (Princeton Univ.
380 Press, 2013).
- 381 [4] D. Smith and L. Moore, The SIR Model for
382 Spread of Disease - The Differential Equation Model.
383 [https://www.maa.org/press/periodicals/loci/joma/the-sir-](https://www.maa.org/press/periodicals/loci/joma/the-sir-model-for-spread-of-disease-the-differential-equation-model)
384 [model-for-spread-of-disease-the-differential-equation-model](https://www.maa.org/press/periodicals/loci/joma/the-sir-model-for-spread-of-disease-the-differential-equation-model).
- 385 [5] S. Flaxman, S. Mishra, A. Gandy, H. J. T. Unwin, H. Coup-
386 land, T. A. Mellan, H. Zhu, T. Berah, J. W. Eaton, P. N. P.
387 Guzman, N. Schmit, L. Cilloni, K. E. C. Ainslie, M. Baguelin,
388 I. Blake, A. Boonyasiri, O. Boyd, L. Cattarino, C. Ciavarella,
389 L. Cooper, Z. Cucunubá, G. Cuomo-Dannenburg, A. Dighe, B.
390 Djaafara, I. Dorigatti, S. van Elsland, R. FitzJohn, H. Fu, K.
391 Gaythorpe, L. Geidelberg, N. Grassly, W. Green, T. Hallett, A.
392 Hamlet, W. Hinsley, B. Jeffrey, D. Jorgensen, E. Knock, D. Lay-
393 don, G. Nedjati-Gilani, P. Nouvellet, K. Parag, I. Siveroni, H.
394 Thompson, R. Verity, E. Volz, C. Walters, H. Wang, Y. Wang,
395 O. Watson, P. Winskill, X. Xi, C. Whittaker, P. G. T. Walker, A.
396 Ghani, C. A. Donnelly, S. Riley, L. C. Okell, M. A. C. Vollmer,
397 N. M. Ferguson, S. Bhatt, Report 13: Estimating the number of
398 infections and the impact of non-pharmaceutical interventions on
399 COVID-19 in 11 European countries (Imperial College London,
400 2020); .doi:10.25561/77731
- 401 [6] N. M. Ferguson, D. Laydon, G. Nedjati-Gilani, N. Imai, K.
402 Ainslie, M. Baguelin, S. Bhatia, A. Boonyasiri, Z. Cucunubá, G.
403 Cuomo-Dannenburg, A. Dighe, I. Dorigatti, H. Fu, K. Gaythorpe,
404 W. Green, A. Hamlet, W. Hinsley, L. C. Okell, S. van Elsland,
405 H. Thompson, R. Verity, E. Volz, H. Wang, Y. Wang, P. G. T.
406 Walker, C. Walters, P. Winskill, C. Whittaker, C. A. Donnelly, S.
407 Riley, A. C. Ghani, Report 9: Impact of non-pharmaceutical in-
408 terventions (NPIs) to reduce COVID-19 mortality and healthcare
409 demand (Imperial College London, 2020); 10.25561/77482.doi
- 410 [7] T. Britton, Basic estimation-prediction techniques for Covid-
411 19, and a prediction for Stockholm. Preprint, April 2020 DOI:
412 10.1101/2020.04.15.20066050
- 413 [8] T. Britton, Basic estimation-prediction techniques
414 for Covid-19, and a prediction for Stockholm.
415 <https://www.medrxiv.org/content/10.1101/2020.04.15.20066050v2>
- 416 [9] T. Britton, P. Trapman, F.G. Ball, The disease-
417 induced herd immunity level for Covid-19 is substan-

- 418 tially lower than the classical herd immunity level.
419 <https://www.medrxiv.org/content/10.1101/2020.05.06.20093336v2>
- 420 [10] W. Bock, B. Adamik, M. Bawiec, V. Bezborodov, M. Bodych,
421 J. P. Burgard, T. Goetz, T. Krueger, A. Migalska, B. Pab-
422 jan, T. Ozanski, E. Rafajlowicz, W. Rafajlowicz, E. Skubalska-
423 Rafajlowicz, S. Ryfczynska, E. Szczurek, P. Szymanski, Mit-
424 igation and herd immunity strategy for COVID-19 is likely
425 to fail. medRxiv 2020.03.25.20043109 [Preprint]. 5 May 2020;
426 .doi:10.1101/2020.03.25.20043109 Abstract/FREE Full Text-
427 Google Scholar
- 428 [11] H. Salje, C. T. Kiem, N. Lefrancq, N. Courtejoie, P. Bosetti,
429 J. Paireau, A. Andronico, N. Hozé, J. Richet, C.-L. Dubost,
430 Y. Le Strat, J. Lessler, D. Levy-Bruhl, A. Fontanet, L. Opa-
431 towski, P.-Y. Boelle, S. Cauchemez, Estimating the burden of
432 SARS-CoV-2 in France. *Science* 10.1126/science.abc3517 (2020).
433 doi:10.1126/science.abc3517
- 434 [12] Public Health Institute of Sweden, Estimates of the
435 peak-day and the number of infected individuals dur-
436 ing the covid-19 outbreak in the Stockholm region,
437 Sweden February–April 2020 [in Swedish] (2020);
438 [www.folkhalsomyndigheten.se/contentassets/2da059f90b90458d8454a04955d1697f/skattning-
439 peakdag-antal-infekterade-covid-19-utbrottet-stockholms-lan-
440 februari-april-2020.pdf](http://www.folkhalsomyndigheten.se/contentassets/2da059f90b90458d8454a04955d1697f/skattning-peakdag-antal-infekterade-covid-19-utbrottet-stockholms-lan-februari-april-2020.pdf).
- 441 [13] Public Health Institute of Sweden: Antikroppstester avviker
442 fran prognoserna – matematikern “ser tva förklaringar”.
443 <https://www.svt.se/nyheter/inrikes/antikroppstester>
- 444 [14] Public Health Institute of Sweden: Yngre personer har
445 högre andel antikroppar. Sverige Radion, June 2, 2020;
446 <https://sverigesradio.se/sida/artikel.aspx?programid=83artikel=7487102>
- 447 [15] T. Britton, F. Ball, P. Trapman, A mathematical model reveals
448 the influence of population heterogeneity on herd immunity to
449 SARS-CoV-2. *Science* 23 Jun 2020: eabc6810
- 450 [16] J. Wallinga, P. Teunis, M. Kretzschmar, Using data on so-
451 cial contacts to estimate age-specific transmission parameters
452 for respiratory-spread infectious agents. *Am. J. Epidemiol.* 164,
453 936–944 (2006).
- 454 [17] R. Pastor-Satorras, A. Vespignani, Epidemic spreading in scale-
455 free networks. *Phys. Rev. Lett.* 86, 3200–3203 (2001).
- 456 [18] M. J. Ferrari, S. Bansal, L. A. Meyers, O. N. Bjørnstad, Network
457 frailty and the geometry of herd immunity. *Proc. Biol. Sci.* 273,
458 2743–2748 (2006).

- 459 [19] A. T. Ogielski, Dynamics on Ultrametric Spaces. PHYSICAL RE-
460 VIEW LETTERS, 55, N 15, 1634-1637.
- 461 [20] M. Mezard, G. Parisi, N. Sourlas, G. Toulouse, and M. Virasoro,
462 Phys. Rev. Lett. 52, 1156 (1984).
- 463 [21] R. G. Palmer, Adv. Phys. 31, 66a (1982); G. Parisi, Phys. Rev.
464 Lett. 50, 1946 (1983).
- 465 [22] A. Ansari, J. Berendzen, S. F. Bowne, H. Frauenfelder, I. E. T.
466 Iben, T. B. Sauke, E. Shyamsunder, and R. D. Young, Proc. Natl.
467 Acad. Sci. USA 82, 5000 (1985).
- 468 [23] R. G. Palmer, D. L. Stein, E. Abrahams, and P. W. Anderson,
469 Phys. Rev. Lett. 53, 958 (1984).
- 470 [24] B. A. Huberman and M. Kerszberg, J. Phys. A 18, L331 (1985).
- 471 [25] Parisi, Phys. Rev. Lett. 43, 1754 (1979).
- 472 [26] Khrennikov A., Human subconscious as a p-adic dynamical sys-
473 tem. Journal of Theoretical Biology, 193(2), 179-96 (1998).
- 474 [27] S. Albeverio, A. Khrennikov, P. E. Kloeden, Memory retrieval as
475 a p-adic dynamical system. Biosystems 49, N 2, 105-115 (1999).
- 476 [28] D. Dubischar, V. M. Gundlach, O. Steinkamp, A. Khrennikov,
477 A p-Adic Model for the Process of Thinking Disturbed by Phys-
478 iological and Information Noise. Journal of Theoretical Biology
479 197(4), 451-67 (1999).
- 480 [29] A. Khrennikov, Information Dynamics in Cognitive, Psychologi-
481 cal, Social, and Anomalous Phenomena. Kluwer, 2004
- 482 [30] A. Khrennikov, Probabilistic pathway representation of cognitive
483 information. Journal of Theoretical Biology 231, N 4, 597-613
484 (2004).
- 485 [31] A. Khrennikov, Toward an adequate mathematical model of
486 mental space: Conscious/unconscious dynamics on m-adic trees.
487 Biosystems 90, N 3, 656-675 (2007).
- 488 [32] Tegnell: "Vi får oroande rapporter om utelivet". SVT, Ny-
489 heter, 20-04-2020; [https://www.svt.se/nyheter/inrikes/tegnell-
490 vi-far-oroande-rapporter-om-utelivet](https://www.svt.se/nyheter/inrikes/tegnell-vi-far-oroande-rapporter-om-utelivet)
- 491 [33] Se hur matematikprofessorn räknar ut när Stockholm uppnar
492 flockimmunitet. SVT Nyheter, 20-04-2020.
- 493 [34] G. Holm, Hoppfulla teorin: Sa når vi flockim-
494 munitet redan i juni. Expressen, 10.05.2020.
495 [https://www.expressen.se/nyheter/coronaviruset/hoppfulla-
496 teorin-sa-nar-vi-flockimmunitet-redan-i-juni/](https://www.expressen.se/nyheter/coronaviruset/hoppfulla-teorin-sa-nar-vi-flockimmunitet-redan-i-juni/)

- 497 [35] A. Obminska, Det här vet vi om coronaviruset. Ny Teknik, 30-
498 06-2020; [https://www.nyteknik.se/samhalle/det-har-vet-vi-om-](https://www.nyteknik.se/samhalle/det-har-vet-vi-om-coronaviruset-6985117)
499 [coronaviruset-6985117](https://www.nyteknik.se/samhalle/det-har-vet-vi-om-coronaviruset-6985117)
- 500 [36] Public Health Institute of Sweden, 18-06-2020;
501 [https://www.folkhalsomyndigheten.se/nyheter-och-](https://www.folkhalsomyndigheten.se/nyheter-och-press/nyhetsarkiv/2020/juni/forsta-resultaten-om-antikroppar-efter-genomgangen-covid-19-hos-blodgivare/)
502 [press/nyhetsarkiv/2020/juni/forsta-resultaten-om-antikroppar-](https://www.folkhalsomyndigheten.se/nyheter-och-press/nyhetsarkiv/2020/juni/forsta-resultaten-om-antikroppar-efter-genomgangen-covid-19-hos-blodgivare/)
503 [efter-genomgangen-covid-19-hos-blodgivare/](https://www.folkhalsomyndigheten.se/nyheter-och-press/nyhetsarkiv/2020/juni/forsta-resultaten-om-antikroppar-efter-genomgangen-covid-19-hos-blodgivare/)