Forcing Seasonality of influenza-like epidemics with daily Solar resonance

Fabrizio Nicastro[1*], Giorgia Sironi[2], Elio Antonello[2], Andrea Bianco[2], Mara Biasin[3], John R. Brucato[4], Ilaria Ermolli[1], Giovanni Pareschi[2], Marta Salvati[5], Paolo Tozzi[4], Daria Trabattoni[3], Mario Clerici[6]

[1] *Italian National Institute for Astrophysics (INAF) – Rome Astronomical Observatory, Rome, Italy*
[2] *Italian National Institute for Astrophysics (INAF) – Brera Astronomical Observatory, Milano/Merate, Italy*
[3] *Department of Biomedical and Clinical Sciences L. Sacco, University of Milano, Milano, Italy*
[4] *Italian National Institute for Astrophysics (INAF) – Arcetri Astrophysical Observatory, Firenze, Italy*
[5] *Regional Agency for Environmental Protection of Lombardia (ARPA Lombardia), Milano, Italy*
[6] *Department of Pathophysiology and Transplantation, University of Milano and Don C. Gnocchi Foundation, IRCCS, Milano, Italy*

Summary

Seasonality of acute viral respiratory diseases is a well-known and yet not fully understood phenomenon. Several models have been proposed to explain the regularity of yearly recurring outbreaks and the phase-differences observed at different latitudes on Earth. Such models take into account known internal causes, primarily the periodic emergence of new virus variants that evade the host immune response. Yet, this alone, is generally unable to explain the regularity of recurrences and the observed phase-differences.

Here we show that seasonality of viral respiratory diseases, as well as its distribution with latitude on Earth, can be fully explained by the virucidal properties of UV-B and A Solar photons through a daily, minute-scale, resonant forcing mechanism. Such an induced periodicity can last, virtually unperturbed, from tens to hundreds of cycles, and even in presence of internal dynamics (host's loss of immunity) much slower than seasonal will, on a long period, generate seasonal oscillations.

Introduction

Several models have been proposed to explain the regularity of yearly recurring outbreaks of acute respiratory diseases, and their phase-differences at different latitudes on Earth (e.g. Logfren et al., 2007 and references therein). Internal biological dynamics that allows influenza viruses to evade host's immunity and become more virulent (Cox and Subbarao, 1999), manifest mainly through "antigenic drift", the property of such viruses to mutate rapidly (Agranovski et al., 2004; 2005). For A and B influenza viruses, such mutations develop at rates of 6.7×10^{-3} and 3.2×10^{-3} nucleotide substitution per site per year, respectively (Logfren et al., 2007; McCullers et al., 1999). These imply, on average, 6 and 3 mutations per RNA segment of each virus, every 6 months, respectively, sufficient (especially for type-A influenza) to trigger recurrences of epidemics on antigenic-drift timescale. However, without an external forcing mechanism, such recurring events are typically of modest intensity, can only last for a few cycles, and most importantly are not able to reproduce the observed phase-shift between north and south hemisphere on Earth.

One or more concurring external mechanisms must be invoked to explain all evidence, and several have been proposed (e.g. Ebi et al., 2001; Grais et al., 2003; Grais et al., 2004; Liao et al., 2005; Viboud et al., 2004; Baker et al., 2020). These include, but are not limited to, weather-related temperature and humidity oscillations on Earth (e.g. Baker et al., 2020), the observed yearly pattern of air-travel (e.g. Grais et al., 2003, but see also Grais et al., 2004), indoor heating during winter (e.g. Liao et al., 2005), bulk aerosol transport of virus through global convective currents or related to weather oscillations like "El Niño" (e.g. Ebi et al., 2001; Viboud et al., 2004).

Most of these, however, modulate on timescales that are either typical of seasons on Earth or longer and thus singularly can only explain some aspects of seasonality of recurring outbreaks, or the onsets of particularly violent outbreaks. For example, the load of air-travel flow is typically larger during northern hemisphere summers, and this is in contrast with most of the outbreaks developing during falls and winters in these countries. On the other hand, vastly populated areas in the north hemisphere lie at relatively low latitudes in temperate regions where indoor heating is not massively widespread: yet, even in these areas, influenza epidemics typically develop during winters, and remain inactive in summers. Similarly, temperature, humidity, as well as air pollution, are not homogeneously distributed through the planet, and

micro-climate can vary substantially at the same latitude, depending on the particular geographical or industrial setting of even contiguous areas. The cross-influence of all these factors could certainly synergize to produce the observed geographical and time modulations, but this would require a hard to obtain perfect fine-tuning of the many parameters at play.

The one mechanism that has been acting constantly at (almost) the same pace every day of the year and every year for the past four billion years, is the irradiation of the Sun on Earth. At a given location on Earth, this is modulated daily by the Earth's spinning and yearly by the Earth's orbit around the sun (Figure 1). Solar ultraviolet photons with wavelength in the range 200-290 nm (UV-C radiation) photo-chemically interact with DNA and RNA and are endowed with germicidal properties that are also effective on viruses (Kowalski, 2010; Rauth, 1965; Kesavan and Sagripanti, 2012; Chang et al., 1985; McDevitt et al., 2012; Welch et al., 2018; Walker and Ko, 2007; Tseng and Li, 2005). Fortunately, UV-C photons are filtered out by the Ozone layer of the upper Atmosphere, at around 35 km (e.g. Walker and Ko, 2007). Softer UV photons from the Sun with wavelengths in the range 290-320 nm (UV-B) and 320-400 nm (UV-A), however, do reach the Earth's surface. The effect of these photons on Single- and Double-Stranded RNA and DNA viruses (e.g. Lytle and Sagripanti, 2005; Lubin and Jense, 1995) and the possible role they play on the seasonality of epidemics e.g Martinez, 2018), are nevertheless little studied. In two companion papers (Nicastro et al., 2020, submitted: hereinafter N20; Bianco et al., 2020, submitted), we present a number of concurring circumstantial pieces of evidence suggesting that the evolution and strength of the recent Severe Acute Respiratory Syndrome (SARS-Cov-2) pandemics (Zhu et al., 2019; Cobey, 2020), might have been modulated by the intensity of UV-B and UV-A Solar radiation hitting different regions of Earth during the diffusion of the outbreak between January and May 2020 (N20), and provide a measurement of the lethal UV-C dose for Covid-19 (Bianco et al., 2020, submitted).

For the solar virus-inactivation mechanism to be effective in modulating the disease's transmission and settling the duration and recurrence of epidemics, both direct contact (e.g. through exposed surfaces) and airborne transmission have to play a role (as is the case for most respiratory diseases, e.g. Prather et al., 2020 and references therein) and the time needed for aerosol particles to gravitationally settle has to be comparable or longer than the typical UV-B/A virus's lethal time (N20). This is probably not true (in still air) for the largest (~100 μm in size) and thus heaviest droplets produced in coughs and sneezes, where the gravitational settling time is of the order of few seconds (Morawska and Cao, 2020), but can certainly be the case for significantly smaller, micron or sub-micron size, aerosol particles (like those that commonly contain the influenza virus, e.g. Tellier et al., 2019) that can survive in still air for longer than 12.4 hours (Tellier et al., 2019).

Here we present a simple mathematical model of epidemics that includes the effect of Solar UV photons ("solar-pump", hereinafter) as an infra-day resonance mechanism that acts systematically every day with strength and on timescales (from minutes to hours) dictated only by astronomical parameters, and explore its parameter space. We demonstrate that such a physically-motivated model is able to naturally reproduce the seasonality of influenza-like epidemics as well its distribution with latitude on Earth, without forcing seasonal oscillations with ad-hoc mathematical prescriptions (e.g. Dushoff et al., 2004; Neher et al., 2020), and matches reasonably well the Italian data of the recent SARS-CoV-2 epidemics (see Methods in Supplemental Information: SI), predicting different short-term future scenarios, depending on the efficiency of the solar-pump.

Results

The Model

Our model considers isolated (i.e. no cross-mixing), zero-growth (i.e. death rate equal to birth rate), groups of individuals in whom an efficacious anti-viral immune responses has (i.e. Susceptible → Infected → Recovered → Susceptible: SIRS) or has not (Susceptible → Infected → Recovered: SIR) been elicited (loss of immunity: LOI, hereinafter), parameterized through a population LOI rate γ_{in} (zero for no LOI). Other parameters of the model are the rate of contacts β (proportional to the reproductive parameter R_t), the rates of recovery γ_{out}, the rate of deaths μ, lockdown/phase-2 halving/doubling times, when considered (see below), and the efficiency ε of the solar-pump (see below).

The solar-pump
In our equations (see Methods in SI) the solar-pump forcing term is introduced as an infra-day (minutes/hours) attenuation factor for the reproductive number R_t and modeled by following the sun radiation curves as a function of Earth's latitude, day of the year, and time of the day. Solar curves (Fig. 1) are functions of the solar height angle α (complement to the Zenith angle: i.e. $\alpha=90$ degrees at local noon) that, in turn, depends on Earth's latitude φ, sun's declination δ, day of the year d and the time of the day h. Solar curves modulate daily with $\sin(\alpha)$ and have maximum amplitudes and widths that depend on φ, d and h, as shown in Fig. 1, where the R_t attenuation factor $(1 - \varepsilon \sin(\alpha))$ is plotted (for $\alpha>0$, i.e. during daytime, and reference solar-pump efficiency $\varepsilon=1$) versus h for the 15[th] day of the first 6 months of the year. In each panel of Fig. 1, different curves correspond to different Earth's latitudes (in steps of 5 degrees), from 25 to 90 degrees north (black), -25 to -90 degree south (blue) and -25 to +25 degrees (orange).

The efficiency of the solar-pump is here parameterized through the parameter ε (in the interval [0,1]) but depends, effectively, on internal virus properties, namely the lethal time τ_0 needed to deliver the UV lethal dose D_0 (defined as the UV-C fluence needed to inactivate 63% of the virus) for the given virus (N20). At local noon (the minima of the daily solar curves of Fig. 1) typical UV-B/A solar intensities on Earth range from 0.9–18 J m^{-2} min^{-1} in the north and south hemispheres between winters and summers, respectively, and are roughly constant at the equator throughout the year with a mean value of ~ 18 J m^{-2} min^{-1}. Folding these intensities through the action spectrum (i.e. the ratio of the virucidal dose of UV radiation at a given wavelength to the lethal dose at 254 nm) of Lytle and Sagripanti (2020), yields to lethal UV-B/A intensities of ~0.05–1 J m^{-2} min^{-1}. For Covid-19 we measured a three orders of magnitude inactivation dose of 37 J m^{-2} (Bianco et al., 2020, submitted), which translates into $D_0=4.8$ J m^{-2}. This value is used to produce the lethal-time dotted curves of Figures 3, 5, 7, 9, 11 and 13, which oscillate between 5-100 minutes at local noon depending on latitude and time of the year, and corresponds to our reference solar-pump efficiency $\varepsilon=1$. Efficiencies lower than the reference value 1 correspond to inversely longer lethal-times. As an example, Figure 2 shows two solar-pump attenuated R_t curves over a solar-year period for a 40 degree north location on Earth and for an epidemic with $R_0=3$ (see figure's caption for details).

Lockdown and Phase-2
Lockdown and phase-2 terms are modeled with exponential functions (see Methods in SI) that act as attenuation (lockdown) and amplification (phase-2) factors of the reproductive number R_t, with, respectively, halving and doubling times (in units of 1 day) that are free parameters of the model. The lockdown attenuation ends when phase-2 starts, and phase-2 amplification ends when R_t reaches the original value R_0 characteristic of the disease.

Our two Study-Cases
We focus on two particular study-cases: (A) typical influenza epidemics, with initial reproductive number $R_0=1.5$, solar-pump efficiency $\varepsilon =1$ and recovery and death rate $\gamma_{out}=0.2$ and $\mu=0.001$, respectively, and (B) a more aggressive and higher-mortality epidemics, with parameters similar to those recorded for the recent SARS-Cov-2 pandemics ($R_0=3$, $\gamma_{out}=0.1$ and $\mu=0.01$) and solar efficiencies $\varepsilon=1$ and 0.3. For each case, we consider three different LOI rates: $\gamma_{in}=0$ (i.e. no LOI), $\gamma_{in}=0.0055$ (180-day period: shorter than seasonality) and $\gamma_{in}=0.00055$ (5-year period: much longer than seasonality).
Finally, for each case, we consider three groups of individuals with equal populations (sixty million) and living at three different Earth's latitude: 40 degrees north (black curves in all figures), 40 degree south (blue curves in all figures) and the equator (orange curves in all figures). Groups are taken as isolated and have no cross-relation between them.

Case A: Influenza Epidemics
Figure 3 shows the curves of growth of the infected individuals in the three latitude groups over the first 20 years of influenza epidemics with (top panel) and without (bottom panel) solar-pump (see caption for details). When the solar-pump is not active ($\varepsilon=0$; bottom panel) the growth (and decline) of epidemics proceeds identically at all latitudes (black, blue and orange curves are superimposed to each others). The period of oscillations, in this case, resonates solely with the inverse of the LOI rate (the time it takes for previously infected individuals to become again susceptible), the strength of epidemics declines from cycle to cycle, and the disease dies out completely in less than 10 years. When the solar-pump is active ($\varepsilon=1$; top panel), instead, the reproductive number undergoes small infra-day variations (whose exact magnitude

depends on the particular time of the year and Earth's latitude, other than the efficiency of the solar-pump mechanism: e.g. onsets of Fig. 2) that do not significantly change its daily average value (Fig. 2), but progressively drive the modulation of epidemic oscillations to match the seasonal solar period (Fig. 3, top panel). The north (black), south (blue) and tropical (orange) groups differentiate immediately, and after a few cycles (lasting a total of about 5 years) are perfectly regulated on biannual periodicity, with phase shifted in time by 6 months in the north versus south hemisphere and the tropical group undergoing fast and continuous moderate bursts of epidemics roughly every six months. The biannual oscillations at latitudes ±40 degrees, last for a few tens of years, and then synchronize with the yearly solar-irradiation seasonal period (see figure's caption for details). The cycles can run unperturbed for centuries, as shown in the top panel of the Susceptible versus Infected (SI) phase-diagram of Fig. 4, where the first 100 years of endemic diseases are drawn (see Figure's caption for details)

Fig. 5–8 show analogous figures for the cases with γ_{in}=0 (no LOI: Fig. 5 and 6) and γ_{in}=0.0055 (5-year period of recurrence: Fig. 7 and 8). Interestingly, when the solar-pump is active and LOI periods are long compared to the seasonal period (Fig. 5 and 6 top panels), oscillations are more erratic and chaotic during the long settling period: destructive resonance between the LOI rate and the Solar cycles acts differently in the three latitude groups, and if no group mixing is introduced the intense bursts of epidemics are initially separated by the inverse of LOI rate. This separation progressively squeezes at all latitudes as the continuous solar-pumping imposes its rhythm, and becomes strictly seasonal at all latitudes in about 40-50 years. The squeezing of the time intervals between outbreaks, is visible also when the solar-pump is not active (Fig. 5, bottom panel). This is because, as new outbreaks develop, the reservoir of susceptible individuals made available by previous LOI cycles and the number of people still infected at any time are sufficient to trigger new bursts at earlier times. However, in this case, the strength of each outbreak decreases smoothly and systematically with years, and in about 70 years the reservoir of susceptible individuals at each time completely empties and the epidemics die out (Fig, 5, bottom panel). Moreover, again, the three latitude groups behave identically, which is not observed.
Finally, when γ_{in}=0 (no antigenic-drift), the solar-pump cannot, alone, sustain stationary epidemic cycles, but strongly modulates the course of epidemics during the first three years in the north and south hemisphere and over a longer period of about 8 years in the tropical area (Fig. 7, top panel). By contrast, when the solar-pump is not active (Fig. 7 and 8, bottom panels) the epidemics develop identically at all latitudes and last for a single 1-year cycle (Fig. 8, bottom panel).

Case B: SARS-CoV-2 Pandemics

Our second study-case aims to model the recent SARS-CoV-2 pandemics outbreak under the different assumptions tested in this work, and allows us to gain insights on possible future scenarios. As for study-case A, we use the three groups at latitudes 40^0-north, 40^0-south and the equator, and run simulations both with ($\varepsilon \neq 0$) and without (ε=0) the solar forcing term and for LOI rates γ_{in}=0.0055, 0.00055 and 0. However, for study-case B we also compare our north-hemisphere simulation with the SARS-CoV-2 data for Italy (central latitude +41.5 degrees) observed from 24 February 2020 through 4 July 2020 (see Methods in SI). To do so, we introduce, for all groups, two additional external forcing terms that modulate the contact rate β (chosen to be 0.3, to reproduce the observed – in virtually every monitored country – initial exponential growth rate): a 'lockdown' term and a 'phase-2' term (see Fig. 9's caption for details).

With the particular initial setting described in the caption of Fig. 9, the simulation with active solar-pump (ε=1) and fast antigenic-drift (γ_{in}=0.0055) reproduces extremely well the curve of growth of the observed daily number of infected in Italy (Fig. 9, top panel, black infected curve and data), with an under-sampling factor of 10 (i.e. observed infected are one tenth of actual infected). Preliminary results from serological tests on a statistically selected sample, confirm this average under-sampling factor in Italy (see e.g. https://www.med4.care/italia-test-sierologici-covid-19-risultati/).
The bottom panel of Fig. 9 shows the same simulation when the solar-pump is not active (ε=0; curves of growth from the three latitude groups are superimposed to each other). However, here the under-sampling factor needed to fit the initial exponential phase of the growth of contagions in Italy is an unrealistic factor of 300. Moreover, the declining phase of contagions, with no solar-pump is too fast to reproduce the Italian data. The much smaller absolute number of daily infected in the simulation shown in the top-panel of Fig. 9, compared with that of the bottom panel (under-sampling factor of 10, versus 300), and the slower decay of daily detected infections, are due solely to the daily effect of the solar-pump. The fast-decay in the inactive

solar-pump case is a direct consequence of the much larger 'consumption' of susceptible individuals during the growth of contagions (compare top and bottom panels of Fig. 10).

The sun has also the effect of reducing by almost one and two orders of magnitude, respectively, the number of infected individuals observed from February through May, in the South and tropical group, with respect to the North group, which is exactly what has generally been observed during the SARS-CoV-2 pandemic (N20; Fig. 9 top panel). The curve of growth of the South group, however, starts climbing again in May (which has generally been observed, e.g. N20) and, if no additional distancing measures are adopted, should have a second and higher peak in October, towards the end of the South winter (Fig. 9, top panel). In the north hemisphere, instead, this simulation (but see below for a for an alternative scenario) predicts that the curves of growth of the epidemics should reach a minimum in July, start climbing again (less aggressively than during the initial January outbreak) from August through October and, in absence of new distancing measures, reach a new intense peak in November, while tropical countries will see a second and much stronger outbreak during the second half of 2020. Subsequently, under these conditions, the epidemics will start adjusting to the solar cycle and less aggressive outbreaks (maximum daily number of new infected individuals between ten and thirty thousand) will repeat cyclically for tens of years (Fig. 10).

Should recovered people preserve their immunity against Covid-19 (γ_{in}=0 case), UV photons from the Sun may still be able to indirectly amplify contagions during autumns/winters (as a consequence of the attenuation during previous summers) and produce one additional wave of the SARS-CoV-2 epidemics at all latitudes (Fig. 11 and 12). However, in less than two years (and only 1 additional cycle), in absence of antigenic-drift or cross-mixing between groups at different latitudes, the epidemics will die out.

Finally, as for study-case A (influenza), at the much slower LOI rate γ_{in}=0.00055 (LOI period of 5 years), the seasonal cycle takes much longer to stabilize, and oscillations proceed at intermittent intensities both in the north and south hemisphere (with different phases) for tens of years, and then finally synchronize to the Solar cycles on timescales longer than 50 years (Fig. 13 and 14).

Discussion

Our model is able to reproduce in a particularly precise way the seasonality of influenza-like epidemics and its phase-latitude dependence, and can explain the qualitative differences generally observed when the strength of the SARS-CoV-2 pandemics is compared in countries located in the north, tropical and south hemisphere. However, accurate predictions of ongoing epidemics depend critically on the initial and boundary conditions of the simulations. For example, the particular set of model parameters we used to match the Italian data of the ongoing SARS-CoV-2 epidemics (see Fig. 9's caption for details), is obviously not unique. In particular, these depend (given initial dates for lockdown and phase-2) on the starting time of the epidemics and the actual efficiency of the solar effect, which, in turn, depends on the exact knowledge of the virus' UV-C lethal dose D_0, the way this is delivered and the time the virus (in aerosol or surfaces) remains exposed to the dose. Accurate modeling of this phenomenon is beyond the scope of this work. However, it is useful to show the comparison with a drastically different solar-pump efficiency (ε=0.3, i.e. ~3 times lower than in the case of Figures 9 and 10: compare, e.g. green and blue curves of Fig. 2) to highlight the potential predictive power of the model for e.g. the SARS-CoV-2 epidemics.

Fig. 15 shows the results of a simulation for the three groups at ±40 degree latitudes and the Equator, a solar-pump efficiency ε=0.3 and a set of model parameters that, as for the case of Fig. 9, matches reasonably well the Italian SARS-Cov-2 curve of growth of contagions (see figure's caption for details) from the start of epidemics through 4 July 2020 and with an under-sampling factor of 25 (probably too high, based on preliminary results from serological tests in Italy). Even with such reduced efficiency the solar-pump is able to produce significant differences in the curves of growth of the epidemics at ±40 degree latitudes and at the Equator (black, blue and orange solid curves in Fig. 15), which, however, start to synchronize with seasons only after a few cycles. Moreover, short-term predictions, at all latitudes, differ significantly from the case with higher solar-pump efficiency (Fig. 9). Italy (and in general, countries located at ~40 degrees north), where distancing measures have now greatly being relaxed, should start seeing a strong increase in the number of new daily infected in September, and this should continue, in absence of distancing measures, through the end of January 2021, to then drop quickly in spring 2021 and restart again a few months later, in summer 2021. In the case of Fig. 9, instead, the epidemic in Italy should

have a second sharp peak in November 2020 and then die out quickly just before the beginning of winter 2020. Such different predictions depend mainly (in absence of further distancing measures), on the relative significance of the solar mechanism, and thus offer the opportunity not only to clearly distinguish between the two scenarios, but also to evaluate the magnitude of the effect.

In general, comparing our model predictions with SARS-CoV-2 data at different latitudes over the next few months, should tell us qualitatively about the importance of the solar-pump effect in modulating the spreading of this disease.

Limitations of the Study

Our model for the diffusion of contagions (see Methods in SI) considers population groups as ideal isolated systems (i.e. no cross-mixing of populations in different groups at different latitudes on Earth). This is condition certainly not verified in today's global world, in which daily exchange of people and goods are massive and, generally, not strictly controlled (unless measures to limit the spread of contagions are taken by governments, as in the case of the recent SARS-CoV-2 pandemics). This type of complexity could easily be introduced in our basic system of differential equations (see Methods in SI) by considering cross-mixing through a matrix of either observed (i.e. based on air- ground- and sea-traffic data) or randomized exchange rates of susceptible and infected individuals among groups. However, the introduction of such an additional degree of complexity in the model (which will be the subject of a forthcoming paper), while could contribute to wash-out and partially homogenize differences between specific groups at different latitudes, could hardly build-up or break the strict seasonality that is typically observed in the recurrence of respiratory diseases and whose explanation is the main scope of this work.

The basic ingredient of our model that generates seasonality and latitude-dependent phase differences among groups, is a physically-motivated resonant forcing mechanism (the solar-pump) that acts as a daily modulator of the reproductive number R_t (see Methods in SI). The mechanism is based on the virus-inactivation power of the solar UV photons that reach the Earth's surface, and is dictated by the shapes of the daily solar irradiation curves on Earth, which, for a given latitude and time of the year, depend only on the Earth's spinning around its rotation axis and whose self-similarity throughout a solar year is determined only by Earth's orbit around the Sun. However, the exact efficiency of this mechanism depends on the quantification of several intrinsic parameters of the specific epidemic, such as the lethal dose of UV photons needed to inactivate the specific virus when exposed to solar-like spectra, the average life-time of the virus in aerosol or surfaces, its transport in wind or low-atmosphere currents, etc. The quantification of these parameters for given viruses, is beyond the scope of this work, whose aim is to show the potential of solar radiation in modulating the short- and long-term evolution of epidemics outbreaks and demonstrate that the, unavoidable, infra-day (minute timescales) resonant mechanism naturally offered by the solar-pump, can efficiently synchronize with internal LOI rates to explain the course of epidemics on much longer timescales.

Finally, and related to the above caveat, the efficiency of the solar-pump mechanism, which can only act on outdoor droplets generated through speaking, coughing and/or sneezing, aerosol and surfaces-deposited particles, depends also on the social habits of different population groups, located even at similar latitudes on Earths, often dictated by climate, and on specific location's weather (e.g. diffuse cloud coverage). For example, outdoor living, even in winters, is much more common in Mediterranean countries than, e.g. in North-American states at similar Earth's latitudes. This is often anti-correlated with the massive use of indoor air-conditioning (offices, shopping malls, private homes, private and public transportation, etc.), which is known to be an efficient mean of diffusion of contagions and could, e.g., contribute to explaining why the SARS-CoV-2 epidemics is currently spreading aggressively during summer in northern locations like the state of Texas in the US. Similarly, the diffuse cloud coverage (especially during the times of the days when the solar irradiation is maximum) in tropical countries, during specific times of the year (e.g. the monsoon season in India, or the rain seasons in the northern of Brazil, or hurricane seasons in central and tropical America) will reduce the efficiency of the solar-pump and could act as an amplifier of the growth of contagions in specific periods of the year. Again, the consideration of such additional degrees of complexity, in the model is certainly possible but would require a somewhat arbitrary (if not based on solid social and weather data) introduction of group-dependent weights in the parameterization of the efficiency of the solar-pump mechanism, which is beyond the scope of this paper.

Acknowledgments

The work presented in this paper has been carried out in the context of the activities promoted by the Italian Government and in particular, by the Ministries of Health and of University and Research, against the COVID19 pandemic. Authors are grateful to INAF's President, Prof. N. D'Amico, for the support and for a critical reading of the manuscript. Authors are also grateful to Dr. M. Elvis for critically reading the paper and providing useful comments and suggestions. A research grant from Falk Renewables partially supported this work.

Author Contributions

F.N. designed the study (together with co-authors P.T., G.P. and M.C.), wrote the program that solves the model's equation and provides the curves of growth of the epidemics, collected and analysed Italian SARS-CoV-2 data and wrote the paper. G.S. and M.S. collected, reduced and analysed TEMIS data. G.P., M.C., M.T., M.B. and A.B., provided the measurement of the UV lethal dose for Covid-19 and, together with co-authors G.S., J.R.B., E.A. and I.E., derived the action-spectrum-folded UV-B/A virus lethal times plotted in Fig. 3, 5, 7, 9, 11, 13 and 15. All authors contributed equally to the discussion of the results and commented on the manuscript.

Declaration of Interests

Authors have no competing interest to disclose.

References

- Agranovski, I. E., A. S. Safatov, A. I. Borodulin, O. V. Pyankov, V. A. Petrish-chenko, A. N. Sergeev, A. P. Agafonov, G. M. Ignatiev, A. A. Sergeev, and V.Agranovski, Inactivation of viruses in bubbling processes utilized for per-sonal bioaerosol monitoring. Appl. Environ. Microbiol, 2004, **70**, 6963–6967.
- Agranovski, I. E., A. S. Safatov, A. I. Borodulin, O. V. Pyankov, V. A.Petrishchenko, A. N. Sergeev, A. A. Sergeev, V. Agranovski, and S. A.Grinshpun, New personal sampler for viable airborne viruses: feasi-bility study. J. Aerosol Sci, 2005; **36**, 609–617
- Baker R. E., Yang W., Vecchi G. A., Metcalf C. J. E., Grenfell, B. T., Susceptible Supply limits the role of climate in early SARS-CoV-2 pandemic. Science, 2020; **10**, 1126–1129.
- Bianco, A. et al, UV-C irradiation is highly effective in inactivating and inhibiting SARS-CoV-2 replication. 2020, submitted; medRxiv 2020.04.16.20068387.
- Chang JC, Ossoff SF, Lobe DC, et al. UV Inactivation of Pathogenic and Indicator Microorganisms. *Appl. Env. Microb*. 1985; **49**: 1361-1365.
- Cobey S. Modeling infectious disease dynamics - The spread of the coronavirus SARS-CoV-2 has predictable features. *Science 2020;* **368**, 713.
- Cox, N. J., and K. Subbarao, Influenza. Lancet, 1999; **354**,1277–1282
- Dushoff J., Plotkin J.B., Levin S.A., Earn, D.J.D., Dynamical resonance can account for seasonality ofinfluenza epidemic. *PNAS*, 2004, **101**, 48, 16915–16916.
- Ebi, K. L., K. A. Exuzides, E. Lau, M. Kelsh, and A. Barnston, Association of normal weather periods and El Nino events with hospitalization for viral pneumonia in females: California, 1983–1998. Am. J. Public Health, 2001; **91**,1200–1208.
- Grais, R. F., J. H. Ellis, and G. E. Glass, Assessing the impact of airlinetravel on the geographic spread of pandemic influenza. Eur. J. Epidemiol, 2003; **18**, 1065–1072.37.
- Grais, R. F., J. H. Ellis, A. Kress, and G. E. Glass, Modeling the spreadof annual influenza epidemics in the U.S.: the potential role of air travel, Health Care Manag. Sci., 2004; **7**, 127–134
- Kesavan J, Sagripanti JL. Disinfection Of Airborne Organisms By Ultraviolet-C Radiation And Sunlight. Research and Technology Directorate, Edgewood Chemical Biological Center, U.S. Army, Aberdeen Proving Ground, Maryland 21010-5424 2012 ECBC-TR-1011.
- Kowalski W. Ultraviolet germicidal irradiation handbook: UVGI for air and surface disinfection. 2010, Springer Science & Business Media.
- Liao, C. M., C. F. Chang, and H. M. Liang, A probabilistic transmis-sion dynamic model to assess indoor airborne infection risks. Risk Anal, 2005; **25**,1097–1107.

- Lofgren E., Fefferman N. H., Naumov Y.N., Gorski J., and Naumova E. N., Influenza Seasonality: Underlying Causes and Modeling Theories. Journal of Virology, 2007; **81**, 5429–5436
- Lubin D, Jensen EH. Effects of Clouds and Stratospheric Ozone Depletion on Ultraviolet Radiation Trends. *Nature* 1995; **377**, 710–713.
- Lytle DC, Sagripanti JL, Predicted Inactivation of Viruses of Relevance to Biodefense by Solar Radiation. *J Virol* 2005*;* **79**, 14244–14252.
- Martinez ME, The calendar of epidemics: Seasonal cycles of infectious diseases. *PLoS Pathog* 2018*;* **14**, e1007327.
- McCullers, J. A., G. C. Wang, S. He, and R. G. Webster, Reassortmentand insertion-deletion are strategies for the evolution of influenza B virusesin nature. J. Virol, 1999, **73**, 7343–7348.
- McDevitt JJ, Rudnick SN, Radonovich L. Aerosol susceptibility of influenza virus to UVC light. *Appl Environ Microbiol* 2012; **78**, 1666–1669.
- Morawska L., Cao J., Airborne transmission of SARS-CoV-2: The world should face the reality. *Environ. Int., 2020*, **139**, 105730.
- Neher R.A., Dyrdak R., Druelle V., Hodcroft E.B., Albert J., Potential impact of seasonal forcing on a SARS-CoV-2 pandemic. *Swiss Med Wkly*, 2020, **150**, w20224.
- Nicastro, F. et al., Modulation of COVID-19 Epidemiology by UV-B and -A Photons from the Sun. 2020, submitted; medRxiv 2020.06.03.20121392: N20
- Prather K. A., Wang C. C., Schooley R. T., Reducing transmission of SARS-CoV-2. *Science, 2020*, **368**, 1422–1424
- Rauth AM. The Physical State of Viral Nucleic Acid and the Sensitivity of Viruses to Ultraviolet. *Light Biophys J* 1965; **5**: 257–273.
- Tellier R. et al., Recognition of aerosol transmission of infectious agents: a commentary. *BMC, Infect. Dis., 2019*, **19**, 101.
- Tseng CC, Li CS. Inactivation of Virus-Containing Aerosols by Ultraviolet Germicidal Irradiation. *Aerosol Science and Technology* 2005; **39**, 1136-1142.
- Viboud, C., K. Pakdaman, P. Y. Boelle, M. L. Wilson, M. F. Myers, A. J.Valleron, and A. Flahault, Association of influenza epidemics with global climate variability. Eur, J. Epidemiol, 2004; **19**, 1055–1059
- Walker CM, Ko G. Effect of ultraviolet germicidal irradiation on viral aerosols. *Environ Sci Technol* 2007; **41**, 5460-5465.
- Welch, D., Buonanno, M., Grilj, V. et al., Far-UVC light: A new tool to control the spread of airborne-mediated microbial diseases. *Scientific Reports* 2018; **8**, 2752.
- Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with Pneumonia in China 2019. *New Engl J Med 2020*; **382**, 727-733.

Figure 1 The Solar Pump. Attenuation functions for the reproductive number R_t, due to the daily solar action. Each panel show solar attenuation curves for 36 Earth's latitudes between -90 and -25 degrees south (blue), -25 − +25 degrees (orange) and 25–90 degrees north (black) at the 15[th] day of each of the first 6 months of the year, as labeled. The plotted function is $(1 − \sin(\alpha))$, where α is the solar height angle (complement to the Zenith angle: i.e. α=90 degrees at local noon) that depends on Earth's latitude φ, sun's declination δ, day of the year d and time of the day h.

Figure 2 Efficiency of the solar-pump. One-day averaged reproductive number R_t as a function of time (in days) over one full solar year and at a 40 degree north location on Earth, attenuated only by the solar-pump mechanism with efficiencies of $\varepsilon=0.3$ (green curves) and $\varepsilon=1$ (blue curves), starting from an epidemic-intrinsic $R_0=3$ value. Onsets show the daily attenuation curves as a function of day-time in hours during the first and the 180^{th} day of the year (January 1^{st} and July 1^{st}, respectively).

Figure 3 Seasonality of Influenza. Curves of growths of the daily new infected for the first 20 years of annual Influenza outbreaks at three different latitudes on Earth, +40 degrees north (black), -40 degrees south (blue) and the Equator (orange), with (ε=1; top panel) and without (ε=0; bottom panel) active solar-pump. Model parameters are t_0=January 1st, t_{end}=t0+100 years, R_0=1.5, γ_{out}=0.2, γ_{in}=0.0055, μ=0.001 and no external lockdown or phase-2. When the solar-pump is active (top panel) constructive resonance builds-in regular sun-modulated oscillations already after the first 2 cycles, and the strength of the seasonal outbreaks adjust to constant values that depend on latitude. With these particular model parameters, north- and south-hemisphere oscillations have initially a 2-year period and are shifted in phase by 6 months (with outbreaks only during local winters), while at the Equator the disease is continuously spread at a low-intensity level throughout the year. It takes almost 50 years before oscillations at ±40 degrees latitudes become annual. This happens at the expenses of the maximum intensity of the biannual outbreaks, which starts declining smoothly as mild outbreaks develop during the initially off-state year. Without solar-pump (bottom panel) the cycles follow the LOI period and are the same at all latitudes: the epidemics quickly die out in a few years. Yearly oscillating dotted curves at the bottom of the top panel are UV-B/A lethal-exposure estimated for Covid-19 and are reported here only to mark winter peaks and summer minima.

Figure 4 From Epidemic to Endemic Influenza. Susceptible versus actual infected phase diagrams of the influenza outbreak run described in Fig. 3's caption, with ($\varepsilon=1$; top panel) and without ($\varepsilon=0$; bottom panel) active solar-pump. The first 100 years of the epidemics are plotted (every other 10 years, in the top panel, for easier visualization). When the solar-pump is not active (bottom panel), the curves are the same at all latitudes and the cycles of epidemics die out quickly (essentially in 3-4 cycles) because of lack of supply among the susceptible population. When the solar-pump is switched on (top panel) the resonant mechanism builds up seasonality quickly. During the first 50 years, and at latitudes of ±40 degrees (black and blue curves), the susceptible population at each time oscillates between 65–98% of the population, and the instant fraction of infected people over susceptible, at the peaks, reach maximum values of about 4.5%. At the equator oscillations are much milder (orange curve): after a first long-duration cycle, the susceptible population at each time settles at a fraction of about 98% of the total while infected people oscillates steadily between 0.06–0.2% of the susceptible population.

Figure 5 Slow LOI rate Influenza Epidemics. Same as Fig. 3, but for a slow LOI rate of γ_{in}=0.00055. As for the fast LOI rate case, when the solar-pump is active (ε=1; top panel) constructive resonance eventually (in about 30-40 years) builds-in regular sun-modulated oscillations, and the strength of the seasonal outbreaks adjust to quasi-constant values that depend on latitude. However, during the first 30-40 years, intensive outbreaks are sporadic at all latitudes and typically repeat for 2-3 consecutive years in the north and south hemispheres. Without solar-pump (ε=0; bottom panel) the cycles are the same at all latitudes and are fed by LOIs only till the available number of susceptible is sufficient. The strength of the outbreaks decreases steadily with time, and time-intervals between peaks squeezes, from the inverse of the LOI rate to ~0.2 times that.

Figure 6 Slowly becoming endemic. Same as Fig. 4, for the model of Fig. 5 and for the entire duration of the simulation (100 years). The LOI mechanism alone is able to support a few cycles of the epidemics ($\varepsilon=0$; bottom panel), but these decreases systematically in strength and eventually die out. Switching the solar-pump on ($\varepsilon=1$; top panel) allows the system to remain active for the entire duration of the simulation and to stabilize after 30-40 years to susceptibility fractions of ~85% and 97%, at all times, at ±40 degrees and at the Equator, respectively. Correspondingly, the fraction of actual infected over the susceptible population, oscillates steadily between $(0.001 - 0.2)$% and $(0.005 - 0.02)$%, at ±40 degrees and at the Equator, respectively.

Figure 7 Non-recurring Influenza. Same as Figure 3 for the first 8 years of an influenza-like epidemics with zero LOI rate. When internal dynamics is absent (no LOI) the epidemics cannot become endemic. However, when the solar-pump is active (ε=1; top panel) UV photons from the sun are able to modulate the outbreak and prolong it for a few cycles (top panel). A couple of bursts are seen at ±40 degrees, whereas a single long, mild and Sun-modulated epidemics develop at the Equator and last for about 6 years. Without solar-pump (ε=0; bottom panel) a single intense episode is seen at all latitudes.

Figure 8 No-LOI Influenza Epidemics. Same as Fig. 4, for the model of Fig. 7 and for the entire duration of the simulation (100 years). The epidemics are not sustained by LOI and die out quickly. However, the total number of people infected before the outbreaks are over, depends on whether the solar-pump is active ($\varepsilon=1$; top panel) or not ($\varepsilon=0$; bottom panel). With epidemics starting during north-hemisphere winters (as in our models), latitude +40 degrees reaches a maximum infected over susceptible fraction of 1%, while this fraction is four times larger at -40 degrees and about ten times smaller at the Equator.

Figure 9 Evolution of SARS-Cov-2 Pandemics. Simulation of the curves of growths of the daily new infected for the first 3 years of the SARS-CoV-2 pandemics at three different latitudes on Earth, +40 degrees north (black), -40 degrees south (blue) and the Equator (orange), with (ε=1; top panel) and without (ε=0; bottom panel) active solar-pump. Model parameters are t_0=January 5th 2020, t_{end}=t_0+100 years, R_0=3.0, γ_{out}=0.1, γ_{in}=0.0055, μ=0.01 and external lockdown and phase-2 starting on 11 March 2020 and 10 May 2020 (official lockdown and reopening dates in Italy), with halving and doubling times of 46 and 180 days, respectively. Black points are SARS-CoV-2 data for Italy from 24 February 2020 through 4 July 2020, smoothed over a 7-day moving average. For the active solar-pump case (top panel) all curves of growth of new daily infected have been scaled down by a factor of 10 to let the +40^0 curve (black) match the Italian data of the current outbreak: the model closely matches observations. In the non-active solar-pump case (bottom panel) model curves needs to be scaled-down by an unrealistically high factor of 300, and only the climbing phase of the outbreak in Italy can be reproduced. Yearly oscillating dotted curves at the bottom of the top panel are UV-B/A lethal-exposure estimated for Covid-19.

Figure 10 Will SARS-CoV-2 leave or stay? Susceptible versus actual infected phase diagrams of the SARS-CoV-2 pandemic outbreak run described in Fig. 9's caption, with ($\varepsilon=1$; top panel) and without ($\varepsilon=0$; bottom panel) active solar-pump. The first 10 years of a possible evolution of the epidemics are plotted. When the solar-pump is not active (bottom panel), the curves are the same at all latitudes and the cycles of epidemics die out in 4 cycles because of lack of supply among the susceptible population. Considering the effect of solar-pump (top panel) makes the epidemics evolve quickly to endemic at all latitude. After the first 3 cycles, and at latitudes of ±40 degrees (black and blue curves), the susceptible population at each time oscillates between 33–55% of the population, and the instant fraction of infected people over susceptible, at the peaks, reach maximum values of about 5%. At the equator the amplitude of oscillations is much smaller (orange curve), the susceptible population at each time settles at a fraction of about 50% of the total while infected people oscillates steadily between 0.2–3.5% of the susceptible population

Figure 11 Secondary wave of no-LOI SARS-CoV-2. Same as Fig. 9, but for a zero LOI rate ($\gamma_{in}=0$). With these parameters, the model is not able to adequately match the Italian data of the outbreak (black points), even when the solar-pump is active ($\varepsilon=1$; top panel). However, only when UV photons are considered (top panel), can the model reproduce the observed differences in geographical spread of the pandemics. Should recovered people preserve their immunity against Covid-19, UV photons from the Sun may still be able to amplify contagions during autumns/winters and produce one additional wave of the SARS-CoV-2 epidemics at all latitudes. This cannot happen in absence of UV photons ($\varepsilon=0$; bottom panel), even after lockdown measures are loosened.

Figure 12 Same as Fig. 10, for the model of Fig. 11 and for the entire duration of the simulation (100 years). Both in the active (ε=1; top panel) and non-active (ε=0; bottom panel) solar-pump cases, the epidemics are not sustained by LOI and die out quickly. However, when UV photons are considered, the epidemics last longer (a total of two cycles at all latitudes). With epidemics starting during north-hemisphere winters (as in our models), and in absence of additional lockdown measures, two cycles develop +40 degrees and at the Equator, while a single, longer and asymmetric cycle runs through the end of 2021 in the South hemisphere. The final fractions of susceptible population are 12%, 20% and 35% at +40, 0 and -40 degrees respectively, and maximum 30% instant fractions of infected over susceptible population are recorded in the north hemisphere and at the Equator, towards the end of 2020, while this fraction stays lower (< 11%) at all time during the epidemics in the south hemisphere.

Figure 13 The nightmare of a recurring long-lasting SARS-CoV-2. Same as Fig. 9, but for the slow LOI-rate scenario (γ_{in}=0.00055, or 5-year period) and for the first 50 years of the simulation. Both with (ε=1; top panel) and without (ε=0; bottom panel) solar-pump, cycles proceed driven by the slow-rate reappearing vulnerability of the immune system, with decreasing periods at all latitudes. When UV photons are considered (top panel) the initial epidemics' bursts are generally intense and erratic, especially at north and south altitudes, and become steadily recurring and stabilize to the seasonal cycle only after 25-50 years depending on latitude (and in absence of cross-mixing population), fortunately with moderate strength.

Figure 14 The Violent cycles of SARS-CoV-2. Same as Fig. 10, but for the model of Fig. 13 and for the entire duration of the simulation (100 years). Should LOI happen on a long timescale (5 years in this simulation), the presence of UV photons from the sun (top panel) would make the first 4-5 epidemics bursts in the north and south hemisphere, particularly intense. The fraction of actual people infected at any time over the susceptible population, would reach peak values of up to 30% during the first 25 years, and would then stabilize, oscillating within the range 0.02–10% (at latitudes ±40) and 0.2–1% (at the Equator), for the next 75 years.

Figure 15 Evolution of SARS-Cov-2 with a low-efficiency solar-pump. Same as Fig. 9, but for a solar-pump with efficiency 3 times lower ($\varepsilon=0.3$, corresponding to three times longer lethal times: dotted curves) and model parameters: t_0=January 15th 2020, $t_{end}=t_0$+100 years, R_0=3.0, γ_{out}=0.1, γ_{in}=0.0055, μ=0.01 and external lockdown and phase-2 starting on 11 March 2020 and 20 April 2020 (20 days earlier than the actual official Phase-2 date in Italy), with halving and doubling times of 25 and 370 days, respectively. All curves of growth of new daily infected have been scaled down by a factor of 25 to let the +40^0 curve (black) match the Italian data of the current outbreak: the model closely matches observation, however, short-term predictions differ significantly from the case plotted in Fig. 9.

Supplemental Information

Transparent Methods

Our Deterministic Model's Equations

We adopt a simple deterministic SIRS model, in which the Susceptible population (S) interacts not linearly with Infected (I) at a rate β, Recovers (R) or dies (D) with rates γ_{out} and μ, respectively, and can become again susceptible (loses immunity) with a rate γ_{in}:

$$\frac{dS}{dt} = -\beta\frac{S\,I}{N} + \gamma_{in}\,R$$
$$\frac{dI}{dt} = \beta\frac{S\,I}{N} - (\gamma_{out} + \mu)I$$
$$\frac{dR}{dt} = \gamma_{out}\,I - \gamma_{in}\,R$$
$$\frac{dD}{dt} = \mu\,I$$

We wrote a simple Fortran-90 routine that solves this system of differential equations (non-linear only in I and S) iteratively, for arbitrary values of the model parameters. We adopt a time-unit of 1 day and, at each iteration, perform infra-day integrations.

We further introduce three optional external forcing terms that act on the reproductive number R_t (effectively on the contact rate β=β(t)) by attenuating or amplifying it on timescales of 1 day (lockdown and phase-2 terms) and minutes (solar-pump) and may be switched on and off during runs:

$$\beta = \beta_0(1 - \varepsilon\sin(\alpha))\begin{cases} 1, & no\ lockdwon/phase2 \\ e^{-(t-t_{LD})/T_{LD}}, & t_{LD} \leq t \leq t_{Ph2} \\ Ae^{(t-t_{Ph2})/T_{Ph2}}, & t \geq t_{Ph2} \end{cases}$$

where, (t_{LD}, T_{LD}), and (t_{Ph2}, T_{ph2}) are initial and e-folding times of lockdown and Phase-2, respectively (when present), and $A = e^{-(t_{Ph2}-t_{LD})/T_{LD}}$.

Solar UV Data

In our study, to compute lethal times τ_0, we use Solar UV data made available by the Tropospheric Emission Monitoring Internet Service (TEMIS) archive (http://www.temis.nl). TEMIS is part of the Data User Programme of the European Space Agency (ESA), a web-based service that stores, since 2002, calibrated atmospheric data from Earth-observation satellite. TEMIS data products include measurements of ozone and other constituents (e.g. NO_2, CH_4, CO_2, etc.), cloud coverage, and estimates of the UV solar flux at the Earth's surface. The latter are obtained with state-of-the-art models exploiting the above satellite data.

SARS-CoV-2 Pandemics Data

We collect SARS-CoV-2 data pandemics for Italy from the on-line GitHub repository provided by the Coronavirus Resource Center of the John Hopkins University (CRC-JHUL https://github.com/CSSEGISandData/COVID-19). Such data are provided daily by the Italian civil-defense department. CRC-JHU global data are updated daily and cover, currently, the course of epidemics in 261 different world countries, by providing the daily cumulated numbers of Confirmed SARS-CoV-2 cases, Deaths and Healings. We use CRC-JHU data from 24 February 2020 (recorded start of the epidemics in Italy) through 4 July 2020.
Black points of Fig. 9, 11, 13 and 15 are the 7-day moving average of the daily number of confirmed SARS-CoV-2 cases in Italy.