Machine learning models to identify patient and microbial genetic factors

associated with carbapenem-resistant *Klebsiella pneumoniae* infection

Zena Lapp BA[a], Jennifer H Han MD[b], Jenna Wiens PhD[c], Ellie JC Goldstein MD[d,e], Ebbing

Lautenbach MD[f], Evan S Snitkin PhD[g#]

Running title: Patient and microbial predictors of infection

[a]Department of Computational Medicine and Bioinformatics, University of Michigan; 1510A

MSRB I, 1150 W. Medical Center Dr., Ann Arbor, MI, 48109-5680

[b]GlaxoSmithKline; 14200 Shady Grove Road, Rockville, MD 20850

[c]Department of Electrical Engineering and Computer Science, University of Michigan; 3765

Beyster, 2260 Hayward Street, Ann Arbor, MI 48109

[d]R M Alden Research Laboratory

[e]David Geffen School of Medicine, University of California, Los Angeles; 2021 Santa Monica

Blvd. #640-E, Santa Monica, CA 90404-2208

[f]Department of Medicine (Infectious Diseases), Department of Biostatistics, Epidemiology, and

Informatics, Perelman School of Medicine, University of Pennsylvania; 502A Johnson Pavilion,

3610 Hamilton Walk, Philadelphia, PA 19104-6073

[g]Department of Microbiology and Immunology, Department of Internal Medicine/Division of

Infectious Diseases, University of Michigan, Ann Arbor, Michigan; 1520D MSRB I, 1150 W.

Medical Center Dr., Ann Arbor, MI, 48109-5680

Corresponding author: Evan Snitkin

Address: 1520D MSRB I, 1150 W. Medical Center Dr., Ann Arbor, MI, 48109-5680

Email: esnitkin@med.umich.edu

Telephone: (734) 647-6472

Abstract

Carbapenem-resistant *Klebsiella pneumoniae* (CRKP) is a critical-priority antibiotic resistance

threat that has emerged over the past several decades, spread across the globe, and

accumulated resistance to last-line antibiotic agents. While CRKP infections are associated with

high mortality, only a small subset of patients acquiring CRKP colonization will develop clinical

infection. Here, we sought to determine the relative importance of patient characteristics and

CRKP genetic background in determining patient risk of infection. Machine learning models

classifying colonization vs. infection were built using whole-genome sequences and clinical

metadata from a comprehensive set of 331 CRKP isolates collected across 21 long-term acute

care hospitals over the course of a year. Model performance was evaluated based on area

under the receiver operating characteristics curve (AUROC) on held-out test data. We found

that patient and genomic features were predictive of clinical CRKP infection to similar extents

(AUROC IQRs: patient=0.59-0.68, genomic=0.55-0.61, combined=0.62-0.68). Patient predictors

of infection included the presence of indwelling devices, kidney disease and length of stay.

Genomic predictors of infection included presence of the ICEKp10 mobile genetic element

carrying the yersiniabactin iron acquisition system, and disruption of an O-antigen biosynthetic

gene in a sub-lineage of the epidemic ST258 clone. Altered O-antigen biosynthesis increased

association with the respiratory tract, and subsequent ICEKp10 acquisition was associated with

increased virulence. These results highlight the potential of integrated models including both

patient and microbial features to provide a more holistic understanding of patient clinical

trajectories.

Importance

Multidrug resistant organisms, such as carbapenem-resistant Klebsiella pneumoniae (CRKP),

colonize alarmingly large fractions of patients in endemic regions, but only a subset of patients

develop life-threatening infections. While patient characteristics influence risk for infection, the

relative contribution of microbial genetic background to patient risk remains unclear. We used

machine learning to determine whether patient and/or microbial characteristics can discriminate

between CRKP colonization vs. infection across multiple healthcare facilities and found that

both patient and microbial factors were predictive. Examination of informative microbial genetic

features revealed features associated with respiratory colonization and higher rates of infection.

The methods and findings presented here provide a foundation for future epidemiological,

clinical, and biological studies to better understand bacterial infections and clinical outcomes.

Introduction

Infections due to multidrug resistant organisms lead to hundreds of thousands of deaths

worldwide each year (1). Carbapenem-resistant Enterobacterales (CRE) are a critical-priority

antibiotic resistance threat that has emerged over the past several decades, spread across the

globe, and accumulated resistance to last-line antibiotic agents (2, 3). In the United States

(U.S.), CRE infections are primarily caused by the sequence type (ST) 258 strain of

carbapenem resistant *Klebsiella pneumoniae* (CRKP), which has become endemic in regional

healthcare networks (3–7). In this background of regional endemicity the risk of patient

exposure to CRKP is high, as evidenced by alarmingly high rates of colonization, especially in

long-term care settings (7, 8). However, even among critically ill patients residing in long-term

care facilities, not all colonized patients develop clinical infections that require antibiotic

treatment (3, 9). Currently, our understanding of the factors that influence whether a colonized

patient develops an infection is incomplete.

In addition to clinical characteristics of patients, the genetic background of the colonizing strain

may also influence the risk of infection, as there is extensive intra-species variation in antibiotic

resistance and virulence determinants harbored by *K. pneumoniae* (3). To date, most studies of

virulence determinants have been carried out in model systems (10, 11) or examined in human

populations without considering patient characteristics or clinical context (11, 12). One recent

study investigated virulence determinants in *K. pneumoniae* clinical isolates while controlling for

patient characteristics (13). However, this was a single-site study with a focus on carbapenem-

susceptible *K. pneumoniae*, thereby not addressing the impact of genomic variation in antibiotic-

resistant lineages that circulate in global healthcare systems.

Here, we sought to understand the importance of both patient factors and CRKP genetic

background in determining whether a patient is infected (vs. colonized) with CRKP, and identify

a set of patient and microbial features that are consistent predictors of CRKP infection across

long-term care facilities. To accomplish this, we compared patients with CRKP colonization and

infection based upon both clinical characteristics and the genomes of their colonizing or

infecting strains. We leveraged a comprehensive set of all clinical isolates collected from 21

long-term acute care hospitals (LTACHs) across the U.S. over the course of a year to obtain an

unbiased assessment of the relative contribution of bacterial and host factors to patient

colonization and infection with circulating CRKP lineages. To do this, we employed a rigorous

machine learning approach that allowed us to perform a direct comparison of patient and

bacterial features, as well as identify the most predictive features across our unbiased study

population.

Results

Of 355 clinical CRKP isolates from 21 LTACHs across the U.S. (14), we classified 149 (42%) of

the isolates as representing infection based on modified NHSN criteria (**Figure S2**, **Tables S1-**

3). Stratified by anatomic site, we classified 29/29 (100%) blood isolates as infection, 69/196

(35%) respiratory isolates as infection, and 51/130 (39%) urinary isolates as infection (**Table**

S3). More than 90% of patient isolates were from the epidemic CRKP lineage ST258 (**Tables**

S1). Patients harboring different sequence types of CRKP showed no significant differences in

infection/colonization status or anatomic site of isolation, and no substantive differences in

clinical characteristics (see supplementary material). Thus, we decided to limit our analysis to

ST258 to improve our ability to discern whether genetic variation within this dominant strain is

associated with infection.

The CRKP epidemic lineage ST258 shows evidence of sub-lineage variation in virulence

and anatomic site of isolation

We next evaluated if there exist sub-lineages of ST258 with altered virulence properties by

looking for clustering of isolates by infection on the whole-genome phylogeny (**Figure 1**; see

supplementary methods) (38). Infection status was non-randomly distributed on the phylogeny

(p=0.002), supporting our hypothesis that the genetic background of CRKP influences infection.

We performed a similar clustering analysis to look at potential niche-specific adaptation to

certain anatomic sites (**Figure 1**), and found that respiratory (p=0.001) and urinary (p=0.013)

isolates cluster on the phylogeny, but blood isolates do not (p=0.21). This analysis indicates

that, in addition to patient features, intra-strain variation in virulence and adaptation to the

urinary and respiratory tract might influence whether patients develop an infection.

Both patient and CRKP genetic characteristics are weakly predictive of infection, with

relative performance being highly facility-dependent

We next performed machine learning to quantify the ability of patient and microbial genetic

characteristics to predict CRKP infection (**Figure S1**). To prevent over- or under-fitting and

control for facility-level biases, we generated 100 train/test data splits, wherein a given LTACH

was only included either in the train or test set. Each LTACH occurred a median of 24 times

(range 13-32) in the test data split. In this way, we were able to identify patient and CRKP strain

characteristics consistently associated with infection or colonization across data splits, and thus

across patient populations in different healthcare facilities.

First, we sought to understand if patient and genomic features were individually predictive of

CRKP infection. To this end, we independently evaluated patient characteristics as well as three

different genomic feature sets for their ability to classify colonization and infection. The three

genomic feature sets were uncurated genomic (including SNPs, indels, IS elements, and

accessory genes), uncurated grouped genomic (variants grouped into genes, akin to a burden

test, e.g (28)), and curated genomic (features identified using Kleborate (16)). Across the 100

different train/test splits, we observed that the average predictive performance was weak, with

each of the genomic and patient feature sets predictive of infection to a similar degree (all 1st

quartile AUROCs > 0.5; median range=0.55-0.68; **Figure 2A**; AUPRC: **Figure S3A**).

Additionally, no one feature set was consistently the most predictive (e.g. **Figures 2B**, **2C**; all

comparisons p > 0.30, see supplementary methods for p-value calculation). Furthermore, for

each feature set the AUROCs were distributed such that the test AUROC ranged from below

0.5 to over 0.7, depending on how the data were split (i.e., which facilities appear in the

train/test sets). This variation in model performance across different train/test sets suggests that

the association of CRKP strain and patient characteristics with infection or colonization varies

across facilities.

Integration of patient and CRKP strain features does not improve discriminative

performance of overall or anatomic site-specific models

To determine if the predictive power of patient and genomic features is additive, and if

combining these disparate feature sets improved validation on held-out facilities, we built

models including both patient and curated genomic features. The discriminative performance of

the models based on the combined feature set was not significantly greater than that of the

individual feature sets (**Figure 2A**, p ≥ 0.20). Thus, despite variation in the predictive capacity of

genomic and patient features across facilities (**Figure 2C**), combining the two sets did not

improve overall performance. Focusing on anatomic site-specific models revealed similar

trends, where classification performances were similar for respiratory and urinary specific

models, and the relative predictive capacity of patient and CRKP strain features varied across

facility subsets (**Figure S4**; AUPRC: **Figure S3B**).

Some patient and genomic features consistently discriminate colonization and infection

After evaluating the predictive capacity of models, we next sought to identify patient and CRKP

strain characteristics that are most associated with CRKP infection or colonization. To this end,

we identified those patient and genomic features that consistently improved model performance

across the 100 different data splits (see methods). Evaluating the importance of features in this

way provides insight into those characteristics that generalize across different facility subsets.

This approach was taken for both overall and anatomic site-specific models to identify features

predictive of different anatomic sites of infection (**Figure 3**, **Figures S5-7**).

Several patient features were consistently associated with infection in the overall analysis,

including presence of a gastrostomy tube, presence of a central venous catheter, acute kidney

injury, and severe chronic kidney disease (**Figure 3**), all markers of critically ill patients. Only a

small number of genomic features were consistently associated with infection (**Figure 3**). No

known virulence factors were positively associated with colonization; all of the genomic features

positively associated with colonization are antibiotic resistance elements. Conversely, all but

one of the genomic features positively associated with infection (3/4) are related to virulence.

The ICEKp10 element is positively associated with infection and carries colibactin and two

different types of yersiniabactin, a previously identified *K. pneumoniae* virulence determinant

(12) (**Figure S8**). Additionally, insertion sequence-mediated disruption of the O-antigen

biosynthetic gene *kfoC* (see supplementary methods and **Figure S9** for insertion sequence

identification (21, 23, 39–41)) was associated with respiratory infection. Colibactin is a toxin (3),

and yersiniabactin is an iron scavenging system that has been identified in previous animal and

human studies as being associated with virulence (3, 10). The O-antigen of lipopolysaccharide

(LPS) is a known antigenic marker, although association with a specific anatomic site has not

been noted (42).

A sub-lineage of ST258 clade II appears to have sequentially evolved enhanced

adaptation for the respiratory tract and increased virulence

We noted that *kfoC* disruption is largely confined to a sub-lineage of ST258 present across 12

LTACHs in California (**Figures 4**, **S10**, **S11**). Consistent with this feature being associated with

respiratory infection, the disrupted *kfoC* lineage is enriched in respiratory isolates (82/118, 69%

of isolates in the disrupted *kfoC* lineage are respiratory isolates vs. 101/213, 47% in all other

isolates; Fisher's exact p=0.0001), suggesting that this lineage is associated with increased

capacity for respiratory colonization. Furthermore, a subset of isolates in the disrupted *kfoC* sub-

lineage harbor the ICEKp10 element containing yersiniabactin. Examination of these genetic

events in the context of the whole-genome phylogeny revealed that disruption of *kfoC* occurred

first, followed by at least two different acquisitions of ICEKp10 (**Figures 4**, **S10**). Within the

disrupted *kfoC* lineage, isolates with ICEKp10 are enriched in infection (31/55, 56% of isolates

with ICEKp10 are infection isolates vs. 16/63, 25% of isolates without ICEKp10, Fisher's exact p

= 0.00065), supporting an increase in virulence after acquisition of ICEKp10. It is important to

note that the observed clinical associations with ICEKp10 and kfoC disruption do not

demonstrate causality, as we cannot rule out the role of correlated genetic variation.

Discussion

There have been numerous studies aimed at identifying risk factors for healthcare-associated infections caused by prominent antibiotic-resistance threats. For the most part, these studies have found the dominant risk factors to be linked to the magnitude of exposure (e.g. length of stay or colonization pressure), use of antibiotics, and overall comorbidity (43). Here we found similar results, where length of stay and having certain comorbidities were positively associated with infection. What remains unclear is, in the critically ill populations heavily exposed to antibiotics that are at greatest risk, if genetic variation in circulating resistant lineages influences patient infection status. Here, we addressed this question by focusing on CRKP infection in a comprehensively sampled cohort of patients from 21 LTACHs across the U.S. To gain a realistic assessment of the predictive capacities of patient and CRKP genetic features, we employed a machine learning framework using multiple facility-level train/test splits. Overall, we found that, while neither patient nor CRKP genetic features have high predictive accuracy on held-out test data, both feature sets were independently associated with infection, with one or the other being more predictive on different facility subsets. Moreover, the integration of clinical and genomic data led to the discovery of an emergent sub-lineage of the epidemic ST258 clone that may have increased adaptation for the respiratory tract, and is more strongly associated with infection.

One strength of our machine learning approach is that we were able to measure the variation in discriminative performance across 100 train/test iterations that differed in which facilities were included in train and test sets. We found that performance varied greatly depending on how facilities were allocated to train and test sets, highlighting how smaller studies could overestimate or underestimate the discriminative ability of both their model and individual features. Variation in model performance across facilities could be due to facility-level heterogeneity leading to differences in the prevalence of predictive patient or genomic features

in the different test sets. For instance, certain facilities may have patient populations skewed

towards individuals with characteristics that are predictive of infection. Alternatively, certain

geographic regions may have CRKP strains that are more virulent than in other regions. These

differences could lead to a higher predictive power for certain facilities compared to others.

Another possible explanation for variation in model performance is that the critically-ill nature of

LTACH patients may be such that most patients are actually highly susceptible to infection (i.e.

many patients colonized with CRKP may ultimately develop an infection). However, it is

noteworthy that despite these potential challenges in creating generalizable models, our

analysis did yield predictors of infection and colonization consistent across test sets, and thus

across LTACHs.

We built classifiers including all genomic features as well as a curated subset of features from

Kleborate (16), and found that both are similarly weakly predictive of infection. However, while

the uncurated feature set presented challenges with downstream interpretation, our analyses on

the curated genomic features (16) facilitated novel insights into potential evolutionary

trajectories of anatomic site-specific adaptation and virulence. For example, we observed that

disruption of the O-antigen biosynthetic gene, *kfoC*, is associated with isolation from the

respiratory tract. While we cannot determine from our machine learning analysis if disruption of

kfoC is directly causal, the biological plausibility of an altered O-antigen structure mediating

evasion of innate immunity and/or other beneficial interactions with the host makes this a strong

candidate for follow-up experiments. Supporting this hypothesis, a previous study found that

absence of O-antigen is associated with decreased virulence, but not decreased intrapulmonary

proliferation, in a murine model (44). In addition, we noted that a number of antibiotic resistance

determinants were associated with colonization. We hypothesize that this observation could be

a consequence of longer duration of residence being associated with increased exposure to off-

target antibiotics (45). Finally, we also saw evidence that, after acquiring the virulence factors

yersiniabactin and colibactin on the ICEKp10 element, the disrupted *kfoC* subclade became

more strongly associated with infection, supporting the idea that circulating ST258 sub-lineages

can evolve to become both hypervirulent and multi-drug resistant (19, 46–48).

It is important to note that the machine learning method we employed does not correct for

microbial population structure. We chose this method instead of alternative bacterial genome-

wide association methods because our primary interest was in quantifying the overall predictive

capacity of bacterial genotype in a patient population that was collected in a comprehensive and

unbiased manner (i.e. all clinical isolates from 21 facilities over one year). While alternative

methods controlling for population structure may yield more precise estimates of the contribution

of individual variants, this would obfuscate the realized contribution in our patient population and

hinder direct comparison to the predictive capacity of patient features. However, this then

presented the challenge of interpreting our finding that certain sub-populations of CRKP may

differ in their predilection for causing infections at different sites. For instance, the *kfoC*

disruption is a lineage defining variant, and in principle other variants that define this lineage

could also be causal. Here, we limited our analysis to a curated set of variants belonging to

pathways known to be associated with antibiotic resistance and virulence, and found that only

kfoC disruption was associated with increased respiratory infection, thus making it a strong

candidate for follow-up in in vitro or in vivo models (10). To identify novel loci whose role in

human infection may not be appreciated both computational and experimental strategies may

be employed to help prioritize putative causal versus passenger variants. Computationally,

investigators may search for evidence of parallelism in genotype/phenotype associations, which

would bolster confidence in causality (49). Alternatively, high-throughput screens of genetic

mutants in relevant model systems can help prioritize candidates. Garnering further genomic or

experimental support for the direct role of a specific genetic variant would in turn increase the

likelihood that those genetic markers would be predictive in new strains and patient populations.

Our study also has several important limitations related to the data available. Specifically, CRKP

colonization vs. infection for non-bloodstream isolates may be difficult to discriminate using

surveillance criteria and the clinical data that were available. However, we based our definitions

on established CDC criteria with modifications used previously (7). Encouragingly, we were still

able to identify consistent predictors of infection, even with potential misclassifications. A

second limitation is that our dataset only included one clinical culture for the majority of patients,

meaning that we were unable to investigate clinical or genomic features that may be associated

with progression from colonization to infection. Furthermore, we do not have CRKP rectal

colonization isolates, and therefore cannot evaluate transition from rectal colonization to other

body sites. However, we hypothesize that comparing rectal colonization to infection may be

asking a subtly distinct question - namely bacterial genetic factors that enable translocation from

the gut to other body sites. In contrast, we hypothesize that our study design is ideal to identify

bacterial genetic factors associated with infection once at a given body site. Additionally, we

chose to focus our analysis on ST258 due to its disproportionate presence in our dataset, but

this makes it possible that our findings may not generalize to other sequence types.

Nevertheless, ST258 is the dominant clone in the U.S., and the methods we employed here can

be used to study other sequence types and other pathogens. Furthermore, our focus on the

ST258 lineage led to the particularly notable finding that even within an established endemic

multi-drug resistant lineage (that emerged circa 2000 (14)), there is continued evolution that

influences the manifestation and outcome of infection. We were also limited in the patient data

included in our model. It is likely that important differences in underlying patient conditions were

not captured by the coarse clinical variables we included, and we also did not account for

differences in genetic variation in the host (50). Other limitations include that our study was

restricted to LTACH patients, and had non-random geographic sampling. While LTACHs have

unique structural features, based on prior studies, we expect that the types of patient risk factors

considered are likely to generalize to other patient populations. Moreover, our restriction to

LTACHs in endemic geographic regions has the benefit of focusing on populations at

disproportionate risk for CRKP infection (8).

In conclusion, we employed a machine learning approach to quantify our ability to discriminate

between CRKP colonization and infection using patient and microbial genomic features. This

approach highlighted the high degree of variation in predictive accuracy across different facility

subsets. Furthermore, despite modest predictive power, we identified several genomic features

consistently associated with infection, indicating that variation in circulating CRKP strains

contributes to infection, even in the context of the critically-ill patient populations residing in

LTACHs. Future work should aim to corroborate our findings with larger cohorts, and follow up

on strong associations to determine whether they are indeed risk factors for infection. This could

ultimately help identify patients at high risk for infection and devise targeted strategies for

infection prevention.

Materials & Methods

Clinical and genomic data

We used whole-genome sequences of clinical (non-surveillance) CRKP isolates and associated

patient metadata from a prospective observational study performed in 21 LTACHs from across

the U.S. over the course of a year (BioProject accession no. PRJNA415194) (14). We included

only the first clinical bloodstream, respiratory, or urinary isolate from each patient (n=355;

Figure S1A), and subset to only ST258 isolates for the majority of analyses (n=331; **Table S1**;

see supplementary material for reasoning). Details about the analysis pipeline (15) genomic

data curation (14, 16–22), and phylogenetic reconstruction (23–26) are provided in the

supplementary material. While most clinical data cannot be shared, the deidentified patient ID,

hospital of sample isolation, and isolation site are included in the Sequence Read Archive

metadata for the BioProject.

Outcome definition

Our outcome of interest was colonization vs. clinical infection (**Figure S1B**). Based on the U.S.

Centers for Disease Control and Prevention's (CDC) established National Healthcare Safety

Network (NHSN) surveillance definitions, we considered all bloodstream isolates as

representative of infection, and used modified definitions to classify urinary and respiratory

cultures as representative of infection versus colonization (**Table S2**) (7, 27). Any isolate that

did not meet the criteria for infection was classified as colonization.

Feature sets

We studied the association between five different feature sets and infection/colonization in

CRKP ST258 (**Figure S1C**); the feature sets are described below. See supplementary methods

for details on feature set creation and processing. Counts below are for confident features from

the entire dataset prior to subsetting for different analyses.

Patient: Clinical features described in Han *et al.* (n=50; **Table S3**) (14).

Uncurated genomic: Single nucleotide variants, indels, insertion sequence elements, and

accessory genes (n=2447).

Uncurated grouped genomic: Variants grouped into genes (i.e. a burden test, e.g. (28)) and

accessory genes (n=3159).

Curated genomic: Features identified by Kleborate (16), a tool designed to identify the presence

of various genes and mutations known to be associated with either CRKP virulence or antibiotic

resistance (n=91).

Patient & curated genomic: Patient features and curated genomic features (n=141).

Machine learning & model selection

We aimed to classify clinical infection (vs. colonization) using each of the different feature sets

(see above); we built classifiers using the first clinical isolate from each patient for all isolates,

only respiratory isolates, and only urinary isolates. We performed L2 regularized logistic

regression using a modified version of the machine learning pipeline presented in Topçuoğlu *et*

al. (29) using caret version 6.0-85 (30) in R version 3.6.2 (31) (**Figure S1D1**). We randomly split

the data into 100 unique ~80/20 train/test splits, keeping all isolates from each LTACH grouped

in either the training set or the held-out test set to control for facility-level differences among the

isolates (e.g., background of circulating strains within each facility, patient population, and

clinician test ordering frequency). For valid comparison, the train/test splits were identical across

models generated with different feature sets. Hyperparameters were selected via cross-

validation on the training set to maximize the average AUROC across cross-validation folds.

See supplementary methods for more details.

Model performance

We measured model performance using the median test area under the receiver operating

characteristic curve (AUROC) and area under the precision recall curve (AUPRC), as well as

the interquartile range, across all 100 train/test splits (**Figure S1D2**).

Features consistently associated with colonization or infection

To determine the importance of each feature in predicting colonization vs. infection, we

measured how much each feature influenced model performance by calculating feature

importance using a permutation test (29) (**Figure S1D3**). For each combination of feature and

data split, we randomly permuted the feature and calculated the 'permuted test AUROC' using

the model generated with the training data. Features with a correlation of 1 were permuted

together. We performed this permutation test 100 times for each feature/data split pair, and

obtained a mean feature importance for each data split. A mean feature importance above zero

indicates that that feature improved model performance for that data split. We highlight features

where the mean permuted test AUROC was above zero in at least 75% of the data splits. In this

way, the permutation importance method allows us to take into account the variation we

observe across the 100 models, which is not possible with standard parametric statistical tests

or odds ratios.

Data analysis & visualization

See supplementary material for details on data analysis and visualization in R version 3.6.2

(31–37). All code and data that is not protected health information is on GitHub

(https://github.com/Snitkin-Lab-Umich/ml-crkp-infection-manuscript).

Acknowledgements

We gratefully acknowledge Kindred Healthcare for their assistance in collecting data and

isolates used in this study. We also thank Sean Muldoon, MD for his support and guidance

throughout the study. Finally, we thank the patients and staff of the long-term acute-care

hospitals (LTACHs) for their gracious participation in this study, Begüm Topçuoğlu for help with

the machine learning pipeline, Ali Pirani for bioinformatics support, and members of the Snitkin

lab, Mike Bachman, and Robert Weinstein for critical review of the manuscript.

Competing interests

JHH was employed at the University of Pennsylvania during the conduct of this study. She is

currently an employee of, and holds shares in, the GSK group of companies. All other authors

declare that they have no competing interests.

Funding

This research was supported by a CDC Cooperative Agreement FOA #CK16-004-Epicenters for

the Prevention of Healthcare Associated Infections, and the National Institutes of Health R01

AI139240-01 and 1R01 AI148259-01. ZL received support from the National Science

Foundation Graduate Research Fellowship Program under Grant No. DGE 1256260. Any

opinions, findings, and conclusions or recommendations expressed in this material are those of

the authors and do not necessarily reflect the views of the National Science Foundation. The

funding bodies had no role in the design of the study or collection, analysis, and interpretation of

data, or in writing the manuscript.

Authors' contributions

ESS, JHH, EL, and ZL conceptualized the study and acquired funding to support the project.

JHH and ZL performed data curation. ZL performed formal analysis, investigation, and

visualization. ESS, JW, and ZL developed methodology. ESS supervised the project. ZL and

ESS wrote the original draft. All authors reviewed and edited the manuscript.

References

1.   Organization WH. 2014. Antimicrobial Resistance: Global Report on Surveillance. World

Health Organization, Geneva.

2.   Munoz-Price LS, Poirel L, Bonomo RA, Schwaber MJ, Daikos GL, Cormican M, Cornaglia

G, Garau J, Gniadkowski M, Hayden MK, Kumarasamy K, Livermore DM, Maya JJ,

Nordmann P, Patel JB, Paterson DL, Pitout J, Villegas MV, Wang H, Woodford N, Quinn

JP. 2013. Clinical epidemiology of the global expansion of Klebsiella pneumoniae

carbapenemases. Lancet Infect Dis 13:785–796.

3.   Wyres KL, Lam MMC, Holt KE. 2020. Population genomics of Klebsiella pneumoniae.

Nature Reviews Microbiology 1–16.

4.   Ansari U, Lawsin A, Campbell D, Albrecht V, McAllister G, Bulens S, Walters MS, Jacob

JT, Satola SW, Wilson LE, Lynfield R, Vagnone PMS, Janelle SJ, Xavier K, Dumyati G,

Hardy D, Phipps EC, Culbreath K, Beldavs Z, Morey K, Kainer MA, Roberts S, Kallen A,

Rasheed JK, Karlsson MS. 2017. Molecular Characterization of Carbapenem-Resistant

Enterobacteriaceae in the USA, 2011–2015. Open Forum Infect Dis 4:S179–S179.

5.   Logan LK, Weinstein RA. 2017. The Epidemiology of Carbapenem-Resistant

Enterobacteriaceae: The Impact and Evolution of a Global Menace. J Infect Dis 215:S28–

S36.

6.   Lee C-R, Lee JH, Park KS, Kim YB, Jeong BC, Lee SH. 2016. Global Dissemination of

Carbapenemase-Producing Klebsiella pneumoniae: Epidemiology, Genetic Context,

Treatment Options, and Detection Methods. Front Microbiol 7.

7.   Han JH, Goldstein EJC, Wise J, Bilker WB, Tolomeo P, Lautenbach E. 2017. Epidemiology

of Carbapenem-Resistant Klebsiella pneumoniae in a Network of Long-Term Acute Care

Hospitals. Clin Infect Dis 64:839–844.

8.   Lin MY, Lyles-Banks RD, Lolans K, Hines DW, Spear JB, Petrak R, Trick WE, Weinstein

RA, Hayden MK, Acuna N, Albright R, Alexander P, Anthony T, Bardowski L, Berends C,

Bonebrake A, Bova J, Braggs A, Burtun S, Byrnes MA, Cagle C, Chavis R, Cienkus S,

Collins-Johnson S, Coomer C, Chou T, Cullen D, DeGuzman D, Donceras O, Feller M,

Fung S, Gasienica JAM, Garcia-Houchins S, Genovese G, Gentile M, Gonzaga G,

Goodwin E, Hernandez E, Kerridge J, Kirk J, Lavin MA, Lee S, Lepinski J, Myrick S, Oats

T, O'Donnell A, Pasinos V, Patton JA, Perez M, Purpura J, Rakoski L, Reiner S, Saiyed A,

Santine M, Schmitt B, Schultz C, Semar RM, Sethuraman H, Silkaitis C, Stutler E,

Thompson A, Wickman K, Bolon M, Cha A, Landon E, Malow J, Mohapatra S, Segreti J,

Weber S, Welbel S, Zembower T. 2013. The Importance of Long-term Acute Care

Hospitals in the Regional Epidemiology of Klebsiella pneumoniae Carbapenemase–

Producing Enterobacteriaceae. Clin Infect Dis 57:1246–1252.

9. Lee BY, Bartsch SM, Wong KF, Kim DS, Cao C, Mueller LE, Gussin GM, McKinnell JA,

Miller LG, Huang SS. 2019. Tracking the spread of carbapenem-resistant

Enterobacteriaceae (CRE) through clinical cultures alone underestimates the spread of

CRE even more than anticipated. Infection Control & Hospital Epidemiology 40:731–734.

10. Bachman MA, Oyler JE, Burns SH, Caza M, Lépine F, Dozois CM, Weiser JN. 2011.

Klebsiella pneumoniae Yersiniabactin Promotes Respiratory Tract Infection through

Evasion of Lipocalin 2. Infection and Immunity 79:3309–3316.

11. Ernst CM, Braxton JR, Rodriguez-Osorio CA, Zagieboylo AP, Li L, Pironti A, Manson AL,

Nair AV, Benson M, Cummins K, Clatworthy AE, Earl AM, Cosimi LA, Hung DT. 2020.

Adaptive evolution of virulence and persistence in carbapenem-resistant Klebsiella

pneumoniae. 5. Nature Medicine 26:705–711.

12. Holt KE, Wertheim H, Zadoks RN, Baker S, Whitehouse CA, Dance D, Jenney A, Connor

TR, Hsu LY, Severin J, Brisse S, Cao H, Wilksch J, Gorrie C, Schultz MB, Edwards DJ,

Nguyen KV, Nguyen TV, Dao TT, Mensink M, Minh VL, Nhu NTK, Schultsz C, Kuntaman

K, Newton PN, Moore CE, Strugnell RA, Thomson NR. 2015. Genomic analysis of

diversity, population structure, virulence, and antimicrobial resistance in Klebsiella

pneumoniae, an urgent threat to public health. PNAS 112:E3574–E3581.

13. Martin RM, Cao J, Wu W, Zhao L, Manthei DM, Pirani A, Snitkin E, Malani PN, Rao K,

Bachman MA. 2018. Identification of Pathogenicity-Associated Loci in Klebsiella

pneumoniae from Hospitalized Patients. mSystems 3.

14. Han JH, Lapp Z, Bushman F, Lautenbach E, Goldstein EJC, Mattei L, Hofstaedter CE, Kim

D, Nachamkin I, Garrigan C, Jain T, Bilker W, Wolford HM, Slayton RB, Wise J, Tolomeo

P, Snitkin ES. 2019. Whole-Genome Sequencing To Identify Drivers of Carbapenem-

Resistant Klebsiella pneumoniae Transmission within and between Regional Long-Term

Acute-Care Hospitals. Antimicrobial Agents and Chemotherapy 63.

15. Köster J, Rahmann S. 2012. Snakemake—a scalable bioinformatics workflow engine.

Bioinformatics 28:2520–2522.

16. Holt K. 2020. katholt/Kleborate. Python.

17. Wyres KL, Wick RR, Gorrie C, Jenney A, Follador R, Thomson NR, Holt KE. 2016.

Identification of Klebsiella capsule synthesis loci from whole genome data. Microbial

Genomics, 2:e000102.

18. Wick RR, Heinz E, Holt KE, Wyres KL. 2018. Kaptive Web: User-Friendly Capsule and

Lipopolysaccharide Serotype Prediction for Klebsiella Genomes. Journal of Clinical

Microbiology 56.

19. Lam MMC, Wick RR, Wyres KL, Gorrie CL, Judd LM, Jenney AWJ, Brisse S, Holt KE.

2018. Genetic diversity, mobilisation and spread of the yersiniabactin-encoding mobile

element ICEKp in Klebsiella pneumoniae populations. Microbial Genomics, 4:e000196.

20. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM.

2012. A program for annotating and predicting the effects of single nucleotide

polymorphisms, SnpEff. Fly 6:80–92.

21. Treepong P, Guyeux C, Meunier A, Couchoud C, Hocquet D, Valot B. 2018. panISa: ab

initio detection of insertion sequences in bacterial genomes from short read sequence

data. Bioinformatics 34:3795–3800.

22. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, Fookes M, Falush D,

Keane JA, Parkhill J. 2015. Roary: rapid large-scale prokaryote pan genome analysis.

Bioinformatics 31:3691–3693.

23. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin

R. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25:2078–

2079.

24. Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J, Harris

SR. 2015. Rapid phylogenetic analysis of large samples of recombinant bacterial whole

genome sequences using Gubbins. Nucleic Acids Res 43:e15–e15.

25. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol

32:268–274.

26. Minh BQ, Nguyen MAT, von Haeseler A. 2013. Ultrafast Approximation for Phylogenetic

Bootstrap. Mol Biol Evol 30:1188–1195.

27. 2020. 2020 NHSN Patient Safety Component Manual 434.

28. Saund K, Lapp Z, Thiede SN, Pirani A, Snitkin ES. 2020. prewas: data pre-processing for

more informative bacterial GWAS. Microbial Genomics, 6:e000368.

29. Topçuoğlu BD, Lesniak NA, Ruffin MT, Wiens J, Schloss PD. 2020. A Framework for

Effective Application of Machine Learning to Microbiome-Based Classification Problems.

mBio 11.

30. Kuhn M. 2008. Building Predictive Models in R Using the caret Package. 1. Journal of

Statistical Software 28:1–26.

31. R: The R Project for Statistical Computing.

32. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes

A, Henry L, Hester J, Kuhn M, Pedersen T, Miller E, Bache S, Müller K, Ooms J, Robinson

D, Seidel D, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H. 2019. Welcome

to the Tidyverse. Journal of Open Source Software 4:1686.

33.   Wilke CO. 2019. cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2."

34.   Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. 2017. ggtree: an r package for visualization

and annotation of phylogenetic trees with their covariates and other associated data.

Methods in Ecology and Evolution 8:28–36.

35.   Yu G, Lam TT-Y, Zhu H, Guan Y. 2018. Two Methods for Mapping and Visualizing

Associated Data on Phylogeny Using Ggtree. Mol Biol Evol 35:3041–3043.

36.   Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics 35:526–528.

37.   grid package | R Documentation.

38.   Popovich KJ, Snitkin ES, Hota B, Green SJ, Pirani A, Aroutcheva A, Weinstein RA. 2017.

Genomic and Epidemiological Evidence for Community Origins of Hospital-Onset

Methicillin-Resistant Staphylococcus aureus Bloodstream Infections. J Infect Dis

215:1640–1647.

39.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.

BLAST+: architecture and applications. BMC Bioinformatics 10:421.

40.   Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, Conrad N, Dietrich EM, Disz

T, Gabbard JL, Gerdes S, Henry CS, Kenyon RW, Machi D, Mao C, Nordberg EK, Olsen

GJ, Murphy-Olson DE, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Vonstein

V, Warren A, Xia F, Yoo H, Stevens RL. 2017. Improvements to PATRIC, the all-bacterial

Bioinformatics Database and Analysis Resource Center. Nucleic Acids Res 45:D535–

D542.

41.   Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler

transform. Bioinformatics 25:1754–1760.

42.   Follador R, Heinz E, Wyres KL, Ellington MJ, Kowarik M, Holt KE, Thomson NR. 2016. The

diversity of Klebsiella pneumoniae surface polysaccharides. Microb Genom 2.

43.   Liu P, Li X, Luo M, Xu X, Su K, Chen S, Qing Y, Li Y, Qiu J. 2017. Risk Factors for

Carbapenem-Resistant Klebsiella pneumoniae Infection: A Meta-Analysis. Microbial Drug

Resistance 24:190–198.

44. Shankar-Sinha S, Valencia GA, Janes BK, Rosenberg JK, Whitfield C, Bender RA,

Standiford TJ, Younger JG. 2004. The Klebsiella pneumoniae O Antigen Contributes to

Bacteremia and Lethality during Murine Pneumonia. Infection and Immunity 72:1423–

1430.

45. Tedijanto C, Olesen SW, Grad YH, Lipsitch M. 2018. Estimating the proportion of

bystander selection for antibiotic resistance among potentially pathogenic bacterial flora.

Proceedings of the National Academy of Sciences 115:E11988–E11995.

46. Marsh JW, Mustapha MM, Griffith MP, Evans DR, Ezeonwuka C, Pasculle AW, Shutt KA,

Sundermann A, Ayres AM, Shields RK, Babiker A, Cooper VS, Tyne DV, Harrison LH.

2019. Evolution of Outbreak-Causing Carbapenem-Resistant Klebsiella pneumoniae

ST258 at a Tertiary Care Hospital over 8 Years. mBio 10:e01945-19.

47. Zhou K, Xiao T, David S, Wang Q, Zhou Y, Guo L, Aanensen D, Holt KE, Thomson NR,

Grundmann H, Shen P, Xiao Y. Novel Subclone of Carbapenem-Resistant Klebsiella

pneumoniae Sequence Type 11 with Enhanced Virulence and Transmissibility, China -

Volume 26, Number 2—February 2020 - Emerging Infectious Diseases journal - CDC

https://doi.org/10.3201/eid2602.190594.

48. Gu D, Dong N, Zheng Z, Lin D, Huang M, Wang L, Chan EW-C, Shu L, Yu J, Zhang R,

Chen S. 2018. A fatal outbreak of ST11 carbapenem-resistant hypervirulent Klebsiella

pneumoniae in a Chinese hospital: a molecular epidemiological study. The Lancet

Infectious Diseases 18:37–46.

49. Saund K, Snitkin ES. 2020. hogwash: Three Methods for Genome-Wide Association

Studies in Bacteria. bioRxiv 2020.04.19.048421.

50. Chapman SJ, Hill AVS. 2012. Human genetic susceptibility to infectious disease. 3. Nat

Rev Genet 13:175–188.

 Main figures

Figure 1: Infection and anatomic site cluster on the phylogeny. Maximum likelihood phylogenetic tree

of all isolates including infection or colonization classification for each isolate and anatomic site of

isolation. The scale bar to the right of the phylogeny shows the branch length in substitutions per site.

Testing for non-random distribution of isolates on the phylogeny (see supplementary methods) revealed

clustering of infection, respiratory, and urinary isolates on the phylogeny, respectively.

Figure 2: Test AUROCs for various classifiers identifying CRKP colonization vs. infection vary

substantially across data splits. (A) Test AUROCs for 100 L2 regularized logistic regression models

different using train/test splits. All isolates from a given LTACH were included in either the training split or

the testing split for each data split. We built models using five different feature sets, keeping the same

100 data splits. AUROCs of different feature sets were not significantly different. In the right two panels,

the curated genomic feature set AUROCs are compared to: (B) the uncurated genomic feature set

AUROCs, and (C) the patient feature set AUROCs. Each point is the resulting pair of AUROCs for models

built with the same data split, but the two respective feature sets. The dotted lines in all 3 panels indicate

the AUROC for choosing an outcome randomly (0.5); anything below the line is worse than random, and

anything above the line is better than random. The solid diagonal line in the right two panels is the line

y=x; points below the line correspond to a higher curated genomic AUROC for that data split, and points

above the line correspond to a higher uncurated genomic AUROC (B), or patient AUROC (C),

respectively. The colors in panels (B) and (C) correspond to the colors in panel (A); the points in a given

colored area indicate that that feature set had the higher AUROC for that data split. In both cases, one

feature set does not consistently outperform the other (p=0.4; see supplementary methods for p-value

calculation). AUROC=area under the receiver operating characteristic curve.

Figure 3: Features consistently associated with colonization or infection sometimes differ

between the overall, respiratory, and urinary models. Feature-specific improvement in model

performance, measured as the mean difference between test and permuted AUROC (see methods), of

features found to be consistently associated with colonization or infection in at least one of the following

analyses: overall, respiratory-specific, urinary-specific. We consider features to be associated with

infection/colonization if the AUROC difference was greater than zero in over 75% of the 100 data splits.

The vertical solid black line indicates a difference of zero (i.e. the feature provides no improvement to

model performance). Horizontal dotted lines separate features associated with urinary but not respiratory

isolates (top), both urinary and respiratory (or all) isolates (middle), or respiratory but not urinary isolates

(bottom). Bla=Beta lactamase, res=confers resistance to that antibiotic class. AUROC=area under the

receiver operating characteristic curve.

Figure 4: Select epidemiologic and genomic features visualized on the phylogeny indicate that a

sub-clade of ST258 clade II may exhibit enhanced niche-specific adaptation and virulence. ST258

maximum likelihood phylogeny with the tip labels colored by KPC gene. The blue box indicates the sub-

lineage with apparent altered niche-specific adaptation that acquires an additional virulence locus. The

heatmap beside the tree indicates information about the isolate. From left to right: if it is a respiratory

isolate, if it is an infection isolate, if *kfoC* is disrupted, and if it contains ICEKp10. Disrupted *kfoC* was

associated with infection in the overall and respiratory machine learning analyses and ICEKp10 presence

was associated with infection in the overall analysis. The scale bar to the top left of the phylogeny shows

the branch length in substitutions per site. ybt=Yersiniabactin; ybt0 and ybt17 are two ybtSTs defined by

Kleborate.

Supplementary materials

Supplementary methods

Pipeline

We created a snakemake pipeline (1) to perform data pre-processing, machine learning analysis, and

figure generation.

Genomic data

We used Kleborate version 0.3.0 (2) to identify sequence type, capsular types (K locus (3) and O locus

(4)), virulence factor (5), and antibiotic resistance genes. In addition, we identified single nucleotide

variants (SNVs) and indels by mapping reads to the KPNIH1 reference genome (BioProject accession

number PRJNA73191; https://github.com/Snitkin-Lab-Umich/variant_calling_pipeline (6)). After variant

calling, we identified the predicted functional impact of variants using SnpEff version 4.3T (7). We also

identified large insertion sequence (IS) elements using panISa version 0.1.4 with the default settings (8),

and accessory genome genes using Roary version 3.12.0 with the default settings (9).

Phylogenetic tree reconstruction

We constructed a phylogenetic tree of all ST258 isolates by first creating a whole-genome alignment by

mapping reads to the KPNIH1 reference genome (GenBank accession number CP008827.1) using bwa

mem version 0.7.12 (10); samtools version 1.9 (11) was used to generate binary alignment files. We then

masked sites identified as recombinant by Gubbins version 2.3.2 (12) and used this masked whole-

genome alignment to build a maximum likelihood phylogeny with IQ-TREE version 1.6.12 (13) using a

GTR model of nucleotide substitution and ultra-fast bootstrap with 1000 replicates (-b 1000) (14). The tree

was midpoint-rooted using the midpoint.root function in phytools version 0.6-99 (15), thus splitting the tree

into ST258 clades I and II (16).

Phylogenetic clustering test

To determine whether infection isolates cluster on the phylogeny more than expected by random chance,

we performed a permutation test (17). To identify the true number of isolates in a pure cluster while

controlling for isolate clustering by LTACH, we enumerated isolates in pure infection or colonization

subtrees that also contain isolates from more than one LTACH. Then we randomized the infection labels

across isolates 1000 times and enumerated the number of isolates in a pure subtree for each

randomization. We then compared the random counts of isolates in a pure cluster to the true number and

calculated an empirical p-value.

Feature set pre-processing

We used five feature sets to study the association between infection and colonization. To determine how

well machine learning models perform using a targeted vs. untargeted approach, we performed machine

learning with uncurated genomic features as well as curated genomic features known to be related to

virulence. In an attempt to increase power, we not only performed machine learning on individual

uncurated genomic features, but also using a burden test where genomic variants were grouped into

genes and features were presence/absence of a variant in a gene. This may help capture variation

associated with infection at the gene level that might not be captured at the variant level (e.g. different

samples can have different variants in the same gene that all lead to a similar function). For the uncurated

grouped genomic analysis, grouped variants included SNVs and indels identified as moderate or high

impact by SnpEff and IS elements in a gene or upstream of a gene. We also grouped modifier variants

identified by SnpEff into intergenic regions. Only features that occurred in more than one isolate and less

than n-1 isolates were included. For the uncurated genomic and uncurated grouped genomic datasets we

collapsed genomic features with an identical pattern across all isolates to reduce machine learning

runtime.

Machine learning rationale

696   We chose to perform L2 regularized logistic regression because it is easily interpretable, often performs

697   just as well as more complicated methods when limited training data are available (18), and has a

698   grouping effect, which means that all associated features are identified even if they are collinear (19).

Machine learning details

701   The machine learning pipeline removes variables with near-zero variance, normalizes all features to be

702   between zero and one, and sets at least 20% of the data aside for testing in each train/test split. We split

703   the training data into train and validate sets by LTACH using the groupKMultiFolds function from the caret

704   R package (caret version 6.0-85) (20). Within each training set, we selected hyperparameters via five-fold

705   cross-validation 100 times, maximizing the average cross-validation AUROC.

Comparing model performance between feature sets

708   To determine whether there was a significant difference in model performance between different feature

709   sets, we calculated a two-sided empirical p-value for each identical train/test split using the formula (2 x

710   min(fraction of AUROC differences ≥ 0, fraction of AUROC differences ≤ 0) (18).

Identification of IS element insertions in certain O2v2 kfoC genes

One of the curated genomic features we identified as associated with infection was what Kleborate called

a missing gene (kfoC) in the O locus operon of certain O2v2 serotypes. On the Kleborate website, they

indicate that a missing gene could represent a truly missing gene, or a gene that is split between two

different contigs. To further investigate this, we used blastn version 2.9.0 (21) to search for kfoC genes in

each of the samples. Additionally, we aligned the reads of all genomes to CP031810, a complete *K.*

pneumoniae genome from PATRIC (22) using bwa mem version 0.7.12 (10), and samtools (11) was used

to generate binary alignment files. Next, we used panISa (8) to identify IS element insertions in kfoC.

Data analysis & visualization

We used Fisher's exact tests for bivariable analyses with categorical variables, and Wilcoxon rank-sum

tests for bivariable analyses with continuous variables. We performed all data analysis and visualization

in R version 3.6.2 (23) using the following packages: tidyverse version 1.3.0 (24), cowplot version 1.0.0

(25), ggtree version 2.0.1 (26, 27), ape version 5.3 (28), phytools version 0.6-99 (15), and grid version

3.6.2 (29).

Supplementary results

Patients with different sequence types show no substantive differences in infection status,

anatomic site of isolation, or clinical characteristics

Over 90% of the isolates in our dataset were ST258 (**Table S2**), the dominant strain in the US (30). At the

level of sequence type, we found no difference in infection prevalence (p=0.44) or anatomic site of

isolation (p=0.66), indicating that at this coarse level there was not evidence of differences in strain

virulence or adaptation to a certain anatomic site. Bivariable comparison of patient factors between

patients with different sequence types revealed only four significant differences (unadjusted p-values <

0.05; previous use of piperacillin/tazobactam, cefepime, ciprofloxacin, or fluoroquinolones). As we found

no substantive differences in infection, anatomic site of isolation, or patient variables when comparing

different sequence types, we chose to focus all subsequent analyses on ST258, the dominant sequence

type in our dataset. The genetic variation of isolates within ST258 is much smaller than the genetic

variation between sequence types, so limiting our analyses to ST258 could improve our ability to identify

associations of interest within ST258.

Supplementary references

1.   Köster J, Rahmann S. 2012. Snakemake—a scalable bioinformatics workflow engine.

Bioinformatics 28:2520–2522.

2.   Holt K. 2020. katholt/Kleborate. Python.

3.   Wyres KL, Wick RR, Gorrie C, Jenney A, Follador R, Thomson NR, Holt KE. 2016. Identification

of Klebsiella capsule synthesis loci from whole genome data. Microbial Genomics, 2:e000102.

4.   Wick RR, Heinz E, Holt KE, Wyres KL. 2018. Kaptive Web: User-Friendly Capsule and

Lipopolysaccharide Serotype Prediction for Klebsiella Genomes. Journal of Clinical Microbiology 56.

5.   Lam MMC, Wick RR, Wyres KL, Gorrie CL, Judd LM, Jenney AWJ, Brisse S, Holt KE. 2018.

Genetic diversity, mobilisation and spread of the yersiniabactin-encoding mobile element ICEKp in

Klebsiella pneumoniae populations. Microbial Genomics, 4:e000196.

6.   Conlan S, Park M, Deming C, Thomas PJ, Young AC, Coleman H, Sison C, Program NCS,

Weingarten RA, Lau AF, Dekker JP, Palmore TN, Frank KM, Segre JA. 2016. Plasmid Dynamics in

KPC-Positive Klebsiella pneumoniae during Long-Term Patient Colonization. mBio

7.   Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012.

A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. Fly

(Austin) 6:80–92.

8.   Treepong P, Guyeux C, Meunier A, Couchoud C, Hocquet D, Valot B. 2018. panISa: ab initio

detection of insertion sequences in bacterial genomes from short read sequence data.

Bioinformatics 34:3795–3800.

9.   Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, Fookes M, Falush D,

Keane JA, Parkhill J. 2015. Roary: rapid large-scale prokaryote pan genome analysis. Bioinformatics

31:3691–3693.

10.   Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler

transform. Bioinformatics 25:1754–1760.

11.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R.

2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25:2078–2079.

12.   Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J, Harris SR.

2015. Rapid phylogenetic analysis of large samples of recombinant bacterial whole genome

sequences using Gubbins. Nucleic Acids Res 43:e15–e15.

13.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol 32:268–274.

14.   Minh BQ, Nguyen MAT, von Haeseler A. 2013. Ultrafast Approximation for Phylogenetic

Bootstrap. Mol Biol Evol 30:1188–1195.

15.   Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other

things). Methods in Ecology and Evolution 3:217–223.

16.   DeLeo FR, Chen L, Porcella SF, Martens CA, Kobayashi SD, Porter AR, Chavda KD, Jacobs

MR, Mathema B, Olsen RJ, Bonomo RA, Musser JM, Kreiswirth BN. 2014. Molecular dissection of

the evolution of carbapenem-resistant multilocus sequence type 258 Klebsiella pneumoniae. PNAS

111:4988–4993.

17.   Popovich KJ, Snitkin ES, Hota B, Green SJ, Pirani A, Aroutcheva A, Weinstein RA. 2017.

Genomic and Epidemiological Evidence for Community Origins of Hospital-Onset Methicillin-

Resistant Staphylococcus aureus Bloodstream Infections. J Infect Dis 215:1640–1647.

18.   Topçuoğlu BD, Lesniak NA, Ruffin M, Wiens J, Schloss PD. 2019. Effective application of

machine learning to microbiome-based classification problems. bioRxiv 816090.

19.   Schreiber-Gregory DN, Waller J, Smith T. 2018. Ridge Regression and multicollinearity: An in-

depth review. Model Assisted Statistics & Applications 13:359–365.

20.   Kuhn M. 2008. Building Predictive Models in R Using the caret Package. 1. Journal of

Statistical Software 28:1–26.

21.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.

BLAST+: architecture and applications. BMC Bioinformatics 10:421.

22.   Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, Conrad N, Dietrich EM, Disz T,

Gabbard JL, Gerdes S, Henry CS, Kenyon RW, Machi D, Mao C, Nordberg EK, Olsen GJ, Murphy-

Olson DE, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Vonstein V, Warren A, Xia F, Yoo

H, Stevens RL. 2017. Improvements to PATRIC, the all-bacterial Bioinformatics Database and

Analysis Resource Center. Nucleic Acids Res 45:D535–D542.

23.   R: The R Project for Statistical Computing.

24.   Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes A,

Henry L, Hester J, Kuhn M, Pedersen T, Miller E, Bache S, Müller K, Ooms J, Robinson D, Seidel D,

Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H. 2019. Welcome to the Tidyverse.

Journal of Open Source Software 4:1686.

25.   Wilke CO. 2019. cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2."

26.   Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. 2017. ggtree: an r package for visualization and

annotation of phylogenetic trees with their covariates and other associated data. Methods in Ecology

and Evolution 8:28–36.

27.   Yu G, Lam TT-Y, Zhu H, Guan Y. 2018. Two Methods for Mapping and Visualizing Associated

Data on Phylogeny Using Ggtree. Mol Biol Evol 35:3041–3043.

28.   Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics 35:526–528.

29.   grid package | R Documentation.

30.   Wyres KL, Lam MMC, Holt KE. 2020. Population genomics of Klebsiella pneumoniae. Nature

Reviews Microbiology 1–16.

Supplementary tables & figures

Table S1: Infection status and anatomic site of isolation by CRKP sequence type.

Sequence Type	Colonization			Infection		Total (%)
	Respiratory	Urinary	Blood	Respiratory	Urinary	
ST15	4	3	1	2	0	10 (2.8)
ST15-1LV	0	0	0	1	0	1 (0.3)
ST17	0	0	0	2	0	2 (0.6)
ST2237	0	1	0	0	0	1 (0.3)
ST258	121	71	28	62	49	331 (93.2)
ST307	1	4	0	2	2	9 (2.5)
ST36	1	0	0	0	0	1 (0.3)

822 **Table S2: Definitions of infection for urinary and respiratory cultures.**

Source	Infection definition
Urinary	≥100,000 colony-forming units/mL on urinary culture with no more than 2 species of microorganisms AND at least one of the following: - Leukocytosis (≥10,000 cells/μL) ≤ 7 days prior to specimen collection - Positive urinalysis demonstrated by at least one of the following findings: - Positive dipstick for leukocyte esterase and/or nitrate - Pyuria (urinary specimen with ≥ 10 white blood cells/mL or ≥ 3 white blood cells/high power field of unspun urinary) - Microorganisms seen on Gram stain of unspun urinary

Respiratory	Chest X-ray with interpretation of pneumonia, probable pneumonia, or infiltrate/consolidation AND leukocytosis (≥10,000 cells/μL) ≤ 7 days prior to specimen collection

Table S3: Bivariable analysis of patient features associated with CRKP ST258 infection in LTACH residents.

Variable	Infection (n=139)	Colonization (n=192)	All		Respiratory		Urinary	
			OR (95% CI)	P value	OR (95% CI)	P value	OR (95% CI)	P value
Blood culture	28 (20%)	0 (0%)	-	-	-	-	-	-
Respiratory culture	62 (45%)	121 (63%)	0.47 (0.3-0.76)	0.0011	-	-	-	-
Urinary culture	49 (35%)	71 (37%)	0.93 (0.57-1.5)	0.82	-	-	-	-
Admission to an LTACH in the year prior to culture[1]	9 (6%)	7 (4%)	1.83 (0.59-5.93)	0.30	2.03 (0.45-9.19)	0.31	2.23 (0.25-27.73)	0.40
Length of stay before culture, median (IQR)[2]	23 (9.0-50)	18 (2.0-34)	-	0.98	-	0.66	-	0.88
Age, median (IQR)[2]	72 (64-79)	73 (64-83)	-	0.029	-	0.0010	-	0.55
Sex (male)	74 (53%)	102 (53%)	1 (0.63-1.59)	1.0	1.15 (0.59-2.29)	0.75	0.94 (0.42-2.1)	1.0

Presence of a central venous catheter	87 (63%)	97 (51%)	1.64 (1.03-2.62)	0.033	3.36 (1.54-7.87)	0.00094	0.79 (0.34-1.81)	0.57
Tracheostomy	83 (60%)	141 (73%)	0.54 (0.33-0.88)	0.0091	1.45 (0.41-6.51)	0.78	0.42 (0.17-0.99)	0.035
Presence of a urinary catheter	87 (63%)	94 (49%)	1.74 (1.09-2.79)	0.019	4.19 (2.06-8.88)	0.000020	0.71 (0.32-1.59)	0.45
Congestive heart failure	27 (19%)	38 (20%)	0.98 (0.54-1.75)	1.0	1.74 (0.8-3.76)	0.14	0.25 (0.04-0.95)	0.035
Acute or chronic respiratory failure	55 (40%)	66 (34%)	1.25 (0.77-2.01)	0.36	1.93 (0.99-3.78)	0.040	0.68 (0.27-1.64)	0.42
Acute kidney injury	76 (55%)	76 (40%)	1.84 (1.16-2.93)	0.0074	2.65 (1.36-5.29)	0.0028	1.18 (0.53-2.62)	0.71
Malignancy (solid or liquid)	16 (12%)	27 (14%)	0.8 (0.38-1.61)	0.51	0.97 (0.31-2.76)	1.0	0.62 (0.18-1.94)	0.45
Brain injury[3]	31 (22%)	31 (16%)	1.49 (0.82-2.7)	0.20	3.37 (1.57-7.32)	0.00082	0.1 (0-0.75)	0.014
Presence of a gastrostomy tube	62 (45%)	59 (31%)	1.81 (1.12-2.93)	0.011	2.98 (1.52-5.94)	0.00083	1.45 (0.61-3.44)	0.42
Obesity[4]	4 (3%)	14 (7%)	0.38 (0.09-1.24)	0.091	0.23 (0.01-1.81)	0.28	0.46 (0.04-2.74)	0.47
Malnourished/underweight [5]	38 (27%)	43 (22%)	1.3 (0.76-	0.30	2.02 (0.94-4.32)	0.065	0.66 (0.24-1.67)	0.39

			2.22)					
Transplant[6]	4 (3%)	3 (2%)	1.86 (0.31-12.93)	0.46	3.01 (0.33-36.91)	0.34	1.45 (0.02-115.98)	1.0
Cirrhosis	8 (6%)	6 (3%)	1.89 (0.56-6.77)	0.28	3.43 (0.64-22.82)	0.12	0.47 (0.01-6.12)	0.64
Severe chronic kidney disease[7]	52 (37%)	55 (29%)	1.49 (0.91-2.43)	0.097	2.12 (1.08-4.19)	0.023	1.06 (0.42-2.63)	1.0
Pulmonary disease[8]	25 (18%)	34 (18%)	1.02 (0.55-1.87)	1.0	1.09 (0.44-2.6)	0.84	0.68 (0.21-1.99)	0.48
Ventilator-dependent respiratory failure	52 (37%)	48 (25%)	1.79 (1.09-2.96)	0.021	3.34 (1.69-6.69)	0.00023	0.49 (0.11-1.78)	0.27
Stage IV/V decubitus ulcer	31 (22%)	32 (17%)	1.43 (0.79-2.58)	0.20	2.02 (0.94-4.32)	0.065	1.54 (0.5-4.81)	0.44
Amikacin[9]	23 (17%)	14 (7%)	2.51 (1.18-5.52)	0.013	1.88 (0.63-5.49)	0.21	2.77 (0.66-13.69)	0.12
Aztreonam[9]	5 (4%)	6 (3%)	1.16 (0.27-4.65)	1.0	1.99 (0.26-15.34)	0.41	0 (0-3.49)	0.27
Trimethoprim/sulfamethoxazole[9]	6 (4%)	5 (3%)	1.68 (0.42-7.13)	0.54	1.99 (0.26-15.34)	0.41	1.46 (0.1-20.84)	1.0
Cefepime[9]	20 (14%)	32 (17%)	0.84 (0.43-1.6)	0.65	1.1 (0.42-2.73)	0.83	0.57 (0.17-1.74)	0.33

Ceftaroline[9]	4 (3%)	3 (2%)	1.86 (0.31-12.93)	0.46	0 (0-4.73)	0.55	Inf (0.61-Inf)	0.066
Ceftazidime[9]	3 (2%)	7 (4%)	0.58 (0.1-2.61)	0.53	0.97 (0.15-4.76)	1.0	0 (0-56.46)	1.0
Ceftriaxone[9]	11 (8%)	8 (4%)	1.97 (0.7-5.82)	0.16	1.68 (0.39-6.91)	0.51	3.04 (0.42-34.89)	0.22
Ciprofloxacin[9]	2 (1%)	6 (3%)	0.45 (0.04-2.59)	0.48	1.96 (0.02-155.49)	1.0	0 (0-1.55)	0.078
Colistin[9]	12 (9%)	11 (6%)	1.55 (0.61-4.02)	0.38	2.4 (0.72-8.22)	0.15	0.35 (0.01-3.7)	0.65
Daptomycin[9]	10 (7%)	3 (2%)	4.86 (1.22-28)	0.018	8.18 (0.79-409.93)	0.046	4.75 (0.8-50.2)	0.062
Ertapenem[9]	6 (4%)	4 (2%)	2.12 (0.49-10.4)	0.33	3.97 (0.2-237.55)	0.27	0.47 (0.01-6.12)	0.64
Metronidazole[9]	25 (18%)	35 (18%)	0.98 (0.53-1.8)	1.0	0.97 (0.43-2.11)	1.0	1.27 (0.33-4.77)	0.77
Gentamicin[9]	5 (4%)	5 (3%)	1.39 (0.31-6.19)	0.75	0.98 (0.09-7.03)	1.0	2.95 (0.15-178)	0.57
Imipenem[9]	3 (2%)	2 (1%)	2.09 (0.24-25.36)	0.65	1.96 (0.02-155.49)	1.0	0 (0-56.46)	1.0
Levofloxacin[9]	18 (13%)	25 (13%)	0.99 (0.49-1.99)	1.0	1.57 (0.63-3.85)	0.29	0.61 (0.13-2.37)	0.56

Linezolid[9]	23 (17%)	22 (11%)	1.53 (0.77-3.03)	0.20	1.47 (0.54-3.83)	0.49	1.1 (0.29-3.91)	1.0
Meropenem[9]	39 (28%)	50 (26%)	1.11 (0.66-1.86)	0.71	1.13 (0.54-2.31)	0.73	0.88 (0.32-2.33)	0.83
Polymyxin[9]	8 (6%)	4 (2%)	2.86 (0.75-13.26)	0.13	0.98 (0.09-7.03)	1.0	Inf (0.27-Inf)	0.16
Tigecycline[9]	25 (18%)	17 (9%)	2.25 (1.11-4.66)	0.019	2.63 (1.04-6.75)	0.026	2.55 (0.68-10.63)	0.14
Tobramycin[9]	7 (5%)	10 (5%)	0.97 (0.3-2.89)	1.0	0.47 (0.05-2.47)	0.50	2.23 (0.25-27.73)	0.40
Piperacillin/tazobactam[9]	18 (13%)	23 (12%)	1.09 (0.53-2.22)	0.87	0.9 (0.29-2.52)	1.0	1.53 (0.46-5.09)	0.43
Intravenous vancomycin[9]	52 (37%)	62 (32%)	1.25 (0.77-2.03)	0.35	1.07 (0.54-2.09)	0.87	1.03 (0.4-2.6)	1.0
Aminoglycoside[9]	32 (23%)	27 (14%)	1.82 (1-3.36)	0.042	1.14 (0.47-2.66)	0.84	2.75 (0.83-9.98)	0.099
Polymyxin or colistin[9]	20 (14%)	15 (8%)	1.98 (0.92-4.34)	0.070	1.92 (0.68-5.33)	0.22	1.09 (0.15-6.78)	1.0
Third-generation cephalosporin[9]	14 (10%)	15 (8%)	1.32 (0.57-3.05)	0.56	1.34 (0.45-3.82)	0.62	2 (0.32-14.33)	0.44
Anti-pseudomonal antibiotic[9,10]	88 (63%)	115 (60%)	1.15 (0.72-1.86)	0.57	1.24 (0.63-2.49)	0.52	0.76 (0.34-1.69)	0.58

Fluoroquinolone[9]	20 (14%)	31 (16%)	0.87 (0.45-1.67)	0.76	1.62 (0.67-3.86)	0.29	0.36 (0.08-1.27)	0.12
Carbapenem[9]	59 (42%)	68 (35%)	1.34 (0.84-2.16)	0.21	1.26 (0.64-2.45)	0.52	0.98 (0.41-2.31)	1.0
Piperacillin/tazobactam or ceftazidime[9]	20 (14%)	29 (15%)	0.94 (0.48-1.82)	0.88	0.86 (0.32-2.14)	0.83	1.34 (0.41-4.28)	0.60

[1]LTACH=long-term acute care hospital
[2]IQR=interquartile range
[3]Anoxic brain injury or cerebrovascular accident (CVA) causing paresis

[4]BMI ≥30 kg/m^2

[5]BMI ≤18.5 kg/m^2

[6]Solid organ transplant or hematopoeitic stem cell transplant
[7]Stage IV or dialysis
[8]Chronic obstructive pulmonary disease (COPD) or chronic bronchitis

[9]Receipt for ≥48 hours in the 30 days prior to culture

[10]Aminoglycosides, cefepime, ceftazidime, fluoroquinolones, polymyxins, carbapenems, or piperacillin-tazobactam

Figure S1: Methods overview. (A) Isolate subsets, based on anatomic site, used for different machine

learning analyses. (B) Outcome used for all machine learning analyses. (C) Feature sets used for

machine learning analyses (see main methods for more details). (D) Machine learning analysis pipeline:

(D1) Machine learning pipeline, (D2) Median AUROC calculated from machine learning pipeline, (D3)

Mean difference in test vs. permuted AUROC for each feature (see methods for more details).

AUROC=area under the receiver operating characteristic curve.

Figure S2: There are differences in sample size and distribution of infection and anatomic site of

samples across LTACHs. Distribution of colonization and infection isolates are split by source and

LTACH of origin. A to U are specific LTACHs and the geographic location by state is in parentheses.

LTACH=long-term acute care hospital.

Figure S3: Test AUPRCs for various classifiers identifying CRKP colonization vs. infection for the overall machine learning model and respiratory and urinary models. Test AUPRCs for 100 train/test splits used to build classifiers with L2 regularized logistic regression. (A) Overall model with various classifiers. (B) Respiratory and urinary models; the curated genomic feature set was used here. AUPRC=area under the precision recall curve.

857 **Figure S4: Test AUROC of respiratory and urinary ST258 only machine learning models for**

858 **various feature sets.** The curated genomic feature set was used here. AUROC=area under the receiver

859 operating characteristic curve.

Figure S5: Differences in test vs. mean permuted AUROC across 100 data splits for each feature in

the overall machine learning analysis. All features included in the model are shown. We consider

features to be associated with infection/colonization if the AUROC difference is greater than zero in over

75% of the 100 data splits (see methods for more details). Features above the dotted line fall into this

category. The solid grey line indicates a difference of zero (i.e. the feature provides no improvement to

model performance). Bla=Beta lactamase, res=confers resistance to that antibiotic class.

Figure S6: Differences in test vs. mean permuted AUROC across 100 data splits for each feature in

the respiratory machine learning analysis. All features included in the model are shown. We consider

features to be associated with infection/colonization if the AUROC difference is greater than zero in over

75% of the 100 data splits (see methods for more details). Features above the dotted line fall into this

category. The solid grey line indicates a difference of zero (i.e. the feature provides no improvement to

model performance). Bla=Beta lactamase, res=confers resistance to that antibiotic class.

Figure S7: Differences in test vs. mean permuted AUROC across 100 data splits for each feature in

the urinary machine learning analysis. All features included in the model are shown. We consider

features to be associated with infection/colonization if the AUROC difference is greater than zero in over

75% of the 100 data splits (see methods for more details). Features above the dotted line fall into this

category. The solid grey line indicates a difference of zero (i.e. the feature provides no improvement to

model performance). Bla=Beta lactamase, res=confers resistance to that antibiotic class.

Figure S8: Presence/absence of genomic variants on the ICEKp10 element. The heatmap beside the

ST258 phylogeny indicates variant presence/absence in a given isolate. Color indicates presence and

white indicates absence. The scale bar above the phylogeny shows the branch length in substitutions per

site.

Figure S9: kfoC contains an IS element in a subset of ST258 isolates. The heatmap beside the

ST258 phylogeny indicates information about the isolate (left to right): the O locus type identified by

Kleborate (i.e. the curated genomic data), if Kleborate identified kfoC as missing, blast results for kfoC

against the genome assemblies (split between 2 assemblies or entirely absent; white indicates that kfoC

was intact), and IS elements identified in kfoC (white indicates no IS element detected). The scale bar to

the top left of the phylogeny shows the branch length in substitutions per site.

902 **Figure S10: Isolates from different LTACHs are distributed across the phylogeny.** Phylogeny of

903 isolates indicating the LTACH the patient was in when the isolate was obtained. Isolates from several

904 LTACHs are present in the disrupted kfoC lineage or the ybt0 lineage, and isolates from these LTACHs

905 are not confined to those lineages. The scale bar above the phylogeny shows the branch length in

906 substitutions per site.

Figure S11: Isolates containing both disrupted kfoC and ICEKp10 are mainly confined to one

ST258 clade. Phylogeny of California LTACH isolates (US-CA) in the context of public isolates from

various locations (EUR=Europe, SA=South America, US-MW=US Midwest, US-NE=US Northeast, US-

S=US South, US-W=US West). Isolate location, as well as ybt0 presence and kfoC absence are

displayed for each sample. The scale bar to the right of the phylogeny shows the branch length in

substitutions per site.