Two Separate, Large Cohorts Reveal Modifiers of Age-Related Change in Visual Reaction Time

Performance

J.S. Talboom[1,2], M.D. De Both[1,2], M.A. Naymik[1,2], A.M. Schmidt[1,2], C.R. Lewis[1,2], W.M. Jepsen[1,2],

A.K. Håberg[3], T. Rundek[4], B.E. Levin[4], S. Hoscheidt[2,5], Y. Bolla[2,5], R.D. Brinton[2,5], N.J. Schork[1,6], M.

Hay[2,5], C.A. Barnes[2,5], E. Glisky[2,5], L. Ryan[2,5], and M.J. Huentelman[1,2,]

[1]The Translational Genomics Research Institute (TGen), Phoenix, Arizona 85004 USA

[2]Arizona Alzheimer's Consortium, Phoenix, Arizona 85004 USA

[3]Norwegian University of Science and Technology, Trondheim 7491 Norway

[4]University of Miami Miller School of Medicine and Evelyn F. McKnight Brain Institute,

Miami, FL 33136 USA

[5]University of Arizona, Tucson, Arizona 85721 USA

[6]City of Hope National Medical Center, Duarte, CA 91010 USA

15   **Abstract**

To help identify factors influencing age-related cognitive decline and disease, a web-based

simple visual reaction time (RT and svRT respectively, index of processing speed) and paired-associate

learning (PAL, indexing of verbal episodic memory) task were implemented. Test results were

combined with 22 survey questions. An analysis of RT and svTR data from the MindCrowd ($n=75,666$)

cohort revealed education and reported stroke as potential modifiers of age-associated processing speed

and memory decline. As a complement, we evaluated complex, recognition, reaction time (cvrRT) in the

UK Biobank cohort and compared the results to a subset of the MindCrowd cohort, creating UKBb

MindCrowd ($n=39,795$), mirroring the UK Biobank's ($n=158,247$) age range (40-70y). Many estimates

from an identical linear model, used for both cohorts, were similar despite noted cohort differences (e.g.,

USA vs. the UK). Results from the UK Biobank that differed from UKBb MindCrowd included a first-

degree family history Alzheimer's disease (FHAD) associated with slower cvrRTs and that sex and

education affected age-related cvrRT slowing. While results from UKBb MindCrowd suggested that

education and smoking status were related to opposite effects on age-related svRT slowing.

MindCrowd's collected RT and PAL findings have begun to pull back the curtain on the intricate

network connecting processing speed, memory, and cognition to healthy and pathological brain aging.

Reaction time (RT), an index of processing speed or efficiency in the central nervous system

(CNS) [1], is an essential factor in higher cognitive function[2,3] and is profoundly affected by age [4]. In fact,

of the studied demographics, age is the main factor known to influence RT[4]. Processing speed is an

important limiting factor for most aspects of cognition during aging, most notably memory[5,6]. In studies

where processing speed was used as a covariate, the age-related variance in a variety of episodic

memory measures was substantially reduced or even eliminated[7,8]. Moreover, studies comparing

different factors and tests of age-related episodic memory deficit, implicate age-related decline in

processing speed as the main mediator[9-11]. Collectively, these findings suggest that RT is an excellent

index of age-related cognitive decline as well as healthy brain aging.

RT can be operationally defined as "simple," which typically involves a non-choice reaction to a

visual stimulus (svRT). RT can also be operationally defined as "complex," which involves a reaction to

one or more visual stimuli after recognition (cvrRT) of correct stimuli, and inhibition to incorrect stimuli

[12]. svRT demonstrates variability between individuals, which akin to PAL, is thought to be influenced by

both genetic and environmental factors[13]. In addition, svRT effects are well noted across the field of

neurology; for example, AD and stroke patients demonstrate slower svRT and higher inter-individual

variability [14,15]. However, due to the limitations of traditional research methods, the body of work

concerning RT mainly examined only limited ranges of demographic, health, medical, and lifestyle

factors in small cohorts. For example, the demographics of prior work consisted mainly of college-aged

students, well-educated older adults[16-18], or athletes [19-21].

Further, with notable exceptions[22-24], many studies had few participants (e.g., $n < 1000$) and were

therefore powered to detect only variables with large effect size and to lead to spurious non-replicable

findings[25]. Consequently, many studies of RT had minimal ability to reveal low frequency factors or

those with subtle effect sizes as well as to conduct more sophisticated analyses (e.g., ANOVA vs.

Growth Modeling) to identify interactions and moderators. Collectively, this suggests that if RT

performance is to be used to inform models of disease or normative and atypical aging, we need a

deeper understanding of the normal variation of RT and the genetic and environmental factors associated

with RT performance.

The goal of this study was to characterize RT across a broad range of demographic, health,

medical, and lifestyle variables commonly associated with cognitive performance and AD risk. To

accomplish this, we utilized both the MindCrowd and UK Biobank cohorts[26,27], comprising over 233

thousand combined participants, to model RT as a function of 11 or more demographic, health, medical,

and lifestyle factors. Based on our prior work[28] and previous RT research, we hypothesized that RT, via

its structure of factor association and modifications, would reveal meaningful connections to healthy

brain aging.

66   **Materials and Methods**

Study Participants

MindCrowd: Overview

In January 2013, we launched our internet-based study at www.mindcrowd.org. Website visitors

18 years or older were asked to consent to our study before any data collection via an electronic consent

form. As of 3-26-2020, we have had 356,674 non-duplicate or unique visitors to the website. Of these

unique visitors, over 194,542 (54% of unique visitors) consented to participate. The final data set

contained 75,666 (54% of consented individuals) participants who completed a simple visual reaction

time (svRT) and paired associates learning (PAL) tasks and answered 22 demographic, lifestyle, and

health questions. Approval for this study was obtained from the Western Institutional Review Board

(WIRB study number 1129241).

MindCrowd: simple visual reaction time (svRT)

After consenting to the study and answering five demographic questions (age, biological sex,

years of education, primary language, and country where residing), participants were asked to complete

a web-based svRT task. The svRT task taps primarily into the nervous system and is influenced by

intelligence and brain injury[29]. Participants were presented with a pink sphere that appeared at random

intervals (between 1-10s) on the screen, and they were instructed to respond as quickly as possible after

the sphere appeared by pressing the enter/return key on their keyboard. Once the participant responded,

the sphere disappeared until the next trial. Each participant received a total of five trials. The sphere

stayed on the screen until the participant responded. The dependent variable, response time

(milliseconds), was recorded from the appearance of the sphere on the screen to the key/screen press

made by the participant.

MindCrowd: paired-associate learning (PAL)

Next, participants were presented the PAL task. For this cognitive task, during the learning

phase, participants were shown 12 word-pairs, one word-pair at a time (2s/word-pair). During the recall

phase, participants were given the first word of each pair and were asked to use their keyboard to type in

(i.e., recall) the missing word. This learning-recall procedure was repeated for two additional trials.

Before beginning the task, each participant received one practice trial consisting of three word-pairs not

contained in the 12 used during the test. Word-pairs were presented in different random orders during

each learning and each recall phase. The same word pairs and order of presentation were used for all

participants. The dependent variable/criterion was the total number of correct word pairs entered across

the three trials (i.e., 12x3=36, a perfect score).

MindCrowd: demographic, medical, health, and lifestyle questions

Upon completing the PAL task, participants were asked to fill out an additional 17 demographic

and health/disease risk factor questions. These questions included: marital status, handedness, race,

ethnicity, number of daily prescription medications, a first-degree family history of dementia, and yes/no

responses to the following: seizures, dizzy spells, loss of consciousness (more than 10 minutes), high

blood pressure, smoking status, diabetes mellitus, heart disease, cancer, reported stroke, alcohol/drug

abuse, brain disease, and memory problems). Next, participants were shown their results and provided

with different comparisons to other test takers based on the average scores across all participants, as well

as across sex, age, and education. On this same page of the site, participants were given the option to be

recontacted for future research (see Supplementary Table 2 for the list of MindCrowd questions asked).

UK Biobank: study design and aims

The UK Biobank (research ethics committee approval [11/NW/0382]) is a long-term study and

research resource in the United Kingdom (UK), which investigates genetic and environmental exposure

to the development of the disease. The stated goal of the UK Biobank is to, "build a major resource that

can support a diverse range of research intended to improve the prevention, diagnosis, and treatment of

illness and the promotion of health throughout society." The UK Biobank began in 2006. The study is

currently following about 500,000 participants in the UK, enrolled at ages 40 to 69. Initial enrollment

took place from 2006 to 2010. All participants are being monitored for at least 30 years after recruitment

and initial assessment (i.e., termed "instance 0" by the Biobank). Potential participants were invited to

visit an assessment center, where they completed a questionnaire. Participants were next interviewed

about lifestyle, medical history, and nutritional habits. Lastly, vital measurements, such as weight,

height, and blood pressure, were measured. One aim of the UK Biobank is to electronically record all

health-related changes and events across the entire 30-year study. Notably, this task is aided by the UK's

integrated health system and corresponding electronic health record-keeping, an approach that is not

possible in the United States.

UK Biobank: data procurement

All UK Biobank data were derived from Application #43036, entitled "Exploring and

Accommodating Heterogeneity in Large-Scale Genetic Analyses" as a "Collaborator Project,"

UK Biobank: Complex visual recognition reaction time (cvrRT) and educational attainment.

Each participant's cvrRT was based on 12 rounds of the card-game 'Snap.' Participants were

shown two cards at a time with a picture on them. Participants pressed a button on a table in front of

them as quickly as possible if the images cards/matched. For each of the 12 rounds, the following data

were collected: the pictures shown on the cards (Index of card A, Index of card B), the number of times

the participant clicked the 'snap' button, and the latency to first click of the 'snap' button. This last

record of "latency to click the button" was used as the UK Biobank's criterion for regression analyses.

For Educational Attainment, the following conversions from UK Biobank (UKBb) answer codes

(see http://biobank.ndph.ox.ac.uk/showcase/coding.cgi?id=100305) to MindCrowd (MC) values were

made: a) "UKBb **-7** None of the above" to "MC No high school diploma," b) "UKBb **2** A levels/AS

levels or equivalent" to "MC High school diploma," c) "UKBb **3** O levels/GCSEs or equivalent" to "MC

High school diploma," d) "UKBb **4** CSEs or equivalent" to "MC High school diploma," e) "UKBb **5**

NVQ or HND or HNC or equivalent" to "MC Some college," f) "UKBb **6** Other professional

qualifications (e.g., nursing and teaching)" to" MC Some college," g) "UKBb **1** College or University

degree" to "MC College degree." All UKBb participants selecting "**-3** Prefer not to answer" were

removed from the final dataset before model selection and analysis. While we did our best to ensure a

similar measure of education across UKBb MindCrowd and the UK Biobank, we realize that there are

fundamental differences between US and UK schools that we cannot control or eliminate (see Table 6).

Data Quality Control

For the MindCrowd analysis, a final data set, including all qualifying participants up to 3-17-

2020, was generated. See Supplementary Figure 9, for a flowchart detailing the following filtering steps.

This dataset removed participants: a) with duplicate email addresses (only first entry kept), b) who did

not complete all three rounds of the PAL test, c) whose primary language was not English, d) who was

not between 18-85 years old, e) whose RT trials were above or below 1.5 x the interquartile range (IRQ)

and f) whose median svRT was above or below 1.5 x the IRQ range of all participants of the same age

(see Supplementary Figure 10). Participants from either study were removed if they were missing any

data (listwise deletion). Lastly, for the UKBb MindCrowd and UK Biobank analysis, participants were

removed if any of their responses to a demographic, medical, health, and lifestyle question did not match

the other study. For example, participants in the UK Biobank who responded to the "Race" question

with "Prefer Not to Answer" were removed. "Prefer Not to Answer" was not a choice MindCrowd

participants were given on the "Race" question. Removing these participants was done to align UKBb

MindCrowd and UK Biobank cohorts as much as possible.

Statistical Methods

Statistical analyses were conducted using R within the RStudio shell. For all analyses,

multivariate linear regression was performed using the general linear model (LM) to model Median

　　8

svRT or Median cvrRT (i.e., criterion or dependent variable) as a function of either 24 (MindCrowd) or

11 predictors (UK Biobank analysis).

For the MindCrowd analysis, svRT was modeled as a function of PAL Performance and Age

raised to the power of three (i.e., to fit and estimate nonlinear associations). Most figures and data

manipulation were created using the various R packages bundled together in R package, *"tidyverse"*[30].

Additional figures were generated using the R packages (see Supplemental Table 1 for a complete list of

resources). Continuous by continuous interactions (i.e., simple slopes) were estimated using the R

packages *"interactions," "sandwich," "jtools"*[31]. Categorical by categorical interactions were estimated

using the R package, *"emmeans"* [32]. Adjustments for multiple comparisons were evaluated using

Tukey's method via the *"emmeans"* package.

Model evaluation and controls

All measurements were taken from distinct samples. Model fit and violations of parametric

assumptions were evaluated separately in each model. Here we evaluated different residual plots,

assessing normality, homoscedasticity, outliers, residual autocorrelation, and multicollinearity. The

MindCrowd LM included all 22 demographic questions, health, medical, and lifestyle questions. These

questions were: Age, Biological Sex, Race, Ethnicity, Educational Attainment, Marital Status,

Handedness, Number of Daily Medications, Seizures, Reported Dizziness, Loss of Consciousness,

Reported Hypertension, Smoking Status, Heart Disease, Reported Stroke, Alcohol/Drug Abuse,

Diabetes Mellitus, Cancer, a First-Degree Family History of Alzheimer's disease, history of brain

disease, whether the test was taken on a mobile device, and the version of the MindCrowd site used. Not

surprisingly, the device used to take the RT portion of MindCrowd had an association with RT

performance. Those on a mobile device (i.e., using a touchscreen) had slower svRTs compared to those

who did not (see Supplementary Figure 8). For the UK Biobank analyses, these 11 variables included:

Age, Biological Sex, Diabetes mellitus, Handedness, Reported Stroke, Reported Hypertension, Smoking

Status, Reported Dizziness, Educational Attainment, and a First-Degree Family History of Alzheimer's

Disease.

197   ***Inclusion of polynomials, automated model selection, and figure generation***

198   The MindCrowd analyses included Age and PAL Performance as first through third-order non-

orthogonal polynomials (i.e., cubic regression). This choice was based on empirical evaluations of

models differing only by order of their Age and PAL Performance polynomial. Bayesian information

criterion (BIC) weights were generated from separate models that include the first through sixth-order

polynomial. Pairwise comparison of BIC weights revealed that the model, containing Age and PAL

Performance as third-order polynomials, was 8.60E-301 times more likely compared to the non-

polynomial model. Similarly, another study of complex and simple RT also included age as a third-order

polynomial and found remarkably similar results[33]. Notably, this study was relatively large (i.e., 7,000

participants), included both sexes, and had ages ranging from 18-94. Using "glmulti" allowed us to

avoid the pitfalls of stepwise methods or unintentional biased manual selection. Like the prior

MindCrowd model, here we used "glmulti" to define the "best" (i.e., lowest BIC) UK Biobank model,

which we also used to evaluate our MindCrowd cohort; thus, permitting us to compare the two cohorts

directly. Two rounds of model selection were run using glmulti to select the optimal MindCrowd model.

For round 1, the optimal model contained only the main effects, due to software limitations when all 22

factors were included. So, in round 2, only factors selected in the optimal main effects model were then

included to select an optimal model, including two-way interactions.

Results

MindCrowd

As of March 13th, 2020, after filtering (see Data Quality Control in Materials and Methods),

MindCrowd, had recruited 75,666 qualified participants (see Table 1 for Sociodemographic

Characteristics and Supplementary Figure 1A- for a histogram of age). We modeled svRT as a function

of Age^3 and PAL $Performance^3$ (i.e., curvilinear associations) as well as 20 other factors (see

Supplementary Figure 2 figure for diagnostic regression plots, and Table 2 for all coefficient ns). The

omnibus model was significant ($F_{omnibus}[58, 73406]=858.20$, $p_{omnibus}=<2.2e-16$, Adjusted $R^2=0.40$).

Age and paired-associate learning (PAL) total correct

The model revealed that two Age polynomials were significantly associated with simple visual

reaction time (svRT) performance, Age ($\beta_{Age1}=7.07$, $p_{Age1}=3.06E-18$), and Age^2 ($\beta_{Age2}=-0.15$,

$p_{Age2}=3.23E-17$). On average, per year of life, svRT was seven milliseconds slower, and the seven

milliseconds slope lowered by 0.15 milliseconds per year (Figure 1A). Like Age, the PAL $Performance^1$

and PAL $Performance^2$ polynomials were significantly associated with svRT performance [PAL

$Performance^1$ ($\beta_{PAL1}=-8.89$, $p_{PAL1}=3.77E-34$), PAL $Performance^2$ ($\beta_{PAL2}=0.32$, $p_{PALl2}=2.35E-14$). These

coefficients suggest that, per word pair correct, svRT was associated with nearly nine milliseconds faster

performance, with the nine milliseconds raised by 0.32 milliseconds per word pair correct (Figure 1B).

glmulti-selected third-order polynomials of Age^3 ($\beta_{Age3}=1.47E-03$, $p_{Age3}=1.46E-34$) and PAL

$Performance^3$ ($\beta_{PAL3}=-4.13E-03$, $p_{PAL3}=7.28E-09$) illustrate the small, but significant, additional bend in

the quadratic shape of each factor's association with svRT (Figure 1A and 1B).

Biological sex, educational attainment, and handedness

Biological Sex was a significant predictor of svRT (β_{Sex}=-34.26, p_{Sex}=1.26E-163). Men were

associated with an average of 34 milliseconds (9.63%) faster svRT response, as compared to women

(Figure 1D). Educational Attainment was also a significant factor associated with svRT. Compared to

"No High School Diploma", participants who had "Some College" ($\beta_{College}$=-14.73, $p_{College}$=8.74E-05),

or a "College Degree" ($\beta_{CDegree}$=-31.78, $p_{CDegree}$=2.95E-17, Figure 2A) were faster. Attending college

and attaining a college degree was associated with a respective near 15 (4.14%) and 32 (8.92%)

millisecond faster svRT compared to not graduating from high school. Handedness was also associated

with svRT. Left-handed participants had a near four milliseconds (1.09%) faster svRT (β_{Left}=-3.87,

p_{Left}=0.03, Figure 2B). This association was present in individuals 20 to 40 years old (β_{40Left}=-3.16,

p_{40Left}<0.01, Figure 2C) but not in individuals 40 to 60 years old (β_{60Left}=-2.69, p_{60Left}=0.07, Figure 2D).

Health, medical, and lifestyle factors

For health and medical factors associated with svRT, we found that Smoking Status

($\beta_{Smoking}$=7.07, $p_{Smoking}$=1.26E-03, Figure 3A) and Reported Dizziness (β_{Dizzy}=4.87, p_{Dizzy}=0.04, Figure

3B) were both significant predictors of svRT. Smoking Status was associated with seven milliseconds

(1.99%) slower svRT, and Reported Dizziness was associated with nearly five milliseconds (1.37%)

slower svRT. When compared to participants reporting "no daily medications," taking "Two"

(β_{Meds2}=5.84, p_{Meds2}=2.00E-03), "Three" (β_{Meds3}=6.24, p_{Meds3}<0.01), and "Four" (β_{Meds4}=17.82,

p_{Meds4}=3.51E-16) Daily Medications were associated with an approximately 5 (1.64%), 6 (1.76%), and

18 (5.01%) milliseconds slower svRT, respectively (Figure 4A). Further, Diabetes Mellitus

($\beta_{Diabetes}$=11.23, $p_{Diabetes}$<3.36E-05, Figure 4B), and Stroke (β_{Stroke}=20.38, p_{Stroke}=3.59E-04, Figure 4C)

were related to 11 (3.16%) and 20 (5.73%) milliseconds slower svRT respectively when compared to

participants not reporting either condition. Of note, in this model, both a first-degree family history of

Alzheimer's disease (FHAD; β_{FHAD}=-0.40, p_{FHAD}=0.78) and Hypertension (β_{Hyper}=1.08, p_{Hyper}=0.52)

were not significant predictors of svRT performance (Supplementary Figure 3A-B).

Two-way interactions

For interactions, we found Age significantly interacted with PAL Performance. Age x PAL

Performance ($\beta_{Age*PAL}$=-0.07, $p_{Age*PAL}$=9.93E-62). Analysis of simple slopes suggests that each word-pair

correct was associated with increasingly faster svRT as participant age increased. That is, at 20

($\beta_{Age20*PAL}$=-5.48, $p_{Age20*PAL}$<0.00), 40 ($\beta_{Age40*PAL}$=-6.41, $p_{Age40*PAL}$<0.00), 60 ($\beta_{Age60*PAL}$=-7.75,

$p_{Age60*PAL}$<0.00), and 80 ($\beta_{Age80*PAL}$=-9.08, $p_{Age80*PAL}$<0.00, Figure 1C) years old. There was a significant

Biological Sex x Age interaction ($\beta_{Age*Sex}$=-0.36, $p_{Age*Sex}$=4.61E-08) indicating that the age-related

slowing of svRT in men, compared to women, was 0.36 milliseconds (0.08%) less per one year of age

(Figure 1C). These data suggest that men's svRT performance declined at a slower rate when compared

to women. Of interest, in both women and men, we found significant Age x Educational Attainment

interactions. Compared to Age x "No High School Diploma", participants reporting having "Some

College" ($\beta_{Age*College}$=-0.65, $p_{Age*College}$=4.20E-04) or a College Degree" ($\beta_{Age*CDegree}$=-1.31,

$p_{Age*CDegre}$=2.07E-12) were associated with slower RTs as they aged. These results suggest that attending

college or getting a college degree was associated with a 0.65 (0.15%) and 1.31 (0.30%) millisecond

faster svRT performance per year of life, respectively (Figure 2A). The MindCrowd model revealed a

significant Age x Reported Stroke interaction ($\beta_{Age*Stroke}$=1.97, $p_{Age*Stroke}$=2.72E-06). Participants who

Reported Stroke were about 2 milliseconds (0.37%) slower on svRT per year of life compared to those

without Reported Stroke (Figure 4C). Lastly, we found a significant Age x Smoking Status interaction

($\beta_{Age*Smoke}$=0.62, $p_{Age*Smoke}$=5.67E-07). This interaction suggests that Smoking Status slows svRT by an

additional 0.57 milliseconds (0.11%) per year of age (Figure 3A). See Table 3 for a complete list of the

significant predictors, covariates, and interactions from the MindCrowd analysis.

UKBb MindCrowd and UK Biobank

Of the total 75,666 MindCrowd participants, 39,759 between the ages of 40 and 70 were

selected, to mirror the UK Biobank. This subset is called UKBb MindCrowd from here on to

differentiate it from MindCrowd. The UK Biobank cohort, after filtering (see Data Quality Control),

contained 158,247 participants, derived from a data-request we received on 9-19-2019 (See Table 1 and

Supplementary Figure 1C-figure for age histogram). We model both the UKBb MindCrowd's svRT (see

Supplementary Figure 4 for regression diagnostic plots) as well UK Biobank's cvrRT (see

Supplementary Figure 5 for regression diagnostic plots) as a function of 11 shared survey questions (see

Table 4 for the ns used to estimate each coefficient). The omnibus UKBb MindCrowd (F_{mcomni}[20,

38871]=1039, p_{mcomni}=<2.2e-16, Adjusted R^2 =0.08) and UK Biobank ($F_{ukbbbomni}$[20, 157903]=1038,

$p_{ukbbomni}$=<2.2e-16, Adjusted R^2 =0.13) LMs were both significant.

Age and biological sex

The UKBb MindCrowd cohort revealed Age as a significant predictor of svRT

(β_{Age}=5.75, p_{Age}=2.00E-16). The parallel analysis (see Statistical Methods) of Age in the UK Biobank

cohort was also significant (β_{Age}=3.40, p_{Age}=2.00E-16) for complex, visual, recognition reaction time

(cvrRT). For the association of Age and RT, UKBb MindCrowd and the UK Biobank both showed

increased RTs or worse RT performance with age, with nearly 6 and 3 milliseconds slower RT per year

of age, respectively (Figure 5A-B). For UKBb MindCrowd, Biological Sex was a significant predictor

of RT (β_{Sex}=-40.00, p_{Sex}=8.03E-71), which was also the case in the UK Biobank (β_{Sex}=-18.28,

p_{Sex}=2.00E-16). Here the effect of Biological Sex on RT between UKBb MindCrowd was 40

milliseconds (20.46%), and the UK Biobank was 18 milliseconds (5.16%). However, in both cohorts,

men had faster RT performance compared to women (Figure 6A-B).

Education and handedness

Akin to the association found in the MindCrowd analysis, the UKBb MindCrowd and UK Biobank

comparison found that Educational Attainment was a significant predictor of RT. Indeed, for UKBb

MindCrowd "Some College" ($\beta_{College}$=-26.08, $p_{College}$=1.61E-03, 13.34%) and a "College Degree"

($\beta_{CDegree}$=-49.07, $p_{CDegree}$=1.91E-09, 25.01%), as well as in the UK Biobank a "High School Diploma"

($\beta_{HSDiploma}$=-9.68, $p_{HSDiploma}$=1.23E-33, 2.73%), "Some College" ($\beta_{College}$=-10.26, $p_{College}$=1.30E-40,

2.90%), or a "College Degree" ($\beta_{CDegree}$=-11.71, $p_{CDegree}$=1.46E-41, 3.31%) were all significantly

different from "No High School Diploma" (Figure 7A-B). Here, both UKBb MindCrowd and UK

Biobank large cohorts reported faster RT was associated with more education. Lastly, unlike the

MindCrowd analyses, both the UKBb MindCrowd $\beta_{Handedness}$=-2.58, $p_{Handedness}$=4.00E-01), and the UK

Biobank ($\beta_{Handedness}$=0.58, $p_{Handedness}$=0.36) cohorts between the ages of 40-70, did not find Handedness

to be a significant predictor of RT performance (Supplementary Figure 6).

Health, medical, and lifestyle factors

In terms of health factors associated with RT, in the UKBb MindCrowd cohort, Diabetes

($\beta_{Diabetes}$=11.48, $p_{Diabetes}$=3.31E-03,   5.87%),   Stroke   (β_{Stroke}=18.47, p_{Stroke}=4.00E-02,   9.45%),

($\beta_{Hypertension}$=7.99, $p_{Hypertension}$=3.16E-03, 3.58%), and Dizziness (β_{Dizzy}=12.13, p_{Dizzy}=2.52E-03, 6.19%)

were all significantly associated with lower svRT performance (i.e., slower svRT). These associations

were recapitulated by the UK Biobank. To that end, Diabetes ($\beta_{Diabetes}$=5.48, $p_{Diabetes}$=4.80E-07, 1.55%,

Figure 8A-B), Stroke (β_{Stroke}=10.61, p_{Stroke}=6.15E-07, 2.99%, Figure 9A-B), Hypertension

($\beta_{Hypertension}$=1.14, $p_{Hypertension}$=0.02, 0.31%, Figure 10A-B), and Dizziness (β_{Dizzy}=3.21, p_{Dizzy}=3.71E-14,

0.91%, Figure 11A-B) were all significantly related to lower cvrRT performance (i.e., slower cvrRT);

however, the association between hypertension and cvrRT was small (Figure 10B). Further, in

agreement with the MindCrowd analysis, Smoking Status was significantly related to svRT

(β_{Smoke}=10.18, p_{Smoke}=0.01, 5.21%) in the UKBb MindCrowd cohort; however, FHAD (β_{FHAD}=-0.07,

p_{FHAD}=0.97) was not. This pattern of associations were reversed in the UK Biobank; that is, Smoking

Status was not a significant predictor of cvrRT (β_{Smoke}=0.47, p_{Smoke}=0.22, Figure 12A-B), but FHAD

(β_{FHAD}=2.36, p_{FHAD}=3.35E-05, 0.69%, Figure 13A-B) was.

Two-way interactions

The "glmulti" R package defined two interactions in the UKBb MindCrowd and UK Biobank

analysis. In UKBb MindCrowd we found a significant Age x Biological Sex interaction ($\beta_{Age*Sex}$=-0.63,

$p_{Age*Sex}$=2.00E-02, 0.33%). We found a comparable significant Age x Biological Sex interaction for

cvrRT ($\beta_{Age*Sex}$=-0.55, $p_{Age*Sex}$=4.18E-28, 0.16%) in the UK Biobank. Across both MindCrowd and the

UK Biobank, these interactions indicated that the trajectory of age-related RT slowing in men, when

compared to women, was over 0.5 ms less per one year of age (Figure 6A and B). In addition, the UK

Biobank analysis revealed a significant Biological Sex x Educational Attainment interaction not found

in UKBb MindCrowd. Here, men with a "High School Diploma" ($\beta_{Sex*HSDiploma}$=-11.24,

$p_{Sex*HSDiploma}$=3.30E-12, 3.35%), "Some College" ($\beta_{sex*College}$=-10.54, $p_{sex*College}$=8.71E-12, 3.14%) or a

"College Degree" ($\beta_{sex*CDegree}$=-12.8, $p_{sex*CDegre}$=1.68E-13, 3.18%) were significantly associated with

faster cvrRT performance when compared to women having "No High School Diploma" (Figure 14).

Follow-up analyses of the simple effects via estimated marginal means (EMM, see Statistical Methods)

revealed that men who did not graduate high school (*EMM*=558.91 ms), compared to men with more

education (*EMMs*=542.99, 542.98, and 540.41 ms), had markedly faster cvrRTs, more in line with the

women's cvrRT performance (*EMMs*=566.80, 562.11, 561.40, and 561.09 ms). The associated

difference in cvrRT for men with "No High School Diploma" compared to men with a "High School

Diploma" (β_{Men}=15.92, p_{Men}=2.00E-16, 1.75%), was more than the difference in cvrRT between women

with "No High School Diploma" compared to women with a "High School Diploma" (β_{Women}=4.68,

p_{Women}=3.12E-04, 1.69%, see Supplementary Figure 7 for a graph of the EMM).

348   **Discussion**

Age-Related Slowing of Reaction Time (RT): Factors and Modifiers

The results of our studies illuminate a portion of the intricate relationship between age and RT

performance by identifying demographic, health, medical, and lifestyle factors associated with either

attenuation or exacerbation of age-related slowing of RT (see figure 15 for a summary of the results). A

large body of work on RT, leading back to Francis Gaulton in 1890[34], has consistently demonstrated an

age-related change in RT[35]. As such, it is not surprising that both MindCrowd and UKBb MindCrowd,

as well as UK Biobank models, revealed age-associated slowing of both in simple visual (svRT,

MindCrowd) and complex, visual recognition RT (cvrRT, UK Biobank). Likewise, the MindCrowd

model found that the relationship between svRT and age was modestly curvilinear (Figure 1A). While

this curvilinear relation between RT and age has been noted earlier[36], the large sample size of both

cohorts combined with our application of an AI-based model definition[37], revealed a notable addition to

this picture. Specifically, in the MindCrowd analysis, we observed an interaction between age and both

education (Figure 2A) and smoking (Figure 3A). Here, less education and smoking were related to the

additional slowing of the already age-related slowing of simple visual RT (svRT). Lastly, the age by

reported stroke interaction modeled in MindCrowd was associated with more slowing of RT

performance (Figure 4C). This study's large participant numbers, as well as broad age and surveyed data

range, lends it to be one of the most substantial evaluations of RT across the aging spectrum.

Importantly, our findings suggest that smoking and stroke (i.e., perhaps cardiovascular health) and

amount of education (i.e., maybe cognitive demand or reserve) as factors, modified across age, affect

age-related RT slowing.

A First-Degree Family History of Alzheimer's Disease (FHAD), Cognition, and RT: Intertwined

systems

372   In the UK Biobank cohort[27], we found an association between having an FHAD and slower (2.43

milliseconds) cvrRT (Figure 13B). This effect of FHAD on RT or more so underlying process speed is

in line with our episodic verbal memory task (i.e., paired-associates learning [PAL]) finding [28], where

we found FHAD was linked to lower PAL performance. Furthermore, a prior functional magnetic

resonance imaging (fMRI) study, examining medial-temporal lobe activation using a cvrRT task, found

a ~100 millisecond RT increase in 68 (mean age of 54) FHAD participants[38]. These data suggest that

some genetic and environmental factors relating to AD risk are present in individuals with an FHAD.

Indeed, the first-degree relatives of identified FHAD participants either consisted of early-onset familial

AD cases, or late-onset AD, which also has a high heritability 79%, suggesting that these individuals are

in a higher risk category for developing AD. Thus, such shared AD and FHAD factors may relate to

sensorimotor function and processing speed (i.e., RT) analogous to changes in cognition and memory

(i.e., PAL).

384   We found a correlation between svRT and PAL performance (Figure 1B). This finding was in

line with many prior studies, the dependence of episodic memory on processing speed, a dependence

that grows with age and incident of age-related disease (e.g., AD)[5-8,39]. The association of svRT with

PAL may highlight distributed systems and networks that underlie RT performance. For example, svRT

performance could depend on the functioning of many cognitive areas that PAL requires and vice versa.

Another possibility is that properly functioning memory and cognitive networks, correspond to better

RT performance. Evidence for this is suggested by the fact that higher intelligence is related to faster

RT[40-42]. However, diverse factors (e.g., exercise, time of day, meal proximity, and individual assessing

RT) are known to affect RT performance[39,42,43]; thus, making accurate estimates of the effect of RT on

intelligence difficult to ascertain fully. Together with our prior results, these findings suggest that RT

performance and variability indexing both sensorimotor and cognitive domains may be indicative of AD

well before the average age of AD diagnosis. However, further study, replication, and corroboration of

RT's link to age-related cognitive decline and disease are necessary to support this notion.

The difference in the effects of FHAD between UK Biobank and MindCrowd (for both primary

and UKBb MindCrowd analyses, Figures 3A & 13A-B), could be caused by the vast difference in the

fraction of FHAD participants in the UK Biobank (FHAD=13% of total) compared to UKBb

MindCrowd (FHAD=35% of total). While noting the fact that we target those with an FHAD for

recruitment into MindCrowd, still this substantial disparity could also be due to the accuracy of the UK

Biobank's FHAD. The UK Biobank was calculated from three separate questions (i.e., mother, father,

and siblings AD diagnosis). Adding to this, the United Kingdom uses a massive, detailed, nation-wide

electronic health record system conceivability facilitating respondent health and medical survey

accuracy. Compare this to our single question in MindCrowd, asking aged participants to remember if a

relative had AD. Another possibility is the different RT paradigm used; that is, MindCrowd's test of

svRT compared to the UK Biobank's use of cvrRT. The fact that complex reaction time, requiring

recognition and the choice to "respond or not respond," rather than just stimulus-response, may underly

this difference. UKBb MindCrowd svRT performance was consistently faster across age compared to

UK Biobank cvrRT performance, an observation noted in a prior study also measuring both simple and

complex RT[33]. These findings are consistent with the idea that complex RT requires more processing

time[12,44], and prior work found the age-associated slowing of RT was higher for choice RT as compared

to simple RT [45]. Lastly, the differences in FHAD associations across MindCrowd and the UK Biobank

are perhaps due to the UK Biobank had over 2X the number of participants compared to MindCrowd

(i.e., 158K vs. 76K). The larger sample is expected to produce better model definitions and increased

statistical power. Increased statistical power may also have enhanced accuracy, validity, and

reproducibility.

Biological Sex and Educational Attainment

Numerous studies of RT have found sex differences in RT performance, which does not appear

to be reduced by practice[16,20,33]. Consistent with others[33,45], across both cohorts and all three models,

compared to women, men exhibited faster RT performance (Figure 1C & 6). In addition, the analysis of

the UKBb MindCrowd and UK Biobank both implicated biological sex affecting age-associated RT

slowing. This age by biological sex interaction suggests that women are associated with additional age-

related RT slowing of decline compared to men (Figure 1C & 6). The model of the UK Biobank

revealed an interaction between biological sex and education on RT performance. The breakdown

revealed that men had similar RT performance if they attained a high school diploma and above.

However, men who did not graduate high school showed markedly slower RTs, which brought them in

line with women's RT performance. However, the associated slowing of RT for less-educated men was

vastly more than that found in less-educated women (reported "No High School Diploma" made up

3.83% of women and 3.11% of men; see Supplementary Figure 7). These results largely replicate a

previous sizable study (i.e., 7,000 participants) evaluating simple as well as visual as well as of RT[33].

Identical to prior research, this research found that men consistently outperformed women on all

measures of RT across age, age-related decline in RT performance was nonlinear, a third-order

polynomial provided the best model fit, and women exhibited a sharper age-related decline[33]. Collected

with our prior study of PAL performance [28], these findings replicate prior work and suggest that

biological sex effects RT and age-related RT slowing. Besides, age interactions, education was

associated with svRT across both MindCrowd and UKBb MindCrowd models as well as the cvrRT

model for the UK Biobank. Higher education milestones were related to faster svRT and cvrRT, or

overall better RT performance. (Figure 2A & 7). It is unclear whether this is due to individuals

possessing faster RT or processing speed seeking out higher education, or due to better cognitive

maintenance in more FHAD. Further work utilizing both cohorts is necessary to shed light on the effects

and modifiers of FHAD as well as cross-cohort discrepancies.

Handedness

Handedness, and more specifically, being left-handed, was previously correlated with faster

svRT performance[19,44,46], which corroborates the findings in the MindCrowd analysis. Hemispheric

asymmetries in spatial processing are generally considered to be responsible for the faster svRT of the

left hand[45,47]. The associated faster svRT in left-handed participants revealed in the MindCrowd analysis

appears to diminish with age. Figure 2B shows a separation of the linear regression lines that eventually

cross around the 4th decade of life. This idea is supported in Figures 2C and 2D as the lines separate in

20 to 40-years-olds (Figure 2C), but not in 40 to 70-year-olds (Figure 2D). This fact may underly why

we did not find associations between handedness in UKBb MindCrowd's svRT or UK Biobank's cvrRT.

Perhaps, a portion of older participants (e.g., 40-70 years old) were forced to be right-handed due to

social conventions of the time, something not as likely in younger participants. In doing so, this may

have introduced variability in an aged individual across both cohorts. As its association with svRT is

minimal.

Health, Medical, and Lifestyle Factors

The MindCrowd analysis incorporated all 13 available health, medical, and lifestyle-

related factors, of which six were present and incorporated into the shared UKBb MindCrowd/UK

Biobank model (Tables 2 and 4). Before the launch of MindCrowd, these factors were carefully selected

based upon their known relation to 1) age-related change, 2) RT performance, and 3) PAL Performance.

Of the 13 health, medical, and lifestyle factors evaluated in the MindCrowd analysis, we found

associations between svRT and the number of daily medications, reported dizziness, smoking status,

reported stroke, and diabetes mellitus. Each health and medical factor were associated with slower

svRTs performance (Figures 3 and 4). We should note that the number of daily medications is a likely

proxy for overall health. That is, the worse one's health, the worse one's performance, the increased

number of medications treating the underlying health conditions. The UKBb MindCrowd (svRT) and the

UK Biobank (cvrRT) analyses found *similar* associations between reported dizziness, reported stroke,

diabetes mellitus, indicating hypertension. Although each association differed in magnitude between the

two older cohorts, each was related to slower RT. The UKBb MindCrowd to the UK Biobank found a

different association for FHAD (UKBb MindCrowd=no association; UK Biobank=2.43 milliseconds

slower), smoking status (UKBb MindCrowd=10 milliseconds slower; UK Biobank=no association).

Interestingly, despite some differences, only a few coefficient signs differed between the UKBb

MindCrowd and UK Biobank; indeed, most estimations were well within an order of magnitude

between the two cohorts (e.g., age, educational attainment, and age by biological sex interaction, Table

5).

Many factors are likely to account for the different associations between smoking and FHAD

between UKBb MindCrowd and UK Biobank, (Figures 5-13). Some of these include differences in

demographics, genetic heterogeneity, and age[26]. However, likely candidates include the fractions of

participants reporting each factor (e.g., for diabetes mellitus: MindCrowd=1%, UKBb MindCrowd=7%,

and UK Biobank=3%), and the UK Biobank participant number being twice the size of MindCrowd and

fourfold higher than UKBb MindCrowd. Despite the size of our study, the observational and cross-

sectional methodology means that we cannot rule out effects due to confounding variables.

Consequently, while numbers may be close, we do not assume that the UKBb MindCrowd is similar and

can be compared to the UK Biobank. Furthermore, we observed that UKBb MindCrowd consistently

reported larger estimates and standard errors when compared to the UK Biobank. For example, the

MindCrowd cohort's estimation of the sex-difference was consistently slower (~40 milliseconds), even

in the UKBb MindCrowd cohort, when looking at the UK Biobank (~19 milliseconds). This difference

demonstrates why the study of neuropsychological traits and disease requires large sizes to provide

accurate estimations driving better predictive validity.

Opportunities and Limitations of Our Research

Opportunities: Unique impact

We strongly advocate for large-scale efforts like ours, the UK Biobank[48], and others[22]. Indeed,

studies of this kind possess characteristics that provide unique impact necessary to move the fields of

aging and age-related diseases forward. These include: 1) statistical control, as our MindCrowd analysis,

incorporated all 24 available factors, 11 of which were utilized in the UKBb MindCrowd and UK

Biobank model. 1) The inclusion of each predictor controlled for its association on RT, which

potentially removed variability (noise), thus enhancing statistical power. 2) The two models used for

each of the analyses were selected with little human input by automated application of specific statistical

criteria (see Inclusion of polynomials and automatic model selection in Statistical Methods); thus, likely

decreasing bias, the probability of overfitting, and multicollinearity. 3) For the present study,

MindCrowd had over 76K and the UK Biobank over 158K participants. Many participants in each

cohort were expected to help reduce variance, enhance estimation, select better models, and in turn,

enhance statistical power. Expanded statistical power may then enhance accuracy, validity, and

reproducibility.

Lastly, a recent genome-wide association study examining associations between RT and single

nucleotide polymorphisms (SNP) in the UK Biobank as well as CHARGE and COGENT consortia,

noted weak correlations between the reported cognitive-associated SNPs among US and UK cohorts [49].

Here, MindCrowd presents a future opportunity to resolve these weak associations and get a better

picture of potential cohort effects. Taken together, these characteristics increase the likelihood of

making accurate inferences regarding associations while boosting predictive validity. These are both

necessary and vital attributes when searching for genetic associations and the structure underlying health

brain aging.

Limitations: Internet self-report

There are potential concerns that arise from web-based studies[50]. However, we believe that the

advantage of meaningfully higher participant numbers and enriched cohort diversity facilitated via

online research remediates some of the disadvantages. For example, the range of error reported in recent

internet-based studies of self-reported quantitative traits like height and weight was between 0.3 and

20%[51-54]. Previously, we ran simulations on the association between FHAD and PAL by randomly

shuffling the FHAD responses (e.g., Yes to No, and No to Yes) introducing increasing sequential

amounts of "error." We found that even with a subtle effect such as FHAD on PAL performance, due to

our cohort size, 24% error would still have only made us commit a Type1 error 50% of the time[28]. In

line with this notion, another recent publication demonstrated that online RT studies produce

reproducible results[55]. Further, we developed an extensive data filtering pipeline (see Data Quality

Control) to address these concerns and to enhance validity and accuracy. For example, we controlled for

svRT delays generated with the usage of the touchscreen or mobile device by adding this information as

a variable to the regression model (see Supplementary Figure 8).

Conclusion

Understanding the modifiable and non-modifiable variables associated with RT and related

cognitive function will begin to deconstruct the underlying architecture of elements accounting for the

vast heterogeneity seen in individual trajectories of age-associated cognitive decline. Only then will it be

possible to develop a model of healthy brain aging that is both valid and reliable[56]. Such a model holds

great potential to attenuate age- and disease-related cognitive deficits, thus enhancing the cognitive

healthspan. Any extension of cognitive healthspan, better aligning it to the human lifespan, would be

invaluable and increasingly vital when aggregated across the aging population. Mitigating age- or

disease-related cognitive decline, allowing maintenance of independence by even only a few years,

would have many benefits. For example, the U.S. would save billions of dollars in health care costs and

lost productivity of caregivers, all the while improving the quality of life for the aging population[48]. In

this study, we revealed several factors related to aging and processing speed. Of those, smoking and

education, potentially modifiable factors throughout life, had opposite effects on age-related slowing of

RT performance. With MindCrowd recruitment ever-increasing, our goal is to continue providing and

refining the knowledge necessary to optimizing cognitive performance throughout life.

References

1. Boisgontier, M.P., Wittenberg, G.F., Fujiyama, H., Levin, O. & Swinnen, S.P. Complexity of central processing in simple and choice multilimb reaction-time tasks. *PLoS One* **9**, e90457 (2014).
2. Deary, I.J. & Der, G. Reaction time, age, and cognitive ability: Longitudinal findings from age 16 to 63 years in representative population samples. *Aging, Neuropsychology, and cognition* **12**, 187-215 (2005).
3. Deary, I.J., Allerhand, M. & Der, G. Smarter in middle age, faster in old age: a cross-lagged panel analysis of reaction time and cognitive ability over 13 years in the West of Scotland Twenty-07 Study. *Psychol Aging* **24**, 40-7 (2009).
4. Welford, A.T. Reaction time, speed of performance, and age. *Ann N Y Acad Sci* **515**, 1-17 (1988).
5. Verhaeghen, P. & Salthouse, T.A. Meta-analyses of age-cognition relations in adulthood: estimates of linear and nonlinear age effects and structural models. *Psychol Bull* **122**, 231-49 (1997).
6. Salthouse, T.A. & Babcock, R.L. Decomposing adult age differences in working memory. *Developmental psychology* **27**, 763 (1991).
7. Salthouse, T.A. Aging and measures of processing speed. *Biol Psychol* **54**, 35-54 (2000).
8. Sliwinski, M. & Buschke, H. Processing speed and memory in aging and dementia. *J Gerontol B Psychol Sci Soc Sci* **52**, P308-18 (1997).
9. Bryan, J. & Luszcz, M.A. Speed of information processing as a mediator between age and free-recall performance. *Psychology and aging* **11**, 3 (1996).
10. Baudouin, A., Clarys, D., Vanneste, S. & Isingrini, M. Executive functioning and processing speed in age-related differences in memory: contribution of a coding task. *Brain Cogn* **71**, 240-5 (2009).
11. Lindenberger, U., Mayr, U. & Kliegl, R. Speed and intelligence in old age. *Psychol Aging* **8**, 207-20 (1993).
12. Balakrishnan, G. *et al.* A comparative study on visual choice reaction time for different colors in females. *Neurol Res Int* **2014**, 301473 (2014).
13. Fan, J., Wu, Y., Fossella, J.A. & Posner, M.I. Assessing the heritability of attentional networks. *BMC Neurosci* **2**, 14 (2001).
14. Evarts, E.V., Teravainen, H. & Calne, D.B. Reaction time in Parkinson's disease. *Brain* **104**, 167-86 (1981).
15. Pearce, S.C., Stolwyk, R.J., New, P.W. & Anderson, C. Sleep disturbance and deficits of sustained attention following stroke. *J Clin Exp Neuropsychol* **38**, 1-11 (2016).
16. Spierer, D.K., Petersen, R.A., Duffy, K., Corcoran, B.M. & Rawls-Martin, T. Gender influence on response time to sensory stimuli. *J Strength Cond Res* **24**, 957-63 (2010).
17. Blough, P.M. & Slavin, L.K. Reaction time assessments of gender differences in visual-spatial performance. *Percept Psychophys* **41**, 276-81 (1987).
18. Lupinacci, N.S., Rikli, R.E., Jones, C.J. & Ross, D. Age and physical activity effects on reaction time and digit symbol substitution performance in cognitively active adults. *Res Q Exerc Sport* **64**, 144-50 (1993).

19. Dane, S. & Erzurumluoglu, A. Sex and handedness differences in eye-hand visual reaction times
in handball players. *Int J Neurosci* **113**, 923-9 (2003).
20. Tonnessen, E., Haugen, T. & Shalfawi, S.A. Reaction time aspects of elite sprinters in athletic
world championships. *J Strength Cond Res* **27**, 885-92 (2013).
21. Kaur, P., Paul, M. & Sandhu, J.S. Auditory and visual reaction time in athletes, healthy controls,
and patients of type 1 diabetes mellitus: A comparative study. *International Journal of Diabetes*
in Developing Countries **26**(2006).
22. Vincent, A.S., Roebuck-Spencer, T., Gilliland, K. & Schlegel, R. Automated
Neuropsychological Assessment Metrics (v4) Traumatic Brain Injury Battery: military normative
data. *Mil Med* **177**, 256-69 (2012).
23. Vincent, A.S. *et al.* Reference data from the automated Neuropsychological Assessment Metrics
for use in traumatic brain injury in an active duty military sample. *Mil Med* **173**, 836-52 (2008).
24. Krieg, E.F., Jr. *et al.* Neurobehavioral test performance in the third National Health and Nutrition
Examination Survey. *Neurotoxicol Teratol* **23**, 569-89 (2001).
25. Button, K.S. *et al.* Power failure: why small sample size undermines the reliability of
neuroscience. *Nat Rev Neurosci* **14**, 365-76 (2013).
26. Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes of a wide range
of complex diseases of middle and old age. *PLoS Med* **12**, e1001779 (2015).
27. Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes of a wide range
of complex diseases of middle and old age. *PLoS medicine* **12**(2015).
28. Talboom, J.S. *et al.* Family history of Alzheimer's disease alters cognition and is modified by
medical and genetic factors. *Elife* **8**(2019).
29. Bashore, T.R. & Ridderinkhof, K.R. Older age, traumatic brain injury, and cognitive slowing:
some convergent and divergent findings. *Psychol Bull* **128**, 151-98 (2002).
30. Wickham, H. *et al.* Welcome to the Tidyverse. *Journal of Open Source Software* **4**, 1686 (2019).
31. Long, J.A. & Long, M.J.A. Package 'jtools'. (2017).
32. Lenth, R., Singmann, H. & Love, J. Emmeans: Estimated marginal means, aka least-squares
means. *R package version* **1**(2018).
33. Der, G. & Deary, I.J. Age and sex differences in reaction time in adulthood: results from the
United Kingdom Health and Lifestyle Survey. *Psychol Aging* **21**, 62-73 (2006).
34. Galton, F. Exhibition of instruments (1) for testing perception of differences of tint, and (2) for
determining reaction-time. *The Journal of the Anthropological Institute of Great Britain and*
Ireland **19**, 27-29 (1890).
35. Bleecker, M.L., Bolla Wilson, K., Agnew, J. & Meyers, D.A. Simple visual reaction time: sex
and age differences. *Developmental Neuropsychology* **3**, 165-172 (1987).
36. Kosinski, R.J. A literature review on reaction time. *Clemson University* **10**, 337-344 (2008).
37. Calcagno, V. & de Mazancourt, C. glmulti: an R package for easy automated model selection
with (generalized) linear models. *Journal of statistical software* **34**, 1-29 (2010).
38. Johnson, S.C. *et al.* The influence of Alzheimer disease family history and apolipoprotein E
epsilon4 on mesial temporal lobe activation. *J Neurosci* **26**, 6069-76 (2006).
39. Moradi, A. & Esmaeilzadeh, S. Simple reaction time and obesity in children: whether there is a
relationship? *Environ Health Prev Med* **22**, 2 (2017).
40. Nettelbeck, T. Factors affecting reaction time: Mental retardation, brain damage, and other
psychopathologies. *Reaction times*, 355-401 (1980).
41. Lee, J.J. & Chabris, C.F. General cognitive ability and the psychological refractory period:
individual differences in the mind's bottleneck. *Psychol Sci* **24**, 1226-33 (2013).

42. Taimela, S. Factors Affecting Reaction-Time Testing and the Interpretation of Results.
Perceptual and Motor Skills **73**, 1195-1202 (1991).
43. Jain, A., Bansal, R., Kumar, A. & Singh, K.D. A comparative study of visual and auditory
reaction times on the basis of gender and physical activity levels of medical first year students.
Int J Appl Basic Med Res **5**, 124-7 (2015).
44. Barthelemy, S. & Boulinguez, P. Manual reaction time asymmetries in human subjects: the role
of movement planning and attention. *Neurosci Lett* **315**, 41-4 (2001).
45. Woods, D.L., Wyma, J.M., Yund, E.W., Herron, T.J. & Reed, B. Factors influencing the latency
of simple reaction time. *Front Hum Neurosci* **9**, 131 (2015).
46. Barthelemy, S. & Boulinguez, P. Orienting visuospatial attention generates manual reaction time
asymmetries in target detection and pointing. *Behav Brain Res* **133**, 109-16 (2002).
47. Gonzalez, C.L., Ganel, T. & Goodale, M.A. Hemispheric specialization for the visual control of
action is independent of handedness. *J Neurophysiol* **95**, 3496-501 (2006).
48. Huentelman, M.J., Talboom, J.S., Lewis, C.R., Chen, Z. & Barnes, C.A. Reinventing
Neuroaging Research in the Digital Age. *Trends Neurosci* **43**, 17-23 (2020).
49. Davies, G. *et al.* Study of 300,486 individuals identifies 148 independent genetic loci influencing
general cognitive function. *Nat Commun* **9**, 2098 (2018).
50. Kraut, R. *et al.* Psychological research online: report of Board of Scientific Affairs' Advisory
Group on the Conduct of Research on the Internet. *Am Psychol* **59**, 105-17 (2004).
51. Elgar, F.J. & Stewart, J.M. Validity of self-report screening for overweight and obesity.
Evidence from the Canadian Community Health Survey. *Can J Public Health* **99**, 423-7 (2008).
52. Shields, M., Connor Gorber, S. & Tremblay, M.S. Estimates of obesity based on self-report
versus direct measures. *Health Rep* **19**, 61-76 (2008).
53. Ikeda, N. Validity of Self-Reports of Height and Weight among the General Adult Population in
Japan: Findings from National Household Surveys, 1986. *PLoS One* **11**, e0148297 (2016).
54. Yoon, K., Jang, S.N., Chun, H. & Cho, S.I. Self-reported anthropometric information cannot
vouch for the accurate assessment of obesity prevalence in populations of middle-aged and older
Korean individuals. *Arch Gerontol Geriatr* **59**, 584-92 (2014).
55. Reimers, S. & Stewart, N. Adobe Flash as a medium for online experimentation: A test of
reaction time measurement capabilities. *Behavior research methods* **39**, 365-370 (2007).
56. Ryan, L. *et al.* Precision Aging: Applying Precision Medicine to the Field of Cognitive Aging.
Front Aging Neurosci **11**, 128 (2019).
669

Data Availability

All programs, software, and other materials described herein are publicly or commercially

available. R code for all analyses and figure generation was included as "Source Data." The R code

includes each model selected and the full list of parameters used. The data that support each analysis,

figure, and table is freely available at Dryad (https://datadryad.org).

Acknowledgments

A substantial porting of this research was conducted using the generous data and resources

provided by the UK Biobank (https://www.ukbiobank.ac.uk/published-papers). We acknowledge the

Mueller Family Charitable Trust and Mike Mueller their significant donations. This work was also

supported by the State of Arizona DHS in support of the Arizona Alzheimer's Consortium (PI; Eric

Reiman), the Flinn Foundation (PI; Matt Huentelman), The McKnight Brain Research Foundation, NIH-

NIA grant R01-AG049465 (PI; Carol Barnes), and NIH-NIA grants UH2 AG064706, U19 AG023122,

U24 AG051129, and U24 AG051129-04S1 (PI; Nicholas J. Schork). We also recognize the support of

many anonymous donors and institutional support of TGen, City of Hope, and the University of

Arizona.

Competing Interest

The authors have no known competing or conflicting interests to declare.

Materials and Correspondence

Matt Huentelman, Ph.D., TGen, 445 N Fifth Street, Phoenix, AZ 85004, 602-343-8730 (office),

mhuentelman@tgen.org.

Author Contributions

J.S.T wrote and edited much of the manuscript, analyzed much of these data, supervised several

areas of project management. J.S.T also contributed greatly to data analysis, interpretation, and

synthesis. M.D.DB oversaw much of the information technology required for this study, analyzed a

portion of these data, and provided critical project support. M.A.N analyzed a portion of these data,

Figure 1. Age, paired associates learning (PAL) performance, and Biological Sex were associated with simple visual reaction time (svRT) in the MindCrowd analysis (ages 18-85). (A) Linear model fits (line fill ± 95% *CI*, error bars ± *SEM*) of the median svRT by Age^3 (curvilinear model). There was a curvilinear relationship between svRT and Age^1 (β_{Age1}=7.07, p_{Age1}=3.06E-18), Age^2 (β_{Age2}=-0.15, p_{Age2}=3.23E-17), and Age^3 (β_{Age3}=1.47E-03, p_{Age3}< 1.46E-34, *n*=75,666). (B) Linear model fits (line fill ± 95% *CI*, error bars ± *SEM*) of the median svRT by Age^3 (curvilinear model). There was a curvilinear relationship between svRT and paired-associates learning (PAL) performance PAL Performance^1 (β_{PAL1}=-8.89, p_{PAL1}=3.77E-34), PAL Performance^2 (β_{PAL2}=0.32, p_{PALl2}=2.35E-14), and PAL Performance^3 (β_{PAL3}=-4.13E-03, p_{PAL3}=7.28E-09, *n*=75,666). (C) Simple slope analysis of the Linear model fits (line fill ± 95% *CI*, error bars ± *SEM*) of the median svRT by PAL Performance ($\beta_{Age*PAL}$=-0.07, $p_{Age*PAL}$=1.26E-59). At 20 ($\beta_{Age20*PAL}$=-5.48, $p_{Age20*PAL}$<0.00, *n*=1985), 40 ($\beta_{Age40*PAL}$=-6.41, $p_{Age40*PAL}$<0.00, *n*=739), 60 ($\beta_{Age60*PAL}$=-7.75, $p_{Age60*PAL}$<0.00, *n*=1789), and 80 ($\beta_{Age80*PAL}$=-9.08, $p_{Age80*PAL}$<0.00, *n*=344) years of age. (D) Linear model fits (line fill ± 95% *CI*, error bars ± *SEM*) of the median svRT by Age^3 (curvilinear model) with lines split by Biological Sex. Women demonstrated slower svRT compared to men across the aging spectrum (β_{Sex}=-34.26, p_{Sex}=1.26E-163, Women *n*=47,700, Men *n*=27,966).

Figure 2. Educational attainment and handedness altered simple visual reaction time (svRT) performance. MindCrowd analysis (ages 18-85). (A) Linear model fits (line fill ± 95% *CI*) of the median svRT by Age^3 (curvilinear model) with lines split by Educational Attainment. Participants who had "Some College" ($\beta_{College}$=-14.73, $p_{College}$=8.74E-05, n=22,950), or a "College Degree" ($\beta_{CDegree}$=-31.78, $p_{CDegree}$=2.95E-17, n=44,140) were faster than those with "No High School Diploma" (n=1,881). (B-D) Linear model fits (line fill ± 95% *CI*) of the median svRT by Age^3 (curvilinear model) with lines split by Handedness. (B) From 18-85 years old, left-handed participants showed slightly faster svRTs (β_{Left}=-3.87, p_{Left}<0.03), (C) an association found in 20-40 years olds (β_{40Left}=-3.16, p_{40Left}<0.01, Left-Handed n=8,449), (D) but not 40-60 years olds (β_{60Left}=-2.69, p_{60Left}=0.07, Right-Handed n=66,903).

Figure 3. Simple visual reaction times (svRT) were slower for those who smoked or Reported Dizziness. MindCrowd analysis (ages 18-85). (A) Linear model fits (line fill ± 95% *CI*) of the median svRT by Age^3 (curvilinear model) with lines split by Smoking Status. Participants identifying as a smoker showed slower svRT ($\beta_{Smoking}$=7.07, $p_{Smoking}$=1.26E-03, Smoker *n*=5,793, Non-Smoker *n*=69,873). (B) Linear model fits (line fill ± 95% *CI*) of the median svRT by Age^3 (curvilinear model) with lines split by Reported Dizziness. Participants reporting Reported Dizziness showed slower svRT performance (β_{Dizzy}=4.87, p_{Dizzy}=0.04, Reported Dizziness Reported *n*=4,749, No Reported Dizziness Reported *n*=70,917).

Figure 4. Health factors such as daily medications taken, diabetes, and Reported Stroke all slowed simple visual reaction time (svRT). MindCrowd analysis (ages 18-85). (A) Linear model fits (line fill ± 95% *CI*) of the median svRT by Age^3 (curvilinear model) with lines split by Daily Medications Taken. Compared to participants reporting "no daily medications" (*n*=33,672), taking "Two" (β_{Meds2}=5.84, p_{Meds2}=2.00E-03, *n*=9,651), "Three" (β_{Meds3}=6.24, p_{Meds3}<0.01, *n*=6,656), and "Four" (β_{Meds4}=17.82, p_{Meds4}=3.51E-16, *n*=10,769) slowed svRT. (B) Linear model fits (line fill ± 95% *CI*) of the median svRT by Age^3 (curvilinear model) with lines split by diabetes mellitus. Participants reporting having diabetes had slower svRT performance ($\beta_{Diabetes}$=11.23, $p_{Diabetes}$<3.36E-05, Diabetes Reported *n*=3,887, No Diabetes Reported *n*=71,779). (C) Linear model fits (line fill ± 95% *CI*) of the median svRT by Age^3 (curvilinear model) with lines split by Reported Stroke. Reporting having had a Reported Stroke impaired (slowed) svRT performance (β_{Stroke}=20.38, p_{Stroke}=3.59E-04, Reported Stroke Reported *n*=765, No Reported Stroke Reported *n*=74,901).

Figure 5. Age was linearly associated with visual reaction time (RT).UK Biobank analysis (ages 40-70). (A) UKBb MindCrowd linear model fits (line fill ± 95% *CI*, error bars ± *SEM*) of median simple visual RT (svRT) across Age. (B) UK Biobank linear model fits (line fill ± 95% *CI*, error bars ± *SEM*) of median complex, visual, recognition RT (cvrRT) across Age. UKBb MindCrowd svRT (β_{Age}=5.75, p_{Age}=2.00E-16, *n*=39,795) and UK Biobank cvrRT (β_{Age}=3.40, p_{Age}=2.00E-16, *n*=158,245) showed similar age-associated slowing, but overall higher RT. The average 50 milliseconds difference between UKBb MindCrowd svRT (*M*=478.66milliseconds) and UK Biobank (*M*=528.74 milliseconds) is likely due to the choice component (i.e., do cards match or not > press button) of the UK Biobank's cvrRT task compared to UKBb MinsCrowd's stimulus-response (i.e., the pink sphere appears > press button) svRT.

Figure 6. Visual reaction time (RT) was faster in men. UK Biobank analysis (ages 40-70). (A) UKBb MindCrowd linear model fits (line fill ± 95% *CI*, error bars ± *SEM*) of median simple visual RT (svRT) across Age with lines split by Biological Sex. (B) UK Biobank linear model fits (line fill ± 95% *CI*, error bars ± *SEM*) of median complex, visual, recognition RT (cvrRT) across Age with lines split by Biological Sex. Both UKBb MindCrowd svRT (β_{Sex}=-40.00, p_{Sex}=8.03E-71, Women [*M*=489.75 milliseconds, *n*=29,640], Men [*M*=446.28 milliseconds, *n*=10,155]) and UK Biobank cvrRT (β_{Sex}=-18.28, p_{Sex}=2.00E-16, Women [*M*=534.98 milliseconds, *n*=89,331], Men [520.66 milliseconds, *n*=68,914) found men consistently outperforming women from 40-70 years of age.

Figure 7. More education was related to faster visual reaction time (RT). UK Biobank analysis (ages 40-70). (A) UKBb MindCrowd linear model fits (line fill ± 95% *CI*) of median simple visual RT (svRT) across Age with lines split by Educational Attainment. Participants who had a "High School Diploma" ($\beta_{HSDiploma}$=-9.68, $p_{HSDiploma}$=2.70E-01, 1.87%, *n*=3,176), "Some College" ($\beta_{College}$=-26.08, $p_{College}$=1.61E-03, 5.06%, *n*=11,139), or a "College Degree" ($\beta_{CDegree}$=-49.07, $p_{CDegree}$=1.91E-09, 9.51%, *n*=24,875) were faster than those with "No High School Diploma" (*n*=605). (B) UK Biobank linear model fits (line fill ± 95% *CI*) of median complex, visual, recognition RT (cvrRT) across Age with lines split by Educational Attainment. Like the UKBb MindCrowd cohort, participants who had a "High School Diploma" ($\beta_{HSDiploma}$=-9.68, $p_{HSDiploma}$=1.23E-33, 1.74%, *n*=46,247), "Some College" ($\beta_{College}$=-10.26, $p_{College}$=1.30E-40, 1.85%, *n*=77,270), or a "College Degree" ($\beta_{CDegree}$=-11.71, $p_{CDegree}$=1.46E-41, 2.11%, *n*=23,750) were faster than those with "No High School Diploma" (*n*=10,978).

Figure 8. Diabetes mellitus slowed visual reaction time (RT). UK Biobank analysis (ages 40-70). (A) UKBb MindCrowd linear model fits (line fill ± 95% *CI*) of median simple visual RT (svRT) across Age with lines split by diabetes mellitus. (B) UK Biobank linear model fits (line fill ± 95% *CI*) of median complex, visual, recognition RT (cvrRT) across Age with lines split by diabetes mellitus. Participant visual RT performance for UKBb MindCrowd svRT ($\beta_{Diabetes}$=11.48, $p_{Diabetes}$=3.31E-03, Diabetes Reported *n*=2,807, No Diabetes Reported *n*=36,988) and UK Biobank cvrRT ($\beta_{Diabetes}$=5.48, $p_{Diabetes}$=4.80E-07, Diabetes Reported *n*=4,969, No Diabetes Reported *n*=153,276), was slowed with Diabetes Mellitus.

Figure 9. Visual reaction time (RT) was impaired if a Reported Stroke was reported. UK Biobank analysis (ages 40-70). (A) UKBb MindCrowd linear model fits (line fill ± 95% *CI*) of median simple visual RT (svRT) across Age with lines split by Reported Stroke. (B) UK Biobank linear model fits (line fill ± 95% *CI*) of median complex, visual, recognition RT (cvrRT) across Age with lines split by Reported Stroke. In both the UKBb MindCrowd svRT (β_{Stroke}=18.47, p_{Stroke}=4.00E-02, Reported Stroke Reported *n*=2,807, No Reported Stroke Reported *n*=39,318) and UK Biobank cvrRT (β_{Stroke}=10.61, p_{Stroke}=6.15E-07, Reported Stroke Reported *n*=1,237, No Reported Stroke Reported *n*=157,008) analysis, experiencing a Reported Stroke was associated with slower visual RT.

Figure 10. Reported Hypertension slowed both types of visual reaction time (RT). UK Biobank analysis (ages 40-70). (A) UKBb MindCrowd linear model fits (line fill ± 95% *CI*) of median simple visual RT (svRT) across Age with lines split by Reported Hypertension. (B) UK Biobank linear model fits (line fill ± 95% *CI*) of median complex, visual, recognition RT (cvrRT) across Age with lines split by Reported Hypertension. Unlike the MindCrowd analysis, hypertension was related to slower svRT in UKBb MindCrowd ($\beta_{Hypertension}$=7.99, $p_{Hypertension}$=3.16E-03, Hypertension Reported *n*=9,676, No Hypertension Reported *n*=30,119) and cvrRT in the UK Biobank ($\beta_{Hypertension}$=1.14, $p_{Hypertension}$=0.02, Hypertension Reported *n*=32,593, No Hypertension Reported *n*=125,652).

Figure 11. Reported Dizziness impaired visual reaction time (RT). UK Biobank analysis (ages 40-70). (A) UKBb MindCrowd linear model fits (line fill ± 95% *CI*) of median simple visual RT (svRT) across Age with lines split by Reported Dizziness. (B) UK Biobank linear model fits (line fill ± 95% *CI*) of median complex, visual, recognition RT (cvrRT) across Age with lines split by Reported Dizziness. UKBb MindCrowd svRT (β_{Dizzy}=12.13, p_{Dizzy}=2.52E-03, Reported Dizziness Reported *n*=2,543, No Reported Dizziness Reported *n*=37,252) and UK Biobank cvrRT (β_{Dizzy}=3.21, p_{Dizzy}=3.71E-14, Reported Dizziness Reported *n*=42,210, No Reported Dizziness Reported *n*=116,035) were slowed if participants reported Reported Dizziness.

Figure 12. Smoking Status is related to impaired MindCrowd simple visual reaction time (svRT), not UK Biobank complex, visual, recognition reaction time (cvrRT). UK Biobank analysis (ages 40-70). (A) UKBb MindCrowd linear model fits (line fill ± 95% *CI*) of median svRT across Age with lines split by Smoking Status. Compared to non-smokers, smokers demonstrated slower svRT (β_{Smoke}=10.18, p_{Smoke}=0.01, Smoker *n*=2,783, Non-Smoker *n*=37,012). (B) UK Biobank linear model fits (line fill ± 95% *CI*) of median cvrRT across Age with lines split by Smoking Status. An association between Smoking Status and cvrRT was not found in the UK Biobank (β_{Smoke}=0.47, p_{Smoke}=0.22, Smoker *n*=91,312, Non-Smoker *n*=66,923).

Figure 13. A first-degree family history of Alzheimer's disease (FHAD) was related to slower UK Biobank complex, visual, recognition reaction time (cvrRT), UKBb MindCrowd simple visual reaction time (svRT). UK Biobank analysis (ages 40-70). (A) UKBb MindCrowd linear model fits (line fill ± 95% *CI*) of median svRT across Age with lines split by reported FHAD. An association between FHAD and svRT was not found in the UKBb MindCrowd cohort (β_{FHAD}=-0.07, p_{FHAD}=0.97, FHAD Reported *n*=13,748, No FHAD Reported *n*=26,047). (B) UK Biobank linear model fits (line fill ± 95% *CI*) of median cvrRT across Age with lines split by reported FHAD. Compared to those reporting No FHAD, FHAD was related to worse cvrRT performance in the UK Biobank (β_{FHAD}=2.36, p_{FHAD}=3.35E-05, FHAD Reported *n*=19,741, No FHAD Reported *n*=138,504).

Figure 14. In the UK Biobank, biological sex modified the association of educational attainment and complex, visual, recognition reaction time (cvrRT) (ages 40-70). Linear model fits (line fill ± 95% *CI*) of the median cvrRT by Age with lines split by Educational Attainment and Biological Sex. Compared to women having "No High School Diploma" (*n*=6,056), men with a "High School Diploma" ($\beta_{Sex*HSDiploma}$=-11.24, $p_{Sex*HSDiploma}$=3.30E-12, *n*=18,505), "Some College" ($\beta_{sex*College}$=-10.54, $p_{sex*College}$=8.71E-12, *n*=34,230) or a "College Degree" ($\beta_{sex*CDegree}$=-12.8, $p_{sex*CDegre}$=1.68E-13, *n*=11,257) were all significantly associated with faster cvrRT. See Supplementary Figure 7, displaying simple effects parsed using estimated marginal means (*EMM*).

Figure 15. An illustrative summary of the results. Data shown across the MindCrowd (MC), UKBb MindCrowd, and the UK Biobank (UKBb). The color (i.e., red=low negative and blue=high positive) indicates the size of the β (beta coefficient) estimate." N.S.," indicates if the estimated β value was not statistically significant (α=0.05).

Table 1
Sociodemographic Characteristics of Each Cohort

Cohort	Category	*n*	%
1. MindCrowd 18-85yr: Biological Sex	Women	46,480	63.08
	Men	27,204	36.91
UKBb MindCrowd 40-70yr: Biological Sex	Women	29,195	74.51
	Men	9,986	25.49
UK Biobank 40-70yr: Biological Sex	Women	89,331	56.45
	Men	68,916	43.55
2. MindCrowd 18-85yr: Race	Native American	445	0.6
	Asian	3,491	4.74
	Black/African American	1,555	2.11
	Native Hawaiian/Pacific Islander	270	0.37
	Mixed	490	0.66
	White	67,433	91.52
UKBb MindCrowd 40-70yr: Race	Asian	743	1.90
	Black/African American	732	1.87
	Mixed	182	0.47
	White	37,235	95.03
UK Biobank 40-70yr: Race	Asian	1,612	1.02
	Black/African American	979	0.62
	Mixed	847	0.54
	White	154,809	97.83
3. MindCrowd 18-85yr: FHAD	TRUE	17,382	23.59
	FALSE	56,302	76.41
UKBB MindCrowd 40-70yr: FHAD	TRUE	13,554	34.59
	FALSE	25,627	65.41
UK Biobank 40-70yr: FHAD	TRUE	19,741	12.48
	FALSE	138,506	87.52
4. MindCrowd 18-85yr: Handedness	Left	8,276	11.23
	Right	65,408	88.77
UKBb MindCrowd 40-70yr: Handedness	Left	4,474	11.42
	Right	34,707	88.58
UK Biobank 40-70yr: Handedness	Left	15,287	9.66
	Right	142,960	90.34
5. MindCrowd 18-85yr: Educational Attainment	No High School Diploma	1,792	2.43
	High School Diploma	6,470	8.78
	Some College	22,335	30.31
	College Degree	43,087	58.48
UKBb MindCrowd 40-70yr: Educational Attainment	No High School Diploma	577	1.47
	High School Diploma	3,108	7.93
	Some College	10,955	27.96
	College Degree	24,541	62.63
UK Biobank 40-70yr: Educational Attainment	No High School Diploma	10,979	6.94
	High School Diploma	46,247	29.23
	Some College	77,271	48.83
	College Degree	23,750	15.01

Table 1. Sociodemographic Characteristics Across Each of the Three Cohorts. Study *n*s and related percentages of commonly reported sociodemographic factors, from each of the three cohorts (i.e., MindCrowd as well as UKBb MindCrowd and the UK Biobank).

Table 2

MindCrowd Cohort Sample Sizes (n)

Characteristic	*n*s
Biological Sex	Women = 47,700 Men = 27,966
Educational Attainment	No High School Diploma = 1,881 High School Diploma = 6,695 Some College = 22,950 College Degree = 44,140
Handedness	Left-Handed = 8,449 Right-Handed = 66,903
Daily Medications Taken	None = 33,672 One = 14,409 Two = 9,651 Three = 6,656 Four =10,769
Reported Dizziness	Dizziness Reported = 4,749 No Dizziness Reported = 70,917
Smoking Status	Smoker = 5,793 Non-Smoker = 69,873
Diabetes mellitus	Diabetes Reported = 3,887 No Diabetes Reported = 71,779
Reported Stroke	Stroke Reported = 765 No Stroke Reported = 74,901

Table 2. Summary of sample size numbers (ns) of each multiple regression coefficient from MindCrowd.

Table 3

MindCrowd Summary of Results

Factor	Value	β	SE	t	p
Intercept		355.88	11.99	29.68	1.89E-192
Age^1	Per Year Change in Slope	7.07	0.81	8.71	3.06E-18
Age^2	Per Year Change in How Much the Slope Is Altered	-0.15	0.02	-8.44	3.23E-17
Age^3	Per Year Change in How Much the Slope Is Altered	1.20E-04	0	12.27	1.46E-34
PAL Performance^1	PAL Word Pairs Correctly Remembered	-8.89	0.73	-12.19	3.77E-34
PAL Performance^2	Per Word Pair Change in Slope	0.32	0.04	7.63	2.35E-14
PAL Performance^3	Per Year Change in How Much the Slope Is Altered	7.14E-04	0	-5.78	7.28E-09
Biological Sex	Men	-34.26	1.25	-27.33	1.26E-163
Educational Attainment	Some College	-14.73	3.75	-3.92	8.74E-05
Educational Attainment	College Degree	-31.78	3.76	-8.45	2.95E-17
Handedness	Left	-3.87	1.76	-2.2	0.03
Daily Medications Taken	Two	5.84	1.89	3.09	2.00E-03
Daily Medications Taken	Three	6.24	2.26	2.76	0.01
Daily Medications Taken	Four	17.82	2.18	8.16	3.51E-16
Reported Dizziness	TRUE	4.87	2.35	2.07	0.04
Smoking Status	TRUE	7.07	2.19	3.23	1.26E-03
Diabetes Mellitus	TRUE	11.23	2.71	4.15	3.36E-05
Reported Stroke	TRUE	20.38	5.71	3.57	3.59E-04
Age x PAL Performance	Trajectory of age-related change in svRT performance is altered buy PAL Performance	-7.56E-04	4.41E-05	-1.72E+01	8.34E-66
Age x Smoking Status	Smoking Status alters the trajectory of age-related changes to svRT	0.58	0.12	4.67	2.95E-06
Age x Reported Stroke	Reported Stroke adds to age-related decrements in svRT	1.87	0.42	4.46	8.17E-06
Age x Biological Sex	Men have a shifted trajectory of age-related svRT changes	-0.37	0.07	-5.59	2.23E-08
Age x Educational Attainment	Having Some College vs. No High School Diploma alters the trajectory of age-related alterations to svRT	-0.76	0.18	-4.09	4.28E-05
Age x Educational Attainment	A College Degree vs. No High School Diploma alters the trajectory of age-related modifications to svRT performance	-1.53	0.19	-8.14	4.08E-16

Table 3. Summary of the main findings from MindCrowd, listing main associations and interactions. β = unstandardized regression coefficient, *SE* = standard error, *t* = value of *t*-statistic, *p* = *p*-value.

Table 4

UKBb MindCrowd and UK Biobank Cohort Sample Sizes (n)

Characteristic	MC*n*s	UK*n*s
Biological Sex	Women = 29,640 Men = 10,155	Women = 8,9331 Men n = 68,914
Diabetes Mellitus	Diabetes Reported = 2,807 No Diabetes Reported = 36,988	Diabetes Reported = 4,969 No Diabetes Reported = 153,276
Handedness	Left-Handed = 4,520 Right-Handed = 35,034	Left-Handed = 15,287 Right-Handed = 142,958
Reported Stroke	Reported Stroke Reported = 2,807 No Reported Stroke Reported = 39,318	Reported Stroke Reported = 1,237 No Reported Stroke Reported = 157,008
Reported Hypertension	Hypertension Rep. = 9,676 No Hypertension Rep. = 30,119	Hypertension Rep. = 32,593 No Hypertension Rep. = 125,652
Smoking Status	Smoker = 2,783 Non-Smoker = 37,012	Smoker = 91,312 Non-Smoker = 66,923
Reported Dizziness	Reported Dizziness Reported = 2,543 No Reported Dizziness Reported = 37,252	Reported Dizziness Reported = 42,210 No Reported Dizziness Reported = 116,035
Educational Attainment	No High School Diploma = 605 High School Diploma = 3,176 Some College = 11,139 College Degree = 24,875	No High School Diploma = 10,978 High School Diploma = 46,247 Some College = 77,270 College Degree = 23,750
FHAD	FHAD Reported = 13,748 No FHAD Reported = 26,047	FHAD Reported = 19,741 No FHAD Reported = 138,504
Educational Attainment x Biological Sex	Women No High School Diploma = 426 High School Diploma = 2,496 Some College = 8,740 College Degree = 17,978 Men No High School Diploma = 179 High School Diploma = 680 Some College = 2,399 College Degree = 12,493	Women No High School Diploma = 6,056 High School Diploma = 27,742 Some College = 43,040 College Degree = 23,750 Men No High School Diploma = 4,922 High School Diploma = 18,505 Some College = 34,230 College Degree = 11,257

Table 4. Summary of sample size numbers (*n*s) of each multiple regression coefficient from the UKBb MindCrowd and UK Biobank.

Table 5
UKBb MindCrowd and the UK Biobank Summary of Results

Variable	Value	UKBb MindCrowd				UK Biobank				Outcome
		β	SE	t	p	β	SE	t	p	
Intercept		189.77	10.65	18.35	6.46E-75	335.93	1.69	209.93	2.00E-16	Agree
Age	Per Year	5.75	0.12	46.57	2.00E-16	3.40	0.03	132.87	2.00E-16	Agree
Biological Sex	Men	-40.00	2.24	-17.83	8.03E-71	-18.28	0.38	-47.70	2.00E-16	Agree
Diabetes mellitus	TRUE	11.48	3.91	2.94	3.31E-03	5.48	1.09	5.03	4.80E-07	Agree
Handedness	Left	-2.58	3.05	-0.85	4.00E-01	0.58	0.63	0.92	0.36	N.S. in both - but MindCrowd subtle effect
Reported Stroke	TRUE	18.47	9.02	2.05	4.00E-02	10.61	2.13	4.99	6.15E-07	Agree
Reported Hypertension	TRUE	6.99	2.37	2.95	3.16E-03	1.09	0.48	2.27	2.00E-02	Agree
Smoking Status	TRUE	10.18	3.87	2.63	0.01	0.40	0.38	1.05	0.29	MindCrowd Finding: Both Primary and UKBb
Reported Dizziness	TRUE	12.10	4.00	3.02	2.52E-03	3.21	0.42	7.57	3.71E-14	Agree
Educational Attainment	High School Diploma	-9.68	8.77	-1.10	2.70E-01	-9.68	0.80	-12.09	1.23E-33	Agree
Educational Attainment	Some College	-26.08	8.27	-3.15	1.61E-03	-10.26	0.77	-13.35	1.30E-40	Agree
Educational Attainment	College Degree	-49.07	8.17	-6.01	1.91E-09	-11.71	0.87	-13.51	1.46E-41	Agree
First-Degree Family History of Alzheimer's Disease	TRUE	-0.07	2.06	-0.04	0.97	2.43	0.57	4.26	2.04E-05	UK Biobank Finding
Age x Biological Sex	Men's vs. Women's per year change in Age slope	-0.62	0.26	-2.38	2.00E-02	-0.55	0.05	-10.99	4.18E-28	Agree
Biological Sex by Educational Attainment	Men with High School Diploma vs. Women with No High School Diploma	21.63	19.62	1.10	0.27	-11.24	1.61	-6.96	3.30E-12	UK Biobank Finding
Biological Sex by Educational Attainment	Men with Some College vs. Women with No High School Diploma	16.25	18.28	0.89	0.37	-10.54	1.54	-6.83	8.71E-12	UK Biobank Finding
Biological Sex by Educational Attainment	Men with a College Degree vs. Women with No High School Diploma	14.79	17.94	0.82	0.41	-12.8	1.74	-7.37	1.68E-13	UK Biobank Finding

Table 5. Summary of the main findings from the UKBb MindCrowd and UK Biobank cohorts. Here factor names, factor definitions, β = unstandardized regression coefficients as well as estimate *SE* = standard error, *t* = value of *t*-statistic, and *p* = *p*-value across both cohorts is shown. The last column displays model outcomes in terms of the agreement between the cohorts.

Table 6
UK Biobank Factor Generation

Index #	Field Description	Field ID	Count (*n*)	Type	Description	Column Label	Used to Generate
6	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=31	31	502,524	Categorical (single)	Sex	Sex	Biological Sex
7	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=34	34	502,524	Integer	Year of birth	Birth Year	Age
264	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=404	404	202,464	Integer	Duration to the first press of snap-button in each round	Snap-Button Time to First Press (Instance 0)	Median cvrRT (8 out 12 Rounds)
395	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=1707	1707	501,626	Categorical (single)	Handedness (chirality/laterality)	Handedness	Handedness
422	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=2443	2443	501,593	Categorical (single)	Diabetes diagnosed by doctor	Diabetes mellitus	Diabetes mellitus
426	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=2453	2453	501,593	Categorical (single)	Cancer diagnosed by doctor	Cancer	Cancer
482	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=2966	2966	134,645	Integer	Age high blood pressure diagnosed	Age Hypertension Diagnosed	Hypertension Diag. via MD
616	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=4056	4056	7,579	Integer	Age Reported Stroke diagnosed	Age Reported Stroke Diagnosed	Reported Stroke Diag. via MD
2104	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=20107	20107	487,790	Categorical (single)	Illnesses of father	(Multiple Columns)	FHAD
2144	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=20110	20110	492,928	Categorical (single)	Illnesses of mother	(Multiple Columns)	FHAD
2188	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=20111	20111	433,922	Categorical (single)	Illnesses of siblings	(Multiple Columns)	FHAD
2325	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=20116	20116	501633	Categorical (single)	Smoking status	Smoking status	Smoking Status
2356	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=21022	21022	502,524	Integer	Age at recruitment	Age at Recruitment	Age (verify)
120	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=401	401	495,702	Categorical (single)	Index for card A in round	Needed to Know if Snap-Button Should Have Been Pressed	Median cvrRT
168	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=402	402	495,702	Categorical (single)	Index for card B in round	Needed to Know if Snap-Button Should Have Been Pressed	Median cvrRT
1635	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=6138	6138	497,883	Categorical (multiple)	Qualifications	Education	Educational Attainment
2357	http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=21053	21053	174774	Categorical (single)	Degree bothered by dizziness in the last three months	Dizziness	Dizziness

Table 6. List of derived factors from the UK Biobank. UK Biobank's "Index Number," "Field Description," "Field ID" and sample size (*n*s) for data field used to generate factors evaluated via multiple linear regression. "Data fields" were chosen to match to the questions posed in MindCrowd. If multiple instances (e.g., 0-4, timepoints) were available for data, only the first (instance 0) was used. For cvrRT and FHAD, multiple columns were compiled to generate the factor. For the UKBb MindCrowd and UK Biobank analyses, participants were removed if any of their responses to a demographic, medical, health, and lifestyle question did not match the other cohort.