

1 **Spatial patterns of child mortality in Nanoro HDSS site, Burkina Faso**

2 Navideh Noori^{1*}‡, Karim Derra^{2‡}, Innocent Valea^{2,3}, Assaf P. Oron¹, Aminata
3 Welgo², Toussaint Rouamba², Palwende Romuald Boua², Athanase SOME²,
4 Eli Rouamba², Edward Wenger¹, Hermann Sorgho², Halidou Tinto^{2,3}, and
5 Andre Lin Ouédraogo¹

6

7 ¹ Institute for Disease Modeling, Building IV, 3150 139th Ave SE, Bellevue,
8 WA, USA, 98005

9

10 ² Institut de Recherche en Sciences de la Sante (IRSS)/Clinical Research Unit
11 of Nanoro (CRUN), Nanoro, Burkina Faso. BP: 218 Ouaga 11, Burkina Faso

12

13 ³ Laboratory of Parasitology and Entomology, Centre Muraz, Bobo-Dioulasso,
14 Burkina Faso. 2054 Avenue Mamadou Konate, 01 B.P. 390 Bobo-Dioulasso
15 01, Burkina Faso

16

17 * Corresponding authors: nnoori@idmod.org

18 ‡ Equally contributed.

19

20

21

22

23

24

25

26

27

28 **Abstract**

29 **Background:** Half of global child deaths occur in sub-Saharan Africa.
30 Understanding child mortality patterns and risk factors will help inform interventions
31 to reduce this heavy toll. The Nanoro Health and Demographic Surveillance System
32 (HDSS), Burkina Faso was described previously, but spatial patterns of child
33 mortality in the district had not been studied. Similar studies in other districts
34 indicated accessibility to health facilities as a risk factor, usually without distinction
35 between facility types.

36 **Methods:** Using Nanoro HDSS data from 2009 to 2013, we estimated the
37 association between under-5 mortality and accessibility to inpatient and outpatient
38 health facilities, seasonality of death, and age group.

39 **Results:** Living in homes 40-60 minutes and >60 minutes travel time from an
40 inpatient facility was associated with 1.52 (95% CI: 1.13-2.06) and 1.74 (1.27-2.40)
41 greater hazard of under-5 mortality, respectively, than living in homes <20 minutes
42 from an inpatient facility. No such association was found for outpatient facilities.
43 Seasonality of death was significantly associated with under-5 mortality, and the wet
44 season (July-November) was associated with 1.28 (1.07, 1.53) higher under-5
45 mortality than the dry season (December-June), likely reflecting the malaria season.

46 **Conclusions:** Our results emphasize the importance of geographical accessibility
47 to health care, and also distinguish between inpatient and outpatient facilities.

48 **Keywords:** Children Under 5; child mortality; Burkina Faso; Spatial Analysis;
49 demographic surveillance; HDSS; Nanoro.

50

51

52

53 **Background**

54 Since the establishment of the Millennium Development Goals in 1990, there
55 has been substantial progress in reducing child mortality globally, from 93
56 deaths in 1990 to 39 deaths in 2017 per 1000 live births. Nonetheless, an
57 estimated 5.4 million children under age five died in 2017, out of which 2.5
58 million died during the first month of their life [1]. About half of child deaths
59 occurred in sub-Saharan Africa [2]. In 2015, the Sustainable Development
60 Goals (SDGs) were defined, aiming to reduce under-five mortality to below 25
61 per 1000 live births by 2030 [3]. To achieve these targets, urgent action in sub-
62 Saharan Africa is needed, as well as higher-quality information to guide this
63 action [4]. Among sub-Saharan countries, Burkina Faso, where our study area
64 is situated, has made great progress in reducing under-5 mortality by about
65 58% from 201 to 84.6 deaths per 1,000 live births between 1990 and 2016,
66 but this rate is still much higher than the SDGs [1].

67 To track progress towards child survival goals and to plan effective
68 interventions for child health, identifying the major drivers of child mortality as
69 well as data-driven estimates of child mortality are necessary [4]. However,
70 countries with the highest child mortality burden lack civil registration and vital
71 statistics (CRVS) systems accounting for all births, deaths and causes of
72 death. In these countries, the location and timing of child deaths and the overall
73 death rates, are highly uncertain. What we know about these crucial public-
74 health questions is informed mostly by nationally representative surveys such
75 as the Demographic and Health Surveys (DHS), conducted every several

76 years.

77 A Health and Demographic Surveillance System (HDSS) is a local CRVS
78 system that routinely monitors the health and demographic characteristics of
79 a population living in a specific area. HDSS data facilitate detailed local studies
80 of public health in general, and child mortality in particular. As of 2020, forty-
81 nine HDSS sites participate in the International Network for the Demographic
82 Evaluation of Populations and Their Health in Developing Countries
83 (INDEPTH), recording the life events of over three million people in 17 African
84 and Asian countries [5]. Several studies have investigated spatial [6, 7, 8, 9,
85 10], temporal [11, 12] and demographic [11, 13, 14] factors affecting child
86 mortality in HDSSs. However, no study to date has analyzed such patterns in
87 the relatively new Nanoro HDSS in rural north-central Burkina Faso.

88 Risk of child mortality varies over space and time, and it is important to identify
89 the areas at the highest risk in order to focus intervention-based efforts in those
90 areas. One source of spatial heterogeneity is accessibility to health facilities
91 [15, 16]. Poor access to health care remains a concern in many low-income
92 countries [17]. A growing number of studies have estimated the effect of
93 distance from a health facility upon child mortality. The first meta-analysis of
94 such studies was published in 2012 [16] and was updated more recently [18].
95 They found that living > 5 km away from a facility is associated with 62% higher
96 neonatal mortality based on 4 studies, and 57% higher under-5 mortality based
97 on 9 studies; both effects were deemed highly significant. In addition, a study
98 aggregating 29 DHSs from 21 countries found that living > 10 km from a facility

99 was strongly associated with 27% higher odds of neonatal mortality. Both the
100 meta-analyses and the DHS-based study did not distinguish between smaller
101 and larger facilities. Most above mentioned studies used simple Euclidean
102 distance, or local expert opinion about distance or travel time, as the exposure
103 variable. More sophisticated approaches to estimate real-life travel distance or
104 time [22] have been published only rarely in this context.

105 Mortality also varies over time as a result of changes in health care-seeking,
106 age and season of birth and death [11, 12], and environmental conditions [19].
107 In the Nouna HDSS, Burkina Faso, infants born during the rainy season were
108 associated with higher mortality risk compared with those born during the dry
109 season [11]. During the rainy season, flooded roads limit the access to health
110 care, especially in the rural region. In most of West Africa, the rainy season
111 also coincides with food shortage until the harvest arrives [11]. Seasonality
112 also drives cause-specific mortality patterns due to malaria, pneumonia and
113 diarrhea, which were the leading causes of child mortality in Burkina Faso in
114 2010 [20].

115 Here we present a quantitative investigation of local under-5 mortality patterns
116 in the Nanoro HDSS site. We were particularly interested in identifying the
117 drivers of spatial heterogeneity in child mortality risk in the area, in view of
118 recent progress on global accessibility estimates capturing inequalities in
119 infrastructural development.

120

121

122 **Methods**

123 **Study Area and Data**

124 Nanoro HDSS site was established in 2009 by the Clinical Research Unit of
125 Nanoro (CRUN), located in the Centre Medicale Saint Camille de Nanoro
126 (CMA), with the goal of evaluating population demography and health living
127 conditions within the health district [24]. Nanoro is located about 85km from
128 the capital city, Ouagadougou. The Nanoro Demographic Surveillance Area
129 (DSA) lies within the health district of Nanoro and includes 24 villages. Initial
130 census started from March to April 2009, and recorded housing and
131 demographic characteristics of 54,781 individuals. Since then, census follow-
132 up has been carried out every four months. Data collected at the individual
133 level include births, deaths, pregnancies, in/out-migrations (temporary or
134 permanent), and relationships (mother, father and head of household). Data
135 from 2009 to end of 2013 were included in this analysis. Nanoro has two main
136 seasons: a rainy season from June to October and a dry season from
137 November to May [24]. In this study, to reflect the malaria mortality seasonality
138 and the potential lag effect of rainy season, the wet season was defined as
139 July to November and the other seven months were defined as the dry season.
140 There are 16 outpatient health facilities in the Nanoro health district and one
141 inpatient health facility close to the village of Nanoro. There is also an inpatient
142 health facility in Bousse just east of the district, which is the closest inpatient
143 facility for some residents in the DSA, and therefore was included in this study
144 (Figure 1).

145 Accessibility to both inpatient and outpatient health facilities was measured as
146 Euclidean distance, travel time, and walking travel time. Travel time to the most
147 accessible health facility was calculated using a global “friction surface”
148 provided by the Malaria Atlas Project (MAP) at a resolution of 1 km for 2015,
149 which estimates the travel time through every 1×1 km grid square on Earth
150 using the fastest feasible surface travel [22]. A companion algorithm
151 calculates the fastest journey time between any two user-provided points. This
152 index may better capture the opportunity cost of travel than Euclidean or
153 network distance, and reflects the information humans use to make transport
154 decisions [22]. We also calculated walking travel time by modifying the friction
155 surface developed by MAP, so that all roads received a fixed walking speed of
156 5 km per hour [22]. Fastest travel time was the main variable used to describe
157 health-facility access in our models. Hereafter we will refer to this variable
158 simply as “travel time.” Models using the other proximity variables are shown
159 in Supplementary Material.

160

161 **Statistical Analysis**

162 We estimated the survival probability of children under age five over the
163 study’s nearly 5-year period, as 1 minus the product of average age-specific
164 monthly survival rates from birth through 60 months, multiplied by 1000. Cox
165 proportional hazards regression models [25] were used to estimate the
166 association between under-5 survival and geographic, and seasonal risk
167 factors. These include physical accessibility to health facilities, seasonality of

168 death events during the survey, and age groups. The relationship between
169 each of these factors and mortality risk was assessed one at a time as both
170 categorical and continuous variables (when possible). The final multivariable
171 model adjusted for risk factors that were significant on a univariate model and
172 available for the entire dataset.

173 For each child, the follow-up time was taken as the time an individual was
174 present within the age group during follow-up, which is the time from the date
175 of first event in the survey, birth or enrollment or in-migration until age 5, out-
176 migration, end of 2013, or death. Village was added as a cluster term to the
177 model to estimate a robust variance. All the analyses and the mapping were
178 performed in R [26].

179

180 **Results**

181 **Demographics and Child Mortality**

182 The key demographic characteristics of the study population are given in Table
183 1. At any given time during the study period, about 8,000 children under 5
184 years old lived in the district. Cumulatively 23,639 children were under age of
185 5 and in the district for at least part of the study period, contributing about
186 37,276 child-years of follow-up. Most children lived within 40 minutes of travel
187 to an inpatient facility. Median household size was smaller among those living
188 near an inpatient facility than those live further away (Table 1).

189 The reported overall mortality rate among children under 5 in Nanoro HDSS
190 during the study period was 64.9 deaths per 1,000 live births. Within the

191 district, the village of Nanoro had the lowest under-5 mortality at 29.4 per 1,000
192 live births, approaching the 2030 SDG of <25. The southern and eastern edges
193 of the DSA had substantially higher mortality rate (91.8 per 1,000 live births)
194 (Supplementary Material, Figures S1-S3).

195

196 **Factors Associated with Child Mortality**

197 The association between child mortality and the variables described in
198 Methods is summarized in Table 2 and Figure 2. As expected, risk of death
199 decreased with increasing age. Children between 1 to 2 years old were at
200 lower risk of mortality by 48% (HR=0.52, 95% CI= 0.41-0.65) than infants, and
201 the risk of mortality was lowest for children 3 to 4 years old (HR=0.39 vs.
202 infants, 95% CI = 0.27-0.57) (Figure 2).

203

204 **Seasonality**

205 The under-5 mortality hazard was higher during the wet season (Jul-Nov)
206 (HR=1.28, 95% CI = 1.07-1.53) than the dry season (Figure 2). Out-migration
207 also had a clear seasonal pattern, and was higher during the dry season,
208 highlighting the potential effect of out-migration on the child mortality pattern
209 (supplementary material, Figure S8).

210

211 **Accessibility to Health Facilities**

212 Under-5 mortality increased significantly with increasing travel time to an
213 inpatient health facility (P<0.001). In particular, children living 40-60 minutes

214 away from an inpatient facility experienced a 1.52 times higher mortality
215 hazard (95% CI = 1.13-2.06) than those living within 20 minutes, and children
216 living > 60 minutes away experienced a relative hazard of 1.74 (95% CI = 1.27-
217 2.40) (Figure 2). Similar associations were found when using Euclidean
218 distance or walking travel time (supplementary material, Figure S4 and Tables
219 S1-S2). By contrast, there was no statistically significant association between
220 accessibility to outpatient health facilities and under-5 mortality
221 (supplementary material, Figure S5 and Tables S1-S2).

222

223 **Discussion**

224 Our study provides insight into child mortality patterns in the Nanoro health
225 district, Burkina Faso by linking it to various demographic, spatial and temporal
226 risk factors. One distinction of our study is the evaluation of accessibility to
227 both inpatient and outpatient health facilities. In the recent meta-analysis by
228 Rojas-Gualdrand and Caicedo-Velazquez [18], the majority of studies included
229 in its under-5 mortality endpoint estimate measured distance from any health
230 center with no distinction between inpatient and outpatient. There were also
231 inconsistencies regarding the effect of accessibility to health care on child and
232 neonatal mortalities. In Malawi, DHS data showed no association between
233 distance to delivery care and early neonatal mortality, and in Zambia, early
234 neonatal survival was higher with increasing distance [27]. On the other hand,
235 analysis of DHS data in Madagascar showed a higher risk of infant mortality
236 among those who lived further from a health facility [28]. In rural western

237 Burkina Faso, rural Ethiopia and Tanzania, accessibility to health facilities was
238 found to be a major risk factor for infant, child and overall under-5 mortality
239 [15, 16, 29]. Our analysis is in agreement with the latter studies, and indicates
240 that impeded access to an inpatient health facility might be a major risk factor
241 for child mortality. Our study also suggests that accessibility to outpatient
242 health facilities does not drive the pattern of child mortality in the study area.
243 We speculate that outpatient health facilities do not provide the level of care
244 children need in a life or death situation. We note the confounding factor that
245 inpatient health facilities are usually located in towns and major villages, with
246 better food, water, and other living conditions for residents, as well as generally
247 higher education and socioeconomic status. Another distinction of our study is
248 the use of the recently developed global accessibility map that accounts for
249 the spatial locations and properties of roads, railroads, rivers, water bodies,
250 topographical characteristics, land cover, and national borders [22].
251 Accounting for these features leads to a more accurate measurement of
252 accessibility than Euclidean or network distance that has been commonly used
253 in previous studies.

254 There was a statistically significant association between seasonality of death
255 and under-5 mortality, with the wet season having a higher mortality rate,
256 reflecting the malaria mortality pattern.

257 Some of the limitations of our work are other risk factors that we have not
258 accounted for and may be important to our outcomes, such as family wealth
259 status, family health-seeking behavior, sanitation and hygiene information, and

260 effects of flooding. Also, the travel time index we used in this study is based
261 on the assumption that everyone could use the fastest travel method possible.
262 However, our analysis using walking travel time showed a similar association
263 with under-5 mortality. An additional limitation is that the friction surface
264 developed by MAP does not account for the seasonal variation, which can
265 affect travel time to health facilities. Last but not least, this is an observational
266 study, and therefore any association is subject to potential confounding
267 factors, as discussed above for inpatient facilities.

268

269 **Conclusions**

270 Our study emphasizes that inequity in mortality rate is not only seen between
271 rural and urban areas, but also within a relatively small rural area. It also
272 highlights the importance of accessibility to health care in rural Burkina Faso.
273 Our findings can help health policy makers and program developers in the
274 health district and similar districts, to understand the potential effect of health
275 infrastructure designs and the most effective locations of health facilities.
276 Novel strategies, such as improved transportation to inpatient facilities during
277 a child health emergency, strengthening of outpatient health facilities, and
278 training community health workers in the rural area, are necessary for
279 mitigating the physical limitations to accessing health care in the area. Also,
280 reducing the socioeconomic inequalities between rural and urban areas as
281 well as within each area, can help enhance access to health services for poor
282 people and reduce child mortality [19, 34].

283

284 **Availability of data and materials**

285 The datasets used and/or analyzed during the current study are available from
286 the authors upon reasonable request. A public version of the dataset with
287 fewer variables, and excluding household location, is available on the
288 INDEPTH website.

289

290

291 **Abbreviations**

292 **SDG:** Sustainable Development Goals

293 **HDSS:** Health and Demographic Surveillance System

294 **CRVS:** Civil Registration and Vital Statistics

295 **DHS:** Demographic and Health Surveys

296 **INDEPTH:** International Network for the Demographic Evaluation of
297 Populations and Their Health in Developing Countries

298 **CRUN:** Clinical Research Unit of Nanoro

299 **CMA:** Centre Medicale avec Antenne Chirurgicale Saint Camille de Nanoro

300 **DSA:** Demographic Surveillance Area

301 **MAP:** Malaria Atlas Project

302

303 **References**

304 1. World Health Organization: WHO — Under-five mortality (2018).
305 <https://www.who.int/gho/child health/mortality/mortality under five text/en/>

- 306 2. UNICEF: Child Mortality - UNICEF Data. doi:10.1016/S2214-
307 109X(14)70280-3. [https://data.unicef.org/topic/child-survival/under-five-](https://data.unicef.org/topic/child-survival/under-five-mortality/)
308 [mortality/](https://data.unicef.org/topic/child-survival/under-five-mortality/)
- 309 3. World Health Organization: Health in 2015: from MDGs, Millennium
310 Development Goals to SDGs, Sustainable Development Goals. Technical
311 report (2015). doi:10.1017/S1744552306002023
- 312 4. You D, Hug L, Ejdemyr S, Idele P, Hogan D, Mathers C, Gerland P, New JR,
313 Alkema L. Global, regional, and national levels and trends in under-5
314 mortality between 1990 and 2015, with scenario-based projections to 2030:
315 A systematic analysis by the un Inter-Agency Group for Child Mortality
316 Estimation. *The Lancet*. 2015; 386(10010), 2275–2286. doi:10.1016/S0140-
317 6736(15)00120-8
- 318 5. Demographic Evaluation of Populations and Their Health in Developing
319 Countries (INDEPTH). <http://www.indepth-network.org/about-us>. Accessed
320 5 June 2020.
- 321 6. Sankoh OA, Y'e Y, Sauerborn R, Mu'llera O, Becher H. Clustering of
322 childhood mortality in rural Burkina Faso. *Int J Epidemiol*. 2001; 30, 485–
323 492.
- 324 7. Becher H. Guest Editorial: Analyses of mortality clustering at member
325 HDSSs within the INDEPTH Network– an important public health issue. *Glob*
326 *Health Action*. 2010; 3(1).
- 327 8. Becher H, Muller O, Dambach P, Gabrysch S, Niamba L, Sankoh O,
328 Simboro S, Schoeps A, Stieglbauer G, Y'e Y, Si'e A. Decreasing child
329 mortality, spatial clustering and decreasing disparity in North-Western
330 Burkina Faso. *Trop Med Int Health*. 2016; 21(4), 546–555.
331 doi:10.1111/tmi.12673
- 332 9. Quattrochi J, Jasseh M, Mackenzie G, Castro MC. Spatial analysis of under-
333 5 mortality and potential risk factors in the Basse Health and Demographic
334 Surveillance System, the Gambia. *Trop Med Int Health*. 2015; 20(7), 941–
335 951.
- 336 10. Alabi O, Baloye D, Doctor HV, Oyedokun OA. Spatial analysis of under-five
337 mortality clustering in northern Nigeria: findings from Nahuiche Health and
338 Demographic Surveillance System, Zamfara state. *Int J Trop Dis Health*.
339 2016; 15(1), 1–10.
- 340 11. Becher H, Muller O, Jahn A, Gbangou A, Kynast-Wolf G, Kouyate B. Risk
341 factors of infant and child mortality in rural Burkina Faso. *Bull World Health*
342 *Organ*. 2004; 82:265–273.
- 343 12. Becher H, Kauerniann G, Khomski P, Kouyat'e B. Using penalized splines
344 to model age and season-of-birth-dependent effects of childhood mortality
345 risk factors in rural Burkina Faso. *Biometrical J*. 2009; 51(1), 110–122.
346 doi:10.1002/bimj.200810496

- 347 13. Desta BN, Assefa N, Damte TD, Hordofa LO. Neonatal mortality and its risk
348 factors in eastern Ethiopia: a prospective cohort Study in Kersa Health and
349 Demographic Surveillance System (Kersa HDSS). *Epidemiol Biostat Public*
350 *Health*. 20016; 13(4), 1–8. doi:10.2427/11938
- 351 14. Ingole V, Kovats S, Schumann B, Hajat S, Rocklöv J, Juvekar S, Armstrong
352 B. Socioenvironmental factors associated with heat and cold-related
353 mortality in Vadu HDSS , western India : a population-based case-crossover
354 study. *Int J Biometeorol*. 2017; 61(10), 1797–1804. doi:10.1007/s00484-
355 017-1363-8
- 356 15. Schoeps A, Gabrysch S, Niamba L, Sie A, Becher H. The effect of distance
357 to health-care facilities on childhood mortality in rural Burkina Faso. *Am J*
358 *Epidemiol*. 2011: 173(5), 492–498. doi:10.1093/aje/kwq386
- 359 16. Okwaraji YB, Edmond KM. Proximity to health services and child survival in
360 low- and middle-income countries: a systematic review and meta-analysis.
361 *BMJ Open*. 2012: 2:e001196, 1–9. doi:10.1136/bmjopen-2012-001196
- 362 17. Karra M, Fink G, Canning D. Facility distance and child mortality: A multi-
363 country study of health facility access, service utilization, and child health
364 outcomes. *Int J Epidemiol*. 2017; 46(3), 817–826. doi:10.1093/ije/dyw062
- 365 18. Rojas-Gualdrón DF, Caicedo-Velázquez B. Distance from healthcare
366 facilities and early life mortality: a systematic review and meta-analysis. *Rev*
367 *Fac Nac Salud Pública*. 2017; 35(3), 420–431.
368 doi:10.17533/udea.rfnsp.v35n3a12
- 369 19. Gruebner O, Khan MMH, Burkart K, Lautenbach S, Lakes T, Kramer A,
370 Subramanian SV, Galea S. Spatial variations and determinants of infant and
371 under-five mortality in Bangladesh. *Health Place*. 2017; 47, 156–164.
372 doi:10.1016/j.healthplace.2017.08.012
- 373 20. Sarrassat S, Meda N, Badolo H, Ouedraogo M, Some H, Bambara R, Murray
374 J, Remes P, Lavoie M, Cousens S, Head R. Effect of a mass radio campaign
375 on family behaviours and child survival in Burkina Faso : a repeated cross-
376 sectional, cluster randomized trial. 2018: 6(3), 330–341. doi:10.1016/S2214-
377 109X(18)30004-4
- 378 21. World Health Organization. Causes of child mortality, by country, 2000-
379 2010.
380 https://www.who.int/gho/child_health/mortality/mortality_causes_text/en/
- 381 22. Weiss DJ, Nelson A, Gibson HS, Temperley W, Peedell S, Lieber
382 A, Hancher M, Poyart E, Belchior S, Fullman N, Mappin B, Dalrymple U,
383 Rozier J, Lucas TCD, Howes RE, Tusting LS, Kang SY, Cameron E,
384 Bisanzio D, Battle KE, Bhatt S, Gething PW. A global map of travel time to
385 cities to assess inequalities in accessibility in 2015. *Nature*. 2018; 553,
386 333–336.
- 387 23. Burstein R, Henry NJ, Collison ML, Marczak LB, Sligar A, Watson S,

388 Marquez N, Abbasalizad-Farhangi M, Abbasi M, Abd-Allah F, Abdoli A,
389 Abdollahi M, Abdollahpour I, Abdulkader RS, Abrigo MRM, Acharya D,
390 Adebayo OM, Adekanmbi V, Adham D, Afshari M, Aghaali M, Ahmadi K,
391 Ahmadi M, Ahmadpour E, Ahmed R, Akal CG, Akinyemi JO, Alahdab F,
392 Alam N, Alamene GM, Alene KA, Alijanzadeh M, Alinia C, Alipour V, Aljunid
393 SM, Almalki MJ, Al-Mekhlafi HM, Altirkawi K, Alvis-Guzman N, Amegah AK,
394 Amini S, Amit AML, Anbari Z, Androudi S, Anjomshoa M, Ansari F, Antonio
395 CAT, Arabloo J, Arefi Z, Aremu O, Armoon B, Arora A, Artaman A, Asadi A,
396 Asadi-Aliabadi M, Ashraf-Ganjouei A, Assadi R, Ataeinia B, Atre SR,
397 Quintanilla BPA, Ayanore MA, Azari S, Babae E, Babazadeh A, Badawi A,
398 Bagheri S, Bagherzadeh M, Baheiraei N, Balouchi A, Barac A, Bassat Q,
399 Baune BT, Bayati M, Bedi N, Beghi E, Behzadifar M, Behzadifar M, Belay
400 YB, Bell B, Bell ML, Berbada DA, Bernstein RS, Bhattacharjee NV, Bhattarai
401 S, Bhutta ZA, Bijani A, Bohlouli S, Breitborde NJK, Britton G, Browne AJ,
402 Nagaraja SB, Busse R, Butt ZA, Car J, C´ardenas R, Castan˜eda-Orjuela
403 CA, Cerin E, Chanie WF, Chatterjee P, Chu D-T, Cooper C, Costa VM, Dalal
404 K, Dandona L, Dandona R, Daoud F, Daryani A, Das Gupta R, Davis I,
405 Davis Weaver N, Davitoiu DV, De Neve J-W, Demeke FM, Demoz GT,
406 Deribe K, Desai R, Deshpande A, Desyibelew HD, Dey S, Dharmaratne SD,
407 Dhimal M, Diaz D, Doshmangir L, Duraes AR, Dwyer-Lindgren L, Earl L,
408 Ebrahimi R, Ebrahimpour S, Effiong A, Eftekhari A, Ehsani-Chimeh E, El
409 Sayed I, El Sayed Zaki M, El Tantawi M, El-Khatib Z, Emamian MH, Enany
410 S, Eskandarieh S, Eyawo O, Ezalarab M, Faramarzi M, Fareed M, Faridnia
411 R, Faro A, Fazaeli AA, Fazlzadeh M, Fentahun N, Fereshtehnejad S-M,
412 Fernandes JC, Filip I, Fischer F, Foigt NA, Foroutan M, Francis JM,
413 Fukumoto T, Fullman N, Gallus S, Gebre DG, Gebrehiwot TT, Gebremeskel
414 GG, Gessner BD, Geta B, Gething PW, Ghadimi R, Ghadiri, K, Ghajarzadeh,
415 M, Ghashghaee, A, Gill, PS, Gill, TK, Golding, N, Gomes, NGM, Gona, PN,
416 Gopalani, SV, Gorini, G, Goulart, BNG, Graetz, N, Greaves, F, Green, MS,
417 Guo, Y, Haj-Mirzaian, A, Haj-Mirzaian, A, Hall, BJ, Hamidi, S, Haririan, H,
418 Haro, JM, Hasankhani, M, Hasanpoor, E, Hasanzadeh, A, Hassankhani, H,
419 Hassen, HY, Hegazy, MI, Hendrie, D, Heydarpour, F, Hird, TR, Hoang, CL,
420 Hollerich, G, Rad, EH, Hoseini-Ghahfarokhi, M, Hossain, N, Hosseini, M,
421 Hosseinzadeh, M, Hostiuc, M, Hostiuc, S, Househ, M, Hsairi, M, Ilesanmi,
422 OS, Imani-Nasab, MH, Iqbal, U, Irvani, SSN, Islam, N, Islam, SMS,
423 Juˆrison, M, Balalami, NJ, Jalali, A, Javidnia, J, Jayatilleke, AU, Jenabi, E,
424 Ji, JS, Jobanputra, YB, Johnson, K, Jonas, JB, Shushtari, ZJ, Jozwiak, JJ,
425 Kabir, A, Kahsay, A, Kalani, H, Kahor, R, Karami, M, Karki, S, Kasaeian, A,
426 Kassebaum, NJ, Keiyoro, PN, Kemp, GR, Khabiri, R, Khader, YS, Khafaie,
427 MA, Khan, EA, Khan, J, Khan, MS, Khang, Y-H, Khatab, K, Khater, A,
428 Khater, MM, Khatony, A, Khazaei, M, Khazaei, S, Khazaei-Pool, M,
429 Khubchandani, J, Kianipour, N, Kim, YJ, Kimokoti, RW, Kinyoki, DK, Kisa,

- 430 A, Kisa, S, Kolola, T, Kosen, S, Koul, PA, Koyanagi, A, Kraemer, MUG,
431 Krishan, K, Krohn, KJ, Kugbey, N, Kumar, GA, Kumar, M, Kumar, P, Kuupiel,
432 D, Lacey, B, Lad, SD, Lami, FH, Larsson, AO, Lee, PH, Leili, M, Levine, AJ,
433 Li, S, Lim, L-L, Stefan, L, Longbottom, J, Lopez, JCF, Lorkowski, S,
434 Magdeldin, S, Abd El Razek, HM, Abd El Razek, MM, Majeed, A, Maleki, A,
435 Malekzadeh, R, Malta, DC, Mamun, AA, Manafi, N, Manda, A-L,
436 Mansourian, M, Martins-Melo, FR, Masaka, A, Massenburg, BB, Maulik, PK,
437 Mayala, BK, Mazidi, M, McKee, M, Mehrotra, R, Mehta, KM, Meles, GG:
438 Mapping 123 million neonatal, infant and child deaths between 2000 and
439 2017. *Nature*. 2019; 574(7778), 353–358. doi:10.1136/bmjopen-2012-
440 001196
- 441 24. Derra K, Rouamba E, Tahita MC, Ouedraogo S, Kazienga A, Sorgho H,
442 Valea I, Tinto H. Profile: Nanoro Health and Demographic Surveillance
443 System. *Int J Epidemiol*. 2012; 41(5), 1293–1301. doi:10.1093/ije/dys159
- 444 25. Andreson PK, Gill RD. Cox's regression model for counting processes: a
445 large sample study. *Ann. Stat.* 1982; 10(4), 1100–1120.
- 446 26. R Core Team: A language and environment for statistical computing; 2013.
447 R Foundation for Statistical Computing, Vienna, Austria. [https://www.r-](https://www.r-project.org/)
448 [project.org/](https://www.r-project.org/) Accessed 2015-09-08
- 449 27. Lohela TJ, Campbell OMR, Gabrysch S. Distance to care, facility delivery
450 and early neonatal mortality in Malawi and Zambia. *PLoS ONE*. 2012; 7(12).
451 doi:10.1371/journal.pone.0052110
- 452 28. Kashima S, Suzuki E, Okayasu T, Louis R, Eboshida A, Subramanian SV.
453 Association between proximity to a health center and early childhood
454 mortality in madagascar. *PLoS ONE*. 2012; 7(6).
455 doi:10.1371/journal.pone.0038370
- 456 29. Kadobera D, Sartorius B, Masanja H, Mathew A, Waiswa P. The effect of
457 distance to formal health facility on childhood mortality in rural Tanzania,
458 2005-2007. *Glob Health Action*. 2012; 5, 1–9. doi:10.3402/gha.v5i0.19099
- 459 30. Si'e A, Louis V, Gbangou A, Müller O, Niamba L, Stieglbauer G, Y'e M,
460 Kouyate B, Sauerborn R, Becher H. The Health and Demographic
461 Surveillance System (HDSS) in Nouna, Burkina Faso, 1993–2007. *Glob*
462 *Health Action*. 2010; 3(1), 5284. doi:10.3402/gha.v3i0.5284
- 463 31. Kynast-Wolf G, Hammer GP, Müller O, Kouyate B, Becher H. Season of
464 death and birth predict patterns of mortality in Burkina Faso. *Int J Epidemiol*.
465 2006; 35(2), 427–435. doi:10.1093/ije/dyi150
- 466 32. Gakidou E, Cowling K, Lozano R, Murray CJL. Increased educational
467 attainment and its effect on child mortality in 175 countries between 1970
468 and 2009: a systematic analysis. *Lancet*. 2010; 376, 959–974.
469 doi:10.1111/tmi.12673
- 470 33. Golding N, Burstein R, Longbottom J, Browne AJ, Fullman N, Osgood-

471 Zimmerman A, Earl L, Bhatt S, Cameron E, Casey DC, Dwyer-Lindgren L,
472 Farag TH, Flaxman AD, Fraser MS, Gething PW, Gibson HS, Graetz N,
473 Krause LK, Kulikoff XR, Lim SS, Mappin B, Morozoff C, Reiner RC, Sligar
474 A, Smith DL, Wang H, Weiss DJ, Murray CJL, Moyes CL, Hay SI. Mapping
475 under-5 and neonatal mortality in Africa, 2000–15: a baseline analysis for
476 the Sustainable Development Goals. *The Lancet*. 2017; 390(10108), 2171–
477 2182. doi:10.1016/S0140-6736(17)31758-0
478 ³⁴. Bado AR, Susuman AS. Determinants of under-5 mortality in Burkina Faso.
479 *J Public Health*. 2018; 41(3):1-11. doi:10.1093/pubmed/fdy168
480

481 **Acknowledgements**

482 The authors wish to thank the individuals of the Nanoro HDSS for their
483 participation and their cooperation in this project, and the Ministry of Health of
484 Burkina Faso for providing support.
485

486 **Funding**

487 This work was supported by Bill and Melinda Gates through Global Good
488 Fund.
489

490 **Author Information**

491 **Affiliations**

492 **¹Institute for Disease Modeling, Building IV, 3150 139th Ave SE, Bellevue, WA,
493 USA, 98005.**

494 Navideh Noori, Assaf P. Oron, Edward Wenger, Andre Lin Ouedraogo

495 **²Institut de Recherche en Sciences de la Sante (IRSS)/Clinical Research Unit of
496 Nanoro (CRUN), Nanoro, Burkina Faso.**

497 Karim Derra, Innocent Valea, Aminata Welgo, Toussaint Rouamba, Palwende
498 Romuald Boua, Athanase SOME, Eli Rouamba, Hermann Sorgho, Halidou Tinto

499 **³Laboratory of Parasitology and Entomology, Centre Muraz, Bobo-Dioulasso,
500 Burkina Faso.**

501 Innocent Valea, Halidou Tinto

502

503 Contributions

504 NN, KD, IV, APO, ALO participated in the design of the analysis. NN and KD
505 conducted the analysis, and NN drafted the original manuscript. All authors revised
506 the manuscript. KD, IV, AW, TR, PRB, AB, ER, HS, HT designed the household
507 survey and collected and cleaned the survey data. IV, APO, ALO supervised the
508 project. HT, EW acquired funding and administered the project. The authors read and
509 approved the final manuscript.

510

511 Corresponding author

512 correspondence to nnoori@idmod.org.

513

514 Ethics declarations

515 The ethical approval was waived by the Burkina Faso Ministry of Health,
516 regarding the general population data of the HDSS (demographics and
517 mortality data) used in this study.

518

519 Consent of publication

520 Not applicable.

521

522 Competing interests

523 The authors declare that they have no competing interests.

524

525 Supplementary Information

526 Supplementary material.pdf

527

528


529

530

531


532

533 **Figure 1:** Nanoro health district is located in the rural center of Burkina Faso.
534 Green dots represent the HDSS households and red crosses represent the
535 health facilities.



536
537
538
539
540
541
542
543
544
545
546

547 **Figure 2** Hazard ratios of multivariable models associated with the probability
548 of mortality of children under-5. Risk factors reducing the probability of death
549 have hazard ratios lower than 1, to the left of the vertical dashed line. Hazard
550 ratios (yellow points), 95% confidence intervals (horizontal lines) and p-values
551 are shown. The variables with the yellow points on the vertical dashed line
552 represent the reference groups.



553
554
555
556
557
558

559 **Table 1** Demographic characteristics of Nanoro HDSS for children under-5, 2009-
560 2013

	Travel time to inpatient health facilities			
	[0-20)min	[20-40)min	[40-60)min	60+min
Number of deaths	142	255	124	94
Number of children	7039	10266	3950	2596
Household size	6 (3,11)*	7 (4,14)	7 (4,13)	9 (4,14)
Number of outpatient health facilities	3	3	6	4
Ratio of female to male children	0.96	0.99	1.04	1.01
Mother's age at birth in year	27.1 (22.3,33)	27.2 (22.4,32.7)	27.2 (22.4,32.7)	27.2 (22.4,32.5)

561 *Median and interquartile range.

562

563

564

565

566 **Table 2.** Results of adjusted and unadjusted Cox regression models for under-5

567 mortality. Hazard Ratios are presented with 95% confidence intervals in parentheses.

Parameters	Mortality per 1,000 live births	Unadjusted Hazard ratio	P-value	Adjusted Hazard ratio	P-value
Age group (month)					
[0 – 12]	327*	-	-	-	-
(12 – 24]	136	0.50 (0.40, 0.63)	< 0.001	0.52 (0.41,0.65)	< 0.001
(24 – 36]	71	0.28 (0.20, 0.39)	< 0.001	0.29 (0.21,0.41)	< 0.001
(36 – 48]	56	0.39 (0.27, 0.56)	< 0.001	0.39 (0.27,0.57)	< 0.001
(48 – 60]	25	0.72 (0.48, 1.06)	0.097	0.75 (0.49-1.13)	0.17
Travel time to inpatient health facilities (min)					
[0 – 20)	50.03	-	-	-	-
[20 – 40)	63.44	1.23 (0.89, 1.68)	0.21	1.24 (0.89,1.71)	0.20
[40 – 60)	79.49	1.51 (1.13, 2.03)	0.006	1.52 (1.13,2.06)	0.005
60+	88.51	1.74 (1.28, 2.38)	< 0.001	1.74 (1.27,2.40)	< 0.001
Seasonality					
Dry Season (Dec-June)	64.49	-	-	-	-
Wet Season (July-Nov)	65.22	1.44 (1.15, 1.81)	0.0018	1.28 (1.07, 1.53)	0.0074
Travel time to outpatient health facilities (min)					
[0 - 5)	58.11	-	-	-	-
[5 – 10)	66.99	1.15 (0.8, 1.65)	0.44	-	-
[10 – 15)	69.72	1.20 (0.87,1.67)	0.26	-	-
[15 – 20)	65.87	1.09 (0.75,1.59)	0.64	-	-
20+	63.41	0.99 (0.58,1.69)	0.99	-	-

568 * Number of deaths within each age group.

569

570