Evolution of DNA methylome from precancerous lesions to invasive lung adenocarcinomas

Xin Hu[1,#], Marcos R. Estecio[2,3,#], Runzhe Chen[4], Alexandre Reuben[4], Linghua Wang[1], Junya Fujimoto[5], Jian Carrot-Zhang[6,7,8], Nicholas McGranahan[9], Lisha Ying[10,11], Junya Fukuoka[12], Chi-Wan Chow[5], Hoa Pham[12], Myrna C.B. Godoy[13], Brett W. Carter[13], Carmen Behrens[4,5], Jianhua Zhang[1], Mara B. Antonoff[14], Boris Sepesi[14], Yue Lu[2,3], Harvey Pass[15], Humam Kadara[5], Paul Scheet[16], Ara A. Vaporciyan[14], John V. Heymach[4], Ignacio I. Wistuba[4,5], J. Jack Lee[17], P. Andrew Futreal[1,&], Dan Su[18,&], Jean-Pierre J. Issa[19,&], Jianjun Zhang[1,4,&]

Affiliations: Departments of [1]Genomic Medicine, [2]Epigenetics and Molecular Carcinogenesis, [3]Center of Cancer Epigenetics, [4]Thoracic/Head and Neck Medical Oncology, [5]Translational Molecular Pathology, The University of Texas MD Anderson Cancer Center, Houston, Texas, 77030, USA.

[6]Broad Institute of MIT and Harvard, Cambridge, MA, 02142, USA.

[7]Department of Medical Oncology, Dana-Farber Cancer Institute, [8]Harvard Medical School, Boston, MA, 02115, USA.

[9]Cancer Research United Kingdom-University College London Lung Cancer Centre of Excellence, London, SW73RP, UK.

[10]Institute of Cancer and Basic Medicine (IBMC), Chinese Academy of Sciences, Hangzhou, 310022, China.

[11]Zhejiang Cancer Research Institute, Cancer Hospital of the University of Chinese Academy of Sciences (Zhejiang Cancer Hospital), Hangzhou, 310022, China.

[12]Department of Pathology, Nagasaki University Graduate School of Biomedical Sciences, Nagasaki, 8528523, Japan.

[13]Thoracic Imaging, [14]Thoracic and Cardiovascular Surgery, the University of Texas MD

Anderson Cancer Center, Houston, Texas, 77030, USA.

[15]Department of Cardiothoracic Surgery, New York University Langone Medical Center,

New York, New York, 10016, USA

[16]Epidemiology, [17]Biostatistics, the University of Texas MD Anderson Cancer Center,

Houston, Texas, 77030, USA.

[18]Department of Pathology, Cancer Hospital of the University of Chinese Academy of

Sciences (Zhejiang Cancer Hospital), Hangzhou, 310022, China.

[19]Coriell Institute for Medical Research, Camden, New Jersey, 08103, USA.

[#]Co-first authors.

[&]Correspondence should be addressed to Jianjun Zhang (jzhang20@mdanderson.org),

Jean-Pierre J. Issa (jpissa@coriell.org), Dan Su (sudan@zjcc.org.cn), or P. Andrew

Futreal (afutreal@mdanderson.org).

Key words: epigenetic intratumor heterogeneity, methylome evolution, immune evasion,

lung precancerous lesions

ABSTRACT

The evolution of DNA methylome and methylation intra-tumor heterogeneity (ITH) during

early carcinogenesis of lung adenocarcinoma has not been systematically studied. We

perform reduced representation bisulfite sequencing of invasive lung adenocarcinoma

and its precursors, atypical adenomatous hyperplasia, adenocarcinoma *in situ* and

minimally invasive adenocarcinoma. We observe gradual increase of methylation

aberrations and significantly higher level of methylation ITH in later-stage lesions. The

phylogenetic patterns inferred from methylation aberrations resemble those based on

somatic mutations suggesting parallel methylation and genetic evolution. De-convolution

reveal higher ratio of T regulatory cells (Tregs) versus CD8+ T cells in later-stage

diseases, implying progressive immunosuppression with neoplastic progression.

Furthermore, increased global hypomethylation is associated with higher mutation

burden, copy number variation burden and allelic imbalance burden as well as higher

Treg/CD8 ratio, highlighting the potential impact of methylation on chromosomal

instability, mutagenesis and tumor immune microenvironment during early

carcinogenesis of lung adenocarcinomas.

INTRODUCTION

Lung cancer remains the leading cause of cancer-related death worldwide, yet it is

curable if treated early. Many cancers, including lung cancers, are preceded by

precancers. Treating precancers to prevent invasive lung cancer is theoretically an

attractive approach to reduce lung cancer–associated morbidity and mortality. However,

developing strategies for lung cancer prevention has been challenging owing to our

limited understanding of neoplastic progression from precancers to invasive lung

cancers[1]. Lung adenocarcinoma (ADC) is the most common histologic subtype,

accounting for more than 50% of all lung cancers[2]. It has been proposed that invasive

lung ADC evolves from atypical adenomatous hyperplasia (AAH), the only recognized

precancer for lung ADC, which could progress to ADC *in situ* (AIS), a pre-invasive lung

cancer, then to minimally invasive ADC (MIA), and eventually frankly invasive ADC[3].

AAH, AIS, MIA and some ADC often present as pulmonary nodules with a unique

radiologic feature termed ground-glass opacity (GGO, hazy nodular opacity with the

preservation of underlying bronchial and vascular margins) on CT scans. The biology

and clinical course of these lesions are not well defined and their management is

controversial. Surgical resection is not the standard of care for treating these lung

nodules and the diagnostic yield of biopsy is often low, particularly for GGO-dominant

nodules[4]. Therefore, these lung nodules are often referred to as indeterminate

pulmonary nodules (IPNs). However, as surgical resection is not often offered, obtaining

adequate tissue for comprehensive profiling of these IPNs is difficult, hindering our

understanding of the biology underlying these lesions.

We initiated an international collaboration for the collection and characterization of these

lung precancers, pre-invasive and early invasive ADC presenting as IPNs. We recently

reported the genomic landscape, including the genomic intra-tumor heterogeneity (ITH),

and revealed evidence of progressive evolution from AAH to AIS, MIA, and ADC[5]. In

addition to mutations, epigenetic alterations such as DNA methylation can also impact

neoplastic transformation and fitness. Recent genome-wide methylation profiling studies

have revealed that certain alterations, such as the silencing of tumor suppressor genes

(TSG) and the activation of genes in stem-like cellular programs[6, 7] may contribute to

carcinogenesis. Our previous study has demonstrated that complex methylation ITH was

associated with larger tumor size and increased risk of postsurgical recurrence in

patients with invasive lung ADC[8]. Methylation aberrations have been reported in AAH

lesions and tumor-adjacent lung tissues suggesting that methylation changes may be

early molecular events[9, 10]. However, these pioneer studies only analyzed small numbers

of genes implicated in lung carcinogenesis. The dynamic changes in methylome at the

genome level and the evolutionary trajectories of methylation ITH during the initiation

and progression of lung precancers have not been studied systematically.

In this study, using a unique cohort of resected IPNs of different histologic stages, we

delineate the evolution of the methylation landscape and reveal increased methylation

ITH in lung ADC than its precursors of early stages, as well as global hypomethylation

correlates with immune infiltration, mutational burden and copy number alterations.

RESULTS

DNA methylation aberrations increase with the progression of precancers

We performed reduced representation bisulfite sequencing (RRBS) of 62 resected IPNs

(14 AAH, 15 AIS, 22 MIA, and 11 invasive ADC) and their paired normal lung tissues

from 39 patients (Supplementary Data 1). There was no significant difference regarding

age (p= 0.6288, Kruskal-Wallis H test), sex (p= 0.6482, Chi-squared test) or smoking

status (p= 0.5696, Chi-squared test) between different histologic groups (Supplementary

Data 2). To minimize the impact of "contamination" from non-malignant cells, each IPN

specimen was reviewed by two lung cancer pathologists to confirm the diagnosis, mark

the areas of diseases and estimate the purity. Only specimens with a minimum of 40%

of premalignant or malignant cells were included and manual macro-dissection was

applied to enrich for premalignant or malignant cells for DNA extraction. Furthermore,

we estimated the tumor purity of these specimens *in silico* based on whole exome

sequencing (WES) data by ABSOLUTE[11]. With the caveat that these IPNs had high level

of genomic ITH leading to underestimation of tumor purity[5], we observed no significant

difference in tumor purity between different stages (Supplementary Fig. 1).

The mean RRBS sequencing coverage was 58.98 reads. With a minimum of 10 reads

shared across all samples, the RRBS profiling allowed the quantification of methylation

status at 751,462 CpG sites mapped to 15,761 known genes. Principal component

analysis (PCA) demonstrated that the DNA methylome of AAH was more similar to that

of normal lung tissue, whereas those of AIS, MIA, and ADC were clearly different from

that of normal lung (Fig. 1a). Similarly, unsupervised hierarchical clustering identified two

separate clusters: one comprising normal lung and AAH, and the other including AIS,

MIA, and ADC (Supplementary Fig. 2). In addition, different regions from the same IPNs

tended to cluster together, suggesting that inter-lesion heterogeneity was more prevalent

than intra-lesion heterogeneity. Similarly, the methylation profiles of different regions

from the same IPNs were highly correlated (median coefficient r=0.884 [range, 0.736-

0.969], p < 2.2 x 10^{-16}, Supplementary Fig. 3).

There was a progressive increase in the number of CpG sites displaying

hypomethylation or hypermethylation (as compared with matched normal lung tissues

from the same patients) from AAH to AIS, MIA, and invasive ADC. Hypermethylation

appeared to emerge as early as AAH, whereas hypomethylation was evident in only AIS,

MIA, and ADC. Meanwhile, the correlation coefficients between the methylation profiles

of IPNs and those of their paired normal tissues progressively decreased (r = 0.885 for

AAH–normal, 0.824 for AIS–normal, 0.796 for MIA–normal, and 0.740 for ADC–normal,

p < 2.2 x 10^{-16}, for all comparisons; Fig. 1b). These decreases were marginal reflecting

the substantial heterogeneity among different patients and suggesting overall

methylation changes may be subtle during early lung carcinogenesis. Nonetheless,

these marginal decreases represent methylation alterations in thousands of CpG sites.

Further quantification of differentially methylated regions (DMRs) in these IPNs

compared with their paired normal lung tissues revealed that later-stage IPNs had

numerically more CpG sites with hypermethylation (p = 0.3635, Kruskal-Wallis test) and

significantly more CpG sites with hypomethylation (p = 0.000287, Kruskal-Wallis test)

(Fig. 1c).

Evolution of methylome shows similar trend across all genomic regulatory regions

The epigenome is composed of regions with various regulatory elements. To further

depict the methylation evolution at different genomic regulatory regions, we first

partitioned genomic regions into promoters, enhancers, transcribed regions

(transcription) and repressed regions (heterochromatin) based on the peaks of histone

marks (H3K04me1, H3K04me2, H3K04me3, H3K09ac, H3K09me3, H3K27ac,

H3K27me3, H3K36me3, and H3K79me2) using ChromHMM[12]. We then calculated the

numbers of DMRs that overlap with these regions. As shown in Fig. 2a-d, the number of

DMRs was higher in later-stage IPNs in all genomic regulatory regions. Next, we selected

DMRs overlapping with repetitive elements (UCSC repeatmasker). Later-stage IPNs

demonstrated increased DMRs in repetitive regions as well as non-repetitive regions (Fig.

2e, f). Furthermore, as partially methylated domains (PMDs) have been reported to

represent a major source of DNA methylation variation in a variety of cancer types[13], we

quantified DMRs inside PMDs versus DMRs outside of PMDs. Similar to the distribution

of DMRs in genomic regulatory regions, there were significantly more DMRs in later

stage IPNs both inside and outside of PMDs (Fig. 2g, h). Taken together, these results

suggested that the majority of methylation alterations during early carcinogenesis of lung

ADC may be stochastic across all genomic regulatory regions.

Enrichment of transcription factor binding sites at DMRs

DNA methylation can impact the DNA binding of transcription factors (TFs) to their target

sequence, termed "motifs"[14]. We searched the motifs that were covered by the DMRs in

the IPNs and identified 50 motifs in AAH, 57 in AIS, 64 in MIA, and 66 in ADC that were

significantly enriched over all DMRs in each histologic stage (Supplementary Data 3).

Many of these motifs were aligned with known binding sites recognized by different TFs

known to be involved in critical biological processes, including cell cycle progression, cell

proliferation, apoptosis, tumor metastasis, angiogenesis and immune response[15, 16, 17, 18]

(Supplementary Data 4). Motifs associated with *BHE40, BMAL1, CLOCK, EPAS1, MAX,*

MITF, MXI1, MYC, MYCN, TFE3, USF1 and *USF2* were significantly enriched at DMRs

across all stages suggesting the methylation changes at these CpG sites may be early

events before the establishment of precancers. On the other hand, some motifs were

significantly enriched in DMRs in later-stage IPNs including *ATF3, E2F1, E2F3, E2F4,*

E2F6, E2F7, EGR1, EGR2, KLF1, KLF12, KLF3, KLF4, KLF6, MYOG, PATZ1, SP1, SP2,

SP3, SP4, TFDP1, ZBT14, ZIC1, ZN281, ZN335 exclusively in ADC; *ASCL1, HEN1,*

KLF9, SNAI2 only in ADC and MIA; *SNAI1* and *TFEB* in ADC, MIA and AIS but not in

AAH, suggesting methylation changes at these CpG sites may be associated with later

processes during carcinogenesis of these IPNs.

Later-stage disease has higher methylation ITH

Molecular ITH can have a profound impact on tumor biology[19, 20], and our previous work

demonstrated that methylation ITH was associated with clinical outcome in patients with

invasive ADC[8]. To assess the evolution of methylation ITH during the progression of lung

precancers, we first calculated the "epiallele shift", the combinatory difference in epiallele

status in all captured loci (>60 reads), for each IPN specimen compared to paired normal

lung tissues from the same patients. MIA and invasive ADC showed more pronounced

epiallele shifts than AAH or AIS did, indicating a higher level of methylation ITH in IPNs

of later histologic stages (Fig. 3a). The corresponding abundance of eloci (loci with

distinct epiallele shifts) was also higher in later-stage IPNs (p = 0.004567, Kruskal–Wallis

test) (Fig. 3b). When focusing on the 15 patients with multiple IPNs of different stages

(IPNs of different stages with the identical genetic background and exposure history),

the abundance of eloci was higher in later-stage IPNs than in early-stage IPNs in 13 of

the 15 patients (Supplementary Fig. 4), further suggesting that later-stage IPNs have a

higher level of methylation ITH. On the other hand, some patients (C23 and J43 for

example) had relative high level of methylation ITH in early stage IPNs. Since the post-

operative follow up was rather short and the majority of patients are doing well without

recurrence, the biological or clinical significance of higher level of methylation ITH in

early stage IPNs remains to be determined. Furthermore, we calculated the frequency

of 16 combinatory DNA methylation patterns at four consecutive CpG sites covered by

the same RRBS reads (termed "epipolymorphism") in each specimen. Accordingly, later-

stage IPNs showed higher epiallele diversity than early-stage IPNs did (Supplementary

Fig. 5). Moreover, ADCs had higher levels of epipolymorphism at eloci than MIA, AIS, or

AAH did (Fig. 3c), indicating that DNA methylation status had undergone a greater extent

of drifting in late-stage IPNs than in early-stage IPNs.

In addition, we sought to assess whether methylation ITH differs at different genomic

regulatory regions. As shown in Fig. 4a, loci at heterochromatic regions show the most

pronounced epiallele shifts indicating a higher level of methylation ITH at genomic

regions of repressed chromatin states. Moreover, there were more eloci inside PMDs

than outside of PMDs across all histologic stages (Fig. 4b-e). PMDs are known to be

more epigenetically plastic[21], which makes CpG sites inside PMDs more prone to

develop methylation ITH. Additionally, PMDs have been reported to associate with

repressive chromatin domains, gene-poor regions and low transcription[21, 22], therefore,

higher level of methylation ITH may be better tolerated inside PMDs.

We next examined the relative distance of eloci to the nearest transcription start sites

(TSSs). Interestingly, the vast majority of the eloci were much closer to TSSs in invasive

ADC than in AAH, AIS, or MIA (Supplementary Fig. 6). To explore the potential

association between methylation ITH and histone modification, we performed Locus

Overlap Analysis (LOLA) [23] of the genomic regions identified as eloci to evaluate the

overlap between genomic regions with methylation ITH and genomic regions targeted

by diverse histone posttranslational modifications[24, 25]. LOLA calculates the number of

overlapping versus non-overlapping regions to assess the significance of the overlap.

After adjustment for false discovery rate considering more eloci in later-stage IPNs, MIA

and ADC had higher incidences of eloci significantly enriched in genomic regions

occupied by H3K27me3, H3K9me3, and H3K9me2 (Supplementary Data 5), histone

modifications strongly associated with transcriptional repression[26, 27, 28], than AAH or AIS,

supporting the concept that histone modifications cooperate with DNA methylation

alterations along the evolution of lung precancers[29, 30].

Genomic and methylation evolution were primarily in parallel during early lung

carcinogenesis

To dissect the evolutionary relationship between the epigenome and genome in lung

ADC, we constructed phylogenetic trees. To avoid overfitting, only 5 patients (C5, J7, J8,

J9 and J43) with IPNs having a minimum of four spatially separated specimens were

included. The overall structure of methylation-based phylogenetic trees was similar to

that of phylogenetic trees based on mutations[5] (Fig. 5a-e). Furthermore, the genomic

distance based on somatic mutations was positively correlated with the methylation

distance between any pair of specimens from the same IPNs (Fig. 5f) suggesting parallel

evolution (either collaboratively or independently) in play during early carcinogenesis of

these lung ADCs. Particularly, 4 out of 5 patients with multi-region specimens (C5, J7,

J8, J9 and J43) included for phylogenetic analysis demonstrated strong correlation

between methylation distance and genomic distance with the only exception of patient

J9 (Supplementary Fig. 7). Of note, J9 was a never smoker with the lowest genomic

aberration burden (the lowest total mutation burden (TMB), lowest allelic imbalance (AI)

burden and the second lowest copy number variation (CNV) burden) and the highest

level of genomic ITH (the lowest proportion of trunk mutations) among these 5 tumors,

but similar methylation aberration burden comparable to the other 4 tumors

(Supplementary Data 6).

Interestingly, in patient C10, promoter hypermethylation of *TSC2*, a candidate TSG

known to inhibit cell growth in lung[31], was identified in AIS specimens, whereas copy

number loss was identified in AAH lesions from the same patient. Similar phenomena of

different putative TSGs were observed in several other patients (Supplementary Data 7).

Taken together, these data suggested convergent evolution, whereby the same genes

or pathways are activated or inactivated by different mechanisms in different cancer cell

clones during lung cancer development and progression.

Global hypomethylation and methylation ITH were associated with increased

chromosomal instability in precancerous IPNs

As an essential chemical modification, the methylation status can directly impact the

chromosomal structure and DNA mutagenesis. It has been well documented that global

hypomethylation is associated with chromosomal instability (CIN) and increased

mutational rates in cancers[32, 33]. To further depict the interaction between genome and

epigenome during early carcinogenesis of lung ADC, we assessed the global

methylation status of these IPNs using long interspersed transposable elements-1

(LINE-1), a widely used surrogate marker for global DNA methylation[34]. As shown in Fig.

6a, we observed a significant decrease of LINE-1 methylation in AIS, MIA and ADC

compared to normal lung tissues or AAH indicating increased global hypomethylation in

IPNs of later histologic stages. Importantly, methylation level of LINE-1 was inversely

correlated with CNV burden (Fig. 6b), AI burden (Fig. 6c) and TMB (Fig. 6d) indicating

global hypomethylation is associated with higher level of CIN. Interestingly, LINE-1

methylation status was also inversely correlated with the proportion of clonal mutations

in each specimen (Fig. 6e). Furthermore, there was a significant positive correlation

between the abundance of eloci and CNV burden as well as AI burden (Supplementary

Fig. 8), indicating higher methylation ITH was associated with increased CIN.

Given the complex interaction between genome and epigenome, we next categorized

CpG sites located in chromosomal regions with various CNV status to assess whether

CNV status impacted the results of methylation analyses. As shown in Supplementary

Fig. 9, the evolutionary patterns of methylation ITH (represented by the number of eloci)

were similar regardless of using CpG sites located in genomic regions with copy number

gain, loss or neutral. We also plotted cumulative distribution of epiallele shifts of AAH,

AIS, MIA, and ADC using only CpG sites located in copy number-neutral chromosomal

regions and observed that later-stage IPNs had higher cumulative distribution of epiallele

shifts than early-stage IPNs (Supplementary Fig. 10), similar to that using all CpG sites

(Fig. 3a). Furthermore, we regenerated methylation-based phylogenetic trees using only

CpG sites located in copy number-neutral chromosomal regions. As shown in

Supplementary Fig. 11, the patterns of these methylation-based phylogenetic trees are

almost identical to those using all CpG sites. Taken together, these results suggested

that in these IPNs, most methylation changes were early molecular events that have

occurred before the copy number changes and once established, these patterns were

inherited without significant changes from one cell generation to the next.

Global hypomethylation was associated with suppressed T cell infiltration

Cancer evolution is shaped by interaction between cancer cells and host factors,

particularly the host immune response. Given T cells' central role in anti-tumor immune

surveillance[20, 35], we depicted the T cell infiltration in AIS, MIA and invasive ADC by de-

convoluting RNAseq data from an independent dataset recently published[36]. Our

analysis demonstrated an increase of CD4+ T regulatory cells (Tregs) (p=4.25e-27) and

decrease of CD8+ T cells (although the difference was not significant, p=0.1374) from

normal lung tissues to AIS/MIA and invasive ADC leading to significantly higher

Treg/CD8 ratio in invasive ADC (p=3.17e-31) (Supplementary Fig. 12a-c). We next

applied MethylCIBERSORT[37] to delineate the T cell infiltration of IPNs in our cohort.

Similarly, we observed higher infiltration of Tregs (Fig. 7a, p=0.009758) and lower

infiltration of CD8+ T cells in later-stage IPNs although the difference did not reach

statistical significance (Fig. 7b, p=0.1472), leading to significantly higher Treg/CD8 ratio

in later-stage IPNs (Fig. 7c, p=0.00194). As increased Treg/CD8 ratio is known to

associate with suppressed anti-tumor immune surveillance[38], these results indicated

more suppressive immune microenvironment in later stage IPNs, consistent with our

previous findings[39]. Interestingly, Treg/CD8 ratio was inversely correlated with LINE-1

methylation level (Fig. 7d) implying the potential association between global

hypomethylation and immunosuppression.

DISCUSSION

The methylation landscape has been studied extensively in various malignancies[40, 41, 42].

However, methylation aberrations in precancers are poorly defined, largely due to lack

of appropriate specimens. Using small panels of genes implicated in lung carcinogenesis,

previous studies have demonstrated gradual changes in DNA methylation in AAH, AIS,

and ADC[9, 10]. However, due to small number of loci, primitive technology and lack of

genomic data, many critical questions regarding methylation evolution during early

carcinogenesis were not addressed. Leveraging a unique collection of resected IPNs of

different stages, we delineated the DNA methylome in lung precancers, preinvasive ADC

and invasive ADC. Our results demonstrated an increase in both hypermethylation and

hypomethylation in later-stage diseases. Interestingly, compared to normal lung tissues,

hypermethylation appeared to emerge as early as at AAH, whereas hypomethylation

only became obvious after AIS (Fig. 1b), implying that somatic hypermethylation may

have preceded hypomethylation during early carcinogenesis of lung ADC.

Different cells within the same tumor can exhibit different molecular and phenotypic

features, a phenomenon termed ITH. ITH may foster tumor evolution by providing

diverse cell populations, and the dynamics of ITH architecture may evolve with

neoplastic progression[8, 43]. Methylation ITH has been observed in various advanced

malignancies and higher levels of methylation ITH have been reported to associate with

inferior clinical outcomes[8, 44, 45, 46]. However, there are only a few reports on methylation

ITH in precancers. For example, methylation ITH in Barrett esophagus, a precursor to

esophageal ADC, was associated with the risk of malignant transformation[46]. In the

current study, we demonstrated higher level of methylation ITH in later-stage IPNs than

in early-stages (Fig. 3, Supplementary Fig. 4, 5). These results are in line with findings in

advanced malignancies suggesting that complex methylation ITH may be associated with

more aggressive tumor biology. As methylation ITH is linked to evolutionary plasticity and

phenotypic diversity[44, 47, 48], higher level of ITH could provide higher probability to survive

and progress. Interestingly, eloci were significantly more abundant around TSSs in ADC

than in AAH, AIS, or MIA. One plausible explanation is that although somatic methylation

ITH may be stochastic during early lung carcinogenesis, some of the methylation

aberrations (particularly those that are close to TSSs and potentially impact gene

expression) may convey survival and/or growth advantages, resulting in selection of cells

with higher densities of eloci around TSSs.

Parallel evolution of genome and methylome has been reported in various advanced

malignancies including lung cancers[8, 49]. The current study demonstrated similar

phylogenetic patterns and correlated genetic and methylation distances in IPNs of

different stages (Fig. 5) suggesting genetic alterations and DNA methylation changes

also evolve in parallel during early carcinogenesis of most lung ADCs. Meanwhile,

promoter hypermethylation and copy number loss or mutations of putative TSGs were

identified in independent IPNs within the same patients. These results were reminiscent

of previous findings showing that distinct mutations of the same cancer genes were

present in different regions of the same tumors[19, 50, 51] or in different primary tumors from

the same patients[52], indicating convergent evolution. Although genetic events (e.g., copy

number loss of TSGs) or methylation changes (e.g. promoter hypermethylation of TSGs)

may independently or cooperatively offer proliferation or survival advantages to cells,

these processes may be constrained around certain genes or pathways (e.g.,

inactivation of *TSC2* in the case of patient C10) that are essential to carcinogenesis in

certain patients.

In our cohort, global hypomethylation was found to associate with significantly higher

TMB, CNV burden as well as AI burden (Fig. 6b-d), consistent with previous reports that

global hypomethylation is associated with CIN[32] and increased rate of somatic

mutations[33]. These data suggest that methylation aberrations have not only evolved in

parallel with genomic aberrations but may have also facilitated accumulating genomic

alterations that may have led to more drastic phenotypic changes in IPNs of later stages.

Interestingly, higher-level of global hypomethylation was associated with higher

proportion of clonal mutations (Fig. 6e). As the progression of lung precancers into

invasive lung ADC predominantly follows a clonal sweeping model with selective

outgrowth of fit subclones[5], one plausible explanation for this association is that cells

within the precancers with higher level of global hypomethylation may be prone to

accumulate genomic aberrations, which subsequently provide growth advantages to

these cell clones to develop into major clones in invasive lung ADC.

Cancer evolution results from accumulation of molecular alterations and is constantly shaped by selection pressure such as anti-tumor immune surveillance and therapeutic interventions. These molecular aberrations such as point mutations may accumulate gradually, a model termed microevolution, or in a punctuated or catastrophic manner through processes such as chromoplexy and chromothripsis, a model termed macroevolution, both of which have been reported in advanced malignancies[53, 54]. Largely due to lack of appropriate study materials, our understanding of the molecular evolutionary pattern during early carcinogenesis of lung ADC is rudimentary. Our previous study on genomic profiling of lung ADC and its precursors has demonstrated progressive accumulation of somatic mutations from AAH to AIS, MIA and ADC in line with the microevolution model. Meanwhile, there was a distinct increase in CNV burden from AAH to AIS and increase in AI burden from AIS to MIA. In the current study, we observed progressive increase of methylation changes from AAH to AIS, MIA and ADC. However, the overall difference in methylation aberrations appeared to be subtle between IPNs of different histologic stages (Fig.1b) and methylation evolution from AAH to ADC was similar across different epigenetically defined genomic regulatory regions. These observations suggest that methylation aberrations may have primarily contributed to the stepwise microevolution during early carcinogenesis of these lung ADCs and most methylation changes were stochastic "passengers", as are the majority of somatic mutations[55]. On the other hand, these seemingly stochastic genomic and epigenetic alterations may give rise to heterogeneous subclones in precancers with various biological features, therefore increase the possibility of establishing fit subclones leading to malignant transformation. These principles may also apply to later neoplastic evolution including invasion, metastasis and development of drug resistance, where large-scale genomic sequencing studies have only depicted the underlying molecular mechanisms

in a small subset of patients[56, 57]. Comprehensive molecular profiling incorporating

genomic, epigenomics and transcriptomic profiling are warranted in future studies to

depict these critical cancer evolutionary processes.

Anti-tumor immune surveillance plays a central role during initiation and progression of

precancers. We have previously reported that the immune microenvironment was

suppressed in invasive lung cancers compared to preinvasive cancers or precancers[58].

In this study, we demonstrated higher Treg/CD8 ratio in later stage IPNs (Fig. 7c,

Supplementary Fig. 12c) implying a more suppressed T cell infiltrate in later-stage

diseases, in line with a concomitant study of immune profiling of the same cohort of

IPNs[59]. These findings are consistent with the concept of immune-editing, whereby the

immunogenicity of cancer cells evolves under the selective pressure from anti-tumor

immune response, resulting in the emergence of immune-resistant cancer clones in later

stage diseases. Interestingly, global hypomethylation was associated with higher

Treg/CD8 ratio (Fig. 7d). Methylation aberrations may affect anti-tumor immune

surveillance directly by regulating the expression of immune-related genes[60] and/or

potential neoantigens or indirectly via modifying chromosomal vulnerability for CNV and

mutations, both of which are well known to influence the tumor immune

microenvironment[61, 62]. However, these impacts are complicated as many processes can

affect anti-tumor immune surveillance both positively and negatively. For example, a high

level of global hypomethylation may lead to a high CNV burden known to associate with

a cold tumor immune microenvironment[61]; meanwhile global hypomethylation is also

associated with increased mutation rate, which may increase tumor immunogenicity[62].

In the end, selection of cancer cell clones under immune pressure is determined by the

cumulative effects of these molecular aberrations and only the cells with the best

combination of molecular features including methylation status, mutation and CNV

burden will survive and develop into dominant clones in invasive cancers.

There has been increasing enthusiasm toward moving interventions successfully applied

to metastatic cancers to early stage cancers and even precancers, a concept called

interception[63]. Compared with invasive cancers, precancers and preinvasive cancers

may exhibit less complexity in aberrant molecular landscapes, as well as better

preserved immune contextures, and thus may be easier to eradicate. Accordingly, we

have launched the IMPRINT-Lung clinical trial (NCT03634241), in which patients with

high-risk IPNs (many of which may be AAH or AIS) are treated with immune checkpoint

inhibitors. In the current study, we demonstrated that DNA methylation aberrations are

less complex in precancers and preinvasive lung cancers than in invasive cancers.

Therefore, therapeutic agents that can modulate methylation by targeting aberrant

methylations and potentially reprogram the immune microenvironment may also have

potential in treating precancers and preinvasive cancers to prevent invasive lung cancers.

We delineated the evolution of genome-wide DNA methylation during the early

carcinogenesis of lung ADC using RRBS on invasive lung ADC precursors of different

stages. Rather than aggregated changes from populations of cancer cells measured by

methylation array, RRBS assesses DNA methylation heterogeneity of single molecules

derived from individual cancer cells, which made it possible to delineate methylation ITH

at the allelic level. Furthermore, the WES data from the same IPNs has provided a unique

opportunity to depict the relationship between methylation and genomic evolution. Due

to the scarcity of IPN materials, our study has inevitable limitations. First, the sample

size was relatively small for each histologic stage. Given the substantial heterogeneity

between IPNs even within the same stages, our data was not powered to address some

essential questions relevant to methylation evolution, such as whether certain

methylation aberrations are significantly more common in early-stage IPNs than later-

stage lesions representing "dead-end" IPNs. Second, most of these IPNs were very

small after pathological assessment, which prevented acquisition of sufficient data in

more patients for in-depth analyses such as multi-region profiling to dissect the

interaction between parallel evolution and CIN; transcriptomic profiling to determine

biological impact of observed methylation changes. Although a substantial number of

DMRs were associated with TF binding sites, which could potentially regulate the

transcriptome of the IPNs, these remained to be speculation without confirmation from

transcriptomic data. Third, follow-up time for all patients in this study was relatively short,

so we were unable to investigate the impact of these methylation changes on recurrence

or survival. Finally, the resected specimens in this study could only provide a single

molecular snapshot of the evolutionary process of IPNs. Whether all AAH will evolve into

AIS, MIA, and ADC; whether all ADC evolve from AAH; and whether the observed

methylation changes in IPNs of different stages represent the true evolutionary dynamics

or simply reflect the distinct methylation patterns of IPNs with different malignant

potentials is unknown. Deciphering the temporal evolution during neoplastic progression

will require specimens obtained over the course of disease progression. Clinical trials

collecting longitudinal biopsy specimens, such as IMPRINT-Lung (NCT03634241), may

provide such opportunities going forward.

METHODS

Sample acquisition

A total of 53 resected pulmonary nodules and paired normal lung tissues from 39 patients

treated at Nagasaki University Hospital or Zhejiang Cancer Hospital between 2014 and

2017 were used in the study. None of the patients received chemotherapy or

radiotherapy before surgery. 29 lung nodules from 15 patients had multiregional

specimens for spatial heterogeneity assessment (Supplementary Data 1). Whole-exome

sequencing (WES) data was available for all specimens (EGAS00001004960)[5].

DNA methylation profiling by RRBS and data processing

DNA was extracted using the QIAamp DNA FFPE Tissue Kit (QIAGEN), and 200ng−1μg

of DNA was subjected to RRBS for genome-wide DNA methylation profiling[64]. Briefly,

TrimGalore v.0.4.3 was used to trim the Illumina adapter sequences (a minimum of 5 bp

in a read was required to overlap with the adapter sequence); then Bismark v.0.18.1

integrating bowtie2 v.2.2.3 was used to align the trimmed reads to the GRCh37 assembly

of the human genome. FastQC v.0.11.7 was used for quality control. The DNA

methylation levels for individual CpGs were calculated using methyKit (v.1.16.0)[65].

Methylated reads (containing Cs) and unmethylated reads (containing Ts) at each

cytosine site were counted, and the percentage of methylated reads among total reads

covering the corresponding cytosine was calculated to quantify the DNA methylome for

each sample at the single-base resolution. The CpG sites, DMRs, and loci of known

genes, as well as genomic features, including CpG islands, were annotated using the R

package "ChIPSeeker (v.1.26.0)"[66] and the toolkit "genomation" (v.1.22.0)[67], which is

based on the "TxDb.Hsapiens.UCSC.hg19.knownGene" annotation database and the

UCSC Genome Browser CpG islands table. To avoid bias, we kept CpG sites mapped

to the autosome and removed CpG sites overlapping with the single-nucleotide

polymorphism positions in dbSNP137. DNA methylation was analyzed at either a single-

CpG resolution or at genomic region bins, in which DNA methylation values were

averaged across 5-kb regions. Promoter methylation was calculated as the averaged

DNA methylation values based on GENCODE promoter regions (i.e., 1 kb upstream to

500 bp downstream of the annotated TSS).

Comparison of methylation profiles between different specimens

To assess the DNA methylome profiles of distinct IPNs of different pathological stages and examine the heterogeneity between IPN samples, we first aggregated the DNA methylation levels of 5-kb tiling regions across the genome in each sample by retaining only the CpG sites with ≥10 sequencing reads. We then applied principal component analysis to identify global DNA methylation patterns between samples. To evaluate consistent clusters of these DNA methylation profiles, we performed unsupervised hierarchical agglomerative analysis of CpG sites covered by ≥50 reads across all samples; these reads were based on single CpG methylation calls without any binning. To evaluate the correlation between overall DNA methylation in all samples from each patient, we used a pairwise approach to compare distance and similarity matrices on the basis for all CpGs with coverage of ≥10 reads.

Differential DNA methylation analysis

Differentially methylated regions (DMRs) encompassing the differentially methylated CpGs between paired disease and normal tissue samples were identified using a triangular kernel to smooth the number of methylated reads and total number of reads by applying the "noise filter" function in the "DMR caller" (v.1.22.0) R package[68]. The differentially methylated CpG of paired samples were identified by selecting CpG sites located in DMRs; only CpG sites covered by ≥10 reads in paired samples were included.

Partition of genomic regulatory regions based on chromatin states model

ChromHMM v.1.21 is applied to build the chromatin states model at 200 bp resolution with default parameters[12] on A549 cell line (Encode Broad) for histone marks: H3K04me1, H3K04me2, H3K04me3, H3K09ac, H3K09me3, H3K27ac, H3K27me3,

H3K36me3, and H3K79me2. The genomic regions including promoter regions, enhancer

regions, transcribed regions and repressed regions (heterochromatin) were partitioned

based on resulting segments inferred as defined chromatin states, which were deduced

from chromatin-state signatures using a multivariate hidden Markov model (HMM) that

explicitly models the combinatorial presence or absence of each histone mark.

Repetitive elements

The repetitive elements were retrieved from UCSC repeatmasker track

(https://genome.ucsc.edu/cgi-bin/hgTrackUi?g=rmsk), which was created using Arian

Smit's RepeatMasker program, through screening human genomic DNA sequences for

interspersed repeats and low complexity DNA sequences.

Identification of partially methylated domains (PMDs)

The boundary of partially methylated domains (PMDs) were inferred using whole

genome shotgun bisulfite sequencing (WGSBS) of lung cells from a healthy donor

(GSM983647) applying MethylSeekR (v.1.30.0) package[69]. Before PMD calling, CpGs

overlapping common SNPs (dbSNP build 137) were removed.

Motif identification and prediction of TF binding sites

The de novo methylated DNA motifs in IPNs of each stage were identified by mEpigram

v.0.07, which discovers motifs by using position-specific weight matrices from the k-mers

that are most enriched in the positive sequences compared with the negative sequences

as "seeds" and extending the motifs in both directions[70]. The Tomtom tool (v.5.2.0) from

the MEME suite was used to select significantly enriched methylated DNA motifs based

on the database of known transcript factors (HOCOMOCO_v11)[71].

Estimation of DNA methylation ITH

To estimate DNA methylation ITH, we applied "methclone" (v.0.1) to identify epigenetic

loci whose distributions of epigenetic allele ("epiallele") clonality differed between paired

tumor and normal samples by quantifying the degree to which the compositions of

epialleles at given loci in the tumor samples were distinct from those in the normal tissue

samples. An epiallele was defined by setting 60 reads in four consecutive CpG sites as

the threshold to consider the epigenetic allele composition of the locus. We then

calculated the differences in epiallele entropy between each IPN sample and its matched

normal tissue sample. Loci with combinatorial entropy changes (ΔS) below −60 between

each IPN sample and its paired normal sample were defined as epigenetic shift loci

(termed "eloci"). To reduce the bias due to the different coverage for each sample, we

then calculated relative epiallele shifts (i.e., the normalized number of eloci) via dividing

the number of eloci by the total number of assessed loci in each sample and then

multiplying that ratio by the average number of total loci across all samples[72]. We also

assessed epigenetic polymorphism ("epipolymorphism") to measure the epiallelic

diversity in each IPN sample by calculating the frequency of each specific epiallele from

multiple stochastic changes in the frequencies of many epialleles[73]. We calculated 16

epiallele status (0000, 0100, 0010, 0001, 1000, 1100, 0110, 0011, 0101, 1010, 1001,

0111, 1011, 1101, 1110, and 1111, where 1 represents a methylated CpG site, and 0

represents an unmethylated CpG site), then epipolymorphism for each loci is defined as

$\sum_{i=1}^{16} Fi^2$, where F is the fraction of each epiallele status.

Locus Overlap Analysis

We applied Locus Overlap Analysis (LOLA) [23] to the genomic regions identified as eloci

($\Delta S < -60$) in all samples of each stage to evaluate the overlap between genomic regions

with methylation ITH and chromatin marks. The genomic regions of all loci with ≥ 60

reads were used as background genomic regions, and the selected genomic regions

were mapped to a compendium of publicly available histone mark profiles, including

CTCF, H2AZ, H3K4me1, H3K4me2, H3K4me3, H3K9ac, H3K9me3, H3K27me3, H3K27ac,

H3K36me3, H3K79me2, H4K20me1 in the A549 lung adenocarcinoma cell line

(http://hgdownload.soe.ucsc.edu/goldenPath/hg19/encodeDCC/wgEncodeBroadHistone) and

H3K27me3, H3K4me3, H3K9me2, H3K9ac, CTCF in an immortalized human bronchial

epithelial cell line (BEAS-2B; GSE56053)[25]. P-values in the enrichment analyses were

calculated using one-sided Fisher's exact test. Adjustment for multiple testing with q-

value (FDR adjusted p-value) was performed using the Benjamini–Yekutieli method.

Construction of phylogenetic trees

To construct methylation-based phylogenetic trees from the RRBS data, we used

promoter CpG sites (≥50 reads per CpG site selected for all samples from each patient)

with the most variable methylation values (mean absolute deviation >10%) shared by all

samples, including a normal tissue sample used as the tree root, to build a Euclidean

distance matrix. We built the phylogenetic trees by applying a neighbor-joining algorithm

from the "ape" (v.5.4.1) package to independently infer phylogenetic relationships

between IPN specimens for each patient from the mutation and methylation profiles. To

assess the phylogenetic similarity between the genetic and epigenetic profiles, we

employed an independent but parallel distance matrix construction from the mutation

and methylation profiles for each IPN, in which the genetic distance was quantified by

Hamming distance based on all filtered mutations and methylation distance was

quantified by Euclidean distance based on methylation values of all CpG sites (≥20 reads,

MAD≥20 for all samples in each IPN), then calculated the Spearman correlation

coefficient for all pairwise samples grouped by IPNs.

Estimation of global hypomethylation

To determine global methylation levels, we chose CpG sites (covered by ≥10 aligned reads) within evolutionarily young subfamilies of LINE-1 repeat elements (L1HS and L1PA). LINE-1 family annotation were obtained from the Repeat-Masker of the UCSC genome browser. We averaged the methylation values of the chosen CpG sites, to represent the global methylation level of each sample.

Deconvolution of T cell profiles

To derive tumor-infiltrating T cell subtypes from transcriptomic data previously published[36], processed RNAseq dataset (normalized and log2 transformed) comprise of 197 normal lung tissues, 98 AIS/MIA and 99 ADC samples were retrieved from EGAS00001004006. ImmuCellAI[74] was applied to infer immune cell components for each sample.

To derive the infiltration of T cell repertoire from RRBS data, we first obtained a reference methylation signature by retrieving the methylation calls of whole-genome bisulfite datasets (WGBS) from the BLUEPRINT epigenome project, including those for samples of regulatory T cells (EGAX00001343016/EGAX00001236257 in EGAD00001002492) and CD8+ T cells (EGAX00001195937/EGAX00001195943 in EGAD00001002486). Only promoter CpG sites that were covered by ≥5 reads in all samples were retained and binned with a 50-bp window by mean methylation values, then only 50-bp binned regions that overlapped with 50-bp binned regions of CpG sites covered by the RRBS methylation profiles of preneoplastic lesions in all samples were used for reference signature extraction by non-negative matrix factorization (NMF) implemented in the "MethylCIBERSORT" (v.0.2.0) package. We then performed DNA methylation

deconvolution (average methylation level by 50-bp bin) using the aforementioned

BLUEPRINT signature and the CIBERSORT webserver (https://cibersort.stanford.edu).

We performed T cell deconvolution in relative mode, running 100 permutations with

quantile normalization disabled[75]. The resulting immune-cellular fractions were used to

compare samples of different pathological stages. The average values of immune cell

infiltration for each sample inferred by whole-genome bisulfite analysis from two patients

independently were used to infer Treg/CD8 ratio.

Statistical analysis

Violin plots were created using the "geom_violin" function in R statistical package ggplot2

(v0.9.1) to represent data point density along the *Y*-axis, and the "stat_summary"

function from ggplot2 (v0.9.1) was used to calculate the mean as the center point.

Differences in DMR numbers, eloci numbers, and immune cell infiltration between IPN

specimens of different stages were assessed using the Kruskal–Wallis chi-square test.

We used quantile regression to assess the differential entropy values at 1th percentile

(low percentile of entropy was chosen to reflect the loci with biological interest) and

epipolymorphism values at 50th percentile between different groups. We used a two-

sided Pearson correlation coefficient to compare methylation profiles between two

samples and between groups of samples of different stages. We used a two-sided

Spearman correlation coefficient to determine the extent to which distance matrices were

correlated with DNA methylation profiles and somatic mutation profiles.

Data availability

The RRBS dataset has been deposited at European Bioinformatics Institute European

Genome–phenome Archive (EGA) (accession number: EGAS00001004610,

https://www.ebi.ac.uk/ega/studies/EGAS00001004610) through controlled access.

Whole-exome sequencing is under the accession code: EGAD00001004960

[https://www.ebi.ac.uk/ega/datasets/EGAD00001004960]. To protect patient privacy, interested

researchers need to apply via data access committee (DAC), which will grant all

reasonable requests. GSM983647 (Whole Genome Shotgun Bisulfite Sequencing of

human normal lung cells) was downloaded from GEO

[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSM983647].

EGAS00001004006 (RNA-seq of human preinvasive and invasive lung adenocarcinoma)

was downloaded from EGA [https://www.ebi.ac.uk/ega/studies/EGAS00001004006].

EGAD00001002492 (Bisulfite-seq of regulatory T cells from human venous blood)

[https://www.ebi.ac.uk/ega/datasets/EGAD00001002492] and

EGAD00001002486 (Bisulfite-seq of CD8+ T cells from human venous blood)

[https://www.ebi.ac.uk/ega/datasets/EGAD00001002486] was downloaded from EGA,

respectively. And source data are provided with this study. All other data may be found

within the main manuscript or supplementary information or available from the authors

upon request.

FIGURE LEGENDS

Figure 1. DNA methylation aberrations increase with progression of precancers. (a)

Principal component analysis of the methylation profiles of the average methylation

values in 39,148 5-kb tiling regions (shared in all samples) composed of autosomal, non-

polymorphism CpG sites supported by at least 10X read coverage in IPNs of different

stages. The solid dots of different colors represent IPNs of different stages and the star

(*) represented the centers of specimens of each histologic stage. (b) Overall genome-

wide DNA methylation correlation between IPNs of different stages and matched normal

tissue from the same patients. The yellow-purple clouded dots in the smooth scatter plot

represent the CpG sites covered by IPN specimens (n = 14 for AAH, n = 11 for AIS, n =

18 for MIA, n = 10 for ADC) and paired normal lung. Two-tailed Pearson's correlation

coefficient r values are shown on the top. P-values < 2. 2 × 10^{-16} in all 4 histologic stages.

The red dots represent CpG sites with hypermethylation in IPNs (within DMRs of

methylation gain in IPNs and methylation ≤20% of CpG sites in the corresponding

normal lung) and the green dots represent CpG sites with hypomethylation (within DMRs

of methylation loss in IPNs and methylation ≥20% of CpG sites in the corresponding

normal lung). (c) The number of CpG sites overlapping with DMRs showing

hypermethylation (left) or hypomethylation (right) in IPNs of different histologic stages.

The green dots represent the mean numbers of CpG sites overlapping with DMRs in

each IPN. The solid blue dots represent the mean numbers of CpG sites overlapping

with DMRs in IPNs of each histologic stage with 95% confidence interval as error bars.

The differences among all stages were assessed using the Kruskal-Wallis H test.

(Source data are provided as a source data file.)

Figure 2. The number of DMRs in IPNs of different stages by genomic regulatory regions

and PMDs. The number of DMRs are shown in promoters (a), enhancers (b), transcribed

regions (c), repressed (heterochromatic) regions (d), repetitive regions (e), non-

repetitive regions(f), inside PMDs (g) and outside PMDs (h). The solid blue dots

represent the means of normalized numbers of DMRs in IPNs of each histologic stage

with 95% confidence interval as error bars. (Source data are provided as a source data

file.)

Figure 3. Increased DNA methylation ITH in later-stage IPNs. (a) Cumulative distribution

curves of epiallele shifts of the DNA methylome in AAH, AIS, MIA, and ADC specimens

compared to normal lung, with 1th percentile of entropy as -21.46 for AAH, -28.88 for

AIS, -30.75 for MIA, -42.01 for ADC by quantile regression. (b) The abundance of eloci

(loci with distinct epiallele shifts) among IPNs of different histologic stages. Each green

dot represents the average number of eloci in each IPN. The solid blue dots represent

the mean numbers of eloci in the IPN of each histologic stage with 95% confidence

interval as error bars. The differences among all stages were assessed using the

Kruskal-Wallis H test. (c) The cumulative distribution curves of epipolymorphism for loci

with significant epiallele shifts (ΔS< -60) in IPN specimens of AAH, AIS, MIA, and ADC,

with 50th percentile of epipolymorphism as 0.295 for AAH, 0.362 for AIS, 0.359 for MIA,

0.463 for ADC by quantile regression. *X*-axis denotes the epipolymorphism in IPN

specimens at each loci. *Y*-axis denotes cumulative fraction of epipolymorphism from all

IPN specimens of each stage. (Source data are provided as a source data file.)

Figure 4. Comparison of methylation ITH by epiallele shifts in different genomic

regulatory regions and PMDs. (a) Cumulative distribution curves of epiallele shifts (from

tumor sample versus its matched normal sample) of consecutive loci located in

promoters, enhancers, transcribed regions, repressed (heterochromatic) regions are

shown, with 1th percentile of entropy as -34.07 for all genomic regions, -30.87 for

promoters, -38.12 for enhancer, -42.44 for transcribed regions, -45.48 for repressed

regions, -29.51 for repetitive regions by quantile regression. Cumulative distribution of

epiallele shifts (from IPN sample versus its matched normal sample) of consecutive loci

located inside PMDs (red) versus outside PMDs (blue), with 1th percentile of entropy as

-29.10 inside PMDs and -20.73 outside PMDs in AAH (b), -37.60 inside PMDs and -

28.05 outside PMDs in AIS (c), -44.74 inside PMDs and -29.25 outside PMDs in MIA (d),

-50.82 inside PMDs and -41.25 outside PMDs in ADC (e) by quantile regression. The

boundary of PMDs are derived from WGBS profiling of normal lung. (Source data are

provided as a source data file.)

Figure 5. The evolutionary relationship between genomic and methylation landscape.

Phylogenetic trees based on mutations (blue) and methylation values (green) in patient

C5 (a), J7 (b), J8 (c), J9 (d), J43 (e). The length of each branch indicates the similarity

of mutational or methylation profiles between any pair of two spatially separated tumor

specimens from each patient. To avoid overfitting, only patients with IPNs having a

minimum of four spatially separated specimens were included for this analysis. (f)

Correlation of genetic distance (Hamming distance based on all mutations) and

methylation distance (Euclidean distance based on methylation values of all CpG sites)

between different spatially separated specimens from the same IPNs assessed by two-

tailed Spearman's correlation analysis ($p = 2.62 \times 10^{-38}$). Each dot represents the

normalized distance between each pair of specimens from the same IPNs. (Source data

are provided as a source data file.)

Figure 6. Correlation of LINE-1 methylation with genomic features in IPNs of different

stages. (a) LINE-1 methylation level in IPNs of different stages. Each green dot

represents LINE-1 methylation level in each IPN and the blue dots represent the means

of LINE-1 methylation level in IPNs of each histologic stage with 95% confidence

intervals as error bars. The differences among all stages were assessed using the

Kruskal-Wallis H test. Correlation between LINE-1 methylation levels and percent of

genes with copy number changes (b), number of events with allelic imbalance (AI) (c),

mutational burden (log2 transformed) (d), proportion of clonal mutations (e), assessed

by two-tailed Spearman's correlation analysis. Each dot represents each IPN specimen.

(Source data are provided as a source data file.)

Figure 7. T cell infiltration in IPNs of different stages. The immune cell fraction of T

regulatory cells (Tregs) (a), CD8+ T-cells (b), Treg/CD8 ratio (c) in IPNs of different

stages. Each green dot represents immune cell fraction in each IPN and the blue dots

represent the means of immune cell fraction in IPNs of each histologic stage with 95%

confidence intervals as error bars. The differences among all stages were assessed

using the Kruskal-Wallis H test. Infiltration of T lymphocytes was inferred by

MethylCIBERSORT. (d) Correlation between LINE-1 methylation and Treg/CD8 ratio

was assessed by Spearman correlation analysis. Each dot represents each specimen.

(Source data are provided as a source data file.)

AUTHOR CONTRIBUTIONS

X.H. and J.J.Z. conceived the study and wrote the manuscript. J.J.Z. and P.A.F. jointly

supervised and financially supported the study. X.H. performed all bioinformatics and

statistical data analyses in consultation with M.R.E., J.P.I., L.H.W., J.J.L., and Y.L.; J.F.J.

supervised pathological assessments and the preparation of specimens. L.S.Y., J.F.K.,

and H.P. collected resected specimens and clinical data. C.W.C., C.B, and R.Z.C.

prepared DNA samples. M.C.G. and B.W.C. performed radiological assessment. M.R.E.

supervised RRBS profiling. X.H., M.R.E., R.Z.C., A.R., N.M., L.H.W., J.C.Z., J.H.Z., P.S.,

A.A.V., J.V.H., I.I.W., H.K., M.B.A, B.S., H.P., J.P.I., D.S., P.A.F., and J.J.Z. interpreted

the data. All authors reviewed the manuscript.

COMPETING INTERESTS

Dr. Zhang reports research funding from Merck, Johnson and Johnson, and consultant

fees from BMS, Johnson and Johnson, AstraZeneca, Geneplus, OrigMed, Innovent

outside the submitted work. Dr. Heymach reports research funding from AstraZeneca,

GlaxoSmithKline, and Spectrum; consultant fees from AstraZeneca, Boehringer

Ingelheim, Bristol-Myers Squibb, Catalyst, EMD Serono, Foundation Medicine, Hengrui

Therapeutics, Genentech, GSK, Guardant Health, Eli Lilly, Merck, Novartis, Pfizer,

Roche, Sanofi, Seattle Genetics, Spectrum, and Takeda; licensing fees from Spectrum.

Dr. Kadara reports funding from Johnson and Johnson and from Janssen

Pharmaceuticals. Dr. Sepesi reports consultant fees from BMS. The other authors

declare neither financial nor non-financial interests in the submitted work.

ACKNOWLEDGMENTS

This study was supported by the MD Anderson Khalifa Scholar Award, the National

Cancer Institute of the National Institute of Health Research Project Grant

(R01CA234629-01), the AACR-Johnson & Johnson Lung Cancer Innovation Science

Grant (18-90-52-ZHAN), the MD Anderson Physician Scientist Program, the MD

Anderson Lung Cancer Moon Shot Program, Sabin Family Foundation Award, Duncan

Family Institute Cancer Prevention Research Seed Funding Program, The University of

Texas MD Anderson Cancer Center Pre-Cancer Atlas Project, EDRN U01

(U01CA214195), the Cancer Prevention and Research Institute of Texas Multi-

Investigator Research Award grant (RP160668). We thank MD Anderson Cancer

Center's Epigenomics Profiling Core and its Science Park Next-Generation Sequencing

Core (supported by CPRIT Core Facility Support Award #RP120348) for performing

RRBS profiling. We thank Sally Boyd, Jinzhen Chen, and Rong Yao, Stan Bujnowski,

Eric Sisson for providing excellent technical support for high-performance cluster

resource; we thank Joe Munch from Scientific Publications in MD Anderson's Research

Medical Library for editing the manuscript.

References

1.   Kadara H, Scheet P, Wistuba, II, Spira AE. Early Events in the Molecular Pathogenesis of
Lung Cancer. *Cancer Prev Res (Phila)* **9**, 518-527 (2016).

827
828   2.   Chen Z, Fillmore CM, Hammerman PS, Kim CF, Wong KK. Non-small-cell lung cancers: a
829   heterogeneous set of diseases. *Nat Rev Cancer* **14**, 535-546 (2014).

830
831   3.   Lee HJ*, et al.* IASLC/ATS/ERS International Multidisciplinary Classification of Lung
832   Adenocarcinoma: novel concepts and radiologic implications. *J Thorac Imaging* **27**,
833   340-353 (2012).

834
835   4.   Chen D, Dai C, Kadeer X, Mao R, Chen Y, Chen C. New horizons in surgical treatment of
836   ground-glass nodules of the lung: experience and controversies. *Ther Clin Risk Manag*
837   **14**, 203-211 (2018).

838
839   5.   Hu X*, et al.* Multi-region exome sequencing reveals genomic evolution from
840   preneoplasia to lung adenocarcinoma. *Nat Commun* **10**, 2978 (2019).

841
842   6.   Jones PA, Baylin SB. The epigenomics of cancer. *Cell* **128**, 683-692 (2007).

843
844   7.   Widschwendter M*, et al.* Epigenetic stem cell signature in cancer. *Nat Genet* **39**, 157-
845   158 (2007).

846
847   8.   Quek K*, et al.* DNA methylation intratumor heterogeneity in localized lung
848   adenocarcinomas. *Oncotarget* **8**, 21994-22002 (2017).

849
850   9.   Kerr KM, Galler JS, Hagen JA, Laird PW, Laird-Offringa IA. The role of DNA methylation
851   in the development and progression of lung adenocarcinoma. *Dis Markers* **23**, 5-30
852   (2007).

853
854   10.   Selamat SA*, et al.* DNA methylation changes in atypical adenomatous hyperplasia,
855   adenocarcinoma in situ, and lung adenocarcinoma. *PLoS One* **6**, e21443 (2011).

856
857   11.   Carter SL*, et al.* Absolute quantification of somatic DNA alterations in human cancer.
858   *Nat Biotechnol* **30**, 413-421 (2012).

859
860   12.   Ernst J, Kellis M. Chromatin-state discovery and genome annotation with ChromHMM.
861   *Nat Protoc* **12**, 2478-2492 (2017).

863   13.   Salhab A, *et al.* A comprehensive analysis of 195 DNA methylomes reveals shared and
864   cell-specific features of partially methylated domains. *Genome Biol* **19**, 150 (2018).

865
866   14.   Yin Y, *et al.* Impact of cytosine methylation on DNA binding specificities of human
867   transcription factors. *Science* **356**, (2017).

868
869   15.   Casey SC, Baylot V, Felsher DW. The MYC oncogene is a global regulator of the
870   immune response. *Blood* **131**, 2007-2015 (2018).

871
872   16.   Wils LJ, Bijlsma MF. Epigenetic regulation of the Hedgehog and Wnt pathways in
873   cancer. *Crit Rev Oncol Hematol* **121**, 23-44 (2018).

874
875   17.   Yu T, *et al.* KLF4 regulates adult lung tumor-initiating cells and represses K-Ras-
876   mediated lung cancer. *Cell Death Differ* **23**, 207-215 (2016).

877
878   18.   Wu C, *et al.* WT1 promotes invasion of NSCLC via suppression of CDH1. *J Thorac Oncol*
879   **8**, 1163-1169 (2013).

880
881   19.   Zhang J, *et al.* Intratumor heterogeneity in localized lung adenocarcinomas delineated
882   by multiregion sequencing. *Science* **346**, 256-259 (2014).

883
884   20.   Reuben A, *et al.* Comprehensive T cell repertoire characterization of non-small cell
885   lung cancer. *Nat Commun* **11**, 603 (2020).

886
887   21.   Gaidatzis D, *et al.* DNA sequence explains seemingly disordered methylation levels in
888   partially methylated domains of Mammalian genomes. *PLoS Genet* **10**, e1004143
889   (2014).

890
891   22.   Lochs SJA, Kefalopoulou S, Kind J. Lamina Associated Domains and Gene Regulation in
892   Development and Cancer. *Cells* **8**, (2019).

893
894   23.   Sheffield NC, Bock C. LOLA: enrichment analysis for genomic region sets and regulatory
895   elements in R and Bioconductor. *Bioinformatics* **32**, 587-589 (2016).

896
897   24.   Consortium EP. An integrated encyclopedia of DNA elements in the human genome.
898   *Nature* **489**, 57-74 (2012).

899
900   25.   Jose CC, *et al.* Epigenetic dysregulation by nickel through repressive chromatin domain
901   disruption. *Proc Natl Acad Sci U S A* **111**, 14631-14636 (2014).

903   26.   Lienert F*, et al.* Genomic prevalence of heterochromatic H3K9me2 and transcription
904   do not discriminate pluripotent from terminally differentiated cells. *PLoS Genet* **7**,
905   e1002090 (2011).

906
907   27.   Kondo Y*, et al.* Gene silencing in cancer by histone H3 lysine 27 trimethylation
908   independent of promoter DNA methylation. *Nat Genet* **40**, 741-750 (2008).

909
910   28.   Hon GC*, et al.* Global DNA hypomethylation coupled to repressive chromatin domain
911   formation and gene silencing in breast cancer. *Genome Res* **22**, 246-258 (2012).

912
913   29.   Ohm JE*, et al.* A stem cell-like chromatin pattern may predispose tumor suppressor
914   genes to DNA hypermethylation and heritable silencing. *Nat Genet* **39**, 237-242 (2007).

915
916   30.   Schlesinger Y*, et al.* Polycomb-mediated methylation on Lys27 of histone H3 pre-marks
917   genes for de novo methylation in cancer. *Nat Genet* **39**, 232-236 (2007).

918
919   31.   Astrinidis A, Cash TP, Hunter DS, Walker CL, Chernoff J, Henske EP. Tuberin, the
920   tuberous sclerosis complex 2 tumor suppressor gene product, regulates Rho
921   activation, cell adhesion and migration. *Oncogene* **21**, 8470-8476 (2002).

922
923   32.   Eden A, Gaudet F, Waghmare A, Jaenisch R. Chromosomal instability and tumors
924   promoted by DNA hypomethylation. *Science* **300**, 455 (2003).

925
926   33.   Chen RZ, Pettersson U, Beard C, Jackson-Grusby L, Jaenisch R. DNA hypomethylation
927   leads to elevated mutation rates. *Nature* **395**, 89-93 (1998).

928
929   34.   Saito K, Kawakami K, Matsumoto I, Oda M, Watanabe G, Minamoto T. Long
930   interspersed nuclear element 1 hypomethylation is a marker of poor prognosis in
931   stage IA non-small cell lung cancer. *Clin Cancer Res* **16**, 2418-2426 (2010).

932
933   35.   Reuben A*, et al.* TCR Repertoire Intratumor Heterogeneity in Localized Lung
934   Adenocarcinomas: An Association with Predicted Neoantigen Heterogeneity and
935   Postsurgical Recurrence. *Cancer Discov* **7**, 1088-1097 (2017).

936
937   36.   Chen H*, et al.* Genomic and immune profiling of pre-invasive lung adenocarcinoma.
938   *Nat Commun* **10**, 5472 (2019).

939
940   37.   Chakravarthy A*, et al.* Pan-cancer deconvolution of tumour composition using DNA
941   methylation. *Nat Commun* **9**, 3220 (2018).

943   38.   Peng GL*, et al.* CD8(+) cytotoxic and FoxP3(+) regulatory T lymphocytes serve as
944   prognostic factors in breast cancer. *Am J Transl Res* **11**, 5039-5053 (2019).

945
946   39.   Chen RZ*, et al.* T cell repertoire evolution from the normal lung to invasive lung
947   adenocarcinoma. *Cancer Research* **78**, (2018).

948
949   40.   Cancer Genome Atlas Research N. Comprehensive molecular profiling of lung
950   adenocarcinoma. *Nature* **511**, 543-550 (2014).

951
952   41.   Saghafinia S, Mina M, Riggi N, Hanahan D, Ciriello G. Pan-Cancer Landscape of
953   Aberrant DNA Methylation across Human Tumors. *Cell Rep* **25**, 1066-1080 e1068
954   (2018).

955
956   42.   Chakravarthy A*, et al.* Pan-cancer deconvolution of tumour composition using DNA
957   methylation. *Nature Communications* **9**, (2018).

958
959   43.   Prasetyanti PR, Medema JP. Intra-tumor heterogeneity from a cancer stem cell
960   perspective. *Mol Cancer* **16**, 41 (2017).

961
962   44.   Landau DA*, et al.* Locally disordered methylation forms the basis of intratumor
963   methylome variation in chronic lymphocytic leukemia. *Cancer Cell* **26**, 813-825 (2014).

964
965   45.   Hua X*, et al.* Genetic and epigenetic intratumor heterogeneity impacts prognosis of
966   lung adenocarcinoma. *Nat Commun* **11**, 2459 (2020).

967
968   46.   Merlo LM*, et al.* A comprehensive survey of clonal diversity measures in Barrett's
969   esophagus as biomarkers of progression to esophageal adenocarcinoma. *Cancer Prev*
970   *Res (Phila)* **3**, 1388-1397 (2010).

971
972   47.   Siegmund KD, Marjoram P, Woo YJ, Tavare S, Shibata D. Inferring clonal expansion and
973   cancer stem cell dynamics from DNA methylation patterns in colorectal cancers. *Proc*
974   *Natl Acad Sci U S A* **106**, 4828-4833 (2009).

975
976   48.   Hansen KD*, et al.* Increased methylation variation in epigenetic domains across cancer
977   types. *Nat Genet* **43**, 768-775 (2011).

978
979   49.   Aryee MJ*, et al.* DNA methylation alterations exhibit intraindividual stability and
980   interindividual heterogeneity in prostate cancer metastases. *Sci Transl Med* **5**,
981   169ra110 (2013).

983   50.   Jamal-Hanjani M*, et al.* Tracking the Evolution of Non-Small-Cell Lung Cancer. *N Engl J*
984   *Med* **376**, 2109-2121 (2017).

985
986   51.   consortium TRR. TRACERx Renal: tracking renal cancer evolution through therapy. *Nat*
987   *Rev Urol* **14**, 575-576 (2017).

988
989   52.   Liu Y*, et al.* Genomic heterogeneity of multiple synchronous lung cancer. *Nat Commun*
990   **7**, 13200 (2016).

991
992   53.   Shen MM. Chromoplexy: a new category of complex rearrangements in the cancer
993   genome. *Cancer Cell* **23**, 567-569 (2013).

994
995   54.   Cortes-Ciriano I*, et al.* Comprehensive analysis of chromothripsis in 2,658 human
996   cancers using whole-genome sequencing. *Nat Genet* **52**, 331-341 (2020).

997
998   55.   Flavahan WA, Gaskell E, Bernstein BE. Epigenetic plasticity and the hallmarks of cancer.
999   *Science* **357**, (2017).

1000
1001   56.   Zheng JB*, et al.* DNA methylation affects metastasis of renal cancer and is associated
1002   with TGF-beta/RUNX3 inhibition. *Cancer Cell International* **18**, (2018).

1003
1004   57.   Le X*, et al.* Landscape of EGFR-Dependent and -Independent Resistance Mechanisms
1005   to Osimertinib and Continuation Therapy Beyond Progression in EGFR-Mutant NSCLC.
1006   *Clin Cancer Res* **24**, 6195-6203 (2018).

1007
1008   58.   Sivakumar S*, et al.* Genomic Landscape of Atypical Adenomatous Hyperplasia Reveals
1009   Divergent Modes to Lung Adenocarcinoma. *Cancer Res* **77**, 6119-6130 (2017).

1010
1011   59.   Dejima H*, et al.* Immune evolution from preneoplasia to invasive lung
1012   adenocarcinomas and underlying molecular features. *medRxiv*,
1013   2020.2007.2011.20142992 (2020).

1014
1015   60.   Liu M, Zhou J, Chen Z, Cheng AS. Understanding the epigenetic regulation of tumours
1016   and their microenvironments: opportunities and problems for epigenetic therapy. *J*
1017   *Pathol* **241**, 10-24 (2017).

1018
1019   61.   Davoli T, Uno H, Wooten EC, Elledge SJ. Tumor aneuploidy correlates with markers of
1020   immune evasion and with reduced response to immunotherapy. *Science* **355**, (2017).

1021
1022   62.   Jung H*, et al.* DNA methylation loss promotes immune evasion of tumours with high
1023   mutation and copy number load. *Nat Commun* **10**, 4278 (2019).

1024

1025   63.   Blackburn EH. Cancer interception. *Cancer Prev Res (Phila)* **4**, 787-792 (2011).

1026
1027   64.   Gu H, Smith ZD, Bock C, Boyle P, Gnirke A, Meissner A. Preparation of reduced
1028   representation bisulfite sequencing libraries for genome-scale DNA methylation
1029   profiling. *Nat Protoc* **6**, 468-481 (2011).

1030
1031   65.   Akalin A*, et al.* methylKit: a comprehensive R package for the analysis of genome-wide
1032   DNA methylation profiles. *Genome Biol* **13**, R87 (2012).

1033
1034   66.   Yu G, Wang LG, He QY. ChIPseeker: an R/Bioconductor package for ChIP peak
1035   annotation, comparison and visualization. *Bioinformatics* **31**, 2382-2383 (2015).

1036
1037   67.   Akalin A, Franke V, Vlahovicek K, Mason CE, Schubeler D. Genomation: a toolkit to
1038   summarize, annotate and visualize genomic intervals. *Bioinformatics* **31**, 1127-1129
1039   (2015).

1040
1041   68.   Catoni M, Tsang JM, Greco AP, Zabet NR. DMRcaller: a versatile R/Bioconductor
1042   package for detection and visualization of differentially methylated regions in CpG and
1043   non-CpG contexts. *Nucleic Acids Res* **46**, e114 (2018).

1044
1045   69.   Burger L, Gaidatzis D, Schubeler D, Stadler MB. Identification of active regulatory
1046   regions from DNA methylation data. *Nucleic Acids Res* **41**, e155 (2013).

1047
1048   70.   Ngo V, Wang M, Wang W. Finding de novo methylated DNA motifs. *Bioinformatics* **35**,
1049   3287-3293 (2019).

1050
1051   71.   Bailey TL*, et al.* MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res*
1052   **37**, W202-208 (2009).

1053
1054   72.   Li S*, et al.* Dynamic evolution of clonal epialleles revealed by methclone. *Genome Biol*
1055   **15**, 472 (2014).

1056
1057   73.   Landan G*, et al.* Epigenetic polymorphism and the stochastic formation of differentially
1058   methylated regions in normal and cancerous tissues. *Nat Genet* **44**, 1207-1214 (2012).

1059
1060   74.   Miao YR*, et al.* ImmuCellAI: A Unique Method for Comprehensive T-Cell Subsets
1061   Abundance Prediction and its Application in Cancer Immunotherapy. *Adv Sci (Weinh)*
1062   **7**, 1902880 (2020).

1063

1064   75.   Chen B, Khodadoust MS, Liu CL, Newman AM, Alizadeh AA. Profiling Tumor Infiltrating
1065   Immune Cells with CIBERSORT. *Methods Mol Biol* **1711**, 243-259 (2018).

1066

1067

Figure 1

a

b

c

Figure 2

Figure 3

a

Cumulative distribution vs Epiallele shifts. Legend: AAH, AIS, MIA, ADC.

b

p-value = 0.004567 (Kruskal–Wallis test)

log2 (Normalized number of eloci) vs Stages (AAH, AIS, MIA, ADC). Stages legend: AAH, AIS, MIA, ADC.

c

Cumulative fraction vs Epipolymorphism at eloci. Legend: AAH, AIS, MIA, ADC.

Figure 4

a

b
AAH

c
AIS

d
MIA

e
ADC

Figure 5

a

C5C1-2B C5C1-2

C5C1-3
C5C1-3B

C5C1-1B

C5C1-1

C5M1-2

Normal

C5C1-2B C5C1-2

C5C1-3
C5C1-3B

C5C1-1B
C5C1-1

C5M1-2

Normal

b

J7C1-2

J7C1-4

J7C1-6

J7C1-3

J7C1-1

J7C1-5

J7H1-1B

Normal

J7C1-2 J7C1-6
J7C1-4
J7C1-1

J7C1-3

J7C1-5

J7H1-1B
Normal

c

J8C1-6 J8C1-3

J8C1-1

J8C1-4

J8C1-5

J8C1-2

J8H1-1B

Normal

J8C1-6 J8C1-3
J8C1-4
J8C1-1
J8C1-5

J8C1-2

J8H1-1B
Normal

d

J9C1-6
J9C1-5

J9C1-3

J9C1-4

J9C1-2

J9C1-1

J9H1-1

J9H1-2

Normal

J9C1-6

J9C1-3

J9C1-5

J9C1-4   J9C1-2

J9C1-1

J9H1-1
J9H1-2

Normal

e

J43M2-1 J43M2-2

J43S1-2

J43M1-2

J43S1-1

J43M1-1

J43S2-2

Normal

J43S1-2 J43S1-1   J43M2-1 J43M2-2
J43S2-2   J43M1-2
                                    J43M1-1

Normal

f

Rho = 0.7283
p-value = 2.62e-38

Methylation distance

Mutation distance

● C5_ADC	● J38_AIS	● J44_MIA	● J7_ADC
● J20_AIS	● J43_AIS	● J45_MIA	● J8_ADC
● J23_MIA	● J43_MIA	● J5_ADC	● J9_ADC

Figure 6

Figure 7