

COVID-19 misinformation: mere harmless delusions or much more?

A knowledge and attitude cross-sectional study among the general public residing in Jordan

Authors

Malik Sallam^{1,2,3,*}, Deema Dababseh⁴, Alaa' Yaseen¹, Ayat Al-Haidar⁴, Duaa Taim⁴, Huda Eid⁴, Nidaa A. Ababneh⁵, Faris G. Bakri^{6,7,8}, and Azmi Mahafzah^{1,2}

Affiliations

¹Department of Pathology, Microbiology and Forensic Medicine, School of Medicine, the University of Jordan, Amman, Jordan.

²Department of Clinical Laboratories and Forensic Medicine, Jordan University Hospital, Amman, Jordan.

³Department of Translational Medicine, Faculty of Medicine, Lund University, Malmö, Sweden.

⁴School of Dentistry, the University of Jordan, Amman, Jordan.

⁵Cell Therapy Center (CTC), the University of Jordan, Amman, Jordan.

⁶Department Internal Medicine, School of Medicine, the University of Jordan, Amman, Jordan.

⁷Department of Internal Medicine, Jordan University Hospital, Amman, Jordan.

⁸Infectious Diseases and Vaccine Center, University of Jordan, Amman, Jordan.

*Corresponding author:

Malik Sallam, M.D., Ph.D., Department of Clinical Laboratories and Forensic Medicine, Jordan University Hospital, Queen Rania Al-Abdullah Street Aljubeiha/P.O. Box: 13046, Postal code: 11942. Amman, Jordan. Tel: +962 79 184 5186. E-mail: *malik.sallam@ju.edu.jo*, ORCID ID: 0000-0002-0165-9670.

Abstract

Since the emergence of the recent coronavirus disease 2019 (COVID-19) and its spread as a pandemic, media was teeming with misinformation that led to psychological, social and economic consequences among the global public. Probing knowledge and anxiety regarding this novel infectious disease is necessary to identify gaps and sources of misinformation which can help public health efforts to design and implement more focused interventional measures. The aim of this study was to evaluate the knowledge, attitude and effects of misinformation about COVID-19 on anxiety level among the general public residing in Jordan. An online survey was used that targeted people aged 18 and above and residing in Jordan. The questionnaire included items on the following: demographic characteristics of the participants, knowledge about COVID-19, anxiety level and misconceptions regarding the origin of the pandemic. The total number of participants included in final analysis was 3150. The study population was predominantly females (76.0%), with mean age of 31 years. The overall knowledge of COVID-19 was satisfactory. Older age, male gender, lower monthly income and educational levels, smoking and history of chronic disease were associated with perceiving COVID-19 as a very dangerous disease. Variables that were associated with a higher anxiety level during the pandemic included: lower monthly income and educational level, residence outside the capital (Amman) and history of smoking. Misinformation about the origin of the pandemic (being part of a conspiracy, biologic warfare and the 5G networks role) was also associated with higher anxiety and lower knowledge about the disease. Social media platforms, TV and news releases were the most common sources of information about the pandemic. The study showed the potential harmful effects of misinformation on the general public and emphasized the need to meticulously deliver timely and accurate information about the pandemic to lessen the health, social and psychological impact of the disease.

Introduction

The entire world is facing an unprecedented challenge in the form of the most recent pandemic caused by severe acute coronavirus 2 (SARS-CoV-2) [1]. Coronavirus disease 2019 (COVID-19) resulted in a massive number of infections throughout the world with a higher mortality rate among high risk groups (elderly, those with comorbidities) [2]. The public is left in a state of disarray due to the socio-economic consequences of the pandemic [3-5]. This global phenomenon dominated the media and became part of everyday conversation [6,7]. The emergence of this virus led to a worldwide lockdown, army enforced rules, disruption of education and a shift in the global economy [4]. These events caused public apprehension that raised questions and spread rumors, which demonstrates the significance of providing correct knowledge and reliable information for proper management of this public health emergency [5,8].

The clinical manifestations of COVID-19 vary, but commonly include: fever, cough, shortness of breath, vomiting and diarrhea [9-11]. The virus is primarily transmitted via respiratory droplets and close contact with an infected person [11]. SARS-CoV-2 can remain active for hours and even days on surfaces, therefore, touching infected surfaces can lead to the spread of infection [12,13]. This is why frequent hand washing and social distancing are the ideal protective measures [14]. To date, there are limited therapeutic options and no vaccine available for COVID-19 infection, and management hinges on supportive therapy [15].

Conspiracy theories regarding the origins of COVID-19 might be a way for the public to make sense of this pandemic. However, dangerous speculations about the virus might diminish the efforts in controlling the spread of infection [8,16]. Thus, it is important to assess the misconceptions and misbeliefs among the public which can reveal defects that should be targeted by awareness tools [17].

The potential negative effects of COVID-19 misinformation have been the subject of active research since the onset of the pandemic [8,18-22]. Our previous investigation of this topic entailed students at the University of Jordan (UJ) with results pointed to

an association between the belief in conspiracy regarding the origin of the virus and a lower COVID-19 knowledge accompanied with higher anxiety level [23].

Deciphering the level of knowledge and attitude toward this unprecedented pandemic can help in identifying the current gaps in knowledge about COVID-19. Resources must be utilized to bridge this gap and promote proper knowledge about COVID-19, which in turn will help in disease control. This project aimed to explore the repercussions of misinformation regarding conspiracy beliefs about this pandemic. Thus, the objectives of the current study were: (1) to assess the overall COVID-19 knowledge and attitude among the general public residing in Jordan, (2) to evaluate the effects of misinformation regarding COVID-19 origins on the anxiety level, and (3) to assess the main sources of knowledge regarding the disease in the country.

Methods

Study design

This cross-sectional study was conducted using an online-based questionnaire that took place between April 11, 2020 (21:00) to April 14, 2020 (00:00), thus spanning 75 hours and targeting residents in Jordan aged 18 years and above. Participation in the study was voluntary and an informed consent was included. The questionnaire was sent through Facebook, WhatsApp, Twitter and other social media platforms. The questionnaire comprised six sections with a total of 39 items addressing various subjects regarding knowledge, attitude, misinformation, sources of knowledge, and anxiety of participants regarding COVID-19. The language used to conduct the survey was Arabic (S1 Appendix).

The questionnaire contained items on socio-demographic information (age, nationality, gender, governorate of residence, marital status, monthly family income, educational level, history of smoking, and the presence of any chronic disease). Four items were used to assess the attitude towards the quarantine period (perception of the danger of the disease, adherence to quarantine measures, spending quality time with the family, and annoyance by the inability to attend religious houses of worship).

Two items were used to assess the sources of information, and three items were included to determine the role of conspiracy theories, biological warfare, and 5G networks in the origin and the spread of the pandemic. An additional item was also included to examine the belief in a divine role in the origin of the disease (S1 Appendix).

Ethical permission

The study was approved by the Department of Pathology, Microbiology and Forensic Medicine and by the Scientific Research Committee at the School of Medicine/UJ, (using WhatsApp conference call) which was later registered under the reference number 2479/2020/67 at the School of Medicine/UJ. Participation in the study was voluntary and anonymous. An informed consent was ensured by the presence of an

introductory section of the questionnaire, with submission of responses implying the agreement to participate. All collected data were treated confidentially.

COVID-19 knowledge score (K-score) calculation

Thirteen items were used to assess the overall COVID-19 knowledge among the study participants. These items included knowledge of symptoms of the disease (fever, cough, vomiting and diarrhea, and shortness of breath), knowledge of virus transmission (touching infected surfaces, close contact with an infected person, and transmission via blood), infectivity of the virus on surfaces for long periods of time, use of antibiotics for treating the disease, availability of a vaccine, remedial effect of garlic, onion and ginger on the infection, ability of summer heat to inactivate the virus, and possibility of reinfection by the virus (S1 Appendix). Each correct answer was recorded as a single point and a valid K-score was considered upon having responses to all 13 items.

Assessment of the anxiety score

The final section of seven items was used to measure the level of anxiety during the government-enforced quarantine period using the General Anxiety Disorder-7 (GAD-7) scale [24]. This scale is a reliable method for anxiety assessment and included four possible responses to each item. A valid anxiety score was considered upon having responses to all seven items. The maximum possible anxiety score was 21 with the minimum being zero.

Statistical analysis

All statistical analyses were conducted in IBM SPSS v22.0 for Windows. Significance was considered for P-values <0.050. Chi-squared (χ^2), Mann-Whitney *U* (M-W) and Kruskal Wallis (K-W) tests were used as appropriate.

Results

Characteristics of the study population

The total number of individuals who participated in the survey and that were included in final analysis was 3150 after filtering out responses from those who were less than 18 years old. This resulted in 1.75% margin of error considering the 95% confidence interval and the current total population of Jordan (10,184,790 people) [25,26]. General features of the study participants are summarized in (Table 1). The median age of the study participants was 27 (mean: 31, interquartile range: 22-37). Females dominated the study population (n=2358, 76.0% with 47 cases of non-response), and residents of the Central region of Jordan represented 84.1% of the participants (with 78 cases of non-response). For educational level and monthly income, the majority of study participants had an undergraduate degree (diploma or bachelor's degrees) representing 73.6% and the majority had a monthly income of less than 1000 JODs (n=2413, 78.3%, Table 1).

Table 1. Socio-demographic characteristics of the study population.


Feature		Number	Percentage
Gender	<i>Male</i>	745	24.0%
	<i>Female</i>	2358	76.0%
Nationality	<i>Jordanian</i>	2894	92.8%
	<i>Non-Jordanian</i>	223	7.2%
Region¹	<i>North</i>	375	12.2%
	<i>Central</i>	2584	84.1%
	<i>South</i>	113	3.7%
Marital status	<i>Single</i>	1617	51.8%
	<i>Married</i>	1414	45.3%
	<i>Divorced</i>	59	1.9%
	<i>Widow/widower</i>	29	0.9%
Monthly income	<i>Less than 500 JOD³</i>	1233	40.0%
	<i>500-1000 JOD</i>	1180	38.3%
	<i>More than 1000 JOD</i>	668	21.7%
Educational level²	<i>High school or less</i>	496	15.8%
	<i>Undergraduate degree</i>	2310	73.6%
	<i>Postgraduate degree</i>	333	10.6%
Smoking	<i>non-Smoker</i>	2328	74.4%
	<i>Smoker</i>	803	25.6%
History of chronic disease	<i>No</i>	2858	91.1%
	<i>Yes</i>	278	8.9%

¹Region: North region includes the following Jordanian governorates: Irbid, Ajloun, Jerash and Mafrq; Central region includes Balqa, Amman (the capital), Zarqa and Madaba; South region includes: Karak, Tafilah, Ma'an and Aqaba. ²Educational level: Undergraduate degree includes diploma and bachelor's degrees; postgraduate degree includes masters and Doctor of Philosophy degrees. ³JOD: Jordanian Dinar. Note: Despite having 3150 as the total number of study participants, the numbers above might not add up to reach this total, due to the existence of partial response to some survey items.

COVID-19 knowledge

The overall knowledge of the study participants regarding COVID-19 is illustrated in (Figure 1). The majority of the study participants correctly responded to eleven out of the thirteen items, with the least percentage of correct responses observed in the following two items: the virus can remain active on surfaces for few hours (49.7%) and reinfection by COVID-19 is not possible (23.7%, Figure 1).


Figure 1. The overall knowledge of COVID-19 among the study participants shown as the percentage of correct responses to each of the thirteen items assessing COVID-19 knowledge.



All items were shown as correct statements. COVID-19: coronavirus disease 2019, SARS-CoV-2: severe acute respiratory syndrome coronavirus 2.

The total number of participants who had a valid K-score was 2988, with a mean K-score of 10.2 (range: 1.0-13.0). Higher level of knowledge regarding COVID-19 was seen among residents of Amman (mean K-score 10.3 vs. 10.1, $p=0.003$; M-W), participants with higher income (10.5 vs. 10.2 vs. 10.0, $p<0.001$; K-W), participants with higher educational level (10.6 vs. 10.2 vs. 9.8, $p<0.001$; K-W) and among non-smokers (10.2 vs. 10.0, $p=0.016$, M-W; Figure 2).

Figure 2. Demographic features correlated with differences in COVID-19 knowledge score (K-score) among the study participants.




Higher K-score was seen among residents of Amman (A), participants with higher monthly income (B), non-smokers (C), and among participants with higher educational level (D). M-W: Mann-Whitney *U* test, K-W: Kruskal-Wallis test, CI: confidence interval of the mean K-score, JOD: Jordanian dinar.

In addition, participants who felt annoyed by the inability to attend places of worship displayed a lower mean K-score (10.1 vs. 10.4, $p < 0.001$; M-W). For the survey items related to COVID-19 misinformation, the participants who believed that the pandemic is related to a conspiracy and those who believed that it was part of a biologic warfare had lower mean K-score (10.0 vs. 10.4, $p < 0.001$ for both comparisons; M-W). In addition, those who believed in the role of 5G networks in COVID-19 spread had a lower mean K-score (9.8 vs. 10.3, $p < 0.001$; M-W). Moreover, those who believed that

COVID-19 is a spiritual divine test showed a lower K-score compared to those who did not hold such a belief (10.1 vs. 10.4, $p < 0.001$; M-W, Figure 3).

Figure 3. Correlation between COVID-19 knowledge score (K-score) and items assessing misinformation regarding origin of COVID-19 pandemic.




Lower K-score was seen among participants who stated that COVID-19 is part of a global conspiracy plot (A), participants who stated that COVID-19 is related to biologic warfare (B), participants who believed in the role of 5G networks in COVID-19 spread (C), and participants who thought that COVID-19 is a divine test (D). P-values were calculated using Mann-Whitney U test, CI: confidence interval of the mean K-score.

Anxiety regarding COVID-19


The total number of participants who had a valid anxiety score was 3035, with mean score of 9.2 (range: zero-21.0). Variables with significant association to higher anxiety level were female gender (mean anxiety score: 9.3 vs 8.7, $p=0.007$, M-W), residence outside Amman (9.5 vs. 9.0, $p=0.006$; M-W), lower educational level (10.1 vs. 9.1 vs. 8.5, $p<0.001$; K-W), lower monthly income (9.9 vs. 9.0 vs. 8.3, $p<0.001$; K-W), and smoking (9.9 vs. 9.0, $p<0.001$; M-W, Figure 4). In addition, those who felt annoyed by the inability to attend places of worship had a higher mean anxiety score (9.7 vs. 8.2, $p<0.001$; M-W). While those who thought that the quarantine gave them an opportunity to spend a quality time with their families had a lower mean anxiety score (8.9 vs. 10.8, $p<0.001$; M-W). Higher anxiety scores were found among those who believed that COVID-19 is related to conspiracy (9.7 vs. 8.7; $p<0.001$; M-W), biological warfare believers (9.6 vs. 8.6, $p<0.001$; M-W), those who believed in the role of 5G networks in facilitating COVID-19 spread (10.3 vs. 8.9, $p<0.001$; M-W), and those who believed that the pandemic is a spiritual divine test (9.4 vs. 8.0, $p<0.001$; M-W, Figure 5).

Figure 4. Demographic features correlated with differences in anxiety score among the study participants.



Lower anxiety score was seen among residents of Amman (A), participants with a higher monthly income (B), non-smokers (C), among participants with a higher educational level (D), and among males (E). M-W: Mann-Whitney *U* test, K-W: Kruskal-Wallis test, CI: confidence interval of the mean anxiety score, JOD: Jordanian dinar.

Figure 5. Correlation between anxiety score and items assessing misinformation regarding origin of COVID-19 pandemic.



Higher anxiety score was seen among participants who stated that COVID-19 is part of a global conspiracy plot (A), participants who stated that COVID-19 is related to biologic warfare (B), participants who believed in the role of 5G networks in COVID-19 spread (C), and participants who thought that COVID-19 is a divine test (D). P-values were calculated using Mann-Whitney *U* test, CI: confidence interval of the mean anxiety.

Perception of COVID-19 danger and attitude towards quarantine

The majority of study population felt that the disease is moderately dangerous (n=1896, 60.3%), followed by 1152 participants who perceived the disease as very dangerous (36.6%). Older participants (more than or equal to 27 [the median age of the whole study population]) perceived the disease as very dangerous (40.6% compared to 31.7%) among the younger participants (less than 27 years old, $p<0.001$; χ^2 , Table 2). Males were more likely to perceive the disease as very dangerous compared to females (41.2% vs. 35.2%, $p=0.001$; χ^2). Participants with the lowest monthly income were more inclined to feel that COVID-19 is very dangerous (41.0% vs. 34.5% vs. 32.5%, $p=0.002$; χ^2). Also, those with the lowest educational level were more likely to believe that the disease is very dangerous (44.0% vs. 34.8% vs. 38.1%, $p=0.001$; χ^2). Smokers (42.8% vs. 34.6%) and those with history of chronic disease (48.2% vs. 35.5%) had higher likelihood to perceive COVID-19 as a very dangerous disease ($p<0.001$ for both comparisons; χ^2). Those who believed that the current pandemic was a spiritual test were more likely to perceive the disease as very dangerous (38.0% vs. 29.8%, $p=0.001$; χ^2). Married participants were more likely to feel that COVID-19 is very dangerous compared to single participants (40.6% vs. 32.5%, $p<0.001$; χ^2 , Table 2). The vast majority of study participants reported adhering to the quarantine measures (n=3072, 97.9%). Variables that were associated with higher likelihood to break the quarantine measures included male gender (4.3% vs. 1.3%, $p<0.001$; χ^2) and history of smoking (3.7% vs. 1.4%, $p<0.001$; χ^2).

Table 2. Perception of COVID-19 danger among study participants.

<i>Characteristic</i>	Self-reported COVID-19 danger	Not dangerous	Moderately dangerous	Very dangerous	P-value³
		N ² (%)	N (%)	N (%)	
<i>Age group</i>	Less than 27	36 (2.5)	939 (65.8)	453 (31.7)	<0.001
	More than or equal to 27	52 (3.4)	849 (55.9)	617 (40.6)	
<i>Gender</i>	Male	31 (4.2)	407 (54.6)	307 (41.2)	0.001
	Female	65 (2.8)	1463 (62.1)	829 (35.2)	
<i>Marital status</i>	Single	49 (3.0)	1042 (64.4)	526 (32.5)	<0.001
	Married	46 (3.3)	794 (56.2)	573 (40.6)	
	Divorced	1 (1.7)	31 (52.5)	27 (45.8)	
	Widow/widower	1 (3.4)	12 (41.4)	16 (55.2)	
<i>Monthly income</i>	Less than 500 JOD ¹	38 (3.1)	690 (56.0)	505 (41.0)	0.002
	500-1000 JOD	37 (3.1)	735 (62.3)	407 (34.5)	
	More than 1000 JOD	21 (3.1)	430 (64.4)	217 (32.5)	
<i>Educational level</i>	High school or less	21 (4.2)	257 (51.8)	218 (44.0)	0.001
	Undergraduate degree	67 (2.9)	1439 (62.3)	803 (34.8)	
	Postgraduate degree	9 (2.7)	197 (59.2)	127 (38.1)	
<i>Smoking</i>	non-Smoker	65 (2.8)	1457 (62.6)	805 (34.6)	<0.001
	Smoker	31 (3.9)	428 (53.3)	344 (42.8)	
<i>History of chronic disease</i>	No	92 (3.2)	1750 (61.3)	1015 (35.5)	<0.001
	Yes	4 (1.4)	140 (50.4)	134 (48.2)	

¹JOD: Jordanian dinar, ²N: Number, ³P-value: Calculated using chi-squared test (χ^2).


Beliefs and misinformation about COVID-19 origin

1. Is COVID-19 part of a global conspiracy?

Overall, a total of 1501 of the study participants believed that COVID-19 is part of a global conspiracy (47.9%, Figure 6). This belief was more common among females compared to males (50.1% vs. 41.2%, $p < 0.001$; χ^2), among married participants compared to single participants (50.5% vs. 45.8%, $p = 0.011$; χ^2) and among smokers compared to non-smokers (52.8% vs. 46.3%, $p = 0.001$; χ^2). The gradual increase in monthly income was associated with a gradual decrease in the belief that COVID-19 is part of a global conspiracy (50.5% in those with income of <500 JOD vs. 48.2% in those with income of 500-1000 JOD vs. 43.8% among those with income of >1000 JOD, $p = 0.019$; χ^2). For educational level, the belief in conspiracy was the highest among

those with a lower educational level (50.4% among those with high school or less degree vs. 48.5% among those with an undergraduate degree, vs. 40.8% among those with postgraduate degrees, $p=0.016$; χ^2).

Figure 6. The overall belief of the study participants with regard to origin and spread of COVID-19.



COVID-19: coronavirus disease 2019, 5G: the 5th generation mobile network.

2. Is COVID-19 part of a biological warfare?

The majority of study participants had a belief that SARS-CoV-2 origin was related to biological warfare ($n=1778$, 57.0%, Figure 6). This belief was more common among females (59.7% vs. 48.6%, $p<0.001$; χ^2), married participants (59.2% vs. 55.0%, $p=0.023$; χ^2), participants with low and middle income ($p=0.001$; χ^2), and lower educational level ($p=0.002$; χ^2).

3. Do 5G networks have a role in COVID-19 spread?

The overall belief in 5G networks role in the spread of COVID-19 was generally less compared to the previously mentioned items (conspiracy and biological warfare) ($n=641$, 21.0%, Figure 6). This misbelief was higher among females (23.6% vs. 12.8% in

males, $p < 0.001$; χ^2), married participants (23.1% vs. 19.4% among singles, $p = 0.017$; χ^2), lower monthly income (24.2%, vs. 20.9% vs. 15.9%, $p < 0.001$; χ^2), and lower educational level (28.0% vs. 20.4% vs. 15.5%, $p < 0.001$; χ^2).

4. Is COVID-19 a spiritual divine test or trial?


The majority of study participants stated that COVID-19 pandemic is a divine spiritual test ($n = 2595$, 82.7%). Variables associated with such a belief included female gender (85.0% vs. 75.8%, $p < 0.001$; χ^2), Jordanian nationality (83.3% vs. 77.1% among non-Jordanians, $p = 0.019$; χ^2), residence outside Amman (86.5% vs. 80.1% among those in Amman, $p < 0.001$; χ^2), marriage (86.5% vs. 79.3, $p < 0.001$; χ^2), lower educational level (88.2%, vs. 82.9% vs. 73.5%, $p < 0.001$; χ^2), lower monthly income (88.4% vs. 82.9% vs. 71.6%, $p < 0.001$; χ^2), and non-smoking (84.2% vs. 78.5%, $p < 0.001$; χ^2).

The main media and other sources of information about the pandemic

The most common source of information for study participants regarding the pandemic were social media platforms ($n = 1075$, 34.4%), followed by TV and news releases ($n = 850$, 27.2%), the official Ministry of Health website on COVID-19 ($n = 771$, 24.6%) and finally scientific journals and opinion of medical doctors ($n = 432$, 13.8%). For social media platforms, Facebook predominated as the main source of information about the pandemic (81.1%), followed by WhatsApp (7.0%), YouTube (4.6%), Twitter (4.5%) and Instagram (2.9%).

The participants who relied on TV and news releases as the main source of knowledge about the virus were older in age compared to those who used other sources combined (33 vs. 30 years old, $p < 0.001$; M-W). Those who relied on doctors and scientific journals had a higher mean K-score (10.5 vs. 10.1, $p < 0.001$; M-W) and a lower mean anxiety score (8.1 vs. 9.3, $p < 0.001$; M-W, Figure 7).

Figure 7. Correlation between the main sources of information regarding COVID-19 and both the anxiety and K-scores.



CI: confidence interval of the mean score, K-score: COVID-19 knowledge score. MoH: Jordanian Ministry of Health.

Participants who depended on medical doctors and scientific journals as their main source of information about COVID-19 were less predisposed to believe in conspiracy (38.4% vs. 49.3%, $p < 0.001$; χ^2), and its related misinformation (5G networks role: 14.2% vs. 22.0%, $p < 0.001$; χ^2 , biological warfare role: 47.7% vs. 58.3%, $p < 0.001$; χ^2 and belief that the pandemic is a spiritual test: 69.1% vs. 48.9%, $p < 0.001$; χ^2). They were also less prone to feel that the disease is very dangerous (30.6% vs. 37.6%, $p < 0.001$; χ^2). Higher reliance on information provided by scientific journals and medical doctors was seen among males (18.8% vs. 12.4%, $p < 0.001$; χ^2), participants with a higher educational level (25.7% vs. 14.0% vs. 4.9%, $p < 0.001$; χ^2), participants with a higher monthly income (21.4% vs. 13.7% vs. 9.6%, $p < 0.001$), residence in Amman (15.2% vs. 12.0%, $p = 0.012$; χ^2), non-Jordanian nationality (19.9% vs. 13.4%, $p = 0.007$) and single marital status (17.0% vs. 10.4%, $p < 0.001$; χ^2).

Discussion

The key results of this study can be summarized as follows: the overall knowledge of COVID-19 among residents in Jordan was satisfactory. More than 87% correct responses were found for eight items and more than 63% correct responses for eleven items out of thirteen total items that were used to assess COVID-19 knowledge in this study. The participants scored less for two items: SARS-CoV-2 can remain active on surfaces for few days rather than few hours (50%) and re-infection by COVID-19 is not possible (24%). The lower knowledge in relation to these two items can be attributed to ongoing research that has not achieved a widespread outreach for the public yet. Such research indicated the stability of SARS-CoV-2 on surfaces for more than 24 hours depending on the nature of the surface [13,27]. For the possibility of re-infection, the current evidence points to unlikely occurrence of such a phenomenon despite the need for more research tackling this aim [28-31].

The high overall knowledge might be attributed to general interest of the public in this pandemic that became a global phenomenon and such high knowledge has been reported recently by several studies around the globe [32-37]. However, upon further dissecting COVID-19 knowledge, in relation to possible origins of the pandemic, severe gaps in knowledge were revealed. This was manifested by high prevalence of belief in conspiracy (48%), biologic warfare role (57%) and 5G networks' role (21%) in the origin and spread of the virus. In our previous work among university students, we found a significant association between higher anxiety during the current pandemic and the belief in conspiracy in COVID-19 origin [23]. The results of the current study clearly delineate the existence of an association between misinformation about COVID-19 and the combination of higher anxiety and lower knowledge about the disease among the public in Jordan.

Even though the natural origin of SARS-CoV-2 was scientifically determined to a large extent, which further discredits the role of a conspiracy in the origin of the disease, many people still grasp to such delusions [38,39]. The current climate of fear and uncertainty seems as a fertile soil from which conspiracy beliefs are born and thrive [40,41]. Thus, the high prevalence of inaccurate beliefs about COVID-19 origins seems

a plausible result. In addition, the link that was demonstrated between higher anxiety levels and conspiracy belief in this study, is not unique and can augment the previous evidence that such belief is harmful [23,42,43].

The clearest variable associated with lower overall knowledge about the disease, belief in conspiracy and higher anxiety level was the lower socio-economic status (lower educational level and lower monthly income). This result is consistent with findings from various recent studies, and points to the importance of targeting such groups with intensified awareness campaigns [33,34,44-46].

The most common main source of information about COVID-19 reported by the participants were the social media platforms. The role of social media in fueling and spreading implausible notions cannot be overlooked [47-49]. The spread of such misinformation via social media outlets is not a recent phenomenon that accompanied the current crisis, but also involved several health-related aspects (e.g. vaccination, AIDS denialism, Zika fever outbreak, etc.) in the last decade [49-51]. To fight against the spread of harmful misinformation, the correct public health messages should be delivered in a user-friendly style with emphasis on fact-checking tools [52]. The prime role relies on experts, physicians and the policy makers to advocate for social media campaigns that can aid in establishing a culture of fact-checking [53].

Regarding the anxiety level of the study participants and taking into account that the survey was conducted in April 2020, the overall mean anxiety score showed a mild anxiety among the study participants. Females showed a higher anxiety level compared to males and this can be partly explained by the differences in physiology which increase females' susceptibility to develop anxiety and stress [54,55]. Also, the lower socio-economic status was associated with a higher anxiety level which can be attributed to the lack of income security during the crisis. Moreover, participants living outside Amman had higher anxiety; this might be due to certain hardships accompanying inhabitants of rural areas and more isolated areas, like financial strains and social isolation. Such results are in line with recent research pointing to similar associations [7,34,56-58]. Furthermore, smokers had higher mean anxiety score compared to non-smokers; this is because of the rising emphasis on how smokers are

more vulnerable to infections including COVID-19, besides the worsened prognosis in case of protracting the disease [59,60].

For the perception of COVID-19 degree of threat, males, participants with a lower socio-economic status, smokers and those with a history of chronic disease were likely to perceive the disease as very dangerous. This result appears plausible, particularly for individuals with comorbidities, considering the high-risk of severe disease and mortality among this group [61].

Finally, one observation in this study warrants further and meticulous exploration. This entails the attitude and belief towards the origin of the disease and the government-enforced public health measures from a religious perspective. Feeling annoyed by the inability to perform religious practices due to forced closure of mosques and churches was seen in the majority of study participants (67.3%). Another observation was that about 83% of the study population believed that the origin of the pandemic is a form of test or trial by God. A higher level of anxiety among the aforementioned groups can be attributed to the inability of a majority of participants to worship in large gatherings in either mosques or churches during the lockdown period in the country. A large sum of previous reports showed that religious belief can reduce anxiety in various health-related conditions [62-65]. Further research is needed to establish the role of religious belief in coping at time of crisis.

Study limitations

Despite the relatively large sample size, bias was observed for gender (a majority of females) and for age. However, age seems to reflect the age distribution among the residents of Jordan at least to some extent. In addition, it was justified to have predominance of residents in the Central region as it harbours roughly two-thirds of population in Jordan including the Capital, Amman. Furthermore, we should clearly state that the results of the current study might not be representative of the Jordanian population. This is partly related to survey distribution via contacts and networks of the authors, which make sampling bias inevitable. Thus, further studies are needed to confirm our findings at different national and cultural levels. The study validity can be another limitation despite having a majority of items adopted from previously

published work [35]. Finally, we have to state the inherent limitations of surveys including the response of the participants in a way they think would please the researchers, in addition to the problem of incomplete response.

Conclusions

COVID-19 poses a crisis that drastically changed the world; this is illustrated by the social, psychological and economic impact of the disease. This pandemic is framed with endless streams of misinformation and fake news, which has its own consequences and spreads even more confusion. The results of this study showed satisfactory knowledge about the disease among residents in Jordan, with lack of knowledge in certain aspects of the disease regarding origin and conspiracy surrounding this pandemic. Individuals with a lower socio-economic status showed higher anxiety, lower COVID-19 knowledge and belief in misinformation. Focused awareness and proper delivery of correct information is mandatory, particularly for this group to reduce the negative impact of the pandemic on their lives. An association was seen between belief in the role of conspiracies, biological warfare, and 5G networks in the origin and spread of the disease with lower levels of knowledge regarding COVID-19 and higher level of anxiety. The spread of misinformation and conspiracies is exacerbated by different media outlets, which is why proper management and close monitoring of posted content is necessary.

Supporting information

S1 Appendix

Consent Form and Questionnaire Translated to English (the original form in Arabic is provided as well).

Acknowledgments

NA

Funding

We declare that we received no funding nor financial support/grants by any institutional, private or corporate entity.

Conflicts of Interest

We declare that we have no competing interest nor conflict of interest.

Author contribution

Conceptualization: MS, DD, AM

Data Curation: MS, DD

Formal Analysis: MS

Investigation: MS, DD, AY, AH, DT, HE, NAA, FGB, AM

Methodology: MS, DD, AY, AH, DT, HE, NAA, FGB, AM

Project administration: MS

Supervision: MS, AM

Visualization: MS

Writing - Original Draft Preparation: MS, DD

Writing - Review & Editing: MS, DD, AY, AH, DT, HE, NAA, FGB, AM

References

- [1]. Lipsitch M, Swerdlow DL, Finelli L. Defining the Epidemiology of Covid-19 - Studies Needed. *N Engl J Med.* **2020**;382;13:1194-6. doi:10.1056/NEJMp2002125
- [2]. Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, et al. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA.* **2020**. doi:10.1001/jama.2020.6775
- [3]. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet.* **2020**;395;10227:912-20. doi:10.1016/S0140-6736(20)30460-8
- [4]. Nicola M, Alsafi Z, Sohrabi C, Kerwan A, Al-Jabir A, Iosifidis C, et al. The Socio-Economic Implications of the Coronavirus and COVID-19 Pandemic: A Review. *Int J Surg.* **2020**. doi:10.1016/j.ijssu.2020.04.018
- [5]. Dubey S, Biswas P, Ghosh R, Chatterjee S, Dubey MJ, Chatterjee S, et al. Psychosocial impact of COVID-19. *Diabetes Metab Syndr.* **2020**;14;5:779-88. doi:10.1016/j.dsx.2020.05.035
- [6]. Hua J, Shaw R. Corona Virus (COVID-19) "Infodemic" and Emerging Issues through a Data Lens: The Case of China. *Int J Environ Res Public Health.* **2020**;17;7. doi:10.3390/ijerph17072309
- [7]. Gao J, Zheng P, Jia Y, Chen H, Mao Y, Chen S, et al. Mental health problems and social media exposure during COVID-19 outbreak. *PLoS One.* **2020**;15;4:e0231924. doi:10.1371/journal.pone.0231924
- [8]. Mian A, Khan S. Coronavirus: the spread of misinformation. *BMC Med.* **2020**;18;1:89. doi:10.1186/s12916-020-01556-3

- [9]. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med.* **2020**;382;18:1708-20. doi:10.1056/NEJMoa2002032
- [10]. Zhou M, Zhang X, Qu J. Coronavirus disease 2019 (COVID-19): a clinical update. *Front Med.* **2020**. doi:10.1007/s11684-020-0767-8
- [11]. Rothan HA, Byrareddy SN. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. *J Autoimmun.* **2020**;109:102433. doi:10.1016/j.jaut.2020.102433
- [12]. Kampf G, Todt D, Pfaender S, Steinmann E. Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. *J Hosp Infect.* **2020**;104;3:246-51. doi:10.1016/j.jhin.2020.01.022
- [13]. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *N Engl J Med.* **2020**;382;16:1564-7. doi:10.1056/NEJMc2004973
- [14]. Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schunemann HJ, et al. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *Lancet.* **2020**;395;10242:1973-87. doi:10.1016/S0140-6736(20)31142-9
- [15]. Sanders JM, Monogue ML, Jodlowski TZ, Cutrell JB. Pharmacologic Treatments for Coronavirus Disease 2019 (COVID-19): A Review. *JAMA.* **2020**. doi:10.1001/jama.2020.6019
- [16]. Bavel JJV, Baicker K, Boggio PS, Capraro V, Cichocka A, Cikara M, et al. Using social and behavioural science to support COVID-19 pandemic response. *Nat Hum Behav.* **2020**;4;5:460-71. doi:10.1038/s41562-020-0884-z
- [17]. West R, Michie S, Rubin GJ, Amlot R. Applying principles of behaviour change to reduce SARS-CoV-2 transmission. *Nat Hum Behav.* **2020**;4;5:451-9. doi:10.1038/s41562-020-0887-9

- [18]. Georgiou N, Delfabbro P, Balzan R. COVID-19-related conspiracy beliefs and their relationship with perceived stress and pre-existing conspiracy beliefs. *Pers Individ Dif*. **2020**;166:110201. doi:10.1016/j.paid.2020.110201
- [19]. Ahmad AR, Murad HR. The Impact of Social Media on Panic During the COVID-19 Pandemic in Iraqi Kurdistan: Online Questionnaire Study. *J Med Internet Res*. **2020**;22;5:e19556. doi:10.2196/19556
- [20]. Li HO, Bailey A, Huynh D, Chan J. YouTube as a source of information on COVID-19: a pandemic of misinformation? *BMJ Glob Health*. **2020**;5;5. doi:10.1136/bmjgh-2020-002604
- [21]. Allington D, Duffy B, Wessely S, Dhavan N, Rubin J. Health-protective behaviour, social media usage and conspiracy belief during the COVID-19 public health emergency. *Psychol Med*. **2020**:1-7. doi:10.1017/S003329172000224X
- [22]. Pennycook G, McPhetres J, Zhang Y, Lu JG, Rand DG. Fighting COVID-19 Misinformation on Social Media: Experimental Evidence for a Scalable Accuracy-Nudge Intervention. *Psychol Sci*. **2020**:956797620939054. doi:10.1177/0956797620939054
- [23]. Sallam M, Dababseh D, Yaseen A, Al-Haidar A, Ababneh NA, Bakri FG, et al. Conspiracy beliefs are associated with lower knowledge and higher anxiety levels regarding COVID-19 among students at the University of Jordan. *International Journal of Environmental Research and Public Health*. **2020**;17;14:2020.04.21.20064147. doi:10.3390/ijerph17144915
- [24]. Lowe B, Decker O, Muller S, Brahler E, Schellberg D, Herzog W, et al. Validation and standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the general population. *Med Care*. **2008**;46;3:266-74. doi:10.1097/MLR.0b013e318160d093
- [25]. CheckMarket. Calculate sample size margin of error. 2020 [cited 2020 25/04/2020]. Available from: <https://www.checkmarket.com/>.
- [26]. Worldometer. Jordan Population. 2020 [cited 2020 25/04/2020]. Available from: <https://www.worldometers.info/world-population/jordan-population/>.

- [27]. Zhang DX. SARS-CoV-2: air/aerosols and surfaces in laboratory and clinical settings. *J Hosp Infect.* **2020**. doi:10.1016/j.jhin.2020.05.001
- [28]. Long QX, Liu BZ, Deng HJ, Wu GC, Deng K, Chen YK, et al. Antibody responses to SARS-CoV-2 in patients with COVID-19. *Nat Med.* **2020**;26;6:845-8. doi:10.1038/s41591-020-0897-1
- [29]. Chandrashekar A, Liu J, Martinot AJ, McMahan K, Mercado NB, Peter L, et al. SARS-CoV-2 infection protects against rechallenge in rhesus macaques. *Science.* **2020**. doi:10.1126/science.abc4776
- [30]. Ota M. Will we see protection or reinfection in COVID-19? *Nat Rev Immunol.* **2020**;20;6:351. doi:10.1038/s41577-020-0316-3
- [31]. Sallam M. Urgent necessity to address the issue of COVID-19 reinfection. *Preprint.* **2020**. doi:10.13140/RG.2.2.26744.39687
- [32]. Al-Hanawi MK, Angawi K, Alshareef N, Qattan AMN, Helmy HZ, Abudawood Y, et al. Knowledge, Attitude and Practice Toward COVID-19 Among the Public in the Kingdom of Saudi Arabia: A Cross-Sectional Study. *Front Public Health.* **2020**;8:217. doi:10.3389/fpubh.2020.00217
- [33]. Azlan AA, Hamzah MR, Sern TJ, Ayub SH, Mohamad E. Public knowledge, attitudes and practices towards COVID-19: A cross-sectional study in Malaysia. *PLoS One.* **2020**;15;5:e0233668. doi:10.1371/journal.pone.0233668
- [34]. Abdelhafiz AS, Mohammed Z, Ibrahim ME, Ziady HH, Alorabi M, Ayyad M, et al. Knowledge, Perceptions, and Attitude of Egyptians Towards the Novel Coronavirus Disease (COVID-19). *J Community Health.* **2020**. doi:10.1007/s10900-020-00827-7
- [35]. Zhong BL, Luo W, Li HM, Zhang QQ, Liu XG, Li WT, et al. Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-sectional survey. *Int J Biol Sci.* **2020**;16;10:1745-52. doi:10.7150/ijbs.45221

- [36]. Reuben RC, Danladi MMA, Saleh DA, Ejembi PE. Knowledge, Attitudes and Practices Towards COVID-19: An Epidemiological Survey in North-Central Nigeria. *J Community Health*. **2020**. doi:10.1007/s10900-020-00881-1
- [37]. Hezima A, Aljafari A, Aljafari A, Mohammad A, Adel I. Knowledge, attitudes, and practices of Sudanese residents towards COVID-19. *East Mediterr Health J*. **2020**;26;6:646-51. doi:10.26719/emhj.20.076
- [38]. Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet*. **2020**;395;10224:565-74. doi:10.1016/S0140-6736(20)30251-8
- [39]. Benvenuto D, Giovanetti M, Salemi M, Prosperi M, De Flora C, Junior Alcantara LC, et al. The global spread of 2019-nCoV: a molecular evolutionary analysis. *Pathog Glob Health*. **2020**;114;2:64-7. doi:10.1080/20477724.2020.1725339
- [40]. van Prooijen JW, Douglas KM. Conspiracy theories as part of history: The role of societal crisis situations. *Mem Stud*. **2017**;10;3:323-33. doi:10.1177/1750698017701615
- [41]. Constantinou M, Kagialis A, Karekla M. COVID-19 Scientific Facts Vs. Conspiracy Theories: 0-1: Science Fails to Convince Even Highly Educated Individuals. *Preprint in Research Square*. **2020**. doi:10.21203/rs.3.rs-33972/v1
- [42]. Jolley D. Are conspiracy theories just harmless fun? *Psychologist*. **2013**;26;1:60-2
- [43]. Jakub Šrol, Eva Ballová Mikušková, Vladimira Cavojova. When we are worried, what are we thinking? Anxiety, lack of control, and conspiracy beliefs amidst the COVID-19 pandemic. *Preprint*. **2020**
- [44]. Lau LL, Hung N, Go DJ, Ferma J, Choi M, Dodd W, et al. Knowledge, attitudes and practices of COVID-19 among income-poor households in the Philippines: A cross-sectional study. *J Glob Health*. **2020**;10;1:011007. doi:10.7189/jogh.10.011007

- [45]. Geldsetzer P. Knowledge and Perceptions of COVID-19 Among the General Public in the United States and the United Kingdom: A Cross-sectional Online Survey. *Ann Intern Med.* **2020**. doi:10.7326/M20-0912
- [46]. Roy D, Tripathy S, Kar SK, Sharma N, Verma SK, Kaushal V. Study of knowledge, attitude, anxiety & perceived mental healthcare need in Indian population during COVID-19 pandemic. *Asian J Psychiatr.* **2020**;51:102083. doi:10.1016/j.ajp.2020.102083
- [47]. Ahmed W, Vidal-Alaball J, Downing J, Lopez Segui F. COVID-19 and the 5G Conspiracy Theory: Social Network Analysis of Twitter Data. *J Med Internet Res.* **2020**;22;5:e19458. doi:10.2196/19458
- [48]. Llewellyn S. Covid-19: how to be careful with trust and expertise on social media. *BMJ.* **2020**;368:m1160. doi:10.1136/bmj.m1160
- [49]. Waszak PM, Kasprzycka-Waszak W, Kubanek A. The spread of medical fake news in social media—the pilot quantitative study. *Health policy and technology.* **2018**;7;2:115-8
- [50]. Gu Z, Badger P, Su J, Zhang E, Li X, Zhang L. A vaccine crisis in the era of social media. *National Science Review.* **2018**;5;1:8-10
- [51]. Kou Y, Gui X, Chen Y, Pine K. Conspiracy talk on social media: collective sensemaking during a public health crisis. *Proceedings of the ACM on Human-Computer Interaction.* **2017**;1;CSCW:1-21
- [52]. Tangcharoensathien V, Calleja N, Nguyen T, Purnat T, D'Agostino M, Garcia-Saiso S, et al. Framework for Managing the COVID-19 Infodemic: Methods and Results of an Online, Crowdsourced WHO Technical Consultation. *J Med Internet Res.* **2020**;22;6:e19659
- [53]. Trethewey SP. Strategies to combat medical misinformation on social media. *Postgrad Med J.* **2020**;96;1131:4-6. doi:10.1136/postgradmedj-2019-137201

- [54]. Donner NC, Lowry CA. Sex differences in anxiety and emotional behavior. *Pflugers Arch.* **2013**;465;5:601-26. doi:10.1007/s00424-013-1271-7
- [55]. Bahrami F, Yousefi N. Females are more anxious than males: a metacognitive perspective. *Iran J Psychiatry Behav Sci.* **2011**;5;2:83-90
- [56]. Moghanibashi-Mansourieh A. Assessing the anxiety level of Iranian general population during COVID-19 outbreak. *Asian J Psychiatr.* **2020**;51:102076. doi:10.1016/j.ajp.2020.102076
- [57]. Vindegaard N, Benros ME. COVID-19 pandemic and mental health consequences: Systematic review of the current evidence. *Brain Behav Immun.* **2020**. doi:10.1016/j.bbi.2020.05.048
- [58]. Mazza C, Ricci E, Biondi S, Colasanti M, Ferracuti S, Napoli C, et al. A Nationwide Survey of Psychological Distress among Italian People during the COVID-19 Pandemic: Immediate Psychological Responses and Associated Factors. *Int J Environ Res Public Health.* **2020**;17;9. doi:10.3390/ijerph17093165
- [59]. Vardavas CI, Nikitara K. COVID-19 and smoking: A systematic review of the evidence. *Tob Induc Dis.* **2020**;18:20. doi:10.18332/tid/119324
- [60]. Patanavanich R, Glantz SA. Smoking is Associated with COVID-19 Progression: A Meta-Analysis. *Nicotine Tob Res.* **2020**. doi:10.1093/ntr/ntaa082
- [61]. Yang J, Zheng Y, Gou X, Pu K, Chen Z, Guo Q, et al. Prevalence of comorbidities in the novel Wuhan coronavirus (COVID-19) infection: a systematic review and meta-analysis. *International journal of infectious diseases.* **2020**
- [62]. Moreira-Almeida A, Neto FL, Koenig HG. Religiousness and mental health: a review. *Braz J Psychiatry.* **2006**;28;3:242-50. doi:10.1590/s1516-44462006000300018

[63]. Alves RR, Alves Hda N, Barboza RR, Souto Wde M. The influence of religiosity on health. *Cien Saude Colet.* **2010**;15;4:2105-11. doi:10.1590/s1413-81232010000400024

[64]. Safara M, Bhatia M. Relationship of religious beliefs with anxiety and depression. *Delhi Psyquiatric Journal.* **2008**;11;2:177-9

[65]. Jansen KL, Motley R, Hovey J. Anxiety, depression and students' religiosity. *Mental Health, Religion & Culture.* **2010**;13;3:267-71