COVID-19 effective reproductive ratio determination: An application, and analysis of issues and influential factors

Luis Alfredo Bautista Balbás[1], Mario Gil Conesa[2], Gil Rodríguez Caravaca[2], Blanca Bautista Balbás[3]

[1]Hospital Virgen de la Poveda, Villa del Prado, Madrid, Spain.

[2]Hospital Fundación Alcorcón. Madrid, Spain.

[3]Hospital Son Llatzer. Palma, Balearic Islands, Spain.

Abstract

An essential indicator of COVID-19 transmission is the effective reproduction number (R_t), the number of cases which an infected individual is expected to infect at a particular moment of time; curves of the evolution of R_t over time (transmission curves) reflect the impact of preventive measures and whether an epidemic is controlled. We have created a Shiny/R web application (https://alfredob.shinyapps.io/estR0/) with user-selectable features: open data sources with daily COVID-19 incidences from all countries and many regions, customizable preprocessing options (smoothing, proportional increment, backwards distribution of negative corrections, etc), different MonteCarlo-Markov-Chain estimates of the generation time or serial interval distributions and state-of-the-art R_t estimation frameworks (EpiEstim, R0). We have analyzed the impact of these factors in the obtained transmission curves. We also have obtained curves at the national and sub-national level and analyzed the impact of epidemic control strategies, superspreading events, socioeconomic factors and outbreaks.

We conclude that country wealth and, to a lesser extent, mitigation strategies, were associated with poorer epidemic control. Dataset quality was an important factor, and sometimes dictated the necessity of time series smoothing. We couldn't find conclusive evidence regarding the impact of alleged superspreading events. In the reopening phase, outbreaks had an impact on transmission curves. This application could be used interactively as a tool both to obtain transmission estimates and to perform interactive sensitivity analysis.

Table of Contents

1   Introduction

The *basic reproduction number* or basic reproductive ratio, R_0, is the average number of secondary cases of disease caused by a single infected individual over his infectious period, when all individuals are susceptible to infection; some authors also include the absence of preventive measures in this definition. It depends on the transmissibility (probability of infection, given contact), the average rate of contacts and the average duration of the disease. Although it can be considered as intrinsic to a particular infectious disease, the contact rate is influenced by demography, population density, cultural habits, seasonality and other factors; so this parameter can vary widely between different studies (1). The R_0 can be also understood in terms of the SIR Epidemic models (and other compartmental models): the R_0 is the ratio between the effective contact rate β and the removal rate v, and this family of models have been used to estimate it (2)(3).

The *effective reproduction number* (R_t) is the actual number of cases that an infected individual is expected to infect, given the current state of the population and the implemented

measures, and it typically smaller than R_0. R_t can be analyzed as a function of mobility restrictions and confinement measures(4). R_t also decreases when the proportion of susceptible individuals declines.

Estimating the COVID-19 reproduction number is important to get direct knowledge about the evolution of the pandemic and implement public health regulations and preventive measures. Public health strategies include infection prevention (masks, hand-washing…), social distancing (including schools closures, teleworking, physical distancing,cancellation of mass gatherings…), travel-related measures (travel advice and/or restrictions, area quarantines/"cordon sanitaire", traveller screening…), environmental cleaning, etc… (5). According to their "intensity" public health strategies can be classified in containment, mitigation and suppression. The **containment** strategy involves quarantine of infected patients and requires significant resources (isolation, airborne protection, etc) and a careful epidemiological investigation; the appearance of cases outside the containment areas renders this approach ineffective(6). The **mitigation** strategy the aims to slow down the transmission in order to prevent healthcare system collapse, without necessarily stopping epidemic spread (reaching R values close to 1). Mitigation measures include case isolation, contacts quarantine and social distancing for vulnerable groups. The **suppression** strategy aims to reduce the R values below 1, including measures such as social distancing for all groups, household quarantine and closing education institutions (7).

On 2-Mar-2020 the European CDC published an update that included a 5-scenario analysis (0 to 4); with scenario-bound proposed measures and objectives ranging from containment to mitigation(5). A report by the Imperial College COVID-19 Response Team on 16-Mar-2020 remarked that the preferred policy option, for countries able to attain it, was suppression instead of mitigation because a mitigated epidemic would result in thousands of deaths(8). In March and April 2020 the epidemic outbreak materialized in Europe; 30 out of 31 European Union/European Economic Area and UK (EU/EEA) countries implemented public space closures, 17 out of 31 had mandatory stay-at-home policies for the general population and all countries canceled mass gatherings and closed higher education or secondary schools (9).

After the outbreak was controlled, many countries have implemented multistage dynamic scenarios (the "New Normality" in Spain(10), the phased recovery in UK(11), the regional semaphore plans in Mexico(12), etc). Both the lockdown and the reopening phases require different transmission indicators, including the reproduction numbers(13). For instance, the French Government has defined three R levels: green (<1), orange (1-1.5) or red (>1.5). In the Russian Federation the R_t is a main indicator, along with bed availability and test capacity; proposed cutoffs for the R_t include ≤ 1 to advance to phase I, ≤ 0.8 to advance to phase II and ≤ 0.5 to advance to phase III (14). The WHO recognized a R_t as the best indication that an epidemic is controlled and declining, and recommended large countries should estimate it at subnational level(15).

An important component of transmission are superspreading patients (or events), those who allegedly transmit an infection to a large number of individuals, due to factors like high viral load, asymptomatic cases and extensive social interactions (16). Mass media has reported several events that could qualify as such: The football match between Atalanta and Valencia FC in February, 19(17)), the 8 March events in Spain (18)(19) and other countries (including International Women's Day demonstrations), the annual gathering of the Christian Open Door Church in France (14-24 February) in Mulhouse, France(20), the Spring Break in the US (varying dates depending on the college/institution, peaking March 7-14)(21) and the riots in USA after the

death of George Floyd. Superspreading events are associated with both explosive growth early in an outbreak and sustained transmission in later stages, and small delays in their prevention can be associated with significant morbidity[22]. In the SARS outbreak in Hong Kong in 2003 the estimated R_t peaks coincided with known superspreading events [23].

Knowing the generation time is particularly important to estimate the reproduction number; the generation time (or generation interval) is the period between infection events in an infection sequence, that is to say, the period between the start of the infection in the primary patient (the infector) and the start of the infection in the individual receiving the infection (the infectee). In epidemic periods the generation time is not easily determined, so the serial interval is often used as a surrogate[24]. The serial interval (si) is the interval between the onset of symptoms in the infector and the onset of symptoms in the infectee; it has the same mean as the generation time, but larger variance due to the presence of asymptomatic transmission during the incubation period[25]. It might become negative if asymptomatic transmission takes place and symptoms start earlier in the infectee than in the infector. Coarse data (i.e. data in which the onset of the symptoms is not precisely known in either the infector or the infectee) can still be used to estimate the incubation period[26].

Several descriptions of the COVID-19 serial interval have been made available. Nishiura et al collected information from 28 research articles and case investigation reports and described a serial interval with median 4.0 that was fitted to several distributions including a log-normal distribution[27]. Zhao et al used data from 21 transmission chains in Hong Kong in late January and February 2020[28]. Du et al used data from 468 online reports from China in order to estimate the serial interval [29]. 12.6% of case reports had negative serial intervals, i.e. the symptoms stared earlier in the infected than in the infectee and the authors attributed this to presymptomatic transmission and symptoms starting in the infectee earlier than in the infector. Estimations also exist for the generation time, obtained from clusters from Singapore (91 cases, 48% asymptomatic transmission) and Tianjin (105 cases, 62% asymptomatic transmission) by Ganyani et al; these estimates are slightly larger than those obtained from the serial interval[25]. It is not clear how this intervals might influence the R_t estimates and the practical information obtained from the R_t curves.

Beautiful mathematical descriptions of the relationship between generation times, epidemic growth rates and reproductive numbers have been described elsewhere[30][31][32]. In time-since-infection-models, the Incidence rate $I(t)$ changes depending on the transmissibility $\beta(t, \tau)$ where t denotes calendar time and tau denotes time since infection. (Transmission is defined as a Poisson process such that an individual has a probability $\beta(t, \tau)\delta$ of infecting another individual). Infectiousness can be decomposed into the product of $\beta(t, \tau) = R(t)w(\tau)$ where $w(\tau)$ is the idealized generation time distribution (i.e. the probability distribution of new infection events as a function of time since infection τ, assuming contact rates are constant and not time-dependent) and $R(t)$ is the instantaneous reproduction number. Different estimates exist for the time-varying effective reproduction numbers[31]: The **case reproduction number** at time t (R_t^c), is the average number of cases an infected individual at time t will infect; the R_t^c can be estimated with the method developed by Wallinga and Teunis[32] and it reflects transmissibility after time t. On the other hand, the **instantaneous reproduction number** R_t is the average number of people some individual infected at time t could be expected to infect if conditions remain unchanged; it is estimated by the ratio of new infections generated at time step t, to the total infectiousness of infected individuals at time t[23], which is defined by $\sum_{\tau=1}^{t} I_{t-\tau} w_\tau$.

In this report we focus on the effective reproduction number estimates, and the curves of time-varying effective reproduction number vs time (**transmission curves**, as an analog of epidemic curves which display daily incidence data). We attempt to evaluate the nuances of calculating the effective reproduction number with modern methodologies and the factors influencing this estimation, and how transmission curves have evolved in the last months.

2   Materials and methods

A web application in R/Shiny has been developed in order to estimate effective reproductive ratios (https://alfredob.shinyapps.io/estR0/). This application leverages and combines state-of-the-art reproductive ratio estimation tools, open COVID-19 incidence datasets, generation time/serial interval descriptions and the Shiny framework for web development, which is a R package that allows users to easily build and host interactive web applications (33).

It uses several COVID-19 open data sources: COVID-19 Data Repository by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University, European CDC COVID-19 data, Narrativa COVID-19 tracking project, the New York Times COVID-19 data source, the Spanish Public Health Surveillance (ReNaVE) COVID-19 dataset, the Brazil Covid dataset(34), the official Ministry of Health/Government COVID-19 datasets from Belgium, Canada, Chile, Colombia, Italy, Spain and Mexico, and the Catalonia Open data catalog COVID-19 sources. In many countries, volunteers perform a compilation and curation of daily reports carried out by health authorities; these unofficial/semiofficial sources of data have been included: United Kingdom (Tom White, Unlicense), Germany (J Gehrcke, MIT license), Perú (JM Castagnetto, MIT license), Portugal (Data Science for the Social Good, GPL 3.0), etc. Proper credit is given to these authors in the app.

COVID-19 cases are defined here by having a positive PCR. Datasets can have many problems or issues: data corrections or consolidations (large peaks or valleys), noise (small peaks and variability, possibly arising from a small number of tests performed), an uncertain and time-dependent proportion of undetected cases, delays between case diagnosis and report, etc. Health authorities often report cumulative incidence every day ("number of cases so far"); in this case a time series of approximate daily incidences is obtained by differentiation. As a consequence; this might introduce spurious peaks if the cumulative report includes both new cases and previous cases that were not reported, or negative values if corrections of previous reports are performed. The data quality of some COVID-19 datasets has been examined, along with the impact on R_t estimations, as explained later in this section.

Graphs and tables have been implemented in the application so that the user can examine the dataset before and after preprocessing, before drawing any conclusion.

Several preprocessing steps were implemented:

- The total number of cases can-be increased by a user-defined constant proportion (ranging from 0-99%) to account for a constant proportion of undetected cases (the **undetected rate**). In some studies this proportion of undetected infections has been described to be 80-90% in Iceland (35), 72% in Italy(36) or 56.5% in Austria (37).

- The epidemic start can be defined as the date with at least an user-defined number of cases per three day period. This is important, as calculating the R_t too early in the epidemic does not provide accurate or useful estimates.

- Smoothing can be performed in a variety of ways: Lowess (locally-weighted polynomial regression) and moving averaging (EMA, WMA, SMA, etc). In the Lowess smoothing, the number of points that influence each point can be adjusted (the smoothing span). Mean averaging has been tried and implemented using the previous points, the future points or both for averaging.

- Negative incidences are invalid by definition and reveal an underlying incidence reporting problem; in some series these values reflect data verification and consolidation, implying that these values are meant to compensate cases counted more than one time. This situation might arise for example if an authority reports the number of positive tests every day and then groups them by individual patients every week; or if an authority aggregates data from several institutions every day without considering whether a positive case was previously tested in two institutions and reported twice. Centralized databases of non-aggregate data could solve these issues. In our application these values can be eliminated or set to 0 before or after smoothing, or they are assumed to be periodic corrections to the case series and a **backwards distribution before smoothing** (BD) is performed. In the BD correction, we propose to smooth them with two assumptions: 1) The corrections C_j, C_l are negative values at time points j, l of a time series (T)); they are meant to correct the data points since the previous correction (or the beginning of the time series, if no previous correction is found). 2) The correction should be applied in a direct proportion to the incidence at each time point. (Therefore, to apply the correction C_l each positive point T_k between the two corrections is divided by the total sum of the points between the corrections, and multiplied by their total sum minus the next correction: $T_k = T_k \frac{\sum_{n=j+1}^{k} T_n}{\sum_{n=j+1}^{k-1} T_n}$. The correction point is set to 0.

- Backwards distribution of spurious peaks can also be chosen by the user. A simple peak detector has also been implemented, and it can be applied before smoothing. We arbitrarily define **spurious peaks** as daily incidences greater than 200; and 300% higher than both the two previous and the two following daily incidences; the definition also requires that the two previous and the two following daily incidences are higher than 75. In this case, the corrected spurious peak S_{jc} at time j is set to the centered mean average of 5 values, including the original peak value S_j, ($S_{jc} = \frac{\sum_{n=j-2}^{j+2} T_n}{5}$) and the value to distribute backwards is the difference between the peak and the mean, $T_k = T_k \frac{\sum_{n=j+1}^{k-1} T_n + (S_{jc} - S_j)}{\sum_{n=j+1}^{k-1} T_n}$).

These parameters are implemented in a Shiny app, so the user can immediately see the effect they have on the daily incidence time series; and choose the ones that better fit the data. Unless otherwise specified, we have used Simple moving average for smoothing, backwards distribution of negative values before smoothing, no specific distribution of positive spurious peaks and no proportional correction for undetected cases; and the epidemic curves after preprocessing are displayed.

Different methods can be used to estimate the time-varying reproduction number. Wallinga and Teunis method estimate the case reproduction number at time t (R_t^c), which is the number of cases an infected individual at time t will infect. Anne Cori et al described a method to estimate the instantaneous reproduction number R_t as the number of new infections at time step t divided by the total infectiousness at that time step(23).

Deriving the reproduction number from incidence data requires knowledge of the generation time or the serial interval. The original serial interval datasets from Nishiura et al (27), from Du et al (29) and from Zhao et al (28) were obtained; providing both exact and censored serial intervals. Cases with non-positive serial interval were excluded from the Du et al dataset, due to estimation requirements and due to the characteristics of the serial interval. These data can be used to apply a Markov Chain MonteCarlo method to estimate the parameters of the lognormal distribution of the serial interval(38); a lognormal distribution was chosen since the original authors reported it provided the best fit. The lognormal distribution can be defined as the distribution of the variable $X = e^{\mu + \sigma Z}$, where Z is a standard normal variable, and μ and σ are two real numbers, which are the mean and standard deviation of the logarithm of X. The implementation in the R package coarseDataTools(26) was used (Metropolis-Hastings algorithm, with 6000 iterations and thinning of 1000) and convergence checks were performed with the Gelman-Rubin diagnostic). We also used the Generation times from Ganyani et al obtained with data from clusters in Singapore and Tianjin. We took the mean and standard deviation estimates and their credible interval, and used them as the defining parameters of a triangular distribution (where the estimate is the mode and the credible interval the limits). From this we randomly sampled 5000 means and standard deviations of the serial interval, and calculated 5000 gamma distributions of the serial interval.

These distributions are, in turn, used to estimate R_t using the R package EpiEstim(38)(23). Point estimates of μ and σ were also obtained using the R package coarseDataTools, and these are used to obtain R_t^c estimates using the R0 package, with the method described by Wallinga and Teunis(32).

This code and application are used to look for answers to several questions:

Question A: What impact does data quality have on R_t estimations? Datasets can have many problems or issues. The effect of data quality on R_t estimates (question A) has been analyzed in the Sup. material: The higher-quality COVID-19 incidence reports from the Spanish Public Health Surveillance National Network (RENAVE) have been compared to the initial and deprecated Ministry of Health/Government COVID-19 cumulative incidence reports (5.8); the impact of a spurious peak on the result has been examined (Sup. material, 5.4), and the impact of weekends and smoothing on daily incidences and the subsequent weekly and daily R_t estimations has been evaluated by using official data from the Catalonia regional Government and Belgium Government (Sup. material, 5.10, 5.9).

Questions B: What impact do preprocessing parameters have on R_t estimations? Transmission curves have been obtained with different settings. A rudimentary one-at-a-time sensitivity analysis is also performed (Sup Material, 5.6) using the average daily Gap between non overlapping credible R_t intervals as an indicator. Sensible parameters are chosen and used in the rest of the study.

Question C: What impact do the different serial intervals/generation times and estimation methods have? Transmission curves for different si/GT distributions and estimation methods (R_t, R_t^c) have been analyzed, sensitivity analysis has been performed with different distributions, also using the average daily Gap.

Question D: What impact do wealth and development have on national transmission curves? We have used the John Hopkins University dataset to obtain R_t curves in order to analyze differences between developed and developing countries(39), section 3.4. National Gross Domestic Product (GDP) per capita (purchasing power parity, constant 2017 international dollars)

data from World Bank for 2019 (40) was obtained and used to colour the transmission curves so that economic impact can be visually evaluated. Hypothesizing that wealth might play a significant role in dataset quality, the relationship between GDP and dataset variability is also evaluated (Sup. material, 5.3)

Question E: Do transmission curves follow a similar pattern on a sub-national/regional level? We have also obtained regional R_t curves by using the data provided by Spain Health Ministry(41); these data include positive PCRs and positive antibodies tests, but only the former was used since the dataset does not report the type of positive antibodies. While a positive IgG result can only be interpreted as a "present/past infection" (prevalence information), a positive IgM result could provide incidence information; however we assume a PCR will be performed in most positive IgM cases, following the standard protocols(42) so the impact of this limitation is expected to be minor. It should be noted that this problem was also present in many other datasets. Additionally, in the Sup. Material, we have compared the R_t between different USA states and between different Perú regions (Sup. material: 5.11, 5.12). Regional GDP per capita (US dollars, current prices, current purchasing power parity) data from OECD for 2018 (if available, USA) or 2017 (Perú) (43) was used to color the transmission curves.

Question F: Are transmission curves altered in suspected superspreding events? The possible impact of superspreading events in different countries has been analyzed by using the JHU dataset: The Atalanta vs Valencia football match (Italy), the 8M/International Women's Day demonstrations and other events in Spain, the Annual Assembly of the Christian Open Door Church in France (February), the March spring break in the USA, the George Floyd killing protests in USA (June). Some of these events have been subjects of political debate; but the question examined here has an epidemiological nature: How did the transmission curves change around these events? Is there any temporal correlation between superspreading event dates and transmission surges? Any other implication will not be discussed.

Question G: Can transmission curves be used to detect outbreaks or to guide epidemiological measures in the reopening phase? Finally, given the importance of Public Health surveillance in the reopening phase, we have analyzed the correlation between outbreaks and R_t changes in the reopening phase, by using incidence data of several German districts and states (dataset compiled by J Gehrcke with information by the John Köch Institute), and the outbreak in Lleida, Catalonia (dataset by IDESCAT, Salut, Generalitat de Catalunya(44)), in the beginning of July (Sup. material, 5.13, 5.14). In these analysis several subnational divisions of varying entity and size are analyzed, in order to evaluate whether transmission peaks/valleys are reflected in outbreaks (or the lack of thereof). In 5.14 the analysis is complimented with a sensitivity analysis that includes several si or GT distributions, and an analysis of overall infectivity, as calculated by the EpiEstim package.

Many of these questions are mutually related: Data quality influences and determines R_t estimations, the impact of superspreading events can also be seen on the regional level, COVID-19 data quality can be influenced by economic factors (Sup. material, 5.3), the smoothing might need to be increased in order to correct significant data quality issues, which might in turn reduce the sensitivity of the estimates.

The analysis of the temporal evolution of the R_t (transmission curves) here is mostly performed qualitatively, considering the clinical and epidemiological implications: Whether or not

the disease transmission is controlled ($R_t < 1$), when does this happen and whether or not the disease transmission is increasing or decreasing.

3   Results

3.1   Question A: Data quality

In the 185 countries in the John Hopkins University COVID-19 dataset, 19 countries had any negative value (Antigua and Barbuda, Benin, Ecuador, France, Guyana, Honduras, Italy, Liechtenstein, Lithuania, Madagascar, Mauritius, Nepal, New Zealand, Portugal, San Marino, Spain, Uganda, Uruguay, Zimbabwe) and 20 countries had spurious positive peaks (Cameroon, Chile, Congo (Brazzaville), Congo (Kinshasa), Ecuador, Equatorial Guinea, Ethiopia, France, Gabon, Ghana, Israel, Kazakhstan, Netherlands, Nicaragua, Qatar, Spain, Sudan, Sweden, Zambia, Kosovo). The peak in Spain has been analyzed in the Sup. Material, 5.4. In the European CDC dataset , 10 countries had any negative value (Benin, Ecuador, France, Italy, Lithuania, Portugal, San_Marino, Spain, Uganda, United_Kingdom). It is very important to start any analysis with the visual exploration of the epidemic curve before and after smoothing.

We have observed the proportion of positive tests has significantly changed over the course of weeks/months (Sup. Material, 5.2). In Castile and León the daily proportion of positive tests in March nearly reached 60%; in Norway this proportion peaked at 17% in March; in USA it peaked at 23% in March and in Canada it peaked at 10% in April. Both the percentage of positives and the trend were different between USA (a developed country) and Perú (a developing one); in both cases the number of performed PCR tests gradually increased over from April to June 2020, providing more precise estimates of the real incidence as time passes. Other issues have been observed in the datasets: The reporting delay and diagnostic delay and associated factors (Sup. material, 5.1), the availability of different data sources for the same country (Sup. material, 5.8), the presence of large spurious peaks or frequent small peaks and valleys or the impact of the weekday bias.

3.2   Serial interval distributions

The point serial distribution estimates are displayed in the table 1, with 95% confidence intervals, and the serial interval distributions are available in Figure 1 .

Figure 1: Lognormal fit of the distributions of the serial interval

Table 1:

Serial Interval distributions as estimated via MCMC				
Source	Number of cases	Location	μ (CI 95 %)	σ (CI 95 %
Du et al	361	China	1.458(1.379,1.543)	0.778(1.379,1.543)
Nishiura et al	28	Articles and reports	1.385(1.167,1.599)	0.558(1.167,1.599)
Zhao et al	21	Hong Kong	1.211(0.852,1.546)	0.722(0.852,1.546)

3.3 Questions B, C. Estimation parameters: Preprocessing, GT/si and estimation method

The different smoothing and preprocessing options did have a minor/moderate impact on the transmission curves (Sup. material, 5.5); briefly speaking, multiplying each daily incidence by

a constant factor to correct the undetected proportion does not have any noticeable effect on the R_t curves or create any gap between transmission curves, as previously described(23). Both Lowess and SMA smoothing showed good results and performed better than one-sided mean averaging. Smoothing influenced the final R_t curves, but their relevant trends (faster or slower decrease and epidemic control) are mostly similar.

A rudimentary one-at-a-time sensitivity analysis has been performed, using the mean daily gap between non-overlapping Credible Intervals of the R_t as the target. (Sup. material, 5.6; it confirms the negligible impact of a constant proportion of undetected cases and shows that smoothing parameters might not be more influential than the Generation Time/serial interval distributions used. Smoothing can be specially important in datasets with spurious peaks (Sup. material, 5.4), and Lowess smoothing can provide the best results providing that the smoothing span in properly tuned. Smoothing can also correct the weekday-bias present in most datasets (Sup. material, 5.9, 5.10).

Figure 2 shows the influence of different Generation Time/serial interval distributions and estimation methods on the transmission curves. The fixed parameters include JHU dataset, incidences between 20-Feb-2020 and 14-Jul-2020, Lowess smoothing with a span of 6 days, backwards distribution of negative values and +0% undetected cases. (See sup. material 5.14 for another sensitivity analysis of several si/GT distributions).

Figure 2: Transmission curves with different methods and distributions

Some differences can be remarked: The credible intervals of the estimated R_t were narrower when the serial interval distribution obtained from the larger Du et al dataset was used, and they were wider when the smaller Zhao et al serial interval dataset was used. A visible delay was observed when estimating the R_t using the EpiEstim package (by Cori et al) compared with the R0 package (Wallinga Teunis method). (By default, the EpiEstim package calculates the R_t over a

window period of seven days; and this paper follows the default package behavior, in the epidemic curves the x-axis value is the last day of the window period).

Apart from this small differences, the results were largely equivalent and both methods deliver similar conclusions. However, given the smooth and continuous estimates obtained with the EpiEstim package, this will be used in the rest of this study.

3.4 Question D. Results in several countries

Daily incidences have been obtained for several large countries (Brazil, Belgium, Germany, Iran, Italy, Japan, Sweden, United Kingdom, Russia, India, South Africa, Spain, Singapore, US, Colombia, Mexico, Chile, Pakistan, Bangladesh, Peru) which exhibited a large number of cases and/or deaths, for the period between 2020-02-20 and 2020-07-14 using the JHU COVID-19 dataset. Figure 3 shows the epidemic curve of daily incidences by country after preprocessing and 4 shows the transmission curve of R_t by country after removing the particularly unstable initial periods. Preprocessing parameteris include assuming a time-constant proportion of 40% undetected cases (therefore, increasing the daily incidences by 66%), Lowess smoothing with 6 day span, backwards distribution of negative values. The Nishiura et al si MCMC distribution has been used. World Bank data (GDP per capita, PPP, in constant 2017 international dollars(40)) from 2019 (2017 for Iran) is used to colour the curves.

Figure 3: COVID-19 incidence in several countries

Daily incidences reflect several phases: All countries experienced **exponential transmission** in March; this happened earlier in some countries (Iran, Italy), and later in others (Russia, South Africa, India, Brazil). The pace was not uniform, and the increase was faster in Italy, Iran, and Spain. Several developed countries (Belgium, Germany, Italy, Spain, France, etc.) managed to "bend the curve", with decreasing daily incidences; but Sweden and USA only tentatively reached this in April/May. Developing countries (Russia, India, Brazil, South Africa) had more difficulties reaching decreasing daily incidences; Iran managed to control the epidemic at the end of March, but increasing incidences appeared again in May.

Italy and in particular Spain COVID-19 had an abrupt and exponential increase in daily incidences in late February / early March, respectively. Italy was the first European country to declare the State of Emergency (on the 31 January). On 6 March the Italian Anesthesiology and Intensive Care Association published their 'Recommendations for treatment admission and suspension in exceptional conditions of disbalance between needs and available resources' (45); this attests to the particular healthcare overload seen in this country around this time.

In late June and July some countries exhibit small increases in their incidences; this was also noted by the European CDC. (46)

Figure 4: R_t estimates for several countries. Highlighted dates: solid line = full scale measures/lockdown; dashed line = smaller scale measures

Several phenomena can be identified in R_t curves. The European developed countries quickly achieved a good control of the epidemic in April/May. Most countries implemented lockdowns,

which in some cases involved non-essential economy halting (Spain, Italy). In some countries the lockdown was established at sub-national levels (USA, Chile, Brazil, Russia), while in others the lockdown was at a national level (India, Spain, France, United Kingdom). The United Kingdom implemented lockdown measures late in March, replacing the mitigation strategy with a suppression strategy. Both Sweden and USA show average control, with R_t close to 1 and a plateau in the epidemic curve; the results of Sweden could be considered successful given the mitigation strategy used.

In Japan after an initial "valley" around the 15th March ($R_t \sim 1$ and $I_t = 31$), transmission increased, reaching Rt~1.5-1.8 at the beginning of April with $I_t = 563$ on the 16th of April). R_t started to decline around the 8th April (on 7 April a one-month state of emergency was decreed in the prefectures of Kanagawa, Saitama, Chiba, Osaka, Hyogo, and Fukuoka, and on 16 April the declaration was extended to the rest of the country for an indefinite period). However, no lockdown was implemented in the country.

Developing countries show worse epidemic control. Brazil and India have not consistently reached $R_t < 1$, Russia finally reached epidemic control and South Africa keeps an R_t slightly above 1. Iran also managed to control the R_t, but in late May the situation worsened.

Figure 5 displays the transmission curves of all world countries, using incidences and population values from the ECDC COVID-19 dataset(47), starting from 15-March-2020, coloured by their GDP per capita (with 2017 data from World Bank, Purchasing Power Parity in Constant 2017 dollars(40)) and using population as curve size. Preprocessing parameters include epidemic start after 50 cases in three days, lowess smoothing with 12 days as smoothing span, backwards distribution of negative values. R_t estimates have been obtained with the EpiEstim package by using the Du et al serial interval MCMC estimations. Countries with less than one million inhabitants and R_t estimates with 95% credible intervals wider than 1 have been excluded. This shows there is an important trend towards better epidemic control in countries with GDP per capita above 30000 USD dollars, especially in April/May (this difference is reduced in June, in the reopening phase). It should be remarked that many countries show later epidemic starts in April/May.

Figure 5: R_t estimates for large countries

3.5 Question E. Spain Regional R_t curves

Spain also experienced an accelerated COVID-19 evolution. The 9 Autonomous Communities with the most cases as of 2020-07-15 were selected and epidemic curves after preprocessing (Figure: 6) and transmission curves (Figure: 7) were obtained, by using the EpiEstim package(23) and the current official Health Ministry dataset (41), between 25-February-2020 and 14-July-2020. The epidemic start was defined as the moment with more than 15 cases in three days. Preprocessing parameters include: centered SMA smoothing, 5 days smooth span, backwards distribution of negative values, no modification of positive values, incidences increased by 40% to account for undetected cases.

Figure 6: COVID-19 incidence in several Spanish Autonomous Communities. Highlighted dates: 2020-03-08 (8M), 2020-03-14 (State of Alarm), 2020-03-28 (Economic stop), 2020-04-13 (Partial economic restart), 2020-05-04 (Return to 'normal', phase 0), 2020-06-21 (End of the state of Alarm)

Figure 7: R_t estimates for several Spanish Autonomous Communities. Highlighted dates: 2020-03-08 (8M), 2020-03-14 (State of Alarm), 2020-03-28 (Economic stop), 2020-04-13 (Partial economic restart), 2020-05-04 (Return to 'normal', phase 0), phases 1 and 2, 2020-06-21 (End of the state of Alarm)

In the case of Spain several periods can be identified: uncontrolled exponential dissemination took place early in March; in this period cases in Madrid grew earlier than other Autonomous Communities, along with the Basque Country and Catalonia. (These three Communities are some of the most densely populated regions, include the two largest cities in Spain, and they represent important economic centers with the 1st, 2nd and 4th highest GDP per capita in the country). In the week after the 8th of March the R_t started to decline, following the implementation of several

contact-prevention measures. The State of Alarm declaration on 14 March was associated with a fast and sharp decline in the virus transmission reaching R_t<1 before the end of April, effectively heralding adequate epidemic control. Although the exponential increase did not occur at the same time or pace, the decrease was quite similar between regions. On 28th of March all economic activity was halted except for essential activities. The R_t peak and the epidemic growing period differ slightly across the different Autonomous Communities: both were reached a few days later in Castilla-La Mancha, a less-densely populated region.

Most non-essential economic activities were resumed on 13th of April. At this point the R_t was below the epidemic threshold in all the regions; and remained controlled for the following weeks, showing some hills and valleys that could be caused by noise in the temporal series. The state of alarm ended on 21st June 2020, after several weeks of gradual reopening with controlled transmission. After this several regions exhibited a growth in the number of cases preceded by increased transmission: Aragón(48), Andalucía(49)... These outbreaks did not seem more prone to take place in wealthier and more densely populated areas (Madrid, Barcelona).

3.6 Question F. Superspreading events

Evaluating the COVID-19 spread in shorter periods requires a careful analysis of the underlying data. As incidence data includes important sources of variability and noise, time series smoothing is required; but too much smoothing can obscure the signal. Additionally, R_t is estimated in weekly window periods by default, providing smoother estimates. We have used data from the JHU-CSSE dataset(50). We have applied the EpiEstim(23) package to obtain daily and weekly R_t estimates. Smoothing was performed by using the centered SMA method with a span of 5 days; no proportion of undetected cases was assumed and the serial interval from the Du et al dataset was used. Figure 8 shows transmission curves of several countries in which events with very high transmission of SARS-CoV-2 has occurred.

Figure 8: Transmission curves of countries with alleged superspreading events

The Football Match between Atalanta and Valencia FC (19 February 2020) took place too early in the epidemic to be included in R_t estimates. In Spain the 8 March demonstrations and events took place several weeks later, in a period of exponential increase in the epidemic curve. A mild increase in R_t is observed around the 9th of March, both when using daily and weekly estimation windows. However the estimated R_t in the first two weeks of March are too unstable to draw any conclusion and the credible intervals are too wide. The events took place too early in the epidemic and definite conclusions cannot be drawn. The three-day R_t estimates from Spanish regions don't exhibit any consistent and relevant increase in the R_t around that date either (Sup. Material, 5.7), a mild but inconsistent R_t increase around that date is seen when using the initial cumulative incidence data (Sup. material, 5.8). In Spanish regions the incidence was too low around that date to perform any estimation.

In the USA there was an increase in the number of cases in late March, but this event did not take place at a particular well-defined date; again conclusions can't be drawn. R_t estimates did not noticeably increase either in the riots after the death of George Floyd on the 25th of May. State-level estimates are provided in the Sup. material, 5.11; no consistent increase in R_t is seen in early June: in some states a surge is seen (Arizona, Florida), but in most of them it is not, including Minnesotta (the state where this tragic event occurred), California, New York...

We have also noticed the apparition of peaks without any real-life event correlation; these peaks could be caused by noise in the time series or by reduced laboratory activity on the weekends (weekday bias, Sup. material, 5.9, 5.10), becoming apparent when increasing the sensitivity.

4   Discussion

Many factors can influence the process of obtaining transmission estimates. To start with, daily incidence data is required to perform the estimations, and data quality determines estimates quality. From the data analysis perspective, and as an answer to the question A (data quality), COVID-19 testing should be regarded a sampling process that delivers incidence data. The number of tests performed every day and the positive rate of the tests has greatly varied over time and by location. The daily incidences are biased downwards in periods of healthcare overload and resource limitations, and mild cases or case contacts have not been tested uniformly. Therefore, any utilization of COVID-19 incidence/prevalence time series data for most purposes, including modeling or estimation, should be preceded by an assessment of this sampling process; including the temporal evolution of the number of tests, the percentage of positive tests and whether or not testing limitations were present. The rate of positivity alone is probably not enough to estimate the real incidence in order to compare different countries. Other key issues that influence the quality of a dataset are the usage of cumulative reports to obtain individual incidences and the case definitions used.

Variability is also significant in some datasets; variability in R_t estimates is greater in locations with few COVID-19 cases, and variability in incidence time series is larger in countries with less GDP per capita. On the other hand, many developing countries have shown strikingly exhaustive, resourceful and complete datasets not seen in many developed countries, a testimony to the effort they are making(51)(52). Smoothing has been explored here as a possible solution to issues related to data variability and spurious peaks; it is particularly important to eliminate noise.

The weekday-bias was often present in the datasets used, so small estimation windows are not recommended.

The heterogeneity in data reports is another important issue that impairs comparisons between countries and regions. In the COVID-19 Tracking Project, each state in the USA is graded according to the quality and completeness of the data they provided. Some countries, states or regions limit their reports to the number of positive cases, and do not inform on the number of tests performed, or do not separate PCR and antibodies tests, or do not separate IgM and IgG antibodies tests. International standardization or consensus recommendations regarding COVID-19 reporting and data definitions could improve the data quality and facilitate comparisons.

4.1 Non pharmacological interventions to control the spread

We have obtained time-varying estimates for the R_t and transmission curves. The R_t values obtained in different countries follow the known pattern of transmission control in an epidemic when control measures are initiated. The reproductive ratio in the initial phase of the epidemic can also be estimated by using a compartmental SEIR model(3): in the UK before the lockdown period the R_0 was estimated as 6.94 (6.52-7.39, credible interval). Stochastic SIR models can also be used to obtain reproduction numbers in the initial phase without preventive measures (R_0) and reproduction numbers after preventive measures have been implemented (R_c, **control reproduction numbers**) which showed some variability between different regions but incontrovertibly demonstrated the efficacy of preventive measures(53). A compartmental model in Belgium also yielded similar estimates: The initial R_0 was 3.334 and the R after measures were taken in March was 0.713. (54)

The experience in Wuhan highlights the importance of Public Health Interventions to reduce the spread of the coronavirus. The epidemic was resulting in congested hospitals and high rates of transmission; but control was achieved through a combination of medical and preventive regulations: increases in healthcare capabilities and medical supplies, adequate patients isolation, transmission routes blocking (public transportation suspension, closure of public places, cordon sanitaire of Wuhan City, etc.) and the prevention of new infections (compulsory face-masks, universal stay-at-home policies and symptom surveys, etc) (55). The R_t values were directly dependent on the application of preventive measures, with a marked decline starting on the 24 of January after with the city lockdown, traffic suspensions and entertainment. In Wuhan in the early phases of the epidemic the R_t reached 4-5 but after epidemiological quarantine and isolation the R_t declined to less than 1 (55). Initial R_0 (February) estimates in Wuhan were slightly lower (average of estimates: 3.28, median: 2.79); this could be (56).

We have observed the R_t reflects the impact of the control measures and other factors, including socio-economical factors and cultural factors; and the incidence curves reflect the exponential effect of the R_t. Both in developed countries and developing countries, the R_t reflects the impact of Public Health Interventions to reduce the spread of the coronavirus. The suppression strategy has proven to be more effective, when widespread community transmission is taking place. In the case of Singapore, the third most densely populated country in the world, and South Korea, a successful containment was achieved in February and March 2020. The hallmarks of this strategy are travel restrictions, early detection and early quarantine of positive cases and their contacts, widespread testing, reduction in number of contacts and no or minor restrictions to economic activity(57). However in April in Singapore a new increase in R_t heralded increase a new wave with particular increase in daily incidences, several orders of magnitude

greater than the previous wave. In Europe, different non-pharmaceutical interventions have been applied by the authorities; in compartmental models the stay-at-home enforcement yielded the strongest reductions in normalized numbers of contacts(58).

In the case of Spain, initial **containment** efforts (such as the quarantine of an hospital) were successful(59), but soon it was apparent that the uncontrolled and exponential community transmission was taking place, as evidenced in both the epidemic curves and the R_t curves. In the 9th of March **reinforced containment** measures were initiated (halting in-person education, workplace contacts avoiding, etc.) for areas with significant community transmission (which were the region of Madrid and the cities of Vitoria and Labastida)(60); transmission started to decrease as education institutions and public places were gradually closed in the different regions. The Government decreed a State of Alarm on the 14th of March, switching to a **supression strategy** imposing a nation-wide lockdown involving commercial places closures, stay-at-home regulations and a ban on internal travels. Thus a consistently decreasing R_t and the effective control of the disease it hallmarks was reached(7), and Spain managed to successfully 'bend the curve'. The reproductive ratio has been shown to be highly dependent on mobility (4)(61).

On the other hand, some countries used a **mitigation-oriented strategy**: The Swedish Government declared that they seeked to "reduce the pace of the virus's spread" in order "to 'flatten the curve' so that large numbers of people do not become ill at the same time"(62). The epidemic curve of Sweden was indeed mostly flattened between April and July 2020. It could be argued that whatever policy was followed by developed countries, they are more likely to achieve it than developing countries.

4.2 Patterns and analysis of transmission curves: Developed and developing countries and regions

All transmission curves analyzed here follow the same pattern: exponential incidence increases in March, with higher but unreliable $R_t \sim 4$, incidence curve "bending" with reductions in R_t to or below 1, a possible apparition of outbreaks (R_t surges and incidence surges in June).

The transmission reduction was steeper in developed countries (Spain), except when they did not attempt a though suppression strategy from the start (United Kingdom). Both in developed and developing countries, richer and more densely-populated regions had earlier starts of the exponential propagation. Developing countries required more time to achieve epidemic control or did not achieve epidemic control at all in the period studied.

Developing countries can face multiple challenges when it comes to estimating transmission parameters, including a lack of testing facilities and lack of compliance with the "social distance" mandates(63). In developing countries the rise in the exponential phase of the epidemic was notably slower, but they had more difficulties than developed countries to control the epidemic. At the time of writing this, the R_t of India was and the R_t of Brazil was . In the case of India initial containment seemed successful, but was soon followed by widespread community dissemination that required more stringent preventive measures(64): social distancing, public places closures, whole country lockdown on the 24th of March... Wealthier and more densely populated regions of developed countries paradoxically exhibited a greater number of cases but they apparently seemed less prone to transmission surges or outbreaks.

After epidemic control, the apparition of outbreaks is effectively associated with R_t surges, as seen in the cases of Iran, Spanish regions, German regions, Singapore, etc. These outbreaks were not associated with more developed and wealthier regions, even though these regions were the first ones to suffer the COVID-19. We have shown both incidence curves and transmission curves can be used to detect these outbreaks, but transmission curves alone are not sufficient since they lack incidence data.

As a summary, in all the studied cases, the control of the transmission is effectively measured by the R_t and controlling the transmission translates into less incidence, both in developed and developing countries, and when this control is lost the incidences increase exponentially.

4.3 Superspreading events

Superspreading events are associated with a large number of new contagions. The scale ranges between small collective events involving hundreds-thousands of participants, being amenable to epidemiological investigation (Diamond Princess cruise, Annual Assembly of the Christian Open Church in France), and large, often closed, collective events involving tens of thousands of participants and not being easily amenable to epidemiological investigation (Atalanta vs Valencia FC match, 8M events in Spain). These took place in February or in early March, before lockdown measures were implemented, or despite preventive measures. It should be noted the 2-March-2020 Rapid Risk Assessment by European CDC recommend mass gatherings cancellation in exceptional cases for scenarios 1 or 2(5). The 8 of March in Spain was marked by several political and sports events; which were the subject of debate in Spain. Although the R_t curves in Spain showed some small peak of transmission around that date, this peak was not seen consistently in Spanish regions and its wide credible intervals make any interpretation questionable; therefore we could not find any conclusive evidence. These kind of events often take place in the early phases of the epidemic so the number of cases is small and the estimations are not stable or precise. Ultimately, this methodology could only show certain temporal correlation, but proper demonstration of a causal relationship or a "superspreading patient or event" requires a careful epidemiological investigation, which is usually performed in small-scale events. Daily estimation of R_t could be prone to methodological errors, as these data might reflect peaks and valleys of daily incidence caused by the increased laboratory and healthcare capabilities in workdays vs weekends.

4.4 Study limitations

This study has several limitations, which could be attributed to the underlying data, the uncertainty in the generation time or serial interval distributions used and the estimation methodology.

The proportion of tested cases is not expected to be uniform between different countries and between different stages in the epidemic; and this disparity has been also translated into clinical recommendations. Asymptomatic cases, close-contacts of a case and mild/home-managed cases of COVID-19 might not be tested due to resources constraints. In Spain, the initial protocol for home-management of positive cases remarks that patients with possible or probable COVID-19 with mild symptoms can be quarantined at home, and that in patients managed at home a confirmation PCR is not required(65); but later government protocols required a PCR should be performed in all suspicious cases(42). Similarly in the UK mild asymptomatic cases were also not tested at the

beginning of the epidemic(3). In the early stages of the pandemic, many laboratories might not have been ready for the exponential increase in PCR requests.

We have not observed relevant differences between the R_t curves obtained with reported cases and the R_t obtained if we increase each daily incidence by a constant proportion. It is expected that a significant proportion of cases is not reported: In Spain 5.0% and 5.2% of the surveyed population had tested positive for IgG antibodies against SARS-CoV-2 in the two phases of the National Seroprevalence Study(66), while only 0.42 and 0.52 of the participants in the study and 0.51 of the population on 03-Mar-2020 had had a positive PCR. Around 33% of infections were asymptomatic, which is consistent with the 43% of asymptomatic participants in the population-screening study in Iceland(67). Some authorities report suspicious cases; in the case of Catalonia these cases increased some weeks after the epidemic peak, when the epidemic was controlled and opening measures were being implemented; therefore they don't seem a good surrogate for non-reported cases (Sup. Material, 5.9).

COVID-19 cases are not reported in the same way in different places. For example, in Spain, the Community of Madrid defines the daily incidence I_t for day t as the number of positives whose sample was taken on day t or the result was obtained(68); this is better suited for temporal series analysis. In other Autonomous Communities, the daily incidence I_t is the number of positives reported on day t, thus introducing laboratory delays, increasing the weekday bias and reducing the series accuracy. In addition to this, negative incidence values or unexpectedly high incidences can be observed in the temporal series of some Autonomous Communities(68), which do not correspond to new cases but to data correction, validation and consolidation processes. This denotes there was room for improvement, probably towards the automation of data collection. When individualized epidemiological case-reports-derived statistics were available in May/June(69)(70), with the incidence defined by the symptoms-onset date, unnecessary delays were eliminated and data accuracy remarkably increased. Smoothing and preprocessing should be emphasized when dealing with irregular data. (As a minor comment, in some datasets, the labor-intensive nature of the data collection processes (71) could also be suspected from the longer reporting intervals seen as the epidemic is controlled; however, no criticism or recommendations can be made, since automated dataset creation has drawbacks as well and in any case this does not disparage open data publication).

An important assumption when using this models to estimate transmission is the isolation of the particular country/region/province in study: The initial cases are considered imported, and there are not any other imported/exported cases. This is reasonable in some cases as mobility restrictions are applied in many countries and regions around the epidemic peak. However, if the transmission is estimated for a small city or area, population mobility could alter the estimates (some cases might be exported and accounted for in another region, and some cases which were infected outside this region might be imported and accounted for in this region). The estimation method can deal with imported cases(38), but information on imported vs local cases was not available for most datasets.

Time intervals exist between the symptoms appearance and the diagnosis (the diagnosis delay), or the case report; this is another nuance of this methodology. Delays have been analyzed in the Colombia dataset (Sup. material, 5.1); a small association with month of the year has been found.

When the daily incidence I_d in a particular day d is defined as the number of new cases **with symptoms onset** on day d, this bias disappears; however most data sources report daily incidences as the number of new cases known to be positive and reported on day d. This problem impairs any analysis of individual dates, including superspreading events, and necessarily introduces a delay in the current transmission estimations. This could be solved in a similar way to asymptomatic transmission(23): If the distribution of the delay between infection and report is known, this could be used to back-calculate the incidence fof infections from the incidence of reports, but this yields oversmoothed estimates (31). Other reproductive number estimation methodologies went further than us by using linedata and bootstrap methods to obtain a delay distribution, which is used to change the incidence data(72).

Some COVID-19 datasets, like the Spanish National Public Health Surveillance Network (RENAVE), attempt to report the symptoms onset date, and thus are less prone to this bias (Sup. Material, 5.8). This is a partial solution because the symptoms onset is still not known for asymptomatic patients.

Using the serial interval as a surrogate for the generation time has a number of potential issues (31), and the serial interval distributions could be biased (overestimation/unestimation) due to several factors(29). For example, the serial interval might be contracted during the epidemic peak due to early quarantine of symptomatic infected individuals. We have not observed marked epidemiologically-relevant differences between the results obtained with any of the five distributions tested; so we believe it is often acceptable in practical terms to use the serial interval as a surrogate for the generation time.

Compartmental models can be used to estimate many infectious disease parameters, including the reproduction numbers(54)(58)(3). These models often divide estimate the initial reproduction number R_0 and the control reproduction number R_c (after control measures have been initiated). The reproduction numbers in the control phase we obtained are somewhat similar to the R_c, but the initial reproductive numbers we obtained are somewhat higher; these initial R_t seem somewhat unstable in the time-since-infection models used here, particularly if no clear distinction is made between local and imported cases when performing the estimations. Both types of models rely on their own assumptions, and can be considered as complementary. In time-since infection models no assumptions are made regarding the shape of the transmission curve.

One of the advantages of this study and the application is the possibility of interactively performing sensitivity analysis and exploring different smoothing and preprocessing parameters (or lack of them), we have shown smoothing had some influence in the R_t curves. However, the main information conveyed by the R_t estimates - namely, epidemic control - did barely change, although in some cases untreated spurious peaks can lead to wrong outbreak conclusions.

There is not a single definite way to obtain estimates of disease transmission to guide public policies; we have explored some of the most common ones. The information obtained with this tool can be combined with many other modeling tools available on the Internet.

Additionally, one of the most relevant implications for the reopening phase are the outbreaks. Robust epidemiological monitoring and rapid detection of increased transmission are essential to avoid resurgeces (46). We have observed outbreaks are amenable to detection with transmission curves, providing that quality datasets are available. Increased testing capacity is observed in many countries in June and July 2020, thus reducing noise and variability and improving incidence

data accuracy. Many online tools have been created to monitor the transmission of Coronavirus; compared to others this application puts more emphasis on interaction and parameter tuning.

Finally, one of the objectives of this task was to bring epidemic information closer to the general population. We hypothesize that population compliance with preventive measures (social distancing and mobility restrictions) might improve when adequate information is provided. In this application the users can interactively tailor several steps (data source, preprocessing/smoothing, estimation window, serial interval, etc) to obtain slightly different R_t estimations but nevertheless consistent with a fundamental idea: epidemic control requires lowering transmission.

5   Supplementary material

5.1 Question A. Dataset quality: Diagnostic delays

In Spain the National Epidemiological Vigilance Network (RENAVE) has described the interval between symptoms onset and diagnosis (median = 6 days, interquantile range (IQR) = 3-11 days, cases = 157885) and the interval between symptoms onset and case report (median=6 days, IQR = 3-11 days, cases = 169797 at 21-05-2020)(73). In the official Chilean dataset(52), the mean difference in days between symptom onset and case notification, as calculated from aggregate data, is 3.7.

The Colombia official positive cases dataset(51) has been used to assess the diagnostic delays (defined as the difference between symptoms onset and diagnosis) and the reporting delay (defined as the difference between diagnosis and web report).

13.15% of the reported cases as of 2020-07-15 were asymptomatic. The diagnostic delay for symptomatic cases has the following characteristics:

- Mean= 10.759

- SD= 10.759

- Percentiles 2.5, 25, 50, 75, 97.5 = 3, 6, 10, 14, 23

- Distribution = As shown in figure 9

Figure 9: Distribution of the diagnostic delay

A linear model has been created with log-transformed diagnostic delay as an dependent variable, and condition, ethnicity and month of diagnosis as predictors (Table: 3). Asymptomatic cases are excluded. This model was chosen even though log-transformed models are less interpretable. With an adjusted R^2 of 0.0160839, the model performance is poor, but many predictors are significant (table 3).

Table 2:

Colombia COVID-19 summary					
Condition	Freq Cond	month	Freq month	Ethnicity	Freq Ethn
Mild	115126	03-2020	905		70297
Asintomático	19670	04-2020	5563	Indigenous	2238
Deceased	5255	05-2020	22934	Black	7019
Severe	953	06-2020	66558	Other	70503
Moderate	9106	07-2020	51027	Palenquero	18
N/A	335			Raizal	73

Colombia COVID-19 summary

Condition	Freq Cond	month	Freq month	Ethnicity	Freq Ethn
				Rom	297

Table 3:

Linear model of diagnostic delay

	Estimate	Standard Error	t value	Pr(>|t|)
(Intercept)	2.217	0.018	122.803	0.000
ConditionDeceased	0.107	0.007	14.733	0.000
ConditionSevere	0.050	0.017	2.977	0.003
ConditionModerate	0.083	0.006	14.840	0.000
ConditionN/A	0.030	0.028	1.074	0.283
month04-2020	0.254	0.019	13.452	0.000
month05-2020	0.079	0.018	4.461	0.000
month06-2020	0.104	0.018	5.878	0.000
month07-2020	0.171	0.018	9.399	0.000
EthnicityIndigenous	0.010	0.012	0.788	0.431
EthnicityBlack	0.041	0.008	5.413	0.000
EthnicityOther	-0.044	0.005	-9.265	0.000
EthnicityPalenquero	-0.132	0.127	-1.041	0.298
EthnicityRaizal	-0.038	0.065	-0.588	0.556
EthnicityRom	-0.037	0.031	-1.202	0.229

*Signif. codes: 0 <= '***' < 0.001 < '**' < 0.01 < '*' < 0.05 < '.' < 0.1 < '' < 1*

Residual standard error: 0.5068 on 127321 degrees of freedom
Multiple R-squared: 0.01619, Adjusted R-squared: 0.01608
F-statistic: 149.7 on 127321 and 14 DF, p-value: 0.0000

Mild cases of March with non-specified ethnicity were used as a reference. Diagnostic delay was greater in April, May and June when compared to March, and greater in patients who died, and greater in black or indigenous patients, but smaller in patients of other ethnicities. However, these differences were mostly small in this dataset.

The following table summarizes the distribution of the reporting delay.

Table 4:

Reporting delay (days)	Number of reports
<0	231
0	135825
1 to 5	9934

Reporting delay (days)	Number of reports
>5	997

Differences in diagnostic and reporting infrastructure might explain these results.

5.2 Question A. Number of tests and rate of positivity

The **rate of positivity**, the number of samples tested positive for COVID-19 in a given location and period, is defined by the WHO as one of the epidemiological criteria of epidemic control: It is met when less than 5% of samples test positive for COVID-19 for at least the previous 2 weeks, but this can only be interpreted when there is thorough surveillance and testing of positive cases(15); this criteria has been translated into national protocols(13) and monitoring tools(74). Herein we use the number of tests and the rate of positivity to assess temporal fluctuations/differences/constraints in the testing process, so the proportion of undetected cases could also vary over time, and the real incidence might particularly be underestimated when the rate of positivity is high.

Five datasets have been used to analyze the temporal distribution of testing efforts and, when available, the daily rate of positivity: The Perú data provided by the Peruvian Ministry of Health (MINSAL) and compiled by JM Castagnetto (75), the Canadian Government Data on COVID-19 in Canada (76), the USA data from the John Hopkins COVID-19 Testing Insights Initiative (CC BY-NC-4.0 license), as reported by USA states(74), dataset containing the number of COVID-19 tests performed in the region of Cantabria, Spain, between 2020-02-29 and 2020-06-24 and the dataset containing the number of COVID-19 tests performed in the region of Castilla y León, Spain(71), until 18-May-2020 (CC BY-3.0 license).

The number of COVID-19 PCR and antibodies tests performed each day are graphically reported, and the proportion of positive results is also reported, when available. A smoothing LOESS trendline is added due to considerable variability between days and due to a "weekday-bias" effect particularly noticeable in antibodies tests. Figure 10 shows the COVID-19 PCR tests performed in the USA, figure 11 shows the information for Peru, figure 12 shows the tests performed and the positive rate in Canada; with a possible nuance in the former dataset: some USA states might not have clarified what kind of tests they are reporting (PCR or antibodies).

Figure 10: PCR tests performed in the USA

Figure 11: COVID-19 tests performed in Peru

Figure 12: COVID-19 tests performed in Canada

Both USA and Canada have shown a steady and almost linear increase in testing capacity over time, reaching 600000 and 40000 daily tests respectively at the end of June. Perú also linearly increased their testing capacity over time. In both countries the daily percentages of positive tests have evolved: in the USA the rate of positivity approached 20-25% in April, but then a marked decline was observed and a plateau of ~5% positive tests was reached in May, but in late June, the rate of positives started to increase mildly. In Canada the peak of positives was reached in mid April, around 9%. In Perú the percentage of positive PCRs remained uniform around 30% between April and June, and no information is available for earlier months.

Figure 13 shows the numbers of PCR and antibodies tests performed in this Northern Spanish region. Only the number of performed tests is reported.

Figure 13: COVID-19 tests performed in Cantabria, Spain

Figure 14 shows the numbers of PCR and antibodies daily tests performed in Castilla y León, and the percentage of positive results for the whole region; figure 15 show the province-level information.

Figure 14: COVID-19 tests performed in Castilla y León, Spain

Figure 15: COVID-19 tests performed in Castilla y León provinces, Spain

In both regions of Spain a considerable **weekday-bias** is observed. The PCR testing capacity of Cantabria increased in March and peaked in May. In Castilla y León the rate of positive tests was higher than 40% in March, which could have implications for underestimation of the daily incidence. Both regions showed a stable number of PCR tests performed in May/June, coinciding with epidemic control and reopening phases.

5.3 Questions A, D. Data variability

We define variability here as the mean absolute error of the original incidences vs the Lowess-transformed incidences, using a smoothing span of 7 observations. (This absolute error definition might overestimate the effect of days with large incidences; however the relative counterparts might have the opposite effect). The European CDC dataset (47) has been used to calculate the variability per country and the total sum of cases by country, using data between 25-Feb-2020 and 14-Jul-2020; this information has been combined with the OECD economic information (GPD per capita, in US dollars, PPP, 2018). Figure 16 shows the relationship between variability and total number of cases; countries with less than 100 cases are excluded.

Figure 16: Variability vs total number of cases

Although the main factor that explains the mean sum of residuals of the Lowess fit, variability is also related to the GDP_per_capita.

A linear model has been fit using log(variability) as the dependent variable and log(sum_of_cases) and log(GDP_per_capita) as predictors, for countries with at least 100 cases as of 2020-07-15. In this model the R^2 was as high as 0.909 and the GDP_per_capita reached statistical significance.

Table 5:

Linear model of variability

	Estimate	Standard Error	t value	Pr(>|t|)
(Intercept)	-2.948	0.028	-106.349	0.000

Linear model of variability

	Estimate	Standard Error	t value	Pr(>|t|)
log(sum_of_cases)	0.741	0.002	470.798	0.000
log(GDP_USD_PC)	-0.103	0.003	-35.152	0.000

Signif. codes: 0 <= '***' < 0.001 < '**' < 0.01 < '*' < 0.05 < '.' < 0.1 < '' < 1

Residual standard error: 0.4964 on 23471 degrees of freedom
Multiple R-squared: 0.9091, Adjusted R-squared: 0.9091
F-statistic: 1.173e+05 on 23471 and 2 DF, p-value: 0.0000

5.4 Questions A, B. The impact of spurious peaks: Catalonia

On the 10th May 2020 Catalonia reported 2721 positive cases respective to the previous day; but only 83 of these cases were new cases(68). (This illustrates a problem that might arise when obtaining daily incidence data from the differences of cumulative total cases: Some health authorities often report cumulative total cases every day; these reports in a particular date might include both the new cases diagnosed that day (incident cases) and cases whose date of symptoms or date of diagnosis is not known). Therefore, some cumulative datasets include a disclaimer against the obtention of incidences by differentiation. An older cumulative version of the Spain Ministry of Health COVID-19 dataset (77), accessed on 21-May-2020, has been used; it should be noted that newer versions correct this issue (Sup. material, 5.8).

Here we assess the impact of the spurious peaks R_t estimates and how different smoothing methods deal with them, including smoothing and proper value replacement. The epidemic curves after preprocessing are shown in figure 17 and the transmission curves are shown in figure 18. Standard parameters are used (backwards distribution of negative values, no proportion is assumed of negative values).

Figure 17: Epidemic curves before and after several smoothing methods

Compared to the SMA, the Lowess smoothing is more tolerant with peaks (and in some settings might not correct them); and the SMA is more influenced by these peaks. This reflects the different philosophy of these methods (mean averaging vs polynomial regression). A second-pass SMA might be useful in some particular cases.

Figure 18: Transmission curves before and after several smoothing methods

The spurious peak causes spurious R_t estimates if it is not properly corrected. Lowess correction seemed a reasonable method when dealing with these peaks, but proper visually-guided tuning of smoother span is recommended, as small spans could left some peaks. When the SMA correction was applied, spurious R_t estimates were obtained.

5.5 Question B. The impact of different preprocessing options

The impact of the different preprocessing options on the transmission curves is evaluated in the figure 19. The following parameters have been used: JHU dataset, dates between 10-March-2020 and 14-July-2020, different smoothing parameters and spans, backwards distribution of negative values, no specific correction for spurious peaks.

Figure 19: Sensitivity analysis for different preprocessing parameters

5.6 Questions B, C. One-at-a-time sensitivity analysis

We numerically evaluate the differences between different conditions by using the mean daily gap between non-overlapping R_t credible intervals. The Gap G_i at time point i between two credible intervals X and Y obtained with the same data but some different parameter is defined as:

$$G_i(X_{i0.025}, X_{i0.975}, Y_{i0.025}, Y_{i0.975}) = \begin{cases} (1 + X_{i0.025} - Y_{i0.975})^2 - 1, & \text{if } X_{i0.025} > Y_{i0.975} \\ (1 + Y_{i0.025} - X_{i0.975})^2 - 1, & \text{if } Y_{i0.025} > X_{i0.975} \\ 0, & \text{otherwise} \end{cases}$$

where $X_{i0.975}$ and $X_{i0.025}$, respectively, are the upper and lower posterior limits of the 95% credible interval of the reproduction number at time i. The Gap defined here increases with the square of the distance between non-overlapping credible intervals. The existence of this Gap

between the limits of the credible intervals reflects there is room for different information and interpretations of the transmission.

The Gap between two transmission curves is the average daily gap, (defined as $G = \frac{\sum_{j}^{N} G_j}{N}$, where N is the total number of days in which both transmission curves are defined).

5.6.1 Generation interval/serial intervals (GT/si), distributions obtained via MCMC

For each of four countries (Spain, Brazil, USA, United Kingdom) with incidences from the JHU-CSSE dataset ([50]) between 10-March-2020 and 14-July-2020, the Gap between pairs of R_t curves obtained with different generation time distributions as the only varying parameter is calculated. As described in 2, the datasets from Du et al, Nishiura et al and Zhao et al are used to obtain estimates of the serial interval distribution via MCMC by using the original data; and the generation time estimated parameters described by Ganyani et al with data from Singapore or Tianjin are used to sample generation time distributions. 250 distributions are used due to computational constraints.

Several estimation parameters are kept constant: 0% of undetected rate, backwards distribution of negative values, no specific modification of spurious positive values, centered SMA smoothing with a period or order of 4 days. The results are displayed as triangular matrix of Gap.

Spain:

Table 6:

Sens. analysis: GT/si. Spain. Mean daily GAP					
	Du	**Nishiura**	**Zhao**	**Ganyani Singapore**	**Ganyani Tianjin**
Du	0				
Nishiura	0	0			
Zhao	0.02	0.021	0		
Ganyani Singapore	0.002	0	0.003	0	
Ganyani Tianjin	0.01	0.006	0.012	0	0

Brazil:

Table 7:

Sens. analysis: GT/si. Brazil. Mean daily GAP					
	Du	**Nishiura**	**Zhao**	**Ganyani Singapore**	**Ganyani Tianjin**
Du	0				
Nishiura	0	0			
Zhao	0.015	0.019	0		
Ganyani Singapore	0.001	0	0.005	0	
Ganyani Tianjin	0.005	0.002	0.009	0	0

United States:

Table 8:

	Du	Nishiura	Zhao	Ganyani Singapore	Ganyani Tianjin
			Sens. analysis: GT/si. USA. Mean daily GAP		
Du	0				
Nishiura					
Zhao	0	0			
Ganyani Singapore	0.045	0.05	0		
Ganyani Tianjin	0.001	0	0.018	0	
Du	0.009	0.008	0.034	0	0

United Kingdom:

Table 9:

	Du	Nishiura	Zhao	Ganyani Singapore	Ganyani Tianjin
			Sens. analysis: GT/si. United Kingdom. Mean daily GAP		
Du	0				
Nishiura	0	0			
Zhao	0.011	0.012	0		
Ganyani Singapore	0	0	0.003	0	
Ganyani Tianjin	0.006	0.004	0.005	0	0

5.6.2 Generation interval/serial intervals (GT/si), point estimates

This section is identical to the previous one, with the difference that for each of the five previously mentioned *GT* or *si* sources a single distribution is obtained by using point estimates (Pt), (instead of a sample of *GT* or *si* distributions obtained via MCMC or bootstrap).

Spain:

Table 10:

	Du Point	Nishiura Pt	Zhao Pt	Ganyani Singapore Pt	Ganyani Tianjin Pt
			Sens. analysis: GT/si. Spain. Mean daily GAP		
Du Point	0				
Nishiura Pt	0	0			
Zhao Pt	0	0	0		
Ganyani Singapore Pt	0.126	0.126	0.126	0	

Sens. analysis: GT/si. Spain. Mean daily GAP

	Du Point	Nishiura Pt	Zhao Pt	Ganyani Singapore Pt	Ganyani Tianjin Pt
Ganyani Tianjin Pt	0.126	0.126	0.126	0	0

Brazil:

Table 11:

Sens. analysis: GT/si. Brazil. Mean daily GAP

	Du Point	Nishiura Point	Zhao Point	Ganyani Singapore Point	Ganyani Tianjin Point
Du Point	0				
Nishiura Point	0	0			
Zhao Point	0	0	0		
Ganyani Singapore Point	0.03	0.03	0.03	0	
Ganyani Tianjin Point	0.03	0.03	0.03	0	0

United States:

Table 12:

Sens. analysis: GT/si. USA. Mean daily GAP

	Du Point	Nishiura Point	Zhao Point	Ganyani Singapore Point	Ganyani Tianjin Point
Du Point	0				
Nishiura Point	0	0			
Zhao Point	0	0	0		
Ganyani Singapore Point	0.171	0.171	0.171	0	
Ganyani Tianjin Point	0.171	0.171	0.171	0	0

United Kingdom:

Table 13:

Sens. analysis: GT/si. United Kingdom. Mean daily GAP

	Du Point	Nishiura Point	Zhao Point	Ganyani Singapore Point	Ganyani Tianjin Point
Du Point	0				
Nishiura Point	0	0			
Zhao Point	0	0	0		

	Du Point	Nishiura Point	Zhao Point	Ganyani Singapore Point	Ganyani Tianjin Point
Ganyani Singapore Point	0.006	0.006	0.006	0	
Ganyani Tianjin Point	0.006	0.006	0.006	0	0

When creating the si / GT distribution with point estimates, there was a Gap between the credible intervals obtained via Generation Time distributions, and the credible intervals obtained via Serial Intervals distributions.

5.6.3 Smoothing

For each of four countries (Spain, Brazil, USA, United Kingdom), the Gap between pairs of R_t curves obtained with different smoothing orders as the only varying parameter is calculated; the smoothing order is the number of points used for each point smoother. Several estimation parameters are kept constant: 0% of undetected rate, backwards distribution of negative values, no specific modification of spurious positive values, centered SMA smoothing, serial interval distribution obtained from Du et al(29). The results are displayed as triangular matrix of Gap.

Spain:

Table 14:

Sens. analysis: Smoothing. Spain. Mean daily GAP

	2	5	7
2	0		
5	0.001	0	
7	0.005	0	0

Brazil:

Table 15:

Sens. analysis: Smoothing. Brazil. Mean daily GAP

	2	5	7
2	0		
5	0	0	
7	0.001	0	0

USA:

Table 16:

Sens. analysis: Smoothing span. USA. Mean daily GAP

	2	5	7
2	0		
5	0	0	
7	0	0	0

UK:

Table 17:

Sens. analysis: Smoothing span. United Kingdom. Mean daily GAP			
	2	5	7
2	0		
5	0	0	
7	0.003	0	0

5.6.4 Undetected ratio

For each of four countries (Spain, Brazil, USA, United Kingdom), the Gap between pairs of R_t curves obtained with different proportions of undetected ratio (0%, 40%, 95%; which is used to calculate the total estimated incidence) as the only varying parameter is calculated; the smoothing order is the number of points used for each point smoother. Several estimation parameters are kept constant: 0% of undetected rate, backwards distribution of negative values, no specific modification of spurious positive values, centered SMA smoothing, serial interval distribution obtained from Du et al(29). The results are displayed as triangular matrix of Gaps.

Spain:

Table 18:

Sens. analysis: Undetected rate. United Kingdom. Mean daily GAP			
	0	**0.4**	**0.95**
0	0		
0.4	0	0	
0.95	0	0	0

Brazil:

Table 19:

Sens. analysis: Undetected rate. Brazil. Mean daily GAP			
	0	**0.4**	**0.95**
0	0		
0.4	0	0	

	0	0.4	0.95
0.95	0	0	0

Sens. analysis: Undetected rate. Brazil. Mean daily GAP

USA:

Table 20:

	0	0.4	0.95
0	0		
0.4	0	0	
0.95	0	0	0

Sens. analysis: Undetected rate. USA. Mean daily GAP

UK:

Table 21:

	0	0.4	0.95
0	0		
0.4	0	0	
0.95	0	0	0

Sens. analysis: Undetected rate. United Kingdom. Mean daily GAP

As apparent from the GAPs, the GT/si distribution chosen is the factor that most influences the differences in transmission curves. However the mean daily differences were small. Undetected rate has a negligible impact when a naïve proportional correction is applied.

5.7 Question C,E. Spain regional estimates, three-day windows

As a counterpart for the weekly estimates for Spanish regions, three-day estimates are obtained and displayed here for reference purposes. The same setting as are used as in the previous case.

R_t estimates for several Spanish Autonomous Communities. Three-day estimation windows. Highlighted dates: 2020-03-08 (8M), 2020-03-14 (State of Alarm), 2020-03-28 (Economic stop), 2020-04-13 (Partial economic restart), 2020-05-04 (Return to 'normal', phase 0),phases 1 and 2, 2020-06-21 (End of the state of Alarm)

Given the significant weekday bias, three day windows are not recommended.

Daily estimates are more susceptible to noise and show a striking weekday/weekend bias, and don't provide much useful information when compared to weekly R_t estimates.

5.8 Question A, E. Spain: Public health surveillance data source

In Spain the New Normality Transition Plan established the daily automated and individual notification of cases to the national Health Surveillance authorities for some Epidemic indicators(69); thus highlighting the importance of quality individualized reports of cases, instead of plain cumulative reports. The switch between cumulative reports and individualized epidemiological reports was not easy and exemplifies the challenges that many countries faced(70). The Spanish Public Health Surveillance Network (RENAVE) collects COVID-19 reports via the SIVIES system(73), and symptoms-onset dates are used to calculate the incidence. The median interval between symptoms onset and report is six days, and in this dataset when symptoms onset date is unknown it is assumed to be the diagnosis date minus 6 days. The purpose of this section is to show the improvements in daily incidences when proper epidemiological reports are used.

Figure 20 compares the incidence reported via the RENAVE and the deprecated previous cumulative report of the government. Figure 21 displays the transmission curve (parameters include SMA smoothing, 4 day smoothing span, estimation start with >15cases in 3 days, 40% proportion of undetected cases).

Figure 20: Incidences for Spain and regions (Initial cumulative reports (Cum_MSC) vs individualized reports(I_RENAVE)). Highlighted dates: 2020-03-08 (8M), 2020-03-14 (State of Alarm), 2020-03-28 (Economic stop), 2020-04-13 (Partial economic restart), 2020-05-04 (Return to 'normal', phase 0).

Figure 21: R_t estimates for Spain and regions (Government vs public health data source). Highlighted dates: 2020-03-08 (8M), 2020-03-14 (State of Alarm), 2020-03-28 (Economic stop), 2020-04-13 (Partial economic restart), 2020-05-04 (Return to 'normal', phase 0).

The epidemiological symptoms-onset report is less prone to weekday bias (which was particularly important when the incidence was decreasing) and exhibits the effect of the diagnosis/reporting delay in the cumulative data. The variability, as defined in 5.3, is greater in the initial cumulative dataset (314) compared to the improved RENAVE dataset (172).

5.9 Question B. Catalonia: Confirmed vs suspected cases, weekend smoothing

The official Catalonia open data COVID-19 repository(44) has been analyzed. This data includes several time series corresponding to different case definitions: Positive PCR, antibodies test, rapid tests, suspected cases; thus providing insights on the evolution of suspected vs confirmed cases over time. Additionally, the impact of smoothing and the **weekday bias** (a decrease in the assumed daily incidence in the weekends due to the laboratories having reduced activity).

The application and code have been used to obtain epidemic and transmission curves. Mild smoothing is defined as **3** days of smoothing with the "centered SMA" method, and Strong smoothing is defined as **7** days of smoothing with the "centered SMA" method; other parameters include: negative values are distributed backwards before smoothing, spurious positive peaks are

not removed and the undetected rate is zero. Figure 22 displays the epidemic curves after preprocessing and figure 23 displays the transmission curves.

Figure 22: Epidemic curve of Catalonia. Cat_Abs: Antibodies. Cat_PCR: PCR tests. Cat_RAP: Rapid tests. Cat_Susp: Suspected cases. Cat_Susp_Conf: Suspected and Confirmed cases. Highlighted dates: 2020-03-14 (State of Alarm); 2020-05-04 (Phase zero, reopening); 2020-06-21 (End of the State of Alarm)

In Catalonia, Spain, epidemic curves show a significant show a pattern seen in other datasets: the incidence decreases in weekends, maybe due to laboratory workdlows and the role of physicians and epidemiologists.

The number of positive PCRs has decreased over time after the epidemic peak, signaling an effective epidemic control. On the other hand suspected cases have kept increasing after the epidemic peak while the confirmed cases declined; this could be explained by many reasons/hypotheses (non-specific symptoms associated with COVID-19, seasonal allergies, increased public awareness, increased Primary Care and Public Health throughput, a shifting from care-at-home to early testing…), but in any case it demonstrates Suspected cases cannot be reliably used as a surrogate for undetected cases or total cases, at least in this dataset). Truly underreported cases should specially increase near the epidemic peak due to both asymptomatic cases and laboratory limitations, but here "suspected" cases increase after the epidemic peak.

Variability exists between regions and countries regarding the management of these suspected cases. In the epidemic peak many countries and guidelines don't require a PCR is performed in all suspected cases due to constrained resources.

Figure 23: R_t estimates for Catalonia. Highlighted dates: 2020-03-08 (8M), 2020-03-14 (State of Alarm), 2020-03-28 (Economic stop), 2020-04-13 (Partial economic restart), 2020-05-04 (Return to 'normal' (phase 0)), gradual reopening, 2020-06-21 (End of State of Alarm).

Both smoothing the time series and using longer time windows for the estimations can reduce the impact of weekend fluctuations in the epidemic curves and in the transmission curves.

5.10 Question B. Belgium: The impact of smoothing

The official Belgium dataset, collected by the Sciensano institute,(78) has been analyzed using the application and code in order to assess the impact of smoothing and the "weekday bias". Mild smoothing is defined as **3** days of smoothing with the "centered SMA" method, and Strong smoothing is defined as **7** days of smoothing with the "centered SMA" method; other parameters: negative values are distributed backwards before smoothing, spurious positive peaks are not removed and the undetected rate is zero. Figure 24 displays the epidemic curves after preprocessing and figure 25 displays the transmission curves.

Figure 24: Epidemic curves of Belgium

Figure 25: Transmission curves of Belgium

As seen in the case of Catalonia, using smaller window periods for the R_t estimation is equivalent to using smaller SMA smoothing periods. The differences between Belgian divisions are small, and the R_t intervals become wider as the incidence decreases.

5.11 Question E. USA: The state estimates of a developed country

The state-level number of cases from the USA between 15-March-2020 and 14-July-2020 has been analyzed; dataset is from The New York Times, based on reports from state and local health agencies(79). Preprocessing options include SMA centered smoothing, 7 day as smoothing span, epidemic start after 15 cases/three days. Figure 26 displays the epidemic curve after preprocessing and figure 27 displays the transmission curves; in the transmission curve the states have been coloured according to their GDP per capita (purchase power parity, current prices) (43) and in the epidemic curve the date of the death of George Floyd has been highlighted, after which riots ensued (section 3.6).

Figure 26: COVID-19 incidence in several USA states

Figure 27: R_t estimates for USA states

Some wealthy and densely-populated states (New York, New Jersey, Massachusetts, Connecticut, Columbia) have shown higher incidences than other states and they kept the epidemic under control from April to June, although most of them reached a R_t close to 1 by the

start of July. In other states (California, Washington, North Dakota, Delaware) epidemic control was slightly worse. In some states with lower GDP per capita the epidemic control was less consistent (Alabama, Arizona, Arkansas, South Carolina, West Virginia). However, this findings are subtle and not very well-defined; so there are other factors influencing between-state differences in transmission.

New York State has arguably reached the best COVID-19 control so far, albeit this could be explained by the peak of daily incidences>10000 in April. It could be speculated that some particular peaks/valleys could be attributed to real-life events, such as the meat packing plants cases in Nebraska in April. Looking for life-events to attribute them to peaks could lead to flawed conclusions due to the confirmation bias.

Many states showed an incipient rebound in transmission in late June: Florida, Oklahoma, South Carolina, Oregon...); small increases in the R_t above 1 in this period were associated with thousands of new cases (Texas, California, Arizona, Florida).

5.12 Question E. Peru: The regional estimates of a developing country

Using the MINSAL incidence data between 15-March-2020 and 14-July-2020, compiled by JM Castagnetto(75), the regional epidemic curves (figure (28)) and transmission curves (figure 29) of Perú have been obtained. Preprocessing parameters include: SMA smoothing, 7 days as smooth span, +40% proportion of undetected cases, backwards distribution of negative values and epidemic start after >15 cases / 3 day period. In the transmission curve the states have been coloured according to their GDP per capita (purchase power parity, current prices) (43).

Figure 28: COVID-19 incidence in several Peruvian regions

Figure 29: R_t estimates for Peruvian regions

In regions with smaller GDP per capita the epidemic apparently started later.

When compared to the regions in developed countries (Spain, Belgium), the regions in Perú show a marked pattern of variability (more peaks and valleys, maybe due to smaller incidences) and more difficulty reaching control; similarly to the infection curves in other developing countries. Many regions exhibit peaks, but not all of them had the same impact on the incidence:

The Moquegua and Pasco peaks reflected an increase from around 5 reported cases per day to 20 reported cases per day; the Piura peak was caused by an increase from less than 200 cases per day to 500 cases per day. This information should be interpreted with caution; the daily reported incidence could be influenced by testing capacity. The subtle pattern seen in USA states (wealthier states have more cases but better epidemic control) is not clearly observed here.

Overall, both regions in USA and regions in Perú follow a similar pattern, and the GDP per capita was not associated or mildly associated to differences the transmission curves.

5.13 Question G: Implications for reopening (Germany)

In Germany, in late June, after the reopening, several cases took place in clusters and the authorities managed them with a containment policy: In the middle-late of June an outbreak took place in an apartment building in Berlin(80), requiring quarantine measures. On the 24th of June the districts of Warendorf and Gütersloh in the state of North Rhine-Westphalia were put in lockdown following an outbreak in a meat processing plan(81). The purpose of section is to evaluate whether these outbreaks were reflected in the R_t of affected areas and whether non-affected areas showed any R_t impact. By using the Germany COVID-19 dataset compiled by J Gehrcke from the information provided by the Robert-Köch-Institute(82), R_t estimates from 20-March-2020 to June have been obtained, for the districts of Gütersloh, Warendorf, the state of Westphalen, three "randomly" selected districts (Eisenach, Leipzig and Weimar), the state of Berlin and the state of Brandenburg. The preprocessing parameters include centerd SMA smoothing with a smoothing span of 4, backwards distribution of negative values, no specific modification of positive spurious peaks, 0% undetected rate and starting after 10 cases/3 days. Figure 30 shows the epidemic curves and figure 31 shows the transmission curves.

Figure 30: Epidemic curves of Germany

We can observe the scale of the epidemic is one-two orders of magnitude higher in states than in districts; the incidence in districts might be too low to obtain R_t estimates.

Figure 31: Transmission curves of Germany

The transmission curves show a significant increase in the districts of Gütersloh and Warendorf and even in the state of Westphalen in late June. An increase was also observed in Berlin and in Germany. On the other hand, these curves exhibit a non-significant increase in Eisenach, Leipzig and Weimar; this could be caused by a few "imported" cases. Three-day windows did not help to detect the outbreaks and showed more noise. Thus a surge in the epidemic curves accompanied by a significant increase in the transmission curves could be used as a signal to detect outbreaks.

5.14Question C, G: Implications for reopening (outbreaks in Catalonia)

On 04-July-2020 in the county of Segrià in Lerida, Catalonia, Spain, was quarantined ("cordon sanitaire"), following a surge in the number of cases(83); this comarca neighbours the Autonomous Community of Aragón, where an outbreak was taking place(48). Using the Official Catalonia Basic Health Areas dataset(44), R_t have been obtained for several Basic Health Areas in Catalonia, using several si/GT distributions and keeping preprocessing settings constant (centered SMA smoothing with a span of 4 days, backwards distribution of negative values, no modification of spurious peaks, no correction for undetected cases). Figure 32 displays the incidences in different Basic Health Areas, with the dates of the end of the national State of Alarm and the start of the "cordon sanitaire", and figure 33 displays the transmission curves. The date when the state of alarm ended (2020-June-21) and the day when new travel restrictions were issues have been marked.

Figure 32: Epidemic curves of Catalonia

Although the initial outbreak of Coronavirus in March-April had a more pronounced in number of cases in regions with more population (Barcelona Metropolitan Area), the outbreak in the end of June/beginning of July affected the less populated Area of Lleida.

Figure 33: Transmission curves of Catalonia

Two R_t surges can be seen in the Lleida area: One around May, in which daily incidences rose from less than 10 cases/day to 40-50 cases/day, and one in July, in which daily incidences rose from around 10-20 cases day in mid June to 80-90 cases/day at the end of June. Therefore, the information provided by the R_t needs to be complimented by the incidence information. This confussion can be solved by using the "overall infectivity" magnitude, as seen in figure 34. Overall infectivity at time t is defined as the sum of all the previously infected individuals, multiplied by their infectivity at time t (given by the serial interval distribution)(23). Overall infectivity has been calculated by using the mean of the 250 *si* or *GT* distribution samples obtained via MCMC or sampling (section 2).

Figure 34: Overall infectivity curves of Catalonia

Overall infectivity can therefore provide an accurate picture of outbreaks. It mostly follows the pattern of daily incidences epidemic curves. But compared to daily incidences, it is less influenced by weekday bias, but it might be influenced by the *si* or *GT* distribution, albeit very slightly.

Author disclosures: Bautista Balbás LA: Theoretical work, programming, scientific analysis, manuscript writing. Gil Conesa, Rodríguez Caravaca, Bautista Balbás, B: Manuscript review.

6   Bibliography

Note: Unless otherwise specified, all datasets have been used in their latest update available as of July 15, 2020.

1. Delamater PL, Street EJ, Leslie TF, Yang YT, Jacobsen KH. Complexity of the basic reproduction number (r0). Emerging Infectious Diseases. 2019 Jan;25(1):1–4. Available from: https://doi.org/10.3201/eid2501.171901

2. Wan H, Cui J-a, Yang G-J. Risk estimation and prediction by modeling the transmission of the novel coronavirus (COVID-19) in mainland china excluding hubei province. 2020 Mar; Available from: https://doi.org/10.1101/2020.03.01.20029629

3. Dropkin G. COVID-19 UK lockdown forecasts and r0. Frontiers in Public Health. 2020 May;8. Available from: https://doi.org/10.3389/fpubh.2020.00256

4. Arenas A, Cota W, Gomez-Gardenes J, Gomez S, Granell C, Matamalas JT, et al. Derivation of the effective reproduction number r for COVID-19 in relation to mobility restrictions and confinement. 2020 Apr; Available from: https://doi.org/10.1101/2020.04.06.20054320

5. European Centre for Disease Prevention and Control. Rapid risk assessment: Outbreak of novel coronavirus disease 2019 (covid-19): Increased transmission globally – fifth update. 2020 Mar 2; [accessed 12 Jul 2020] Available from: https://www.ecdc.europa.eu/en/publications-data/rapid-risk-assessment-outbreak-novel-coronavirus-disease-2019-covid-19-increased

6. Parodi SM, Liu VX. From containment to mitigation of COVID-19 in the US. JAMA. 2020 Apr;323(15):1441. Available from: https://doi.org/10.1001/jama.2020.3882

7. Kassem AM. COVID-19: Mitigation or suppression? Arab Journal of Gastroenterology. 2020 Mar;21(1):1–2. Available from: https://doi.org/10.1016/j.ajg.2020.04.010

8. Ferguson N, Laydon D, Nedjati Gilani G, Imai N, Ainslie K, Baguelin M, et al. Report 9: Impact of non-pharmaceutical interventions (npis) to reduce covid19 mortality and healthcare demand. 2020; Available from: http://spiral.imperial.ac.uk/handle/10044/1/77482

9. European Centre for Disease Prevention and Control. Rapid risk assessment: Coronavirus disease 2019 (covid-19) in the eu/eea and the uk– ninth update. 2020 Apr 23; [accessed 12 Jul 2020] Available from: https://www.ecdc.europa.eu/en/publications-data/rapid-risk-assessment-coronavirus-disease-2019-covid-19-pandemic-ninth-update

10. Ministerio de Sanidad. Plan para la transición hacia una nueva normalidad. 2020. [accessed 5 Jun 2020] Available from: https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov-China/documentos/PlanTransicionNuevaNormalidad.pdf

11. United Kingdom Cabinet Office. Our plan to rebuild: The uk government's covid-19 recovery strategy. Command paper number cp 239. 2020. [accessed 5 Jun 2020] Available from:

https://www.gov.uk/government/publications/our-plan-to-rebuild-the-uk-governments-covid-19-recovery-strategy/our-plan-to-rebuild-the-uk-governments-covid-19-recovery-strategy

12. Varela JCA, Mexico G de. ACUERDO por el que se establece una estrategia para la reapertura de las actividades sociales, educativas y económicas, así como un sistema de semáforo por regiones para evaluar semanalmente el riesgo epidemiológico relacionado con la reapertura de actividades en cada entidad federativa, así como se establecen acciones extraordinarias. 2020. [accessed 21 Jun 2020] Available from: https://dof.gob.mx/nota_detalle.php?codigo=5593313&fecha=14/05/2020

13. Gouvernement de la France. 2ème étape du plan de déconfinement. 2020. [accessed 7 Jun 2020] Available from: https://www.gouvernement.fr/sites/default/files/document/document/2020/05/presentation_-_2eme_etape_du_plan_de_deconfinement_-_28.05.2020.pdf

14. СФедеральная служба по надзору в сфере защиты прав потребителей и благополучия человека (Federal Service for Surveillance on Consumer Rights Protection and Human Wellbeing). Профилактика инфекционных болезней методические рекомендации мр 3.1.0178-20. 2020. [accessed 13 Jun 2020] Available from: https://xn--80aesfpebagmfblc0a.xn--p1ai/ai/doc/214/attach/mr_poetapnoe_snyatie_ogranich._08.05.2020.pdf

15. WHO Team. Department of Communications. Annex: Public health criteria to adjust public health and social measures in the context of covid-19. World Health Organization; 2020. [accessed 27 Jun 2020] Available from: https://www.who.int/publications/i/item/public-health-criteria-to-adjust-public-health-and-social-measures-in-the-context-of-covid-19

16. Al-Tawfiq JA, Rodriguez-Morales AJ. Super-spreading events and contribution to transmission of MERS, SARS, and SARS-CoV-2 (COVID-19). Journal of Hospital Infection. 2020 Jun;105(2):111–2. Available from: https://doi.org/10.1016/j.jhin.2020.04.002

17. Bengel C. Coronavirus: How a champions league match contributed to italy's covid-19 outbreak. CBS Sports. 2020 Mar 26; [accessed 1 Jun 2020] Available from: https://www.cbssports.com/soccer/news/coronavirus-how-a-champions-league-match-contributed-to-italys-covid-19-outbreak/

18. Parra E. Sanidad defiende marcha del 8M en madrid porque el aumento de casos de coronavirus "se concretó el lunes por la mañana". Europa Press. 2020 Mar; [accessed 1 Jun 2020] Available from: https://www.europapress.es/epsocial/igualdad/noticia-sanidad-defiende-marcha-8m-madrid-porque-aumento-casos-coronavirus-concreto-lunes-manana-20200311132735.html

19. Gomez MV, Quesada JD. Women's day marches in spain attract mass numbers despite coronavirus fears. El Pais. 2020 Mar; [accessed 1 Jun 2020] Available from: https://english.elpais.com/society/2020-03-09/womens-day-marches-in-spain-attract-mass-numbers-despite-coronavirus-fears.html

20. Salaün T. Special report: Five days of worship that set a virus time bomb in france. Reuters. 2020 Mar 30; [accessed 30 Jun 2020] Available from: https://www.reuters.com/article/us-health-coronavirus-france-church-spec/special-report-five-days-of-worship-that-set-a-virus-time-bomb-in-france-idUSKBN21H0Q2

21. Zdanowicz C. More than 40 spring breakers who ignored public health advice test positive for coronavirus. CNN travel. 2020 Apr 2; [accessed 1 Jun 2020] Available from: https://edition.cnn.com/travel/article/austin-spring-breakers-coronavirus-trnd/index.html

22. Frieden TR, Lee CT. Identifying and interrupting superspreading eventsImplications for control of severe acute respiratory syndrome coronavirus 2. Emerging Infectious Diseases. 2020 Jun;26(6):1059–66. Available from: https://doi.org/10.3201/eid2606.200495

23. Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to estimate time-varying reproduction numbers during epidemics. American Journal of Epidemiology. 2013 Sep;178(9):1505–12. Available from: https://doi.org/10.1093/aje/kwt133

24. Griffin JM, Collins AB, Hunt K, McEvoy D, Casey M, Byrne AW, et al. A rapid review of available evidence on the serial interval and generation time of COVID-19. 2020 May; Available from: https://doi.org/10.1101/2020.05.08.20095075

25. Ganyani T, Kremer C, Chen D, Torneri A, Faes C, Wallinga J, et al. Estimating the generation interval for coronavirus disease (COVID-19) based on symptom onset data, march 2020. Eurosurveillance. 2020 Apr;25(17). Available from: https://doi.org/10.2807/1560-7917.es.2020.25.17.2000257

26. Reich NG, Lessler J, Cummings DAT, Brookmeyer R. Estimating incubation period distributions with coarse data. Statistics in Medicine. 2009 Sep;28(22):2769–84. Available from: https://doi.org/10.1002/sim.3659

27. Nishiura H, Linton NM, Akhmetzhanov AR. Serial interval of novel coronavirus (COVID-19) infections. International Journal of Infectious Diseases. 2020 Apr;93:284–6. Available from: https://doi.org/10.1016/j.ijid.2020.02.060

28. Zhao S, Gao D, Zhuang Z, Chong M, Cai Y, Ran J, et al. Estimating the serial interval of the novel coronavirus disease (COVID-19): A statistical analysis using the public data in hong kong from january 16 to february 15, 2020. 2020 Feb; Available from: https://doi.org/10.1101/2020.02.21.20026559

29. Du Z, Xu X, Wu Y, Wang L, Cowling BJ, Meyers LA. Serial interval of COVID-19 among publicly reported confirmed cases. Emerging Infectious Diseases. 2020 Jun;26(6):1341–3. Available from: https://doi.org/10.3201/eid2606.200357

30. Wallinga J, Lipsitch M. How generation intervals shape the relationship between growth rates and reproductive numbers. Proceedings of the Royal Society B: Biological Sciences. 2006 Nov;274(1609):599–604. Available from: https://doi.org/10.1098/rspb.2006.3754

31. Fraser C. Estimating individual and household reproduction numbers in an emerging epidemic. Galvani A, editor. PLoS ONE. 2007 Aug;2(8):e758. Available from: https://doi.org/10.1371/journal.pone.0000758

32. Wallinga J. Different epidemic curves for severe acute respiratory syndrome reveal similar impacts of control measures. American Journal of Epidemiology. 2004 Sep;160(6):509–16. Available from: https://doi.org/10.1093/aje/kwh255

33. Chang W, Cheng J, Allaire J, Xie Y, McPherson J. Shiny: Web application framework for r. 2020. Available from: https://CRAN.R-project.org/package=shiny

34. Cota W. Monitoring the number of COVID-19 cases and deaths in brazil at municipal and federative units level. SciELOPreprints:362. 2020 May; Available from: https://doi.org/10.1590/scielopreprints.362

35. Stock J, Aspelund K, Droste M, Walker C. Estimates of the undetected rate among the sars-cov-2 infected using testing data from iceland. 2020 Jun;23. doi: 10.1101/2020.04.06.20055582

36. Tuite AR, Ng V, Rees E, Fisman D. Estimation of COVID-19 outbreak size in italy. The Lancet Infectious Diseases. 2020 May;20(5):537. Available from: https://doi.org/10.1016/s1473-3099(20)30227-9

37. Böhning D, Rocchetti I, Maruotti A, Holling H. Estimating the undetected infections in the covid-19 outbreak by harnessing capturerecapture methods. International Journal of Infectious Diseases. 2020 Aug;97:197–201. Available from: https://doi.org/10.1016/j.ijid.2020.06.009

38. Thompson RN, Stockwin JE, Gaalen RD van, Polonsky JA, Kamvar ZN, Demarsh PA, et al. Improved inference of time-varying reproduction numbers during infectious disease outbreaks. Epidemics. 2019 Dec;29:100356. Available from: https://doi.org/10.1016/j.epidem.2019.100356

39. International Monetary Fund. World economic outlook, october 2019. Global manufacturing downturn, rising trade barriers. WorldBank; 2019. [accessed 13 Jul 2020] Available from: https://www.imf.org/en/Publications/WEO/Issues/2019/10/01/world-economic-outlook-october-2019

40. International Comparison Program, World Bank | World Development Indicators database, World Bank | Eurostat-OECD PPP Programme. WorldBank; [accessed 1 Jul 2020] Available from: https://data.worldbank.org/indicator/NY.GDP.PCAP.PP.KD

41. Ministerio de Sanidad. Evolución de enfermedad por el coronavirus (covid-19). 2020. [accessed 15 Jul 2020] Available from: https://datos.gob.es/es/catalogo/e05070101-evolucion-de-enfermedad-por-el-coronavirus-covid-19

42. Ponencia de Alertas y Planes de Preparación y Respuesta and Comisión de Salud Pública del Consejo Interterritorial. Ministerio de Sanidad. Estrategia de diagnóstico, vigilancia y control en la fase de transición de la pandemia de covid-19. Indicadores de seguimiento. 2020. [accessed 5 Jun 2020] Available from: https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov-China/documentos/COVID19_Estrategia_vigilancia_y_control_e_indicadores.pdf

43. OECD. Regional economy - gross domestic product, large regions tl2. OECD; [accessed 25 Jun 2020] Available from: https://stats.oecd.org/Index.aspx?DataSetCode=REGION_ECONOM

44. Departament de Salut. Registre de casos de covid-19 realitzats a catalunya. Segregació per sexe i àrea bàsica de salut (abs). dades obertes catalunya; [accessed 15 Jul 2020] Available from: https://analisi.transparenciacatalunya.cat/Salut/Registre-de-casos-de-COVID-19-realitzats-a-Catalun/xuwf-dxjd

45. Società Italiana di Anestesia Analgesia Rianimazione e Terapia Intensiva, Grupo di Lavoro. Raccomandazioni di etica clinica per l'ammissione a trattamenti intensivi e per la loro sospensione, in condizioni eccezionali di squilibrio tra necessità e risorse disponibili. 2020. [accessed 24 May 2020] Available from: http://www.siaarti.it/SiteAssets/News/COVID19%20-%20documenti%20SIAARTI/SIAARTI%20-%20Covid19%20-%20Raccomandazioni%20di%20etica%20clinica.pdf

46. European Centre for Disease Prevention and Control. Rapid risk assessment: Resurgence of reported cases of covid 19 in the eu/eea, the uk and eu candidate and potential candidate countries. 2020 Jul 2; [accessed 12 Jul 2020] Available from: https://www.ecdc.europa.eu/en/publications-data/rapid-risk-assessment-resurgence-reported-cases-covid-19

47. Disease Prevention EC for, Control. COVID-19 coronavirus data. EU Open Data Portal; 2020. [accessed 15 Jul 2020] Available from: https://data.europa.eu/euodp/es/data/dataset/covid-19-coronavirus-data

48. Pérez R. Temor por el crecimiento de contagios en málaga tras confirmarse tres nuevos brotes de covid en andalucía. ABC. 2020 Jun 23; [accessed 3 Jul 2020] Available from: https://www.abc.es/espana/aragon/abci-agravan-brotes-aragon-66-casos-covid-solo-dias-diez-veces-mas-hace-semana-202006231332_noticia.html

49. El Mundo. Temor por el crecimiento de contagios en málaga tras confirmarse tres nuevos brotes de covid en andalucía. El Mundo. 2020 Jun 26; [accessed 3 Jul 2020] Available from: https://www.elmundo.es/andalucia/2020/06/26/5ef5e0ec21efa0580e8b4617.html

50. Center for Systems Science and Engineering (CSSE) at Johns Hopkins University. COVID-19 data repository. John Hopkins University; 2020. [accessed 15 Jul 2020] Available from: https://github.com/CSSEGISandData/COVID-19

51. Instituto Nacional de Salud, Colombia Government. Casos positivos de covid-19 en colombia. [accessed 15 Jul 2020] Available from: https://www.datos.gov.co/Salud-y-Protecci-n-Social/Casos-positivos-de-COVID-19-en-Colombia/gt2j-8ykr

52. Ministerio de Ciencia, Tecnología, Conocimiento e Innovación. Datos-covid19. [accessed 15 Jul 2020] Available from: https://github.com/MinCiencia/Datos-COVID19

53. You C, Deng Y, Hu W, Sun J, Lin Q, Zhou F, et al. Estimation of the time-varying reproduction number of COVID-19 outbreak in china. International Journal of Hygiene and Environmental Health. 2020 Jul;228:113555. Available from: https://doi.org/10.1016/j.ijheh.2020.113555

54. Herzog S, Bie JD, Abrams S, Wouters I, Ekinci E, Patteet L, et al. Seroprevalence of IgG antibodies against SARS coronavirus 2 in belgium: A prospective cross-sectional study of residual samples. 2020 Jun; Available from: https://doi.org/10.1101/2020.06.08.20125179

55. Pan A, Liu L, Wang C, Guo H, Hao X, Wang Q, et al. Association of public health interventions with the epidemiology of the COVID-19 outbreak in wuhan, china. JAMA. 2020 May;323(19):1915. Available from: https://doi.org/10.1001/jama.2020.6130

56. Liu Y, Gayle AA, Wilder-Smith A, Rocklöv J. The reproductive number of COVID-19 is higher compared to SARS coronavirus. Journal of Travel Medicine. 2020 Feb;27(2). Available from: https://doi.org/10.1093/jtm/taaa021

57. Lee VJ, Chiew CJ, Khong WX. Interrupting transmission of COVID-19: Lessons from containment efforts in singapore. Journal of Travel Medicine. 2020 Mar;27(3). Available from: https://doi.org/10.1093/jtm/taaa039

58. European Centre for Disease Prevention and Control. Projected baselines of covid-19 in the eu/eea and the uk for assessing the impact of de-escalation of measures. 2020 May 26; [accessed 27 Jun 2020] Available from: https://www.ecdc.europa.eu/en/publications-data/projected-baselines-covid-19-eueea-and-uk-assessing-impact-de-escalation-measures

59. Gironés-Bredy CE, Posca-Maina M, Pinto-Plasencia RG, Mahtani-Chugani V, Grupo de Trabajo Hotel Medicalizado Tenerife Sur COVID 19. First public health measures for the containment of covid-19: A hotel in quarantine. Emergencias. 2020 Jun;32(03):194–6. Available from: http://emergencias.portalsemes.org/descargar/primeras-medidas-de-salud-pblica-para-la-contencin-del-covid19-cuarentena-de-un-hotel/

60. Gabinete de Prensa. Ministerio de Sanidad y Consumo. Gobierno de España. El consejo interterritorial del sns acuerda medidas concretas para zonas con transmisión comunitaria significativa de coronavirus. 2020. [accessed 12 Jun 2020] Available from: https://www.mscbs.gob.es/gabinete/notasPrensa.do?id=4806

61. Bergman NK, Fishman R. Correlations of mobility and covid-19 transmission in global data. 2020 May; Available from: https://doi.org/10.1101/2020.05.06.20093039

62. The Swedish Government. Strategy in response to the covid-19 pandemic. Government Offices of Sweden; 2020. [accessed 9 Jul 2020] Available from: https://www.government.se/articles/2020/04/strategy-in-response-to-the-covid-19-pandemic/

63. Torres-Roman JS, Kobiak IC, Valcarcel B, Diaz-Velez C, Vecchia CL. The reproductive number r0 of COVID-19 in peru: An opportunity for effective changes. Travel Medicine and Infectious Disease. 2020 Apr;101689. Available from: https://doi.org/10.1016/j.tmaid.2020.101689

64. Varghese G, John R. COVID-19 in india: Moving from containment to mitigation. Indian Journal of Medical Research. 2020;0(0):0. Available from: https://doi.org/10.4103/ijmr.ijmr_860_20

65. Dirección General de Salud Pública, Calidad e Innovación. Ministerio de Sanidad. Manejo domiciliario del covid-19. 17.03.2020. 2020. [accessed 24 May 2020] Available from: https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov-China/documentos/Manejo_domiciliario_de_COVID-19.pdf

66. Centro Nacional de Epemiología, ISCIII. Consejo Interterritorial. Ministerio de Sanidad. Estudio ene-covid19: Segunda ronda. Estudio nacional de seroprevlaencia de la infección por sars-cov-2 en españa. 2020. [accessed 5 Jun 2020] Available from: https://portalcne.isciii.es/enecovid19/ene_covid19_inf_pre2.pdf

67. Gudbjartsson DF, Helgason A, Jonsson H, Magnusson OT, Melsted P, Norddahl GL, et al. Spread of SARS-CoV-2 in the icelandic population. New England Journal of Medicine. 2020 Jun;382(24):2302–15. Available from: https://doi.org/10.1056/nejmoa2006100

68. Dirección General de Salud Pública, Calidad e Innovación. Ministerio de Sanidad. Actualización nº 102. Enfermedad por el coronavirus (covid-19). 11.05.2020. 2020. [accessed 10 Jun 2020] Available from: https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov-China/documentos/Actualizacion_102_COVID-19.pdf

69. Ministerio de Sanidad. Estrategia de diagnóstico, vigilancia y control en la fase de transición de la pandemia de covid-19. Indicadores de seguimiento. 2020. [accessed 30 Jun 2020] Available from: https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov-China/documentos/Anexo_I_PANEL_DE_INDICADORES.pdf

70. Llaneras K. The problems spain's outdated data methods have caused during a 21st-century pandemic. El País. 2020 Jun 24; Available from: https://english.elpais.com/society/2020-06-24/the-problems-spains-outdated-data-methods-have-caused-during-a-21st-century-pandemic.html

71. Dirección General de Sistemas de Información, Calidad y Prestación Farmacéutica - Consejería de Sanidad. Pruebas realizadas coronavirus. Junta de Castilla y León; [accessed 15 Jul 2020] Available from: https://datosabiertos.jcyl.es/web/jcyl/set/es/salud/pruebas-coronavirus/1284947954927

72. Abbott S, Hellewell J, Thompson RN, Sherratt K, Gibbs HP, Bosse NI, et al. Estimating the time-varying reproduction number of SARS-CoV-2 using national and subnational case counts. Wellcome Open Research. 2020 Jun;5:112. Available from: https://doi.org/10.12688/wellcomeopenres.16006.1

73. Equipo COVID-19. RENAVE. CNE. CNM (ISCIII). Informe sobre la situación de covid-19 en españa. 2020. [accessed 11 Jun 2020] Available from: https://www.isciii.es/QueHacemos/Servicios/VigilanciaSaludPublicaRENAVE/EnfermedadesTransmisibles/Documents/INFORMES/Informes%20COVID-19/Informe%20n%C2%BA%2032.%20Situaci%C3%B3n%20de%20COVID-19%20en%20Espa%C3%B1a%20a%2021%20de%20mayo%20de%202020.pdf

74. John Hopkins Testing Hub. Johns hopkins covid-19 testing insights initiative. Coronavirus Resource Center (John Hopkins University of Medicine); 2020. [accessed 15 Jul 2020] Available from: https://coronavirus.jhu.edu/testing

75. MINSAL, M. CJ. Jmcastagnetto/covid-19-peru-data: Datos actualizados al 2020-06-19. Zenodo; 2020. [accessed 15 Jul 2020] Available from: https://zenodo.org/record/3901865

76. Public Health Agency of Canada. Public health infobase - data on covid-19 in canada. Government of Canada; [accessed 15 Jul 2020] Available from: https://open.canada.ca/data/en/dataset/261c32ab-4cfd-4f81-9dea-7b64065690dc

77. Ministerio de Sanidad. Evolución de enfermedad por el coronavirus (covid-19). 2020. [accessed 21 May 2020] Available from: https://datos.gob.es/es/catalogo/e05070101-evolucion-de-enfermedad-por-el-coronavirus-covid-19

78. Sciensano, Belgium. COVID-19. Sciensano; 2020. [accessed 15 Jul 2020] Available from: https://epistat.wiv-isp.be/covid/

79. Data from The New York Times, based on reports from state and local health agencies. Coronavirus (covid-19) data in the united states. The New York Times; 2020. [accessed 15 Jul 2020] Available from: https://github.com/nytimes/covid-19-data

80. Immanuel Marcus. Berlin: Authorities fight coronavirus outbreak in apartment building. Berlin Inspectator; 2020. [accessed 24 Jun 2020] Available from: https://berlinspectator.com/2020/06/15/berlin-authorities-fight-coronavirus-outbreak-in-apartment-building/

81. Deutsche Welle and dj/rc (Reuters, dpa, AFP). Germany puts two western districts on lockdown. Deutsche Welle; 2020. [accessed 24 Jun 2020] Available from: https://p.dw.com/p/3eEdt

82. J Gehrcke, based on reports from Robert-Koch-Institut, and health agencies. COVID-19 case numbers in germany by state, over time. The New York Times; 2020. [accessed 15 Jul 2020] Available from: https://github.com/jgehrcke/covid-19-germany-gae

83. Sànchez G, Pérez B. Catalunya decreta el confinamiento de lleida y su comarca por los rebrotes. elPeriódico. 2020 Jul 4; [accessed 4 Jul 2020] Available from: https://www.elperiodico.com/es/sociedad/20200704/el-govern-decreta-el-confinamiento-de-la-comarca-del-segria-lleida-8026251