

1 **Moderate to vigorous physical activity and sedentary behavior change in self-**
2 **isolating adults during the COVID-19 pandemic in Brazil: A cross-sectional survey**
3 **exploring correlates**

4

5 Felipe B. Schuch^{1,a}, Rugero A. Bulzing¹, Jacob Meyer², Guillermo F. López-Sánchez³,
6 Igor Grabovac⁴, Peter Willeit^{5,6}, Davy Vancampfort⁷, Cristina M. Caperchione⁸, Kabir
7 P. Sadarangani^{9,10}, André O. Werneck¹¹, Philip B. Ward^{12,13}, Mark Tully^{14*}, Lee
8 Smith^{15*}

9

- 10 1- Federal University of Santa Maria, Santa Maria, Brazil.
11 2- Iowa State University, Ames, IA, United States of America.
12 3- Faculty of Sport Sciences, University of Murcia, Murcia, Spain.
13 4- Department of Social and Preventive Medicine, Centre for Public Health,
14 Medical University of Vienna, Vienna, Austria.
15 5- Department of Neurology, Medical University of Innsbruck, Innsbruck, Austria.
16 6- Department of Public Health and Primary Care, University of Cambridge,
17 Cambridge, UK.
18 7- KU Leuven Department of Rehabilitation Sciences, Leuven, Belgium.
19 8- School of Sport, Exercise and Rehabilitation, Human Performance
20 Research Centre, University of Technology Sydney, Sydney, NSW, Australia.
21 9- Universidad Autónoma de Chile, Chile.
22 10- Escuela de Kinesiología, Facultad de Salud y Odontología, Universidad Diego
23 Portales, Santiago 8370057, Chile.
24 11- Department of Nutrition, School of Public Health, University of São Paulo
25 (USP), São Paulo, Brazil.
26 12- School of Psychiatry, UNSW Sydney, New South Wales, Australia;
27 13- Schizophrenia Research Unit, Ingham Institute for Applied Medical Research,
28 Liverpool, New South Wales, Australia.
29 14- Institute of Mental Health Sciences, School of Health Sciences, Ulster
30 University, Newtownabbey, UK.
31 15- The Cambridge Centre for Sport and Exercise Sciences, Anglia Ruskin
32 University, Cambridge, CB1 1PT, UK.

33

34

35 *Joint final author

36

37

38

39

40 a=Corresponding author:

41 Felipe Barreto Schuch (felipe.schuch@ufsm.br)

42 Federal University of Santa Maria, Santa Maria, RS, Brazil.

43 Av. Roraima, 1000, Prédio 51 (CEFD), sala 1025.

44 CEP:07105-900 Phone: +55 55 3220 7788

45 **Abstract:**

46 **Background:** Self-distancing measures imposed major changes in daily life. This study
47 aimed to (i) evaluate the changes (pre-versus during pandemic) in time spent in
48 moderate to vigorous physical activity (MVPA) and sedentary behavior (SB) in self-
49 isolating Brazilians during the COVID-19 pandemic, and (ii) to explore correlates of
50 MVPA and SB.

51 **Methods:** A cross-sectional self-report online survey, evaluating the time spent in
52 MVPA and SB pre and during the COVID-19 pandemic. Sociodemographic,
53 behavioral, clinical, variables, and time in self-isolation were also obtained. Changes in
54 MVPA and SB and their correlates were explored using generalized estimating
55 equations (GEE).

56 **Results:** A total of 877 participants (72.7% women, 53.7% young adults [18-34 years])
57 were included. Overall, participants reported a 59.7% reduction (95%CI:35.6 to 82.2) in
58 time spent on MVPA during the pandemic. Time spent in SB increased 42.0%
59 (95%CI:31.7 to 52.5). Greater reductions in MVPA and/or increases in SB were seen in
60 younger adults, those not married, those employed and those with a self-reported
61 previous diagnosis of a mental disorder.

62 **Conclusions:** People in self-isolation significantly reduced MVPA levels and increased
63 SB. Public health strategies should be implemented during epidemic times to mitigate
64 the impact of self-isolation on MVPA and SB, particularly in vulnerable groups.

65

66

67

68

69

70

71

72

73

74

75

76

77 **Introduction**

78

79 The coronavirus disease 2019 (COVID-19) pandemic, caused by the spread of
80 the severe acute respiratory coronavirus 2 (SARS-CoV-2), has infected more than 13
81 million people in more than 200 countries around the world resulting in nearly 570
82 thousand deaths on the 14th of July 2020 according to the World Health Organization
83 (WHO)¹. As a response to reduce the virus spread, the WHO recommended national
84 governments to adopt non-pharmacological strategies based on social and physical
85 distancing, such as lockdown, quarantine, and self-isolation .

86 In Brazil, the epidemiological report number 05 of the Ministry of Health has
87 recommended the adoption of social distancing measures, including self-isolation in
88 areas with community transmission². When self-isolating, people were advised to stay at
89 home, and only go out in public for essential activities, such as going to the
90 supermarket, pharmacy, or to use essential services, such as medical assistance. All
91 other non-essential services, including gyms, parks, stadiums, and other places where
92 people exercised were closed.

93 Self-isolation measures impose a drastic and sudden disruption of daily life
94 routine, resulting in limited physical and social mobility, and fewer opportunities to be
95 active³. Moreover, the emotional burden^{4, 5} of the pandemic likely resulted in additional
96 barriers to remaining focused and motivated to be and/or stay physically active,
97 potentially reducing the time spent in physical activity (PA), defined as any bodily
98 movement produced by skeletal muscles that result in energy expenditure⁶. The negative
99 impact of the pandemic and self-isolation measures on PA levels, both on light PA and
100 moderate to vigorous PA (MVPA), was noted in many countries including Australia^{7, 8},

101 Canada⁹, Croatia¹⁰, France¹¹, Italy¹², Spain¹³, the UK^{14, 15}, and the USA¹⁶⁻¹⁸. However,
102 some moderating factors on PA changes in this period were identified. For example,
103 age^{12, 14}, sex¹³ and the presence of chronic physical diseases or mental disorders
104 moderated the pandemic impact on PA levels¹⁴. Further, increases in time spent in
105 sedentary behavior (SB), defined as any waking behavior characterized by an energy
106 expenditure ≤ 1.5 metabolic equivalents (METs), while in a sitting, reclining or lying
107 posture¹⁹, were noted during the pandemic in the US^{17, 18}, France and Sweden²⁰, and
108 Spain¹³. In Spain, sex moderated the increase in time spent in SB¹³.

109 To the best of our knowledge, no study has evaluated how the COVID-19
110 pandemic changed PA and SB during self-isolating Brazilians. The present study aimed
111 (i) to examine the changes (pre-versus during COVID-19 pandemic) on PA and SB in
112 self-isolating Brazilians, and (ii) to evaluate whether sociodemographic (sex, age,
113 ethnicity, marital status, employment, monthly household income), behavioral
114 (smoking, current alcohol consumption), clinical (presence of chronic physical diseases
115 or mental disorders), and contextual factors (i.e., number of days of self-isolation)
116 moderated these changes.

117

118 **Methods**

119 This paper presents pre-planned interim analysis of data from a cross-sectional
120 study. Data collection was performed through an online survey (www.qualtrics.com).
121 The study was launched on 11 April 2020 and data collection continued until 05 May.
122 The study was approved by the Federal University of Santa Maria Research Ethics
123 Committee and by the National Commission of Ethics in Research [CONEP]
124 (30244620.1.0000.5346).

125 Participants were recruited through social media (Facebook, Instagram, Twitter),
126 and by distributing an invitation to participate through existing researcher networks.
127 The inclusion criteria were: (1) Brazilians adults (18-65 years), (2) currently residing in
128 Brazil, and (3) in self-isolation due to the COVID-19 pandemic. Self-isolation was
129 defined as staying-at-home and only leaving for essential activities such as food
130 shopping, visiting the pharmacist or other health professionals. Participants who self-
131 reported the presence of COVID-19 symptoms, assessed through a list of symptoms
132 (fever, cough, dry mouth, coriza, sore throat), were removed from this analysis (n=65).

133

134 Moderate to vigorous physical activity (MVPA) and sedentary behavior (SB)
135 assessment

136 Participants were asked to recall the amount of time in vigorous and moderate
137 physical activity, and sedentary behavior they undertook on an average day, separately
138 both pre- and since self-isolation¹⁵. Participants were asked: (1) “How much time on an
139 average day have you spent in vigorous activity before/since social distancing?”; (2)
140 “How much time on an average day have you spent in moderate activity since/before
141 social distancing?” and (3) “How much time on an average day have you spent sitting
142 since/before social distancing?” Responses were given in hours and minutes. MVPA
143 and SB were analyzed as continuous variables (minutes per day). We also categorized
144 PA levels (≥ 30 minutes/day or < 30 minutes/day of MVPA), which is in accordance with
145 the WHO recommendations. Next, four categories were derived to identify patterns of
146 change: (1) persistent inactive (< 30 minutes/day of MVPA pre and during the
147 pandemic), (2) decreased PA (≥ 30 minutes/day of MVPA pre and < 30 minutes/day of
148 MVPA during the pandemic), (3) increased PA (< 30 minutes/day of MVPA pre and ≥ 30

149 minutes/day of MVPA during the pandemic) and (4) persistent active (≥ 30 minutes/day
150 of MVPA pre and during the pandemic).

151 Covariates

152 Demographic data were collected, including sex (men or women), age (in 10-
153 year age bands), ethnicity (Caucasian, Black, Asian, mixed, others), marital status
154 (single, divorced, widowed or married), employment (employed, student, military,
155 unemployed), monthly household income \leq R\$1254, R\$1255-R\$2004, R\$2005-R\$8640,
156 R\$8641-R\$11261 \geq R\$11262). Health behaviors data included current smoking (y/n) or
157 alcohol consumption (y/n). Clinical data included self-reported previous diagnosis of
158 physical diseases or mental disorders, such as: obesity, hypertension, myocardial
159 infarction, angina pectoris, and other coronary diseases, other cardiac diseases, varicose
160 veins of lower extremities, osteoarthritis, chronic neck pain, chronic low back pain,
161 chronic allergy (excluding allergic asthma), chronic bronchitis, emphysema or chronic
162 obstructive pulmonary disease, type 1 diabetes, type 2 diabetes, diabetic retinopathy,
163 cataract, peptic ulcer disease, urinary incontinence or urine control problems,
164 hypercholesterolemia, chronic skin disease, chronic constipation, liver cirrhosis and
165 other hepatic disorders, stroke, chronic migraine/others, depression, anxiety disorders,
166 bipolar disorder and schizophrenia/others. Number of days (extension) in self-isolation
167 was registered with a single question.

168 Statistical analyses

169 Data were analyzed using mean and standard deviation (SD) or 95% confidence
170 interval (95%CI) for continuous data and the raw numbers and % for categorical
171 variables. Normality was checked with the Kolmogorov-Smirnov test. Due to the non-
172 normal distribution, the mean changes (pre-versus during the pandemic) of MVPA

173 levels and SB were evaluated using two generalized estimating equations models
174 (GEE), one with changes in MVPA and one with change in SB as the dependent
175 variable. The models were run testing the time effects (pre/during) and the interactions
176 between time and the factors included in the model. The factors included in the models
177 were (sex [male versus female], age [young adults {18-34 years} versus middle-age
178 adults {35-54 years} versus older adults {55-65 years}], ethnicity [Caucasian versus
179 Asian/Black/mixed/others], marital status [married versus single/divorced/widowed],
180 employment [employed/students/military versus unemployed/retired], monthly
181 household income [$<R\$2,005$ versus $R\$2,005$ - $R\$8,640$ versus $R\$8,641$ - $R\$11,261$
182 versus $>R\$11,261$], current smoking [yes versus no], alcohol consumption [yes versus
183 no], self-reported previous diagnosis of any chronic diseases [yes versus no] or any
184 mental disorders [yes versus no]). When the interaction between time and any factor
185 was significant, the Bonferroni test was applied. The results of the models are presented
186 using estimated marginal means and 95%CI. We also calculated the delta% change (pre
187 to during), together with 95% CI as the effect size measure. The associations between
188 the time (in days) in self-isolation and the change in MVPA and SB were tested using
189 linear regression models. Days in self-isolation were collected as a continuous variable,
190 and linear regression models were used to test the association of time in self-isolation
191 with changes in MVPA and SB. The level of statistical significance was set at p-value $<$
192 0.05. All analyses were performed using SPSS (v. 21).

193

194 **Results**

195 A total of 877 participants participated. The sample was predominantly
196 comprised of women (n=635, 72.7%), young adults ranging from 18-34 years (n=471,

197 53.7%), Caucasians (n=669, 76.3%), singles (n=442, 50.9%), currently
198 employed/students/military (n=723, 92.6%), with a monthly income ranging from
199 R\$2005 to R\$8640 (n=364, 41.5%), non-smokers (n=836, 95.3%), currently consuming
200 alcohol (n=605, 69.1%), with a self-reported previous diagnosis of a physical chronic
201 disease (n=824, 94%) and without a self-reported previous diagnosis of a mental
202 disorder (n=523, 59.6%). Participants were on average of 27.13 (6.57) days in self-
203 isolation. The full details of the sample can be seen at Table 1.

204 Mean changes in MVPA and SB (pre versus during pandemic)

205 A total of 432 (49.3%) of participants persisted to be active during the pandemic,
206 32 (3.6%) increased MVPA levels during the pandemic, 306 (34.9%) reduced MVPA
207 levels during the pandemic, and 107 (12.2%) persisted to be inactive. We found an
208 overall reduction of 64.28 (95% CI: 36.06 to 83.33) minutes per day on time spent in
209 MVPA, corresponding to a reduction of 59.7% (95%CI: 35.64 to 82.21, $p<0.001$). The
210 average time spent in SB increased 42.0% (95% CI: 31.74 to 52.50, $p<0.001$) during the
211 pandemic, corresponding to additional 152.3 (95% CI: 111.9 to 192.7) minutes per day
212 in SB. The mean time spent in MVPA and SB at baseline and during the pandemic can
213 be found at figure 1.

214 Correlates of MVPA change

215 Significant interactions in MVPA change were found for age ($p=0.013$), marital
216 status ($p=0.006$) and employment ($p=0.008$). Bonferroni post hoc test found that young
217 adults (mean change=-71.37, 95%CI: -99.76.51 to -42.98) and middle age adults (mean
218 change=-66.76, 95%CI: -94.50 to -39.01) had greater decreases in time spent in MVPA
219 when compared to older adults (mean change=-54.70, 95%CI: -86.25 to -23.16). Also,
220 those not married (single/divorced/widowed) had greater reductions (mean change=-

221 75.50, 95%CI:-102.00 to -49.00) when compared to those married (mean change=-
222 53.05, 95%CI: -79.36 to -26.75), and those with an occupation
223 (employed/student/military) had greater reductions (mean change=-78.69, 95%CI: -
224 105.21 to -52.16) when compared to those without occupation (unemployed/retired.
225 Mean change=-49.87, 95%CI: -78.83 to -20.90). The detailed results of the MVPA
226 model with mean changes can be seen in Table 2. Those not married (mean
227 difference=22.08, 95%CI: 6.79 to 37.36, p=0.005), and with no occupation (mean
228 difference=38.55, 95%CI: 14.08 to 63.03, p=0.002) had greater MVPA levels at
229 baseline (pre and during pandemic means are shown in supplementary material 1). The
230 number of days in self-isolation was not associated with changes in MVPA
231 (unstandardized beta coefficient=0.234, 95%CI: -0.816 to 1.284, p=0.662, R²=0.00).

232 Correlates of SB changes

233 The interactions found for the SB model were age (p<0.001), marital status
234 (p=0.024), employment (p=0.03), and self-reported previous diagnosis of mental
235 disorders (p=0.003). Young adults had greater increases (mean change=190.48,
236 95%CI: 149.65 to 231.30) in time spent in SB when compared to middle age (mean
237 change=143.35 95%CI: 99.48 to 187.21) or older adults (mean change=136.71, 95%CI:
238 77.88 to 195.54). Also, greater increases in time spent in SB were found in those not
239 married (mean change=176.15, 95%CI: 133.74 to 218.56) compared to those married
240 (mean change=137.96, 95%CI: 96.85 to 178.24), in those with an occupation (mean
241 change=179.85, 95%CI: 142.15 to 217.55) compared to those with no occupation (mean
242 change=133.84, 95%CI: 79.26 to 188.42), and in those with a self-reported previous
243 diagnosis of mental disorders (mean change=173.16, 95%CI: 129.97 to 216.34)
244 compared to those with no history of mental disorders (mean change=140.53, 95%CI:
245 101.07 to 180.00). The detailed results of the SB model with interactions can be seen in

246 Table 2. Younger adults spent more time in SB than middle-age adults (mean
247 difference=39.99, 95%CI: 5.27 to 74.71, $p=0.017$), but not more than older adults (mean
248 difference=43.28, 95%CI: -13.15 to 99.75, $p=0.20$). Those with a self-reported previous
249 diagnosis of a mental disorder spent more time in SB at baseline (mean
250 difference=24.09, 95%CI: 0.41 to 48.66, $p=0.046$) than those without (pre and during
251 pandemic means are shown in supplementary material 2). The number of days in self-
252 isolation was not associated with changes in SB (unstandardized beta coefficient=0.306,
253 95%CI: -1.732 to 2.345, $p=0.77$, $R^2=0.00$).

254 **Discussion**

255 The present study is, to the best of our knowledge, the first study demonstrating
256 the impact of the self-isolation during the COVID-19 pandemic on Brazilians for self-
257 reported MVPA and SB. Approximately 35.0% of the sample became insufficiently
258 active during the self-isolation period. Only 3.6% became active with self-isolation. On
259 average, there was a reduction of about 1 hour/day of time spent in MVPA, which
260 corresponds to a reduction of 60.0% of their MVPA pre-pandemic levels. Participants
261 reported spending 2.5 hours/day more in SB during the pandemic than before the
262 pandemic, corresponding to an increase of 40.0%.

263 The reduction in MVPA levels in Brazilians is consistent with the findings of
264 previous studies in other countries. For example, we found that roughly 35.0% of
265 respondents became inactive during the self-isolation period as did about 50.0% in
266 similar studies collected in France¹¹, the USA¹⁶ and Australia⁸. Our results included
267 reductions of 60.0% of the time spent in MVPA in Brazil, which is comparable to the
268 reductions found in the USA, where there was a decrease of 47.0% on time spent in
269 moderate PA¹⁶. Additionally, we observed an increase of about 2.5 hours/day on time

270 spent in SB, which is consistent with other studies that have found an increase of about
271 2 to 3 hours/day of SB in multiple countries^{13, 21}. These findings highlight the urgent
272 need for public health strategies to mitigate the impact of self-isolation on MVPA and
273 SB.

274 Greater reductions in MVPA and increases in SB were found among younger
275 adults, which is line with the findings from Italy and the UK^{12, 14}. It is possible that this
276 age group had fewer resources and greater difficulty coping with emotional responses to
277 this situation²². Also, those not married and currently working had higher MVPA levels
278 at baseline, but decreased their MVPA to similar levels to those not married and with no
279 work during the pandemic. Of note, those currently employed might have reduced their
280 commuting-related PA and have likely increased their SB time due to online meetings
281 and activities. Lastly, those with a self-reported previous pandemic diagnosis of a
282 mental disorder spent more time in SB and reported the greatest increases in time spent
283 in SB during the pandemic. This finding is in accordance with a study in the UK¹⁴ that
284 found a greater reduction in MVPA in people with depression. This finding is also
285 consistent with previous studies showing that people with mental disorders have higher
286 SB levels than people without mental disorders²³⁻²⁵, and suggests that self-isolation
287 during the pandemic might be specifically detrimental to people with a previous
288 diagnosis of a mental disorder.

289 There is ample evidence to justify making PA promotion a global public health
290 priority during the coronavirus pandemic³. The COVID-19 pandemic appears to have
291 impacted mental health globally, increasing rates of depression and anxiety symptoms
292 and disorders⁵. On the other hand, those more active are less likely to develop mental
293 disorders²⁶⁻²⁸ and reported fewer symptoms in comparison to those with mental
294 disorders²⁹⁻³². During the pandemic, cross-sectional evidence suggests that those with

295 higher PA or lower SB levels are less likely to present depressive symptoms³³.
296 Promoting MVPA and reducing SB during the pandemic is also essential for physical
297 health. Higher mortality due to COVID-19 is seen in those with clinical comorbidities
298 such as hypertension, diabetes, and coronary heart disease³⁴. Increasing time spent in
299 MVPA and reducing time spent in SB seems to reduce the risk of development of
300 multiple chronic diseases, including those associated with a higher risk of COVID-19
301 mortality³⁵. Lastly, research is beginning to show that physical inactivity may be a risk
302 factor for hospitalization due to COVID-19³⁶, further underlining the potential
303 importance of promoting PA during the pandemic.

304 The strengths of the manuscript are the large sample size of self-isolating
305 Brazilians and the possibility to explore a variety of moderators. The present study has
306 some limitations. First, MVPA and SB were assessed using self-reported questionnaires.
307 Self-reported questionnaires are commonly associated with overestimations of MVPA³⁷
308 and underestimation of SB³⁸. Second, pre-pandemic MVPA and SB were assessed
309 retrospectively, and both can be susceptible to memory bias. Third, the
310 representativeness of the sample is limited. However, participants were drawn from 24
311 of the 27 federative units of Brazil, with more than half of participants from the Rio
312 Grande do Sul, 14% from the Rio de Janeiro, and about 6% from Ceará. Also, some
313 groups such as adults aging 55-64, Asian and Black people, and those with a household
314 income lower than <R\$1,254 are poorly represented. Fourth, we could not explore
315 changes in light PA, such as walking. Also, we could not explore the changes in time
316 spent on MVPA across the different PA domains (work/occupational, leisure,
317 transportation, household). It is possible, for example, that some participants reduced
318 the time spent in leisure, transportation, or work/occupational activities, but increased
319 the time spent in household activities. This is important since we know that some

320 mental health benefits are more likely to be associated with time spent in leisure
321 activities³⁹.

322

323 **Conclusion**

324 Self-isolation during the pandemic significantly reduced time spent in MVPA
325 and increased time spent in SB in Brazilian adults, particularly in younger adults, those
326 who were single, and those who were employed. These findings highlight the urgent
327 need of the adoption of public health strategies to address the impact of self-isolation
328 during the COVID-19 pandemic on MVPA and SB.

329

330

331 **Acknowledgements**

332 This study was part financed in part by the Coordenação de Aperfeiçoamento de Pessoal de
333 Nível Superior - Brasil (CAPES) - Finance Code 001. André Werneck is supported by a São
334 Paulo Research Foundation PhD scholarship (FAPESP process: 2019/24124-7). Joseph
335 Firth is supported by a University of Manchester Presidential Fellowship (P123958) and a UK
336 Research and Innovation Future Leaders Fellowship (MR/T021780/1). Brendon Stubbs is
337 supported by a Clinical Lectureship (ICA-CL-2017-03-001) jointly funded by Health Education
338 England (HEE) and the National Institute for Health Research (NIHR). Brendon Stubbs is part
339 funded by the NIHR Biomedical Research Centre at South London and Maudsley NHS
340 Foundation Trust. The views expressed are those of the author(s) and not necessarily those of
341 the (partner organisation), the NHS, the NIHR or the Department of Health and Social Care.
342 Mark Tully is partly supported by funding as Director of the Northern Ireland Public Health

343 Research Network by the Research and Development Division of the Public Health Agency
344 (Northern Ireland).

345

346

347

348 **References**

- 349 1. World Health Organization. Coronavirus disease 2019 (COVID-19): situation report,
350 179. 2020.
- 351 2. Ministério da Saúde. Boletim epidemiológico 05 - 20/03/2020. In: Saúde Md, ed.
352 http://maismedicos.gov.br/images/PDF/2020_03_13_Boletim-Epidemiologico-05.pdf;
353 2020.
- 354 3. Sallis J, Adlakha D, Oyeyemi A, Salvo D. An international physical activity and public
355 health research agenda to inform COVID-19 policies and practices. *Journal of Sport and*
356 *Health Science*. 05/01 2020.
- 357 4. Ornell F, Schuch JB, Sordi AO, Kessler FHP. Pandemic fear and COVID-19: mental
358 health burden and strategies. *Brazilian Journal of Psychiatry*. 2020.
- 359 5. Brooks SK, Webster RK, Smith LE, et al. The psychological impact of quarantine and
360 how to reduce it: rapid review of the evidence. *The Lancet*. 2020.
- 361 6. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical
362 fitness: definitions and distinctions for health-related research. *Public Health Rep*. Mar-
363 Apr 1985;100(2):126-131.
- 364 7. Gallo LA, Gallo TF, Young SL, Moritz KM, Akison LK. The impact of isolation measures
365 due to COVID-19 on energy intake and physical activity levels in Australian university
366 students. *medRxiv*. 2020:2020.2005.2010.20076414.
- 367 8. Stanton R, To Q, Khalesi S, et al. Depression, Anxiety and Stress during COVID-19:
368 Associations with Changes in Physical Activity, Sleep, Tobacco and Alcohol Use in
369 Australian Adults. *Int J Environ Res Public Health*. 06/07 2020;17:4065.
- 370 9. Lesser I, Nienhuis C. The Impact of COVID-19 on Physical Activity Behavior and Well-
371 Being of Canadians. *Int J Environ Res Public Health*. 05/31 2020;17:3899.
- 372 10. Sekulic D, Blazevic M, Gilic B, Kvesic I, Zenic N. Prospective Analysis of Levels and
373 Correlates of Physical Activity During COVID-19 Pandemic and Imposed Rules of Social
374 Distancing; Gender Specific Study Among Adolescents from Southern Croatia.
375 *Sustainability*. 2020;12(10):4072.
- 376 11. Deschasaux-Tanguy M, Druesne-Pecollo N, Esseddik Y, et al. Diet and physical activity
377 during the COVID-19 lockdown period (March-May 2020): results from the French
378 NutriNet-Sante cohort study. *medRxiv*. 2020:2020.2006.2004.20121855.
- 379 12. Giustino V, Parroco A, Gennaro A, Musumeci G, Palma A, Battaglia G. Physical Activity
380 Levels and Related Energy Expenditure during COVID-19 Quarantine among the Sicilian
381 Active Population: A Cross-Sectional Online Survey Study. *Sustainability*. 05/26
382 2020;12:4356.
- 383 13. Castañeda A, Arbillaga-Etxarri A, Gutiérrez-Santamaria B, Coca A. *Impact of COVID-19*
384 *confinement on the time and intensity of physical activity in the Spanish*
385 *population*2020.

- 386 14. Rogers N, Waterlow N, Brindle H, et al. *Behavioural change towards reduced intensity*
387 *physical activity is disproportionately prevalent among adults with serious health issues*
388 *or self-perception of high risk during the UK COVID-19 lockdown*2020.
- 389 15. Lee Smith JM, Guillermo Felipe López-Sánchez, Felipe Schuch, Igor Grabovac, Nicola
390 Veronese, Ahmad Abufaraj, Cristina Coperchiona, Mark Tully. Prevalence and
391 Correlates of Physical activity in a sample of UK adults observing social distancing
392 during the COVID-19 pandemic. *BMJ Open Sport & Exercise Medicine*. 2020;in press.
- 393 16. Dunton G, Wang S, Do B, Courtney J. Early Effects of the COVID-19 Pandemic on
394 Physical Activity in US Adults. 2020.
- 395 17. Dunton G, Do B, Wang S. Early Effects of the COVID-19 Pandemic on Physical Activity
396 and Sedentary Behavior in US Children. 2020.
- 397 18. Meyer J, McDowell C, Lansing J, et al. *Changes in physical activity and sedentary*
398 *behaviour due to the COVID-19 outbreak and associations with mental health in 3,052*
399 *US adults*2020.
- 400 19. Tremblay MS, Aubert S, Barnes JD, et al. Sedentary behavior research network (SBRN)–
401 terminology consensus project process and outcome. *International Journal of*
402 *Behavioral Nutrition and Physical Activity*. 2017;14(1):75.
- 403 20. Cheval B, Sivaramakrishnan H, Silvio M, et al. *Relationships between changes in self-*
404 *reported physical activity, sedentary behaviours and health during the coronavirus*
405 *(COVID-19) pandemic in France and Switzerland*2020.
- 406 21. Ammar A, Brach M, Trabelsi K, et al. Effects of COVID-19 Home Confinement on Eating
407 Behaviour and Physical Activity: Results of the ECLB-COVID19 International Online
408 Survey. *Nutrients*. 05/28 2020;12.
- 409 22. Scott SB, Sliwinski MJ, Blanchard-Fields F. Age differences in emotional responses to
410 daily stress: the role of timing, severity, and global perceived stress. *Psychol Aging*. Dec
411 2013;28(4):1076-1087.
- 412 23. Schuch F, Vancampfort D, Firth J, et al. Physical activity and sedentary behavior in
413 people with major depressive disorder: a systematic review and meta-analysis. *Journal*
414 *of affective disorders*. 2017;210:139-150.
- 415 24. Vancampfort D, Firth J, Schuch F, et al. Physical activity and sedentary behavior in
416 people with bipolar disorder: A systematic review and meta-analysis. *J Affect Disord*.
417 Sep 1 2016;201:145-152.
- 418 25. Vancampfort D, Firth J, Schuch FB, et al. Sedentary behavior and physical activity levels
419 in people with schizophrenia, bipolar disorder and major depressive disorder: a global
420 systematic review and meta-analysis. *World Psychiatry*. Oct 2017;16(3):308-315.
- 421 26. Brokmeier LL, Firth J, Vancampfort D, et al. Does physical activity reduce the risk of
422 psychosis? A systematic review and meta-analysis of prospective studies. *Psychiatry*
423 *Res*. Nov 14 2019;112675.
- 424 27. Schuch FB, Vancampfort D, Firth J, et al. Physical Activity and Incident Depression: A
425 Meta-Analysis of Prospective Cohort Studies. *Am J Psychiatry*. Jul 1 2018;175(7):631-
426 648.
- 427 28. Schuch FB, Stubbs B, Meyer J, et al. Physical activity protects from incident anxiety: A
428 meta-analysis of prospective cohort studies. *Depression and anxiety*. Sep
429 2019;36(9):846-858.
- 430 29. Ashdown-Franks G, Firth J, Carney R, et al. Exercise as medicine for mental and
431 substance use disorders: A meta-review of the benefits for neuropsychiatric and
432 cognitive outcomes. *Sports Medicine*. 2020;50(1):151-170.
- 433 30. Schuch FB, Vancampfort D, Richards J, Rosenbaum S, Ward PB, Stubbs B. Exercise as a
434 treatment for depression: a meta-analysis adjusting for publication bias. *Journal of*
435 *psychiatric research*. 2016;77:42-51.
- 436 31. Stubbs B, Vancampfort D, Hallgren M, et al. EPA guidance on physical activity as a
437 treatment for severe mental illness: a meta-review of the evidence and Position

- 438 Statement from the European Psychiatric Association (EPA), supported by the
439 International Organization of Physical Therapists in Mental Health (IOPTMH). *European*
440 *psychiatry : the journal of the Association of European Psychiatrists*. Oct 2018;54:124-
441 144.
- 442 **32.** Stubbs B, Vancampfort D, Rosenbaum S, et al. An examination of the anxiolytic effects
443 of exercise for people with anxiety and stress-related disorders: A meta-analysis.
444 *Psychiatry Res*. Mar 2017;249:102-108.
- 445 **33.** Schuch FB, Bulzing RA, Meyer J, Vancampfort D, Firth J. Associations of moderate to
446 vigorous physical activity and sedentary behavior with depressive and anxiety
447 symptoms in self-isolating people during the COVID-19 pandemic: A cross-sectional
448 survey in Brazil.
- 449 **34.** Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients
450 with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet*.
451 2020/03/28/ 2020;395(10229):1054-1062.
- 452 **35.** Pedersen BK, Saltin B. Exercise as medicine—evidence for prescribing exercise as
453 therapy in 26 different chronic diseases. *Scandinavian journal of medicine & science in*
454 *sports*. 2015;25:1-72.
- 455 **36.** Hamer M, Kivimäki M, Gale CR, Batty GD. Lifestyle risk factors, inflammatory
456 mechanisms, and COVID-19 hospitalization: A community-based cohort study of
457 387,109 adults in UK. *Brain Behav Immun*. Jul 2020;87:184-187.
- 458 **37.** Lee PH, Macfarlane DJ, Lam TH, Stewart SM. Validity of the international physical
459 activity questionnaire short form (IPAQ-SF): A systematic review. *International Journal*
460 *of Behavioral Nutrition and Physical Activity*. 2011/10/21 2011;8(1):115.
- 461 **38.** Prince S, Cardilli L, Reed J, et al. A comparison of self-reported and device measured
462 sedentary behaviour in adults: a systematic review and meta-analysis. *International*
463 *Journal of Behavioral Nutrition and Physical Activity*. 03/04 2020;17:1-17.
- 464 **39.** Teychenne M, White RL, Richards J, Schuch FB, Rosenbaum S, Bennie JA. Do we need
465 physical activity guidelines for mental health: What does the evidence tell us? *Mental*
466 *Health and Physical Activity*. 2020;18:100315.

467

468

469

470

471

472

473

474

475

476

477

478

479

480
481
482
483
484
485
486
487

Table 1. Sample characteristics

	Category/Mean (Standard deviation)	Overall N=877* (%)
Sex	Male	238 (27.3)
	Female	635 (72.7)
Age	18-24 years	131 (14.9)
	25-34 years	340 (38.8)
	35-44 years	220 (25.1)
	45-54 years	108 (12.3)
	55-64 years	78 (8.9)
Ethnicity	Asian	3 (0.3)
	Black	24 (2.7)
	Mixed	163 (18.6)
	Caucasian	669 (76.3)
	Others	16 (1.8)
Marital Status	Married	363 (41.4)
	Widowed	4 (0.5)
	Divorced	60 (6.9)
	Single	442 (50.9)
Employment	Employed	558 (63.6)
	Unemployed	43 (1.6)
	Student	235 (26.8)
	Military	19 (2.2)
	Retired	22 (2.5)
Monthly household income	<R\$1254	30 (3.4)
	R\$1255-R\$2004	94 (10.7)
	R\$2005-R\$8640	364 (41.5)
	R\$8641-R\$11261	139 (15.8)
	>R\$11262	250 (28.5)
Current smoking	No	836 (95.3)
	Yes	41 (4.7)
Current alcohol consumption	No	271 (30.9)
	Yes	605 (69.1)
Self-reported previous diagnoses of physical disease	No	53 (6)
	Yes	824 (94)
Self-reported previous diagnoses of mental disorder	No	523 (59.60)
	Yes	354 (40.4)
Days in self isolation	Mean (standard deviation)	27.07 (6.71)

488

489 * Total sample with available data. Number of cases can be different for each variable due to missing
490 cases (minimum=869)

Table 2. Moderate to vigorous physical activity change (pre-post pandemic) in self-isolated adults during the SARS-CoV-2 pandemic in 2020 in Brazil

Characteristics	Category	MVPA change (pre versus during)						Interaction p value
		Mean change in minutes	95% CI		Delta%	95% CI		
Sex	Male	-65.86	-92.26	-39.52	-61.24%	-85.79%	-36.75%	0.778
	Female	-62.71	-89.37	-36.05	-58.18%	-82.91%	-33.44%	
Age	Young adults (18-35 years)	-71.37a	-99.76	-42.98	-72.68%	-101.51%	-43.77%	0.013
	Middle age adults (36-55 years)	-66.76a	-94.50	-39.01	-64.53%	-91.34%	-37.07%	
	Older adults (55-64 years)	-54.70b	-86.25	-23.16	-45.08%	-71.08%	-19.08%	
Ethnicity	Black/Asian/Mixed/Others	-66.00	-95.52	-36.48	-62.67%	-90.70%	-34.64%	0.216
	White	-62.55	-86.72	-38.38	-56.59%	-78.87%	-34.98%	
Marital status	Single/divorced/widowed	-75.50a	-102.00	-49.00	-63.60%	-85.93%	-41.28%	0.006
	Married	-53.05b	-79.36	-26.75	-54.91%	-82.14%	-27.68%	
Employment	Employed/students/military	-78.69a	-105.21	-52.16	-61.99%	-82.88%	-41.09%	0.008
	Unemployed/retired	-49.87b	-78.83	-20.90	-56.42%	-89.19%	-23.64%	
Monthly household income	<R\$2005	-82.58	-114.49	-50.66	-63.52%	-88.15%	-39.00%	0.647
	R\$2005-R\$8640	-57.53	-82.31	-32.75	-56.57%	-80.94%	-32.20%	
	R\$8641-R\$11261	-56.72	-89.26	-24.19	-58.06%	-91.38%	-24.76%	
	>R\$11261	-60.28	-88.92	-31.63	-59.44%	-87.69%	-31.19%	
Current smoking	Yes	-70.76	-112.95	-28.57	-67.02%	-106.9%	-27.06%	0.090
	No	-57.80	-77.29	-38.30	-52.77%	-70.43%	-34.90%	
Current alcohol consumption	Yes	-58.39	-84.85	-31.93	-57.09%	-82.97%	-31.22%	0.378
	No	-70.16	-97.22	-43.10	-62.06%	-85.29%	-38.12%	
Self-reported previous diagnosis of mental disorders	Yes	-67.46	-95.88	-39.05	-63.37%	-90.07%	-36.68%	0.112
	No	-61.09	-85.60	-36.58	-56.11%	-78.63%	-33.60%	
Self-reported previous diagnosis of physical diseases	Yes	-67.42	-89.62	-45.23	-59.64%	-79.28%	-40.01%	0.803
	No	-61.13	-96.90	-25.63	-59.77%	-94.74%	-25.63%	


Different letters mean significant differences according the Bonferroni post-hoc test (p<0.05).

Table 3. Sedentary behavior change (pre-post pandemic) in self-isolated adults during the SARS-CoV-2 pandemic in 2020 in Brazil

Characteristics	Category	SB change (pre versus during)						
		Mean change in minutes	95% CI		Delta%	95% CI		Interaction p value
Sex	Male	161.41	119.48	203.35	42.51%	31.47%	53.56%	0.380
	Female	152.28	111.87	192.69	41.72%	30.61%	52.78%	
Age	Young adults (18-35 years)	190.48a	149.65	231.30	47.62%	37.42%	57.81%	<0.001
	Middle age adults (36-55 years)	143.35b	99.48	187.21	39.87%	27.62%	51.99%	
	Older adults (55-64 years)	136.71b	77.88	195.54	38.34%	21.82%	54.80%	
Ethnicity	Black/Asian/Mixed/Others	155.43	109.04	201.81	41.42%	29.07%	53.81%	0.940
	White	158.26	121.24	195.29	42.81%	32.80%	52.83%	
Marital status	Single/divorced/widowed	176.15a	133.74	218.56	46.87%	35.58%	58.15%	0.024
	Married	137.54b	96.85	178.24	37.29%	26.18%	48.32%	
Employment	Employed/students/military	179.85a	142.15	217.55	45.58%	36.02%	55.14%	0.030
	Unemployed/retired	133.84b	79.26	188.42	38.22%	22.63%	53.81%	
Monthly household income	<R\$2005	171.17	118.64	223.70	47.39%	32.85%	61.94%	0.632
	R\$2005-R\$8640	166.74	125.52	207.95	45.81%	34.49%	57.14%	
	R\$8641-R\$11261	150.63	102.05	199.21	40.36%	27.34%	53.38%	
	>R\$11261	138.85	92.10	185.59	35.50%	23.54%	47.45%	
Current smoking	Yes	158.87	108.25	209.49	42.72%	29.10%	56.33%	0.984
	No	154.82	116.98	192.66	41.53%	31.38%	51.68%	
Current alcohol consumption	Yes	173.39	134.25	212.53	46.86%	36.28%	57.44%	0.091
	No	140.30	96.75	183.85	37.44%	25.82%	49.06%	
Self-reported previous diagnosis of mental disorders	Yes	173.16a	129.97	216.34	45.02%	33.79%	56.25%	0.003
	No	140.53b	101.07	180.00	39.03%	28.07	49.99%	
Self-reported previous diagnosis of physical diseases	Yes	139.60	107.20	172.00	37.85%	29.07%	46.64%	0.358
	No	174.09	117.49	230.69	46.31%	31.25%	61.37%	

Different letters mean significant differences according the Bonferroni post-hoc test (p<0.05).

Figure 1. Moderate to vigorous physical activity and sedentary behavior, pre and during COVID-19 pandemic in 2020 in Brazil



Values are shown as the estimate marginal means, in minutes per day, of moderate to vigorous physical activity (MVPA) and sedentary behavior (SB) together with their standard error. Significant changes across time were found for MVPA ($p < 0.001$) and SB ($p < 0.001$).