Different adrenal response of critically ill neonates; An Egyptian model

Suzan Abd Razik , Wafaa Osman A

Pediatrics' and neonatology Department - Faculty of Medicine – Ain Shams University, Egypt.

Corresponding author: Wafaa osman A   Address:-Pediatrics' and neonatology Department - Faculty of Medicine -Ain Shams University, Egypt. Tel: +20225117993;

E-mail: wafaaosman83@gmail.com

 wafaaosman83@med.edu.asu.eg

Short running head: Adrenal response configuration in ill preterm

Keywords: ACTH, Cortisol, critically, ill, neonates

Abstract

Background: Adequate adrenocortical function is essential for survival of critically ill neonates. Although most of them display elevated plasma cortisol concentrations, which reflects activation of the hypothalamic pituitary adrenal axis (HPA), yet; adrenocortical insufficiency is seen in septic shock. **Objectives**: Evaluate the HPA response in critically ill neonates with shock. **Subjects**:this prospective observational study included a total of 60 neonates divided into 3 groups;(A)30 critical ill neonates with septic shock on inotropic support, (B)15 patients with sepsis who did not need inotropic support and(C) control group(n=15). **Methods**: Adrenal functions were evaluated by taking a single diurnal ACTH reading and two readings for serum cortisol level (diurnal and nocturnal). **Results**: Gram negative organism was more prevalent among the patients; 53%, 63% in groups A and B respectively. Group A showed Significant statistical hypotension before vasopressor drug administration (p<0.001) as compared to both groups. Group A showed Significant statistical improvement of blood pressure after vasopressor drug administration (p<0.001) as compared to both

groups B, C. Serum cortisol was sigificantly higher in group A(57.21±24.31) and B (48.01±18.27), while it was lower in group C(19.57±16.05) . A highly statistically significant rise of serum cortisol level(p=0.000) and ACTH(p=0.000) in group A when was compared to the other two groups**Conclusion:** This study introduced a new pattern of serum cortisol response in neonates ranging from very high cortisol level to a near normal values; highlighting a state of glucocorticoid resistance in neonates and relative adrenal insufficiency.

Introduction

During stressful conditions and shock; the patient tends to respond by release of hypothalamic hormone including corticotropin releasing hormone (CRH) and arginine vasopressin 1 that stimulate release of adrenocorticotropin (ACTH) from the pituitary gland; with subsequent secretion of cortisol from the adrenal glands. Toxemia triggers inflammatory cytokine release with subsequent activation of the clotting cascade to form microthrombi. Following that vasodilation of arteries and arterioles occur and lower the vascular resistance and increase cardiac output that lead to warm shock. If no treatment received and the condition was not reversed; Blood pressure and cardiac output decrease leading to cold shock *(Constantine et al., 2016).* Microthrombi cause poor capillary perfusion of kidneys, adrenal glands, brain and heart. Inflammatory mediators can cause both hypothalamic pituitary adrenal axis (HPA) activation and blunting ; by the effect of TNF, which cause relative adrenal insufficiency(RAI). RAI can result from dysfunction of HPA at any point from hypothalamus down to the adrenal, blunting the effect of glucocorticoids causing tissue resistance and irreversibility of shock *(Asha et al., 2009).*

Many mechanisms were postulated to explain the microcirculatory changes that occur during sepsis. Among those; the mechanism that postulate dysregulation of nitric oxide production together with functional impairment of body cells including endothelial cells damage, the lost adrenergic sensitivity of the smooth muscle cells, increased platelets aggregation and subsequent increased leukocyte endothelial reactions(*(Marcos et al.,2016)* .)

Subjects and methods

Subjects

The 60 patients were further divided into three groups

Group A: included 30 critically ill patients who are proved clinically and laboratory to have septic shock (15 preterm and 15 fullterm); thus furtherly subdivided into 22(73.3%) patients received double inotropic drug(dopamine and dobutamine on a dose up to 20 mcg /kg/min) and the other 8 patients received both drugs plus noradrenaline on a dose of 0.1 to 0.5 mcg /kg/min.

Group B: Included 15 patients proved to have sepsis and they did not need inotropic support; (9 preterm, 6 full term)

Group C: 15 health control neonates (8 pretrm, 7 full term), both cases and controls were sex and age matched.

Inclusion criteria: hemodynamically unstable neonates. The enrolled babies who developed hypotension and needed inotropic support; the mean arterial pressure < 2 SD for the gestation.

Exclusion criteria: suspected cause of adrenal insufficiency, example patients proved to have shock due to: cardiac cause, adrenal cause, ambiguous genitalia, central

cause(example; brain malformations, encephalitis, panhypopituitarism), postoperative, postnatal exogenous steroid treatment.

Methods

This study was a prospective descriptive study, that included 60 patients aged below one month (gestational age 32-36weeks), admitted to neonatal intensive care (NICU), Ain Shams university hospital over period of six month from january to june, 2019. Written consents were obtained from care givers of the admitted neonates. The study protocol was approved by Ain Shams University NICU Ethical Committee, Fullfills the criteria of the Helsinski Declaration. Informed consent was taken from mother or legal guardians of included neonates in the study.

History taking and clinical examination

All patients were thoroughly examined after taking full history with emphasis on history of maternal illnesses, gestational age, diagnosis on admission to NICU. The need for respiratory support and inotropes were recorded to facilitate further subgrouping of patients

IV- Laboratory investigations

A- Sepsis indicators: Total Leucocytic Count (TLC), neutrophil count, CRP, bacterial blood culture and other cultures

C- Metabolic profile: Liver function tests, kidney function tests, serum electrolytes, calcium, magnesium and phosphorus, random blood sugar, serum albumin

D- ACTH and serum cortisol levels: During sepsis total serum cortisol level was measured in neonates with sepsis on inotropic support (double or triple inotropic drugs) twice in the morning (8-9 am) and and in the evening (8-9.30 pm).

ACTH was measured once in the morning (8-9 am)

Measurement of serum cortisol:

Using enzyme chemiluminescent technique on the immulite (DPC).

a) Specimen collection: 2mL of venous blood were withdrawn and left to clot. The separated sera were stored at -20°C.

b) Principle of the procedure: It is a solid phase competitive chemiluminescent enzyme immunoassay. The solid phase is a polystyrene bead enclosed within an immulite test unit, it is coated with murine anti cortisol

antibody. The patient sample and alkaline phosphatase conjugated cortisol are simultaneously introduced into the test unit, and incubated for approximately 30 minutes at 37C with intermittent agitation.

During this time, cortisol in the sample competes with the enzyme labeled cortisol for a limited number of antibodies binding sites on the head. Unbound enzyme-conjugate is then removed by a centrifugal wash, after which substrate is added and the test unit is incubated for a further 10 minutes.The chemiluminescent substrate, a phosphate ester of adamantly dioxetane, undergoes hydrolysis in the presence of alkaline phosphatase to yield an unstable intermediate. The continuous production of the intermediate results in the sustained emission of light. The bound complex and thus also the photon output, as measured by the luminometer is inversely proportional to the concentration cortisol in the sample.

c) Reagents

1-Cortisol test units (100 units): Each barcode-labeled unit contains one head. 1 Kco: 100 units.

2- Cortisol reagent wedge (LCO,) with barcode 6.5 ml alkaline phosphatase (bovine calf intestine) conjugated to cortisol in buffer. Store capped and refrigerated.

3- Cortisol adjustors (LCOL, LCOH): Two vials (low and high), 3mL each, cortisol in processed human serum.

Measurement of serum adrenocortical hormone (ACTH

a) Specimen collection:5 cc of venous blood were collected by vein puncture (heamolysis was avoided), the tubes were immersed in an ICP bath. Following collections; plasma was separated from the cells

by centrifugation in refrigerated centrifuge, then aliquoted and freezed immediately in plastic tubes at -20C. ACTH test units (LAC1) (100 units). Each barcode labeled unit contains one bead.

b) Principle of the procedure:

IMMULITE ACTH is a solid phase, two-site chemiluminescent immunometric assay. The solid phase, a polystyrene bead enclosed within an immulite test unit, is coated with a monoclonal antibody specific for ACTH. While the patient serum sample and alkaline phosphatase-conjugated

polyclonal antibody are incubated for approximately 30 minutes at 37C in the test unit with intermittent agitation, ACTH in the sample is bound to form an antibody sandwich complex. Unbound complex is then removed by a centrifugal wash, after which the substrate is added and the Test Unit is incubated for further 10 minutes.

The chemiluminescent substrate, a phosphate ester of adamantly dioxetane, undergoes hydrolysis in the presence of alkaline phosphatase to yield an unstable intermediate. The continuous production of these intemediate results in the sustained emission of light, thus improving precision by providing a window for multiple readings. The bound complex and thus also the photon output, as measured by the luminometer is proportional to the concentration of ACTH in the sample.

The immulite system automatically handles sample and reagent additions, the incubation and separation step, and measurement of the photon output via the temperature controlled luminometer. It calculates the test results for controls and patient samples from the observed signal, using a stored

master curve, and generates a printed report.

c) ACTH reagent wedges (LACA, LACB) LACA: 6.5 mL of an alkaline phosphatase (bovine calf intestine) conjugated to polyclonal rabbit anti-ACTH antibody in buffer. ACTH adjustors (LACL, LACH)Two vials (low and high) of lypophilized ACTH inabovine protein-based matrix

Results:

Demographic data

   Group A had 15 preterm males and 15 full term females, while group B had 6 full term and 9 preterm babies; 11 of them were females and group C had 7 full term females and 8 preterm males. Three groups were age and sex matched. There was no significant difference between the 3 studied groups regarding mode of delivery, sex and age.

Laboratory results

Both groups A and B had significantly high total leukocytic count, neutrophil and CRP as compared with the control group.

Reverse correlation between neutrophil count and cortisol level was documented.

Both ACTH and cortisol are not affected by addition of inotropes in group A.

Culture proved sepsis

Sepsis proved cultures was evident in 27 out of 45 patients of both groups A(19) and B(8), where Gram positive staph aureus and E coli had the highest proportion of culture yield (25% each), followed by both streptococci and pneumococci (7.1% each) and finally was the klebsiella proved cultures in (3.6%) of patients.

Blood pressure in the 3 groups

Systolic blood pressure was nearly equal in group B and C (mean=70.2± 6.44), (71.93±9.6) respectively, while it was lower in group A (46.6±6.58). So as the diastolic pressure; that was nearly in the same range; (41.6±7.3), (41.93±5.39) in groups B and C respectively, while group A had lower diastolic range (23.39±3.88). Although blood pressure was improved after addition of inotropes (group A); yet both diastolic and systolic were lower than other 2 groups.

Cortisol level assessment in the 3 groups

Serum cortisol (both readings) was higher in both groups A, B as compared with group C. While ACTH was nearly in the same range in both groups B and C; yet it was found to be higher (double time) in group A rendering a significant statistical difference. A characteristic pattern of cortisol release was documented in group A; 73.33% had quantitative folding (am>pm values with concomitant rise), 50% had

quantitative folding and disturbed rhythm (am>pm values with inconcomitant rise), 23.33% had quantitative folding with reversed rhythm (pm>am) and 10% had quantitative low level (am was within normal range, pm slightly increased)

Discussion

culture proved sepsis

Group A had culture proved sepsis in 19(63.3%); of which 42% was gram negative, while group B had culture proved sepsis in 8(53%); of which all were gram negative. E coli was the most prevalent followed by the klebsiella species. This is supported by Macharashivii *et al.* (2009) and Couto *et al.* (2007) who reported 18-78% of neonatal sepsis were due to gram negative organism.

Blood pressure difference among the three groups

Non statistical difference in blood pressure was found comparing group B and C. While hypotension was evident in group A before and after inotropic support in comparison to the other two groups showing a highly statistical significant difference, indicating a state of a peripheral tissue resistance expressing a vasopressor resistant hypotension that was more pronounced in group A especially preterm patients. **Table 1** To demonstrate the response of premature babies versus full term ones regarding vasopressor drugs (groupA); a total number of 30 ill neonates was equally divided into preterm and full term, where 11 of each group received double inotropic support while the other 4 received triple inotropic support. The difference of response of full term and preterm neonate, in face of vasopressor drugs, can be shown by improved systolic blood pressure in full term post inotropic support as compared to preterm of

same group(p<0.05). So full term babies showed a statistical significant rise in systolic blood pressure compared to preterm patients.

In addition, after starting inotropic support, we found preterm patients of group A had statistically significant hypotension as compared to those in group B, while on the other hand a non statistically significant difference was observed among full term patients of same groups(A,B), indicating a more poor response to vasopressors and immature hemodynamic compensatory mechanisms in pre terms. This goes on with Biniwale *et al.* 2013 who described a vasopressor resistance hypotension in pre terms. **Table 2**

   This confirmed the poor response to vasopressor and an immature hemodynamic compensatory mechanisms, also Higgins *et al.* (2010) stated that some preterms are either vasopressor resistant and/ or dependant due to the relative adrenal insufficiency plus , down regulation of cardiovascular adrenergic receptors due to both critical illness and chronic stimulation. As one of the genomic actions of corticosteroids is the upregulation of those mentioned receptors and the down regulation thought to be due to RAI

   Although absolute adrenal insufficiency is quite rare in critically ill neonates, relative adrenal insufficiency is relatively common. Together with the presence of hemodynamic instability despite adequate fluids and vasopressors; other diagnostic clues include hyponatremia, hyperkalemia and hypoglycemia (Sprung *et al.* 2008).

In the present study, on monitoring the laboratory clues for adrenal insufficiency, hyperkalemia was statistically significant in group A as compared to the other two groups, while hyponatrmia and hypoglycaemia showed a non statistical difference.

This agrees with Fernandez *et al.,* 2009 who reported a hyperkalemic state in ill preterm presenting with inadequate adrenal insufficiency. But it is important to take into consideration that electrolyte assessment was not taken in serious evaluation because the ill groups (A,B) were receiving continuous fluid therapy in NICU. So in this study, laboratory evidence of adrenal insufficiency was assessed basically on measuring total serum cortisol level.

Assessing serum cortisol level in the 3 groups

In assessing the total serum cortisol level in all the studied groups, the following data were recorded; First, normal neonates (group C) showed a circadian rhythm of an (am to pm) ratio (3:1) in both preterms and full terms, with a mean cortisol level (19.57ug/dl) and a range of (3.5-35ug/dl) in am values and range of (2.9-6.9ug/dl) in pm values. **Table 3.** In addition, there was non signficant difference in serum cortisol and ACTH levels comparing normal preterm and normal full terms. Opposite to this, Vermes *et al.* (2001), mentioned that diurnal rhythms were still absent in first 2 months, and the typical circadian rhythms of the plasma cortisol levels were present in infants aged 3 months. Second, total serum cortisol level shows a highly statistical significant rise in both group A and B together with loss of the recorded rhythm in group C (normal neonates). In agreement with that Hvenkatesh *et al.* (2005); Kristenson, 2011) mentioned that, the normal circadian rhythm of cortisol secretion, (characterized by achievement of the highest concentration in the morning and subsequently decreased nadir in the evening), is lost during critical illness. This also goes on with study done by Asare (2007) who stated that for critically ill patients, it is not

necessary to obtain cortisol levels at a specific time of the day because most patients lose the diurnal variation in their cortisol levels.

In the current study it was also found that in group A (septic neonates with inotropic support), a quantitative folding of mean total serum cortisol level together with dysrythmic pattern were detected, categorizing this group into four response patterns. Category I: 15/30(50%) quantitative folding shows of both (am and pm) values with a concomitant rise. Category II: 5/30(16.6%) quantitative and reversed rhythm shows quantitative folds of both (am and pm) values but with a pm values higher than am values. Category III: 3/30(10%) quantitative and disturbed rhythm shows quantitative folds than normal of am values while pm values showed a relatively lower folding values compared to the previous two categories. Finally, Category IV: 7/30(23%) (relative low) where am values fall within normal range values despite severe sepsis, i.e; relative low serum cortisol. In this category preterm represents (71%) and full term 28.5%.

Passing through the adrenal gland response in the four categories gives an idea about the state of the hypothalamic pituitary axis (HPA) activation behaviour in sepsis and septic shock, where a stage of initial rise occurs in serum cortisol level exemplified in category (I&II), followed by exhaustion of the adrenal gland, exemplified in category (III&IV). Fernandez *et al.* (2009) reported that patients with severe sepsis have cortisol values three fold higher than normal and that very preterm neonates would be anticipated to be at increased risk for cortisol insufficiency in the face of acute illness and stress.

Relative adrenal insufficiency is present when the cortisol response is inadequate for the patient's degree of stress, despite a cortisol level that may appear normal. This is due to dysfunction at any point in the HPA axis or due to resistance to glucocorticoids (Gomez, 2013).

Asare (2007) defined Relative adrenal insufficiency as a random cortisol level of less than 25 ug/dl. That random cortisol testing is a useful diagnostic threshold for the diagnosis of adrenal insufficiency.

In the present study, aiming at assessing the severity of the vasopressor resistant shock based on adrenal gland response (serum cortisol level), it was found that triple inotropic therapy was used in (4/8) 50% in category I (highest folding), (2/8) 25% in category IV (relatively low) followed by 12.5% in other two categories (III&IV). Very high levels in category I also indicates a state peripheral tissue resistance to glucocorticoids, while the relatively low category (IV) indicates a state of an adrenal exhaustion.

This goes on with the study done by Annane *et al.* (2008), who mentioned that, the hypothalamus-pituitary-adrenal axis is activated, initially leading to increased free cortisol. Next, the entire axis is exhausted and peripheral cortisol resistance leads to the development of adrenal insufficiency.

This also agrees with Arnildo *et al.* (2011), who described a state of adrenal exhaustion in chronic critical illness due to catecholamine receptor desensitization or down regulation and/or chronic secretion of cytokines and other substances that suppress the HPA axis.

In the present study, it was concluded that relative adrenal insufficiency can be either in a quantitative form as presented in (relative low) category IV (mean serum cortisol 19 ug/dl) "adrenal exhaustion", or as a functional adrenal insufficiency despite high cortisol levels indicating "peripheral tissue cortisol resistance". Conforming the state of 'tissue resistance", mean serum cortisol level in preterms (55.13ug/dl) and full terms (59.29 g/dl) in group A (septic neonates with inotropic support) showed a significant and highly significant reversed correlation respectively to both systolic and diastolic blood pressure. This goes on with Gomez (2013), who found that relative adrenal insufficiency is present when the cortisol response is for the patient's degree of stress, despite a cortisol level that ever that may appear normal and was explained by dysfunction at any point in the HPA axis or due to resistance to glucocorticoids. Prigent *et al.* (2004) and Indyk *et al.* (2013) also reported that an inadequate HPA axis response to stress can aggravated by peripheral resistance to glucocorticoids. In addition to reduction of glucocorticoid receptors and an increase in the conversion of cortisol to inactive cortisone by enhanced activity of 11-B hydroxysteroid dehydrogenase stimulated by IL-2, IL-4, and IL-13. Mark (2003) and Arnildo *et al.* (2011) believed that a random cortisol level in severely stressed patients (ie. with hypotension) should be above 25 mcg/dl and that higher levels may be appropriate in patients with septic shock due to "tissue cortisol resistance. Fernandez *et al.* (2009) and Cooper *et al.* (2007) recorded that random cortisol value of <15 mcg/dl is considered to be a clear evidence of adrenal insufficiency. Also Soliman *et al.* (2004) reported that 11 of 30 (36.6%) septic term newborns had basal cortisol values of <15 mcg per 100 ml.

Dellinger *et al.* (2008), reported that relative adrenal insufficiency is seen in 16.3 to 55% of the patients with septic shock. While, Annane *et al.* (2006) , Hebbar *et al.* (2012) and Shraga (2013), reported that prevalence rates of adrenal insufficiency (AI) in critically ill patients vary according to the characteristics of the study population and the diagnostic criteria used and rates as high as 60-90% have been documented in patients with septic shock. Elfaramawy (2012), reported that the proportion of relative adrenal insufficiency in children with sepsis (66.6%) was much higher than that without sepsis (25%), which declares the strong association between sepsis and septic shock with relative adrenal insufficiency.

Cortisol versus blood pressure in both preterm and full term

There was reversed significant correlation between serum cortisol level and both systolic and diastolic blood pressure in preterm and fullterm of group A. This finding could demonstrate the intact HPA axis while reflects the resistance at the tissue levels. While we reported a significant correlation between ACTH and serum cortisol in full term (p=0.00 7) ; the same correlation was not found in preterm group(p=0.68).

Studying serum ACTH behaviour in all the studied groups, it was concluded that ACTH showed a highly statistically significant rise in group A, (septic neonates with inotropic support) compared to the other two groups (group B septic neonates not receiving inotropes and group C normal neonates). **Table 3**. The high mean ACTH (29.3pg/dl) together with the high mean cortisol level (57.2ug/dl) in group A (septic neonates with inotropic support) (a positive correlation) indicates either a direct stimulation of the HP (hypothalamic pituitary axis) provoked by the septic inflammatory mediators in case of severe sepsis and septic shock (a hypothalamic pituitary depend-

ent mechanism), loss of the normal negative feedback mechanism (negative feedback block) between the adrenal and pituitary glands, indicating a central resistance of the hypothalamic/pituitary to the adrenal glucocorticoids. A poor adrenal response to ACTH can be also a contributing factor. These factors collectively lead to the creation of a pseudo positive feedback loop between pituitary and adrenals.

The non significant levels of ACTH in group B (septic neonates without inotropic support) reaching (14.17 pg/dl) nearly the normal values in group C (14.1 pg/dl), despite the rise in serum cortisol level (mean 48ug/dl), indicates that the initial rise m serum cortisol in response to sepsis may occur as "a pituitary independent mechanism" in cases not reaching the septic shock state, or may be in part due to reduction in negative feedback effect of cortisol on hypothalamic pituitary axis (negative feedback block).

This goes on with Sharaga (2013), who mentioned that although severe stress activates the HPA axis, dissociation may occur between plasma ACTH and cortisol concentrations, as demonstrated by suppressed ACTH and elevated cortisol concentrations. Addingly, a study done by Boonen *et al.* (2013), suggested a pituitary independent mechanism for increased cortisol production during critical illness. In group A, number of inotropes used, sex and mode of delivery showed non statistical significant correlation with serum cortisol level and ACTH. In group A; studying the drugs that may be aggravating the state of relative adrenal insufficiency: antiepileptic drugs (pheytoin) and antifungal (fluconazole) usage was reported. 5/30(16.6%) received antifungal therapy (fluconazole) and 8/30(26.6%) received anti epileptic (phenytoin), 2 out of these 8 were preterm and showed a relatively low serum cortisol level (catego-

ry IV). Rock *et al.* (2007), mentioned that, phenytoin, barbiturates, and rifampicin are drugs that accelerate the metabolism of cortisol and most synthetic glucocorticoids by inducing hepatic mixed- function oxygenase enzymes, causing adrenal insufficiency in patients with limited pituitary or adrenal reserve and in agreement with this Nicolas *et al.* (2011), also, mentioned an association between high-dose fluconazole and adrenal insufficiency in critically ill patients.

Both serum cortisol and ACTH level were not affected by mode of delivery, sex or the inotropic support that was added in group A.

conclusion

Septic shock causes a state of relative adrenal insufficiency that contributes to the higher mortality rate. Early steroid measurement and replacement can improve vassopressor resistance and hypotension leading to a better clinical outcome.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this paper

Funding

This research did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

Suzan A Razek acquisition, analysis and interpretation of data, drafting of paper;
…… conception and design, revision for critical intellectual content;
……………, final approval;
Wafaa Osman : final approval, agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved

Reference:

Tsigos, Ioannis Kyrou, Eva Kassi, Kenneth Feingold and George P Chrousos:
Stress, Endocrine Physiology and Pathophysiology. In: Endotext [Internet].
South Dartmouth (MA): MDText.com, Inc.; 2000–.2016 Mar 10.

Asha Tyagi, Ashok Kumar Sethi, Gautam Girotra, and Medha Mohta: The Mi
crocirculation in Sepsis. Indian J Anaesth. 2009 Jun; 53(3): 281–293.

Marcos Miranda, Michelle Balarini, Daniella Caixeta, and X Eliete Bouskela,
Oswaldo Cruz Foundation, Fiocruz, Rio de Janeiro, RJ, Braz: Microcirculatory
dysfunction in sepsis: pathophysiology, clinical monitoring, and potential thera
pies.Am J Physiol Heart Circ Physiol 311: H24–H35, 2016. doi:10.1152.

Annane D, Maxime V, Ibrahim F, Alvarez JC, Abe E,Boudou P 2006 Diagnosis of
adrenal insufficiency in severe sepsis and septic shock. *Am J Respir Crit Care
Med*; **174**:1319-1326. (doi: 10.1164/rccm.200509-1369)

Arnildo LJ, Eduardo AR, Luiza KM, Jefferson RC 2011 Adrenal insufficiency in
children with sepsis. *Pediatric Intensive Care Unit*; **23(4)**:478-483.
(doi.org/10.1590/S0103-507X2011000400013).

Asare K 2007 Diagnosis and Treatment Adrenal PatientInsufficiency in the Critical-
ly
ill patient. *Pharmacotherapy*; **27(11)**:1512-1528. (doi: 10.1592/phco.
27.11.1512)

Biniwale M, Sardesai S and Seri I 2013 Steroids andVasopressor-Resistant Hypoten

sion in Preterm Infants.*Current Pediatric Reviews*, **9(1)**: 75-83.

Boonen E, Vervenne H, Meersseman P, Andrew R, MortierL, Declercq PE, Vanwijn

gaerden YM, Spriet 1,Wouters PJ, Vander Perre S, Langouche L,

Vanhorebeek

I, Walker BR, Van den Berghe G 2013 Reduced cortisol metabolism during

critical illness. *N Engl J*; **368**:1477-1488.L.(doi: 10.1056/NEJMoa1214969.

Epub 2013 Mar 19.).

Cooper MS and Stewart PM 2007 Adrenal insufficiency in critical illness. *J Inten*

sive Care Med; **22(6)**:348-62.(doi.org/10.1177/0885066607307832).

R. Phillip Dellinger, Mitchell M. Levy, Jean M. Carlet, Julian Bion, Margaret M.

Parker, Roman Jaeschke, Konrad Reinhart, Derek C. Angus, Christian Brun-

Buisson, Richard Beale, et al. 2008 Surviving Sepsis Campaign: International

guidelines for management of severe sepsis and septic shock: 2008. *Intensive*

Care Med **34**, 17–60. (doi.org/10.1007/s00134-007-0934-2)

Fernandez EF and Watterberg KL 2009 Relative adrenal insufficiency in the pre-

term

and term infant. Division of Neonatology, Department of Pediatrics, University of

New Mexico School of Medicine, Albuquerque, NM, USA *Journal of Perinatol*

ogy; official Journal of the California Perinatal; **29**:S44-S49. (doi: 10.1038/jp.

2009.24).

Gomez-Sanchez CE 2013 Adrenal dysfunction in criticallyill patients. *N Engl J Med*;

368:1547-1549. (doi: 10.1056/NEJMe1302305).

Hebbar KB, Petrillo T, Fortenberry JD 2012 Adrenal insufficiency and response to

corticosteroids in hypotensive critically ill children with cancer. *J Crit Care*. Epub 2012 Jun 12; **27**:480-487. (doi: 10.1016/j.jcrc.2012.03.013).).

Virginie M and Djillali A 2004 Clinical review: *Corticotherapy in sepsis Critical Care*, **8**:122-129. (doi: 10.1186/cc2374).

Higgins S, Friedlich P, Seri I 2010 Hydrocortisone for hypotension and vasopressor dependence in preterm neonates: a meta-analysis. *Journal of Perinatology*, **30(6)**: 373-378. (doi: 10.1038/jp.2009.126)

Indyk JA, Candido-Vitto C, Wolf IM, Venkataraman S,S, Munoz R, Saladino RA, Witchel SF, Defranco DB 2013 Reduced glucocorticoid receptor protein ex pression in children with critical illness. *Horm ResPaediatr*; **79**: 169-178. (doi: 10.1159/000348290. Epub 2013 Mar 8).

Macharashvili N, Kourbatova E, Butsashvili M,N.Tsertsvadze T, McNutt LA, Leonard MK 2009 Etiology of neonatal blood stream infections in Tbilisi, *Re public of Georgia. Int J Infect Dis*;**13(4)**:499-505.(doi: 10.1016/j.ijid. 2008.08.020).

Mark 2011 Endocrine Development - Adrenal Glands UNSW. *Embryology* www.embryology.med.Unsw.edu.com

Nicolas CN, Evangelia C, George PC 2011 AdrenalInsufficiency ch13 endotext, the most accessed source on endocrinology for medical professionals.

Shraga LY 2013 Critical illness-related corticosteroidinsufficiency in children. *Hormone Research in Paediatrics*, **80(5)**: 309-317.

Soliman AT, Taman KH, Rizk MM, Nasr IS, AI-RimawyH and Hamido MS 2004 Cir

culaing adrenocorticotropic hormone (ACTH) and cortisol concentrations in nor mal, appropriate for gestational age newborns versus those with sepsis and respi ratory distress: cortisol response to low-dose and standard-dose ACTH tests. *Me tabolism*; **53**:209-214.

Sprung CL, Annane D, Keh D, Moreno R, Singer M,Freivogel K, et al. 2008 Corti cus Study Group.Group.Hydrocortisone therapy for patients with septic shock. *NEngl J Med*; **358**:111-124.(doi: 10.1056/NEJMoa071366).

Vermes I and Beishuizen A 2001 The hypothalamic-pituitary-adrenal response to critical illness. *Best Pract Res Clin Endocrinol Metab*; **15**:495-511. (doi: 10.1053/ beem.2001.0166)

Vrezas I, Willenberg HS, Bornstein SR 2013 (www.Endotext) Chapter 1, Adrenal Cortex. Development Anatomy, Physiology USMLE Step 1 *Physiology Lecture Notes*; **pp**. 263-289

Stress, Endocrine Physiology and Pathophysiology

Constantine Tsigos 1 , Ioannis Kyrou 2 , Eva Kassi 3 , George P Chrousos 4

1 , Bradley Anawalt 2 , Alison Boyce 3 , George Chrousos 4 , Kathleen Dungan 5 , Ashley Grossman 6 , Jerome M Hershman 7 , Gregory Kaltsas 8 , Christian Koch 9 , Peter Kopp 10 , Márta Korbonits 11 , Robert McLachlan 12 , John E Morley 13 ,

Maria New [14], Leigh Perreault [15], Jonathan Purnell [16], Robert Rebar [17], Frederick Singer [18], Dace L Trence [19], Aaron Vinik [20], Don P Wilson [21], editors.

Blood pressure	neonates with inotropes group A	neonates without inotropes group B	Normal neonates group c	One way ANOVA
	N=15	N=15	N=30	(p value) significance
Systolic (before inotropes)	46.68 6.58	6.44 70.2	9.6 71.93	(0.000) HS
Diastolic (before inotropes)	23.39 3.88	7.37 41.6	5.39 41.93	(0.000) HS
Systolic (after inotropes)	61.54 8.53	6.44 70.2	9.6 71.93	(0.001) HS
Diastolic (after inotropes)	35.43 5.6	7.37 41.6	5.39 41.93	(0.002) HS

Table(1):Comparison between the studied groups regarding blood pressure parameters before and after inotropic support

This table shows a highly statistically significant decrease in mean value of the systolic and diastolic blood pressure for group A as compared to the other two groups (B,C) even after inotropic support.

Table (2): Comparison between all the studied groups regarding serum cortisol level and ACTH .

	neonates with inotropes group A N=15	Neonates without inotropes group B 15=N	Normal neonates group c N=30	p value significance
Cortisol am	57.21 t 24.31	48.01 t 18.27	19.57 16.05	0.000 HS
Cortisol pm	44.72 t 24.08	31.52 15.36	4.96 +2.00	0.001 HS
ACTH am	29.31 t 17.12	14.11 3.79	14.17 +3.28	0.000 HS
POST HOC tests :LSD				
Parameters	Normal vs Sepsis with inotropes.	Normal vs Sepsis without Inotropes	Sepsis with inotropes vs sepsis without inotropes	
Cortisol am	0.000 HS	0.002 HS	0.245 NS	
Cortisol pm	0.000 HS	0.027 S	0.097 NS	
ACTH am	0.001 HS	0.990 NS	0.002 HS	

Normal level of ACTH=10-60 pg/ml. Normal serum cortisol level =5-25 ug/dl

This table shows a highly statistically significantdecrease in mean value of the systolic and diastolic blood pressure for group A compared to the other two groups (B, C) even after inotropic support..