Quantifying SARS-CoV-2 infection risk within the Google/Apple exposure notification framework to inform quarantine recommendations

Amanda M. Wilson, PhD[1]; Nathan Aviles[2], James I. Petrie[3], Paloma I. Beamer, PhD[1] ; Zsombor Szabo[3], Michelle Xie[3], Janet McIllece[4], Yijie Chen[5], Young-Jun Son, PhD[5]; Sameer Halai[3], Tina White[3], Kacey C. Ernst, PhD[1]; Joanna Masel, DPhil[6,*]

[1] Mel & Enid Zuckerman College of Public Health, University of Arizona, 1295 N Martin Ave, Tucson, AZ 85724, USA
[2] Graduate Interdisciplinary Program in Statistics, University of Arizona, 617 N. Santa Rita Ave. University of Arizona, Tucson, AZ 85721, USA
[3] Covid Watch, 433 S Stratford Drive, Tucson, AZ 85716, USA
[4] World Wide Technology, 1 World Wide Way, St. Louis, MO 63146, USA
[5] Systems and Industrial Engineering, University of Arizona, 1127 E. James E. Rogers Way, CIMS Lab, 85721, USA
[6] Ecology & Evolutionary Biology, University of Arizona, 1041 E Lowell St., Tucson AZ 85721
* corresponding author masel@email.arizona.edu 520 626 9888

ORCIDs: AMW 0000-0003-3259-8169, NA 0000-0001-9998-4406, JIP 0000-0001-7959-7687, PIB 0000-0001-5287-2183, ZS 0000-0002-3653-1378, MX 0000-0001-9512-3684, YC 0000-0001-5316-4103, YS 0000-0002-4004-2155, KCE 000-0002-3346-7788 JM 0000-0002-7398-2127

Abstract

Background: Most Bluetooth-based exposure notification apps use three binary classifications to recommend quarantine following SARS-CoV-2 exposure: a window of infectiousness in the transmitter, ≥15 minutes duration, and Bluetooth attenuation below a threshold. However, Bluetooth attenuation is not a reliable measure of distance, and infection risk is not a binary function of distance, nor duration, nor timing.

Methods: We model uncertainty in the shape and orientation of an exhaled virus-containing plume and in inhalation parameters, and measure uncertainty in distance as a function of Bluetooth attenuation. We calculate expected dose by combining this with estimated infectiousness based on timing relative to symptom onset. We calibrate an exponential dose-response curve on the basis of the infection probabilities of household contacts. The conditional probability of current or future infectiousness, conditioned on how long post-exposure an exposed individual has been free of symptoms, decreases during quarantine, with shape determined by the distribution of incubation periods, proportion of asymptomatic cases, and distribution of asymptomatic shedding durations. It can be adjusted for negative test results using Bayes Theorem.

Findings: We capture a 10-fold range of risk using 6 infectiousness values, 11-fold range using 3 Bluetooth attenuation bins, ~6-fold range from exposure duration given the 30 minute duration cap imposed by the Google/Apple v1.1, and ~11-fold between the beginning and end of 14 day quarantine. Imposing a consistent risk threshold for the probability of infection can recommend quarantine with weaker Bluetooth signal, even when not recommended for the entirety of the infectious period.

Interpretation: The Covid-Watch app is currently programmed either to use a threshold on initial infection risk to determine 14-day quarantine onset, or on the conditional probability of current and future infectiousness conditions to determine both quarantine and duration. Either threshold can be set by public health authorities.

Funding: University of Arizona

Research in Context:

Evidence before this study

Epidemiological models suggest a benefit from smartphone-based automatic notification of
exposure to Covid-19 infected individuals. Google and Apple have rolled out a privacy-
preserving API with which apps can offer such notifications based on the degree of Bluetooth
attenuation of a signal issued by an infected individual, its duration, and its timing relative to
symptom onset in and test date for that infected individual. Apps deployed so far have focused
on approximating close contact definitions of 15 minutes and a given distance such as 1 meter
or 2 meters. However, Bluetooth attenuation is not a reliable measure of distance, and infection
risk is not a binary function of distance, nor duration, nor timing.

Added value of this study

We integrate all sources of information (including our own new experiments on the relationship
between distance and Bluetooth attenuation) to provide a measure of infection risk that is as
accurate as possible given the data on exposure that is available to an app based on the
Google/Apple system. We also provide a method to calculate the probability of current or future
infectiousness, which is a function not only of initial infection risk, but also of the number of
symptom-free days since exposure, and any negative test results.

Implications of all available evidence

Our risk scoring system can guide quarantine recommendations in exposure notification apps,
targeting them to individuals who are the most likely to be infected. It can also recommend
release from quarantine on the day that the probability of current or future infectiousness falls
below a threshold, leading to different quarantine durations for different individuals.

Background

Manual contact tracing followed by quarantine of known contacts is a critical method for containing or mitigating the spread of communicable diseases. It is, however, extremely resource and time-intensive and relies on case recall of contacts. New technologies can supplement this approach.[1] Manual contact tracing can be effective for COVID-19[2–4], however, a significant challenge is the extremely short window of time between an infected individual presenting for testing and the contacts that they infected beginning to shed infectious virus.[5,6] Automatic exposure notification approaches based on Bluetooth proximity have the potential to achieve many of the benefits of contact tracing, while also providing more rapid notification, greater privacy[7,8], more objective recall of contacts including those whose identity is unknown to the case, and greater scalability.[5,9] The two approaches of contact tracing and exposure notifications are complementary and may directly interact e.g. when those receiving digital exposure notifications are referred to human contact tracers for the information and support needed for quarantine adherence and further investigation.[10]

Apps have access to data on timing, duration, and Bluetooth attenuation. Determining the threshold for entering quarantine based on probability of infection should yield better results than from combining three binary thresholds for duration, distance, and the infectious period of the transmitter. A threshold for exiting quarantine based on the conditional probability of current or future infectiousness could also be used. Both would help optimize the reduction in disease transmission per day of quarantine recommended.

Here we lay out a framework for doing so using the decentralized protocol of the Google/Apple Exposure Notification (GAEN) Application Programming Interface (API). When a user reports positive infection status, the GAEN v1.1 framework (Figure 1) allows apps to assign a "Transmission Risk Level" to each day that they might have been shedding, and to communicate this level to the receiver's phone via a Temporary Exposure Key (TEK). On the receiver's device, the GAEN framework records Bluetooth attenuation as a rough estimate of distance, and the duration of exposure.

The risk of infection depends on viral dose,[11] which in turn depends on the shedding rate of the infected individual, and on the duration and distance of the interaction. As days go by without onset of symptoms, the probability of future infectiousness decreases, because the probability is conditioned on lack of symptoms for an increasing stretch of time. We parameterize calculations of both probabilities using both past literature and new experiments and illustrate what different risk thresholds imply for quarantine recommendations. We are piloting and evaluating the Covid Watch app using portions of this scheme on the campus of the University of Arizona.

129
130
131 **Figure 1.** Assessment of the probability of infection following a single exposure. The calibration
132 work is reported in this manuscript, and the procedures on the Transmitter's and Receiver's
133 phones are part of the Covid Watch app.

134
135
136 **Methods**

137
138 The overall approach to calculating infection risk is summarized in Figure 1. Parameter values
139 and their descriptions and sources are summarized in Supplementary Table 1 for calculations
140 performed by the app and in Supplementary Table 2 for parameters we used during calibration.

141
The GAEN v1.1 framework records exposure durations only up to 30 minutes, in order to protect
anonymity of COVID-positive patients by limiting the risk that users will be able guess the
source of their exposure, while still meeting contact definitions that invoke minimum exposure
duration of 15 minutes. Durations can be recorded separately within three bins of attenuation,
thus allowing a total of 90 minutes if the attenuation varies in the necessary manner over the
exposure. We calculate a weighted sum of the three durations, using the weights to capture the
differences in expected dose (number of inhaled particles over an exposure time). Note that
GAEN v1.5 lifts the duration cap but restricts our ability to record levels of infectiousness. We
focus here on GAEN v1.1, but our scheme can be adapted for later versions.

151
We measured Bluetooth attenuation under GAEN v1.1 for a range of distances, phones, and
scenarios of possible signal interference with the potential to affect the attenuation – distance

relationship (Supplementary Materials Section 1).[12] We rebalanced this dataset to form a
pseudo dataset that is more representative of the distribution of barriers and scenarios in the
real world (Supplementary Materials Section 1).

From this pseudo dataset, we first sample a datapoint that falls within the attenuation bin in
question. If this is a zero-risk barrier scenario, we assign an infection risk of 0. Otherwise, we
record the distance ρ in meters. Note that our method is not based on mapping thresholds in
distance to thresholds in Bluetooth attenuation, but instead on resampling from the probability
distribution of distance as a function of attenuation.

We feed this distance into a microbial exposure model that estimates the airborne spread of
viral particles from an emitter's mouth following a Gaussian plume formation, and their
subsequent inhalation by contacts (Supplementary Materials Sections 2 and 3). We use a
Monte Carlo approach to sample angle, exhalation rate of the transmitter, cross-section of the
transmitter's open mouth, and inhalation rate of the exposed individual, to obtain a mean
dose/time for that attenuation bin. For distances ≤ 1 meter, we assume face to face interactions,
consistent with distances measured for "interpersonal" interactions.[13] We choose thresholds
between attenuation bins, and relative risks for time spent in each bin (Supplementary Materials
Section 4).

We use an exponential dose-response curve, which is derived from the assumption that each
host is susceptible and that each virus has an independent probability of survival and
subsequent initialization of infection.[11] In our case, this probability k, multiplied by a constant C
to convert from arbitrary units to number of viruses, sets the parameter $\lambda = kC$ in the equation

$$P(\text{infection}) = 1 - e^{-\lambda D},$$

where expected dose D comes from a shedding rate multiplied by a weighted sum of time spent
within 3 attenuation ranges. An exponential dose-response curve is superior to the approximate
beta-Poisson for some other viruses (http://qmrawiki.org/content/recommended-best-fit-
parameters, accessed 09/07/2020). These viruses include adenovirus, enterovirus, poliovirus,
and SARS-CoV-1.

Our weighted sum of durations and our estimates of shedding rates S in the Results are both in
arbitrary units. We therefore fit λ to obtain infection probabilities that are compatible with
household spread. Asymptomatic infection and low test sensitivity can both deflate estimated
household infection risks, while indirect chains of infection via a third household member can
inflate them. A meta-analysis by Curmei et al.[14] attempted to correct for these complications and
estimated a secondary attack rate of household contacts of 30%. We assumed exposure is
equivalent to 8 hours with the maximum shedding rate in the lowest attenuation and calculated
λ for this dose that would result in a 30% infection risk.

Our scheme can be used either to 1) set a threshold on the initial probability of infection to
trigger 14-day quarantine, or 2) set a threshold for the probability of current or future
infectiousness to determine both who should quarantine and for how long. Our method for
calculating the probability of current or future infectiousness, conditioned on lack of symptoms
so far and incorporating the possibility of asymptomatic infection, is described in Supplementary
Materials Section 5. The parameters used are summarized in Table 1. This method can be
extended to inform the timing of testing and the effect of a negative test result on a
recommended duration of quarantine (Supplementary Materials Section 6). Note that strictly

speaking when using this latter threshold, our "quarantine" recommendations are, through their
treatment of the possibility of undiagnosed asymptomatic infection, a combination of quarantine
and isolation.
207
208 Our scheme, by expressing exposures in terms of probabilities of infection and infectiousness,
209 naturally lends itself to combining risks over multiple exposures (Supplementary Materials
210 Section 7).
211

212
Figure 2. Expected dose and corresponding probability of infection for a 30-minute exposure,
estimated using our Monte Carlo procedure as a function of distance from an infected individual.
The discontinuity at 1 meter indicates our assumption that this distance threshold indicates face-
to-face interaction. Faded points show doses and infection risks that would be estimated if a
face-to-face or non-face-to-face interaction assumption were consistent across distances. The
bolded points indicate what we assumed in our framework. Note that Bluetooth information likely
contains more risk information regarding whether an interaction was face-to-face than it does
about risk as a function of the distance at which either a face-to-face or a non-face-to-face
interaction takes place. The WHO close contact definition invoking 1 meter also invokes face-to-
face interaction.[15] The same is true, only with 2 meters, for European guidance (available at
https://www.ecdc.europa.eu/en/covid-19-contact-tracing-public-health-management, accessed
on 7/1/2020).[16] The Centers for Disease Control and Prevention (CDC)'s definition departs from
this in omitting reference to face-to-face when referring to interactions occurring within 6 feet
(available at https://www.cdc.gov/coronavirus/2019-ncov/php/public-health-
recommendations.html, accessed on 7/1/2020).[17]

Results

Our Gaussian plume model of microbial exposure produces the relationship between distance and infection risk shown in Figure 2. Training on both this and our distance-attenuation measurements (as summarized in Methods), we chose the 3 attenuation bins of ≤ 50dB (close range), 50-70dB (medium range), and >70 dB (far range), with weights 2.15, 0.79, and 0.19, respectively. Using these weights, we calibrate λ = 3.50 x 10^{-6} (see Methods) to obtain an infection probability of 0.30 for household contacts. Note that the best way to calibrate both weights and λ would be after the app is rolled out, with manual contact tracers compiling exposure characteristics and relating them to the rate of subsequent infection. While Eq. 5 calculates the function of an expectation rather than an expectation of a function, treating variance in dose amounts to using an "effective" value of λ (Supplementary Materials Section 8).

Bluetooth attenuation thus only distinguishes a 2.7-fold difference in dose and hence risk between close and medium range, and only 11-fold between close and far range. In contrast, informed both by TCID50 data[18] and by epidemiological evidence[19], we assign a 10-fold higher risk to exposures to individuals during peak shedding than during the margins of the infectious period (Supplementary Materials Section 9, illustrated in Figure 3A). The magnitude of shedding (transmission risk value) has received less attention than attenuation and exposure duration and is not to our knowledge widely used by other GAEN apps.

Apps can measure duration far more accurately than distance, but the GAEN v1.1 duration cap of 30 minutes, combined with the use of 5 minute or similar windows, mean that only ~6-fold differences in the risk from low-attenuation exposures are captured in this way, or up to 8.6-fold when stochastic variation in attenuation over time also shows up at lower attenuations, or much longer in GAEN v1.5 where there is no cap on duration.

The relatively low predictive power of Bluetooth attenuation gives rise to diagonal patterns in the quarantine recommendations in Figure 3B. These diagonal patterns mean that quarantine will sometimes be recommended following prolonged exposure to a high shedder, even if the interaction took place at well beyond the estimated 2 m. distance. However, these exposures are not risk-free either, in particular if taking place in an indoor environment, especially in cases with heavy breathing, such as exercise environments[20] or choir rehearsals,[21] where aerosols may mix throughout the room and also deposit on surfaces. The diagonal pattern reflects the compelling evidence that exposure timing and duration also significantly contribute to infection risk. We therefore sometimes recommend quarantine recommendation even when Bluetooth attenuation, which is a poor proxy for distance, is not low. However, Bluetooth attenuation is nevertheless critical to concluding that an interaction occurred at all.

Figure 3. Examples of quarantine recommendations using a threshold for infection risk (B) vs. for current or future infectiousness (C). Shaded cells indicate that a 30-minute interaction is insufficient to trigger quarantine. A) Transmission risk levels 1-6 are used to capture the 10-fold range of relative infectiousness on different days as a function of timing relative to symptom onset. Evidence from both transmission pairs and TCID50 measurements is reviewed in the Supplementary Materials Section 9. B) The minimum length interaction needed to trigger 14-day quarantine is a function both of Bluetooth signal attenuation and of Transmission Risk Level. Approaches that neglect the latter correspond to a single row of 15 minutes, and potentially a second row of 30 minutes. C) Number of quarantine days recommended following a 30-minute interaction.

So far, we have estimated the probability of infection from an exposure. Each day that passes
without symptoms provides more information to make infection less likely, and eventually also to
increase the probability that shedding from an asymptomatic infection has ended. To calculate
the probability of current or future infectiousness on a subsequent day, conditional on no
symptoms until that day, we apply a discount factor based both on time elapsed without
symptoms and also any negative test results (Supplementary Materials Sections 5 and 6). We
multiply the probability of infection from an exposure by this discount factor to determine the
remaining risk of infectiousness from a given exposure.

Traditional quarantine guidelines are binary (either 14 days from date of last exposure, or no
quarantine required). However, a consistent approach to risk, combined with a desire to impose
quarantine days in the most efficient manner possible to combat disease spread, suggests that
individuals should quarantine for longer following a higher-risk exposure (Figure 4A) (although
see Supplementary Materials Section 5 for a caveat with very high doses). This approach
calculates the number of days post-interaction that would be needed to drop below a given
threshold of probability of current or future infectiousness. Exposure scenarios of 30 minutes are
illustrated in Figure 3C.

We used a 0.13% threshold in Figure 3C, because it recommends a 14-day quarantine for 15
minutes in close range with a high shedder. Such an interaction has a 1.12% infection risk,
which falls below a 0.13% probability of current or future infectiousness after 14 days of
quarantine during which no symptoms appear. Note that this initial infection risk is broadly
compatible with the attack rate reported in Taiwan (1.0%, 95% CI: 0.6-1.6%) for those
interacting with infected individuals in the first 5 days of symptom onset,[22] which is similar to the
1.9% attack rate (95% CI 1.8%–2.0%) reported in South Korea.[23]

Current advice treats the larger risk of longer exposure the same, making a 0.13% threshold
more conservative because it is calculated to generate a 14-day quarantine for a minimal
duration of exposure. However, this is offset by our assuming maximal shedding in calculating
this benchmark example. In other words, while this threshold approximates the risk tolerance of
current advice, the details of who is recommended for quarantine and for how long will be
different in our quantification of total risk than it would be if we were to combine independent
binary thresholds for infectious period of transmitters, duration of exposure, and distance to
produce a quarantine duration of uniform length. This leads to more consistent treatment of risk
to yield the greatest benefit in terms of transmission prevented per day of quarantine
recommended. Shorter quarantines might significantly reduce the harms imposed by
quarantine,[24] and increase compliance[25] (although see McVernon et al[26]). Quarantining for 14
days post-exposure may be exceptionally challenging for essential workers, individuals without
sick leave, or those who would endure significant financial hardship due to lost income.

The assumed fraction of asymptomatic infections affects the discounting of risk. The
symptomatic fraction is discounted according to the distribution of incubation periods from
exposure to symptom onset, while releasing the asymptomatic fraction from quarantine is not
safe until not only onset, but also shedding is over (Supplementary Materials Section 5). Our
calculations so far assume that 20% of infections are asymptomatic. If we instead assume that
50% infections are asymptomatic, e.g. in a young age group, even a 15-minute contact
registered as low attenuation and with peak shedding in the transmitter would require a 16-day
quarantine to meet a 0.13% threshold (Figure 4B). However, if an individual were to test

negative during their quarantine, their conditional probability of current or future infectiousness
would drop, shortening their quarantine to 13 days for a test with 70% sensitivity (Figure 4C).
334
335

336
Figure 4. Applying a consistent risk tolerance for current or future infectiousness causes
quarantine duration to be a function of initial risk, of the tolerated degree of risk, of the fraction of
infections that are assumed to be asymptomatic, and of any negative test results. A) Initial
infection risk is 1.12% following 15 minutes of close contact with an individual around the time of
symptom onset. With a 20% asymptomatic fraction, a 14-day quarantine is recommended under
a 0.13% risk threshold, but only a 7-day quarantine under a 0.5% threshold. Following a lower
risk exposure with 0.2% infection risk, quarantine would be 5 days with the stricter threshold,
and there would be no quarantine with the less strict. B) Quarantine must be longer to mitigate a
high likelihood of asymptomatic infection in the exposed individual. C) A negative test result,
shown here as taking place on Day 5, can shorten quarantine, in particular mitigating the risk of
asymptomatic infection. We apply Bayes theorem with 70% sensitivity and 100% specificity.
Note that widespread availability of testing would allow much stricter risk thresholds to be used.
Day 0 is included in the total quarantine times.
350
351 **Discussion**
352
353 Here we quantify relative risk of infection using experiments to inform the noisy distance-
354 attenuation relationship, and Monte Carlo simulations to inform both this and other sources of
355 variability and uncertainty that affect risk. We roughly calibrate relative infection risk to absolute
356 probability of infection based on limited information from the infection probability of household
357 contacts.
358
359 Errors in calibration are likely but will generally not affect the rank order of risks. For example,
360 adjusting the risk threshold of 0.13% for quarantine will have similar effects to adjusting the
361 value of λ. Knowledge of absolute vs. relative risk does have some effect once some saturation
362 in risk begins to occur, little of which will occur unless much longer durations are recorded.
363
With 20% cases being asymptomatic and no testing, the risk of current or future infectiousness
falls ~11-fold over the first 14 days of quarantine. Under GAEN v1.5, risk will sometimes differ
more between two individuals entering quarantine than when comparing the same individual
before vs. after a 14-day quarantine. For this reason, our scheme could recommend
quarantines longer than 14 days. Variation in quarantine length is to be expected – if total risk is
scored consistently, some quarantines will be longer and others shorter, in order for residual
infection probability, conditional on time elapsed without symptoms, to fall below a threshold.
371

The Covid Watch app is currently programmed either to use a threshold on infection risk to
determine 14-day quarantine onset, or on risk of current and future infectiousness to determine
both quarantine and duration. Either threshold can be set by public health authorities flexibly in
the light of external factors such as level of community transmission, jurisdictional comfort with
uncertainty related to digital exposure notifications, and current public health science and
recommendations. Communities that have achieved containment might choose to set a stricter
threshold, testing individuals once or twice to lower their risk following each negative test.
379
Communities with high prevalence might raise the threshold if it seems likely that the number of
quarantine recommendations being issued by the app will cause it to fall out of use. Note that
the maximum possible initial infection risk in our v1.1 scheme is 3.23%; relaxing the 30-minute
cap on durations, while maintaining infectiousness information, would be necessary in order to
get resolution among higher risks. The effects of this problem are partly ameliorated by using
the variable quarantine duration approach (Figure 3C). When a threshold is set well below the
probability that a randomly chosen member of the population is currently infected, it should be
recognized that individuals agreeing to download and comply with the recommendations of the
app are implicitly agreeing to adhere to higher standards than those implied by the current
absence of a general stay-at-home order. At the time of writing (September 5), the rate of
current infection is ~1% in Arizona.[27]
391
392   When the infection risk of the average person in the population is high, we believe that the best
393   solutions are population-level restrictions and closures. Under these circumstances, a GAEN
394   app might still have utility for essential workers. However, a GAEN app could be an inferior but
395   still useful option should the political will for population-level restrictions not exist.
396
As the conditional probability of current or future infectiousness (conditioned on the exposed
individual being asymptomatic) falls throughout their quarantine period, messaging can also
change. E.g., during the initial high risk days, users might be offered concrete resources such
as grocery delivery, or the option to quarantine in a specialized facility in order to protect other
household members, before transitioning to self-quarantine once risks falls. Even with self-
quarantine, an app might identify the days on which staying home is the highest priority (I.e.
days where the potential infectivity may be highest). Messaging considerations are discussed in
Supplementary Material Section 10.
405
We caution that our derived relationship between Bluetooth attenuation and infection risk is
extremely approximate and model-dependent. We have more confidence in our settings of
Transmission Risk Levels for symptomatic cases, but very little for asymptomatic cases. These
parameters need to be calibrated with real world data on app users who report their app-
recorded exposures to manual contact tracing efforts, who then track which users go on to test
positive, and who are therefore able to mine the data to quantify the quantitative relationship
between exposure details (duration, attenuation, Transmission Risk Level) and probability of
infection. Transfer of this data to manual contact tracers' contact management databases is
critical to improve the targeting of quarantine recommendations to those at highest risk of being
infected, which will make most efficient use of each day of quarantine recommended to reduce
transmission.
417

Short of this, more quantitative data on infectivity would be extremely valuable. Our
determination of shedding duration partly relies on the prospective sampling of all individuals in
a skilled nursing facility[28], where many patients subsequently got sick. Daily samples during
similar outbreaks could be used to quantify how shedding varies both among individuals and as
a function of time relative to symptom onset. TCID50 data would be ideal, but even Ct values
can be valuable for this purpose. However, the fact that the settings we originally chose based
on infectivity data agreed with later and improved epidemiological approaches[19,29] is
encouraging.
426
Without extended durations, our default calibrations will not recommend quarantine (Figure 3B)
or extended quarantine (Figure 3C) for lower Transmission Risks. However, with the duration
cap lifted in a GAEN v1.5 or later, 43 minutes in the ≤50 dB range, 1.92 hours in the 50-70 dB
range, or 7.93 hours in the >70 dB range with an individual of transmission risk level 2 would be
required. However, GAEN v1.5 does not quantify infectiousness, and GAEN v1.6 re-introduces
only two levels of infectiousness. Limited durations and infectiousness information have been
driven by privacy concerns, but this must be weighed against the significant ethical
considerations in favor of efficient allocation of quarantine.[30] In Supplementary Section 10, we
suggest an alternative method to preserve anonymity, which is to conceal all exposure details
from the user's view. When using variable quarantine duration, this also effectively conceals the
date of exposure.
438
Our framework can be used not only to guide recommendations for who should quarantine and
for how long, but also to allocate associated resources including quarantine facilities, grocery
delivery and other social support, and priority for access to scarce tests. Both manual contact
tracing and digital exposure notification require rapid testing to be effective. Given limited tests,
targeting those at highest risk of infection will do the most good in finding new positive cases
who are early enough in the course of infection for these approaches to stem transmission the
most.
446
447   **Data Availability:** Supplementary Data Table 1 provides the alpha test data used to calibrate
448   our weights.
449
450   **Code Availability:** Code and necessary data are accessible under a Creative Commons
451   license at https://github.com/awilson12/risk_scoring
452

453   **Acknowledgements**
454
We thank the many testers who participated in alpha testing of the app for their time and
feedback, including those attempting to test in Androids who were stymied by technical issues.
We also thank Marm Kilpatrick for literature pointers, Jacob Steinhardt and Bruce Helming for
comments on a draft manuscript, Dave Lines for writing instructions for executing test scenarios
and providing participant support, Ami Buczek for installing the app on loaner phones, Danielle
Zhou for helping with testing instructions, setting up test logs, and making test assignments, all
of the Covid Watch team for agreeing to include our risk scoring scheme into their app, Yuanxia
Li and Sudha Ram for agreeing to let us publish alpha testing data ahead of and separately
from their analysis, Luca Ferretti for sharing a manuscript in preparation that helped confirm our

choice of pre-symptomatic Transmission Risk Scores, Gary Windham for doing the code
reviews that led the University of Arizona to adopt Covid Watch, Darcy Van Patten for
organizing the details of that adoption, and Joyce Schroeder for her leadership of the Covid
Watch pilot at the University of Arizona. Funding was provided by the University of Arizona.

Competing Interests

Author Contributions

JM conceived the overall risk scoring approach and oversaw all aspects of the project. AMW,
PIB, and JM designed the Gaussian plume component, and AMW implemented it. JM and AMW
reviewed the literature to choose Transmission Risk Levels. JM designed and implemented the
variable quarantine approach. MX and JMc designed and managed the alpha testing. JIP
preprocessed and cleaned the alpha test data. YC and YJS designed and performed people
traffic simulations to inform a realistic distribution of distances. NA implemented the rebalancing
procedure to reflect that distribution, and performed sampling to choose weights. AMW and JM
calibrated the dose-response curve and NA explored its sensitivity to variance in dose. JIP, ZS,
SH, and TW designed and implemented the core components of the Covid Watch app used in
the alpha testing. JIP modified the app to obtain high resolution attenuation data. ZS provided
up-to-date guidance about what is possible under GAEN, and checked calculations. PIB and
KCE provided input on all public health aspects of the research. AMW and JM wrote the first
draft of the manuscript, followed by significant input from PIB and KCE.

References

1   Armbruster B, Brandeau ML. Contact tracing to control infectious disease: When enough
     is enough. *Health Care Manag Sci* 2007; **10**: 341–55.
2   Kucharski AJ, Klepac P, Conlan A, *et al.* Effectiveness of isolation, testing, contact tracing
     and physical distancing on reducing transmission of SARS-CoV-2 in different settings.
     medRxiv 2020; **3099**: 2020.04.23.20077024.
3   Aleta A, Martin-Corral D, Pastore Y Piontti A, *et al.* Modeling the impact of social
     distancing, testing, contact tracing and household quarantine on second-wave scenarios
     of the COVID-19 epidemic. *medRxiv Prepr Serv Heal Sci* 2020; : 1–31.
4   Bi Q, Wu Y, Mei S, *et al.* Epidemiology and transmission of COVID-19 in 391 cases and
     1286 of their close contacts in Shenzhen, China: a retrospective cohort study. *Lancet
     Infect Dis* 2020; **3099**: 1–9.
5   Ferretti L, Wymant C, Kendall M, *et al.* Quantifying SARS-CoV-2 transmission suggests
     epidemic control with digital contact tracing. *Science (80-)* 2020; **368**: eabb6936.
6   Kretzschmar ME, Rozhnova G, Bootsma MCJ, van Boven M, van de Wijgert JHHM,
     Bonten MJM. Impact of delays on effectiveness of contact tracing strategies for COVID-
     19: a modelling study. *Lancet Public Heal* 2020; **5**: e452–9.
7   Fraser C, Abeler-Dörner L, Ferretti L, Parker M, Kendall M, Bonsall D. Digital contact
     tracing: comparing the capabilities of centralised and decentralised data architectures to
     effectively suppress the COVID-19 epidemic whilst maximising freedom of movement and

maintaining privacy. Introduction 2. 2020.

8   Von Arx S, Becker-Mayer I, Blank D, *et al.* Slowing the Spread of Infectious Diseases
Using Crowdsourced Data. Covid Watch. 2020. https://blog.covidwatch.org/en/covid-
watch-whitepaper-using-crowdsourced-data-to-slow-virus-spread (accessed Aug 19,
2020).

9   Salathé M, Althaus CL, Anderegg N, *et al.* Early Evidence of Effectiveness of Digital
Contact Tracing for SARS-CoV-2 in Switzerland. Github. 2020; : 1–18.

10   Webster RK, Brooks SK, Smith LE, Woodland L, Wessely S, Rubin GJ. How to improve
adherence with quarantine: rapid review of the evidence. *Public Health* 2020; **182**: 163–9.

11   Haas CN, Rose JB, Gerba CP. Quantitative Microbial Risk Assessment. John Wiley and
Sons, 1999.

12   Farrell S, Leith DJ. Pairwise Handset Types and Orientations Are Sufficient to Blur
Exposure Notification Thresholds. Trinity Coll. Dublin, Irel. 2020.
https://down.dsg.cs.tcd.ie/tact/posorient.pdf (accessed Aug 6, 2020).

13   Zhang N, Su B, Chan PT, Miao T, Wang P, Li Y. Infection spread and high-resolution
detection of close contact behaviors. *Int J Environ Res Public Health* 2020; **17**.
DOI:10.3390/ijerph17041445.

14   Curmei M, Ilyas A, Evans O, Steinhardt J. Estimating Household Transmission of SARS-
CoV-2. *medRxiv* 2020. DOI:10.1101/2020.05.23.20111559.

15   World Health Organization. Considerations for quarantine of individuals in the context of
containment for coronavirus disease (COVID-19). 2020.
https://www.who.int/publications/i/item/considerations-for-quarantine-of-individuals-in-the-
context-of-containment-for-coronavirus-disease-(covid-19) (accessed July 1, 2020).

16   European Centre for Disease Prevention and Control. Public health management of
persons, including healthcare workers, having had contact with COVID-19 cases in the
European Union - second update. ECDC. 2020. DOI:10.2807/1560-
7917.ES.2020.25.9.2000202.

17   Centers for Disease Control and Prevention. Coronavirus Disease 2019 Community-
Related Exposures. 2020. https://www.cdc.gov/coronavirus/2019-ncov/php/public-health-
recommendations.html (accessed Jan 7, 2020).

18   Bullard J, Dust K, Funk D, *et al.* Predicting infectious SARS-CoV-2 from diagnostic
samples. *Clin Infect Dis* 2020; : 1–18.

19   Ashcroft P, Huisman JS, Lehtinen S, *et al.* COVID-19 infectivity profile correction. *arXiv*
2020. http://arxiv.org/abs/2007.06602.

20   Jang S, Han SH, Rhee J-Y. Cluster of Coronavirus Disease Associated with Fitness
Dance Classes, South Korea. *Emerg Infect Dis* 2020; **26**. DOI:10.3201/eid2608.200633.

21   Hamner L, Dubbel P, Capron I, *et al.* High SARS-CoV-2 Attack Rate Following Exposure
at a Choir Practice - Skagit County, Washington, March 2020. *Morb Mortal Wkly Rep*
High 2020; **69**: 606–10.

22   Cheng HY, Jian SW, Liu DP, Ng TC, Huang WT, Lin HH. Contact Tracing Assessment of
COVID-19 Transmission Dynamics in Taiwan and Risk at Different Exposure Periods
before and after Symptom Onset. *JAMA Intern Med* 2020.
DOI:10.1001/jamainternmed.2020.2020.

23   Park YJ, Choe YJ, Park O, *et al.* Contact Tracing during Coronavirus Disease Outbreak,
South Korea, 2020. *Emerg Infect Dis J* 2020; **26**. DOI:10.3201/eid2610.201315.

24   Brooks SK, Webster RK, Smith LE, *et al.* The psychological impact of quarantine and
how to reduce it: rapid review of the evidence. *Lancet* 2020; **395**: 912–20.

25   Soud FA, Cortese MM, Curns AT, *et al.* Isolation compliance among university students
during a mumps outbreak, Kansas 2006. *Epidemiol Infect* 2009; **137**: 30–7.

26   McVernon J, Mason K, Petrony S, *et al.* Recommendations for and compliance with

social restrictions during implementation of school closures in the early phase of the
influenza A (H1N1) 2009 outbreak in Melbourne, Australia. *BMC Infect Dis* 2011; **11**.
DOI:10.1186/1471-2334-11-257.
27   covid19-projections.com. 2020. https://covid19-projections.com/us-az (accessed Sept 5,
2020).
28   Arons MM, Hatfield KM, Reddy SC, *et al.* Presymptomatic SARS-CoV-2 infections and
transmission in a skilled nursing facility. *N Engl J Med* 2020; **382**: 2081–90.
29   Ferretti L, Ledda A, Wymant C, *et al.* The timing of COVID-19 transmission. *medRxiv*
2020. DOI:10.1101/2020.09.04.20188516.
30   Singer P, Masel J. How (Not) to Fight COVID 19. Proj. Synd. 2020; published online Sept
3. https://www.project-syndicate.org/commentary/tech-solutions-to-targeting-covid19-
quarantine-by-peter-singer-and-joanna-masel-2020-08.
574

Supplemental Materials for

Quantifying SARS-CoV-2 infection risk within the Google/Apple exposure notification framework
to inform quarantine recommendations

Amanda M. Wilson[1], Nathan Aviles[2], James I. Petrie[3], Paloma I. Beamer[1], Zsombor Szabo[3], Michelle Xie[3], Janet McIllece[4], Yijie Chen[5], Young-Jun Son[5], Sameer Halai[3], Tina White[3], Kacey C. Ernst[1], Joanna Masel[6,*]

[1] Mel & Enid Zuckerman College of Public Health, University of Arizona
[2] Graduate Interdisciplinary Program in Statistics, University of Arizona
[3] Covid Watch
[4] World Wide Technology
[5] Systems and Industrial Engineering, University of Arizona
[6] Ecology & Evolutionary Biology, University of Arizona
* corresponding author masel@email.arizona.edu

ORCIDs: AMW 0000-0003-3259-8169, NA 0000-0001-9998-4406, JIP 0000-0001-7959-7687, PIB 0000-0001-5287-2183, ZS 0000-0002-3653-1378, MX 0000-0001-9512-3684, YC 0000-0001-5316-4103, YS 0000-0002-4004-2155, KCE 000-0002-3346-7788 JM 0000-0002-7398-2127

Contents

43 **Supplementary Table 1. Parameter values used by app to calculate risk**

Parameter	Description	Distribution or Point Value		References
S	Viral shedding rate in arbitrary units that are proportional to viral copies/m^3	10^1	5 days pre- or 8-9 days post-symptom onset	10-fold range informed by TCID50 measures[1], timing informed by[1–4]
		$10^{1.2}$	6-7 days post-symptom onset, or asymptomatic within 2-4 days of test	
		$10^{1.4}$	4 days pre- or 5 days post-symptom onset, or asymptomatic within 1 day of test	
		$10^{1.6}$	3 days pre- or 4 days post-symptom onset	
		$10^{1.8}$	2 days pre- or 3 days post-symptom onset	
		10^2	1 day pre- to 2 days post-symptom onset	
T_{low}, T_{med}, T_{high}	Duration of exposure	Durations for Bluetooth attenuations ≤50dB, between 50dB and 70dB, and >70dB are multiplied by weights 2.15, 0.79, 0.19, respectively		This study
λ	Probability that one viral particle establishes infection × conversion from arbitrary units	3.50×10^{-6}		Calibrated from secondary attack rate of household contacts = 30%.[5]
Fraction of asymptomatic infections	Higher values lead to longer quarantine	20% for a population, but depends on age		[6]
Incubation period	Days until symptom onset	Probabilities for {0,1,2…} days = {0,4E-05, 0.011842, 0.088541, 0.181965, 0.207344, 0.174797, 0.123761, 0.081488, 0.051057, 0.031469, 0.018734, 0.011235, 0.006786, 0.00422, 0.002518, 0.001626, 0.000978, 0.000592, 0.000364, 0.000231, 0.00014, 0.000093, 0.000062, 0.00004, 0.000025, 0.000017, 0.000011, 0.000008}		[7]
Asymptomatic shedding duration	We assume that asymptomatic shedding begins 3 days before what would have been the day of symptom onset if symptomatic, or else immediately upon infection, whichever occurs later, and that shed viral particles are nonviable beyond 12 days	Cumulative probabilities for days {5,6,7…}={0.054054054,0.094594595,0.12162162,0.148648649, 0.189189189,0.21621621,0.256756757,1}		[8,9]
Risk threshold	Set by public health agency, cognizant of limitations in current calibration of λ	Benchmark of 0.5% for probability or infection or 0.13% for probability of current or future infectiousness		This study

 Supplementary Table 2. Parameter values used by us to calibrate parameter values in Table 1

Parameter	Description	Distribution or Point Value	References
X	Exhalation rate (m³/day)	Normal(16.3, 4.15), left-truncated at 9	[10]
A	Cross-sectional area of the mouth. Used to calculate the breath velocity, U from X	Uniform(23, 59) (cm²)	[11]
I_y	"Lateral intensity" of plume deviation	Uniform(0.08, 0.25)	[12]
I_z	"Vertical intensity" of plume deviation	Uniform(0.03, 0.07)	
I	Inhalation rate (m³/day)	Normal(16.3, 4.15), left-truncated at 9	[10]
ρ	Distance	Sampled from attenuation-distance dataset to inform weights.	This study
φ	Angle between the z-axis and the xy-plane	Used while informing weights. If $\rho \leq 1m$, $\varphi = \pi/2$, If $\rho > 1m$, φ randomly sampled from Triangular(min= $\pi/4$, mode= $\pi/2$, max= $3\pi/4$)	1 m. cutoff for face-to-face interaction informed by [13]
θ	Angle between x and y axes	Used while informing weights. If $\rho \leq 1m$, $\theta = 0$, If $\rho > 1m$, θ randomly sampled from Uniform(0, 2π)	1 m. cutoff for face-to-face interaction informed by [13]

46
47

1.0 Experiments on Distance-Attenuation Relationship

Using a developer version of the Covid Watch app, we called the API multiple times with different attenuation thresholds in order to achieve resolution of 3dB in the 30dB-99dB range. The API appears to round up durations to 5-minute increments, each with its own attenuation value; we consider each of these to be a datapoint. Our tests were all short, e.g. a 12-minute test would yield 3 datapoints.

There were 7 testers and 14 phones, representing a variety of models, all of iPhones – handset type and orientation can affect signal.[14] 49 measurements were taken with specific phone orientations, while for the remaining 986 the devices were side-by-side and facing upwards if not otherwise specified by the barrier type (e.g. pocket). 203, 222, 199, 374, 17, 20 measurements were at 0.5m., 1m., 1.5m., 2m., 3m., and 5m., respectively. We also used the 28, 28, 29, 27, and 16 zero-risk barrier measurements at 0.5m., 1m., 1.5m., 2m., and "N/A", respectively. Note that improvements made in GAEN v1.5 are expected to improve our ability to estimate distance as a function of attenuation. While useful, we note that new experimental data would be required for recalibration. The imbalanced nature of the collected data, with few long distance measurements and no Android measurements, reflects the fact that the decision was made to recommence once v1.5 was available, whose calibration will be both qualitatively and quantitatively different.

1163 out of the total 1558 datapoints were used for attenuation weight and attenuation threshold setting, with exclusions of data described in Figure S1. The 1163 non-excluded datapoints are supplied in Supplementary Dataset 1. Of the included 1035 attenuation measures that involved infection risk, 747 did not include a deliberate barrier, while 288 includes barriers such as pockets, backpacks, nearby laptop, and human body. 925 measures were taken inside homes, 49 were taken inside an elevator, and 61 outside.

Figure S1. Attenuation data cleaning

*A few attenuation values were implausibly low, always representing one such increment per device per series of attenuation values corresponding to a single test. We believe this is because these testers used the dropdown menu to turn off Bluetooth rather than going to Settings, and this only disables existing Bluetooth connections, causing an anomalously strong signal to be recorded during the period in which the test was being set up. We manually annotated these, totaling 144 datapoints, and excluded them from further analysis, yielding 1414 datapoints.

**Distance is not needed for the 128 datapoints taken in the presence of a risk-blocking barrier (e.g. closed car doors or walls), so points marking "N/A" for distance were not excluded for any of these.

To rebalance the distance measurements, we created a pseudo-dataset with different multiples of the data collected at each of the distances. To inform the desired distribution of distances, we analyzed the time-weighted pairwise distance in traffic flow simulations of a classroom.[15] These indicate a roughly uniform distribution over possible distances, with a reduction in close contact due to attempts to adhere to social distancing rules. Since close contact might be more common in other settings, and distances beyond 5m. can also register Bluetooth signal, we made 5, 5, 6, 3, 132, and 168 copies of the non-zero-risk data at distances of 0.5m., 1m., 1.5m., 2m., 3m., and 5m, respectively, yielding a data ratio of 1015 : 1100 : 1194 : 1122 : 2244 : 3360 (as a rough approximation of a target ratio of 1:1:1:1:2:3) prior to the sampling described below. To this, we added 4 copies of the zero-risk barrier measurements, so that they made up 4.85% of the total pseudo-dataset.

Our calibration code holds shedding rate and exposure duration constant at 50 arbitrary units/m^3 and 30 minutes, in order to isolate the effect of distance on differences in dose between attenuation buckets.

2.0 Estimation of Exposure Concentrations

It is well-acknowledged that both distance from an infected individual and duration of "close proximity interactions"[16] are important parameters in estimating the probability of infection of those exposed.[17–20] However, there is little quantitative information about the relationship between distance and risk of infection. Chu et al. (2020)[18] quantified risk in terms of answers to binary survey questions about whether the respondent came within distance X of an infected person. They found that the value of the threshold distance X in the survey question predicts the degree to which the answer predicts risk, but this relationship cannot easily be converted into one between actual distance and risk.

For this reason, we instead model the dose inhaled at different distances. Exhaled breath is a likely source of infection.[21,22] Accordingly, we model a Gaussian plume[23] of virus-containing aerosols originating from the emitter's face at (0,0,0). The x axis represents the direction that the transmitter is facing and breathing toward with breath velocity U (m/s). Diffusion causes spread away from y=0 or z=0. The viral concentration is then

$$C(x, y, z) = \frac{Q}{U} \frac{1}{2\pi\sigma_y\sigma_z} e^{\frac{-y^2}{2\sigma_y^2}} e^{\frac{-z^2}{2\sigma_z^2}} \tag{1}$$

$$Q = SX \tag{1.1}$$

$$U = X/A \tag{1.2}$$

where Q is virus emitted per second and is equal to the product of shedding rate, S, (in arbitrary units proportional to copies/m³) and an exhalation rate, X, (taken from measured inhalation rates in m³/s), yielding arbitrary units proportional to copies per second being generated (eq 1.1). We sample our exhalation rates from a normal distribution of inhalation rates with a mean and standard deviation of 16.3 and 4.15 m³/day, respectively. These were informed by the 16-21 year old range from Table 6-1 in the Exposure Factors Handbook (2011).[10] To avoid negative exhalation rates, this distribution was left-truncated at 9 m³/day, the smallest fifth percentile of inhalation rates for males and females in age ranges overlapping with the 16-21 year old range.[10] The velocity of breath U (m/s) was determined by dividing the exhalation rate (m³/s) by the cross-sectional area of an open mouth A (m²), which is the area over which air is assumed to be exhaled at the plume source. The cross-sectional area was informed by a uniform distribution with minimum and maximum cross-sectional areas measured for an open mouth with a "large bite" configuration, ranging from 23 to 59 cm².[11] Note that for a steady-state plume assuming a continuous output of virus, the effects of the exhalation rate (volume of air per second) on amount of virus emitted, and on the velocity with which they disperse, cancel out. For an abrupt exhalation such as a cough, rather than steady state, a higher exhalation rate would affect viral airborne concentration.

For interactions ≤ 1m, we assumed two people interacting are directly in front of each other along the x-axis ($\varphi = \pi$ /2, θ=0). For interactions beyond the close range (>1m), we sample θ from a uniform 360 degrees (min=0, max=2π), and the angle between the z axis and the xy-plane, φ, was randomly sampled from a triangular distribution (min=π/4, mode=π/2, max=3π/4). We then convert from spherical units to (x,y,z) to apply Eq. 1. We assumed that scenarios where the person exposed was behind the emitter (x<0) resulted in a zero dose.

To capture the shape of the plume, we use:

$$\sigma_y = I_y x \tag{2}$$

$$\sigma_z = I_z x \tag{3}$$

Assuming moderately stable conditions, I_y and I_z were randomly sampled from uniform distributions with minimums and maximums of 0.08-0.25 and 0.03-0.07, respectively.[12]

We note that inhalation and exhalation rates are both likely important to risk. For example, one infected dance
instructor spread COVID-19 to 7/26 other instructors at a four hour workshop,[24] representing a similar risk as for
household contacts, despite the presumption that most were at >2 m. distance for most of this time. Limited air
circulation or increased respiratory rates are important factors that cannot be captured in the current GAEN
approach, but the four-hour duration of the workshop could be, combined with considerable uncertainty in the
relationship between Bluetooth attenuation and distance, in order to correctly capture the high risk of such a
scenario.
152
While wind velocity and relative humidity are important factors for determining droplet and fine aerosol dispersion
and deposition,[25,26] as is mask usage, these are uncertain factors that are not recorded by the app, especially
considering that interactions may occur indoors or outdoors. By not accounting for deposition, and by assuming that
masks are either not worn or not worn effectively, we will tend to overestimate dose at greater distances, and in the
presence of masks. This will implicitly lower the app-imposed risk tolerance of individuals who comply with public
health guidelines that recommend masks and physical distancing, and who might therefore also be more inclined to
comply with quarantine recommendations. The 2-meter rule was based on the assumption that most transmission is
via droplets (large aerosols) for which deposition occurs over this distance. However, there is increasing evidence
for transmission via smaller aerosols,[27–31] supporting our assignment of some risk to greater distances, reflecting
short- to medium-distance airborne transmission.
163

164 **3.0 Inhalation Dose per Interaction**

165
166 An inhaled dose of viral particles due to person-to-person interactions was estimated based on the duration of the
167 interaction (minutes) (T), the concentration of virus in the air at this {x,y,z} coordinate during the interaction
168 (arbitrary units of viral particles/m^3) $C(x, y, z)$, and inhalation rates (m^3/minute) (I),
169
170 $$D = T \cdot I \cdot C(x, y, z) \qquad (4)$$
171
172 Inhalation rates were randomly sampled from the same distribution as exhalation rates but allowing for a different
173 value per iteration. As with exhalation rates, we left-truncated the distribution to avoid negative inhalation rates and
174 therefore negative doses. Figure 2 shows the expected dose as a function of distance, with a discontinuity at 1m.
175 arising from our assumption that this distance or below indicates face-to-face interaction.

176 **4.0 Setting attenuation bin thresholds and corresponding weights**

177
178 To select the threshold values (a, b) demarcating 3 attenuation bins, we optimized the differences in mean dose
179 between two randomly sampled attenuation measurements. Specifically, we maximized the value of
180
181 $$d(a,b) = \sqrt{2p_A p_B (A - B)^2 + 2p_B p_C (B - C)^2 + 2p_A p_C (C - A)^2}$$
182
183 where A, B and C are the average doses D from Eq. 4, averaged across Monte Carlo sampling described above,
184 corresponding to bins [0, a], (a, b], and (b, +), and p_A, p_B, and p_C are the probabilities that an attenuation will fall
185 within that bin in our pseudo dataset.
186
187 We examined multiple local maxima of this distance measure before choosing a partition pair. We also investigated
188 alternative versions of a distance metric and alternative rebalancing schemes, to confirm that this is a relatively
189 robust partition pair.

5.0 Probability of Current or Future Infectiousness

To calculate residual risk of infection as a function of initial risk plus time since exposure, we use the probability distribution of incubation periods from Lauer et al.[7], available at https://iddynamics.jhsph.edu/apps/shiny/activemonitr/. Note that it is possible that incubation periods are even more dispersed than reported here[32]; this would lengthen quarantine recommendations.

To calculate risk of current or future infectiousness, we assume a fraction of symptomatic vs. asymptomatic cases and take an average of the discount factors applying in each case. Across a population, 20% of infections are estimated to be asymptomatic.[34] Younger users are more likely to be asymptomatic[33], so the fraction of asymptomatic cases could be personalized on the basis of user age if that information is collected on a voluntary basis. For the symptomatic cases, we discount according to the probability of subsequently developing symptoms, given that symptoms have not appeared yet.

For the asymptomatic cases, we combine the incubation periods from Lauer et al.[7] with a distribution of shedding durations. Long et al.[8] report slightly longer shedding durations for asymptomatic than symptomatic shedding, but other studies[34–38] for which we were unable to obtain the data, report the opposite, or no difference. Shedding declines in magnitude post symptom onset and is considered by the CDC to have reached negligible levels by 10 days post symptom onset. We assume that asymptomatic shedding begins 3 days before what would have been the day of symptom onset if symptomatic, or else immediately upon infection, whichever occurs later.

Using this assumption, we calculated the probability distribution of the day that shedding ends, given both the distribution of incubation periods and a distribution of shedding durations. For the latter, we combine the asymptomatic and symptomatic shedding durations of Long et al. but on the basis of CDC advice for isolation, we truncate the distribution so that all shedding periods longer than 12 days are recorded as exactly 12 days.

Note that low dose exposures, e.g. to asymptomatic individuals, may result in longer incubation periods,[32] suggesting that low initial risk scores should have longer rather than the shorter quarantines we calculate using this method. We currently ignore this by assuming that risk scores primarily capture uncertainty in the likelihood of infection with a minimal dose, and not variation in the infecting dose once above the minimal. To see how this assumption arises from our model, note that the exponential dose-response curve we use assumes that each virus has an independent probability of initiating infection. Under the resulting Poisson distribution for the number of viruses responsible for the initial infection, then even for the 30% infection rate of household contacts, the probability that infection is initiated with two or more viruses is only 5%, and with three or more viruses is only 0.6%. However, the higher variance in dose explored in Figure S2 could make initiation with multiple viruses common enough to matter for high infection probabilities. In this case, our simplifying assumption might require overly long quarantines following very high risk exposures. However, unless the variance is extreme, it might not significantly distort estimated probabilities among the range of lower risk exposures. We note that lognormal distributions of incubation periods with substantial variance occur even in the absence of variation in dose, due both to variance in within-host replication rate and to the stochastics of establishing infection in the first cells.[39]

6.0 Negative Test Results and Recommending a Day for Testing

Incorporation of negative test results can help exclude asymptomatic infection and hence allow for earlier release. From Bayes Theorem, and taking the false positive rate as negligible, a negative test result changes the probability of infection from p to $\frac{Ep}{(1-(1-E)p)}$, where E is the false negative rate. This could be taken as 0.3 [40,41] or made dependent on the timing of the test relative to exposure,[42] which would require distinct application to each of multiple exposures.

Kucirka et al.[42] report a false negative rate as a function of the timing of a PCR test relative to symptom onset, but
most of the data is post-symptom onset, with only a single patient's data informing false positive rates prior to
symptom onset. If sufficient data on false negative rates pre-symptom onset became available on a larger number of
patients, it could be combined with the distribution of incubation periods to calculate the false negative rate as a
function of time since exposure. Before use as E in Bayes Theorem above, it can be adjusted in a form of
survivorship analysis; if testing is conditional on no symptoms to date, then the ratio of pre-symptomatic :
asymptomatic individuals in the reference group will depend on time since exposure. Time-dependent false negative
rates could then be used to recommend an optimal day for testing. For example, we could choose the days on which
the probability of obtaining positive test results is highest.
247
We calculated this probability as the sensitivity on that day × the conditional probability of infection on that day
given no symptoms to date. Given exposures on multiple days, the sum of this product can be computed to choose a
date for testing. Alternative criteria for optimization could also be devised, e.g. explicitly prioritizing highly exposed
individuals whose calculated quarantine in the absence of a test would exceed 14 days, or assigning a value to
testing immediately after quarantine begins, in the service of rapid tracing of that individual's contacts should they
test positive.

7.0 Multiple Exposures and Total Risk

255
256 GAEN v1.1 will not record either a single long exposure or multiple exposures from a single individual beyond the
257 30 minutes cap for each of three attenuation bins, although this has changed in v1.5. However, it will record
258 exposures happening on different days (considered to change at midnight UTC) as independent exposures. To
259 calculate total risk, we combine the probabilities p_i of each exposure i, each discounted as described in the section
260 above, as $1 - \prod_i (1 - p_i)$.
261
Figures 3 and 4 illustrate scenarios of a single exposure. When there are multiple exposures, quarantine durations are
determined with respect to total risk. The risk threshold for initiation and completion of quarantine are the same. In
other words, risk is treated in an internally consistent fashion to achieve the maximum possible benefit from a given
number of recommended quarantine days across a population. When fixed quarantine durations are used, exposure
must be significant on a single day, from which the 14 days are then calculated, and risks are not integrated across
multiple days.

8.0 Variance in Dose

269
270 Our dose response curve is actually the probability of infection as a function of the expectation of dose, rather than
271 dose. To consider the effect of variance in dose, we note that while $E\left[1 - e^{-\lambda D}\right]$ is analytically intractable with
272 respect to a lognormally distributed dose, there is a viable saddle point approximation described by Rojas-
273 Nandayapa (2008) While Eq. 5.6 in Rojas-Nandayapa contains a typo, we use the method given to derive
274
275 $$1 - E\left[e^{-\lambda D}\right] = 1 - \left[\frac{1}{\sqrt{1 + LW(\theta \sigma^2 e^\mu)}} exp\left(\frac{LW^2(\theta \sigma^2 e^\mu) + 2LW(\theta \sigma^2 e^\mu)}{2\sigma^2}\right)\right], \qquad (6)$$

276
277 where $LW(x)$ is the Lambert-W function.
278
Using this approximation, or the associated importance sampling method (Asmussen et al., 2016) which yields
similar results, we can compare the shape for the dose response curves and consider whether an "effective" λ will
perform acceptably. Ct counts in Long et al. (2020) have a standard deviation ~4, representing an upper bound of a
16-fold difference in viral load, because a difference of 1 Ct in PCR measures represents at most a 2-fold difference
in underlying viral load, and because individuals are not sampled at exactly comparable times with respect to the
timecourse of shedding. However, variance in dose from causes other than infectiousness, e.g. the intimacy of

contact, is not included. With a standard deviation representing a 16-fold difference, we find that by using a value
$3/10^{th}$ of $\lambda = 3.50\,E-06$, we can super-impose the two curves up to 20% infection probability (Figure S2).
287

288
289 Figure S2. Although our dose-response curve takes the function of an expectation, for low infection probabilities the
290 effect of this is to change the interpretation of the value of λ, which is 3.50 x 10^{-6} when using Eq. 5 but 0.3 times this
291 value when using Eq.6 with a distribution of log-dose with standard deviation corresponding to a 16-fold difference
292 in dose.

293 **9.0 Transmission Risk Levels**

294
295 Transmission risk level in the GAEN API is a proxy for the magnitude of viral shedding and can be set by the app
296 on the basis of a simple questionnaire administered to users reporting a positive diagnosis. We use this to inform
297 expected shedding S.
298
The first question we ask to inform transmission risk levels is, "What day did your symptoms start"? A curve fit to
known transmission events suggests peak transmission the day before symptom onset, and essentially no
transmission more than three days before symptom onset.[43] However, this might be confounded with behavioral
changes with symptom onset, with ascertainment bias in terms of which transmission events are easiest to document,
with onset of mild symptoms being overlooked and the later onset of more severe symptoms instead being reported,
with possible errors in the direction of transmission given the variability of incubation periods, and with assumptions
regarding the shape of the curve. Worse, the analysis appears to have significant technical flaws.[4]
306
307 A second source of information is quantitative polymerase chain reaction (PCR). However, this may reflect non-
308 infectious viral remnants, especially late in the course of disease, where the proportion of culture-positive PCR
309 results tends to decrease.[1,2] However, this decline is also expected from a simple dose-response curve, where the
310 probability of culture-positivity decreases as the amount of shedding decreases late in infection, i.e. the decline in
311 infectivity might be quantitative rather than qualitative.
312
While informed by the two sources of data above, we focus the most on data in which virus was successfully
cultured from patient samples, as a clear indication of infectivity. Arons et al. [2] took prospective samples throughout
a nursing home, and were able to culture virus from six days before symptom onset until nine days after symptom
onset, with little quantitative trend in shedding rate conditional on a positive test. In hospitalized patients, Wölfel et
al. [3] were unable to isolate live virus from cultures more than 8 days post symptom onset, despite PCR evidence of
high shedding. In one case report, live virus has been isolated 18 days after symptom onset, but this seems to be an
outlier.[44] Bullard et al. [1] quantified both TCID50 and PCR for 7 days post symptom onset, and saw an approximately
10-fold decline in infectious dose. We note that culture methods may not be sensitive enough to capture low
concentrations.[2] We nevertheless use this TCID50 data as motivation in assuming a 10-fold dynamic range in

shedding rates as a function of day relative to symptom onset. More studies measuring infectivity in a quantitative
manner are needed, particularly in populations that represent a broader base of cases of varying ages and health
status. Encouragingly, our settings based on culture data agreed with those from epidemiological modeling after
corrections to the latter were made.[4] We slightly tweaked our settings for pre-symptomatic Transmission Risk
Levels to agree with later epidemiological work.[45]
327
328 A final source of information comes from detailed Taiwanese contact tracing[46], who found a 1.0% symptomatic
329 attack rate (95% CI 0.6-1.6%) for those exposed within five days of symptom onset, and 0% (95% CI 0–0.4%) for
330 those exposed after. Risk from exclusively pre-symptomatic exposure was 0.7% (95% CI 0.2%-2.4%). German
331 contact tracing also points to highest transmission risk around the time of symptom onset.[47]
332
We use 6 of the 8 Transmission Risk Levels in the GAEN API to capture this range, evenly spaced on a log scale
between 10 and 100 in arbitrary units, reserving the use of levels 7 and 8 for individuals for testing purposes and any
future functionality. Transmission risk levels could also be manipulated for testing purposes, e.g. to help learn, if
individuals voluntarily enter exposure details into a manual contact tracing database, how transmission risk varies in
the real world, rather than just in TCID50 studies, as a function of a symptomatic status and time. We note that
Transmission Risk Levels have been deprecated in the announced GAEN v1.5, and replaced with only two possible
levels of infectiousness in v1.6. While the functionality of v1.1 should be preserved, we note that usage of
Transmission Risk Levels would need to be standardized across different apps in order to ensure interoperability. In
the description below, we outline not just our use of v1.1, but also the use we would make of six levels of
infectiousness if they were to again be made available in a future API version. The assignments we use in this
framework can of course be improved as more data become available. Our call for more than two levels of
infectiousness comes from the fact that a systematic 10-fold difference in TCID50 has been observed, and that such
a large difference seems to warrant more levels.
346
347 Based on a holistic reading of the four sources of evidence described above, we assign the maximum level of 6 from
348 one day pre-symptom onset to two days post-symptom onset. Five days before symptom onset we assign level 1,
349 four days before we assign level 3, three days before level 4, and two days before level 5. Three days after symptom
350 onset we assign level 5, four days after level 4, five days after level 3, 6-7 days after level 2, and 8-9 days after level
351 1. Our termination at 9 days is based on current CDC guidance.[9]
352
For users who report a positive test but no symptoms, there is likely a reason they were tested, and so we ask for the
most likely day of exposure, if known. If provided, we assume that shedding did not begin until two days after
exposure, at the earliest. We also ask for the date of sampling for the positive test (which can be reported by the
healthcare provider rather than the app user) and assume peak shedding at around this time. Subject to the constraint
from day of exposure, we assign Transmission Risk Level 3 to dates within with one day of the test, and level 2 to
dates between 2 and 4 days of the test, although as discussed below, one study reports substantially longer shedding
than these 9 days.[8] There is some evidence that viral shedding is lower in asymptomatic vs. symptomatic cases,[48]
while another study indicates the shedding magnitudes may be similar.[49] Note that we assume that those with no
symptoms at the time they receive a positive test result are asymptomatic rather than pre-symptomatic – should test
turnaround times be sufficiently fast, it would be useful for users to be able to report symptom onset after the fact
and trigger a change to previously reported Transmission Risk Levels, and we recommend that this functionality be
added, together with restoring a greater number of Transmission Risk Levels, in future versions of the GAEN
framework.

10.0 Considerations in recommending and messaging variable quarantine durations
367
The need for consistent guidance to the public is an important consideration for implementing tailored risk scoring
and modified quarantine recommendations. If for the sake of a consistency, a public health authority is not willing to
authorize variable quarantine recommendations, as is currently the case in Arizona, but only 0 or 14 day quarantines
from time of the last individually significant exposure, then the threshold for going into quarantine at all would need

to become more strict in order to maintain the same overall risk among the population under quarantine. In other
words, retaining the same average probability of current or future infectiousness among the quarantined population
would require some exposed individuals to no longer go into quarantine at all, in addition to others lengthening their
quarantine out to 14 days. With a binary 0 or 14 day quarantine, the amount by which disease transmission is
prevented per day of quarantine will be lower.

Alternatively, to avoid mixed messaging regarding quarantine even while the app recommends quarantine of
variable duration, one option is to suppress all details about individual exposures from the user's view, including
their date. This has the additional advantage of decreasing the risk that users will be able to guess who exposed
them, further preserving privacy.

The app can communicate the risk of infectiousness either as a simple recommendation for which days to
quarantine, or also as a quantitative score in order to "game-ify" the process of quarantine and give users positive
feedback for each day they succeed in remaining at home until risk falls to a lower level. Further research is needed
to assess the most effective messaging strategies. E.g., the app could display both current and projected risk of
infectiousness on a simple scale of 1 to 10, so users can see how that risk will fall with each day of quarantine. This
visualization might change perceptions. E.g., an individual who wants to comply with a 14-day quarantine, but does
not feel able to, might rush out to get groceries before starting their quarantine in earnest, while shedding virus pre-
symptomatically. Visualizing projected risk into the future would then give the message that if the exposed
individual can only make do for one more day before leaving the home for essentials, that will help, because if they
do not develop symptoms, their risk will be lower even after a single day longer. Risk communication in an app
could focus on day to day coaxing of this form.

Conflicting messages can still arise if manual contact tracers trace an individual who also received an exposure
notification. In this case, it is likely that the two recommend different end dates for quarantine. While this is to be
expected from our procedure for recommending variable quarantine durations, we note that even if the app were to
issue 14 day quarantine recommendations only, it could still arise because the individual has been exposed more
than once, on different days, and the manual contact tracer is following up an infected individual who may not have
used the app. Until there is reliable data on app performance, we recommend that the manual contact tracer's
protocol should override whatever the app says. Should the app turn out to perform well, an alternative procedure
might eventually be to go with whichever protocol recommends the longer quarantine. An intermediate possibility is
for the manual contact tracer to ask for exposure notification details, to determine whether it may be a different
exposure to the one being manually traced. There may also be conflicts in protocols for the timing of testing.

Note that with symptom onset sometimes as early as two days after exposure, and given the possibility of pre-
symptomatic shedding, and the possibility of confusion regarding who infected whom, we currently ignore the
possibility that shedding might not yet have begun. Current testing turnaround times are mostly long, making this
reasonable. However, if same-day tests become more widely available, our approach could be extended to directly
communicate the risk of current infectiousness, rather than as is currently the case, the risk of current or future
infectiousness. A significantly lower risk of infectiousness will be present on the day of exposure and perhaps also
the day after. Delays in going into public to prepare for a long quarantine could inadvertently lead to pushing
individuals past the latent period before they go into public; displaying a full projected timeline of the projected risk
of infectiousness could avert this, at the risk of significantly more complex messaging than "stay home until Friday".

We currently consider the harm from the release of any infectious individual to be equivalent to that from the release
of any other. Extensions of our approach could take into account greater harm from someone never quarantined and
hence shedding for the full duration relative to someone released prematurely partway through asymptomatic or pre-
symptomatic shedding, for whom a portion of the harm has already been mitigated. This would lead to steeper
discounting and might also prevent the app from occasionally recommending quarantines of less than 5 days when
initial risk is already very near the threshold.

References

Bullard J, Dust K, Funk D, *et al.* Predicting infectious SARS-CoV-2 from diagnostic samples. *Clin Infect Dis* 2020; : 1–18.

Arons MM, Hatfield KM, Reddy SC, *et al.* Presymptomatic SARS-CoV-2 infections and transmission in a skilled nursing facility. *N Engl J Med* 2020; **382**: 2081–90.

Wölfel R, Corman VM, Guggemos W, *et al.* Virological assessment of hospitalized patients with COVID-2019. *Nature* 2020; **581**: 465–9.

Ashcroft P, Huisman JS, Lehtinen S, *et al.* COVID-19 infectivity profile correction. *arXiv* 2020. http://arxiv.org/abs/2007.06602.

Curmei M, Ilyas A, Evans O, Steinhardt J. Estimating Household Transmission of SARS-CoV-2. *medRxiv* 2020. DOI:10.1101/2020.05.23.20111559.

Buitrago-Garcia DC, Egli-Gany D, Counotte MJ, *et al.* Asymptomatic SARS-CoV-2 infections: a living systematic review and meta-analysis, version 3. *medRxiv* 2020. DOI:10.1101/2020.04.25.20079103.

Lauer SA, Grantz KH, Bi Q, *et al.* The incubation period of coronavirus disease 2019 (CoVID-19) from publicly reported confirmed cases: Estimation and application. *Ann Intern Med* 2020; **172**: 577–82.

Long Q, Tang X, Shi Q, *et al.* Clinical and immunological assessment of asymptomatic SARS-CoV-2 infections. *Nat Med* 2020. DOI:10.1038/s41591-020-0965-6.

Centers for Disease Control and Prevention. Duration of Isolation & Precautions for Adults. 2020. https://www.cdc.gov/coronavirus/2019-ncov/hcp/duration-isolation.html#cecommendations (accessed July 27, 2020).

U.S. Environmental Protection Agency. Exposure Factors Handbook 2011 Edition (EPA/600/R-09/052F). Washington, DC, 2011 https://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=236252.

Leckie JO, Naylor KA, Canales RA, *et al.* Quantifying Children's Microlevel Activity Data from Existing Videotapes" by Exposure Research Group at Stanford University for the U.S. Environmental Protection Agency. 2000 https://cfpub.epa.gov/ols/catalog/advanced_full_record.cfm?&FIELD1=AUTHOR&INPUT1=FERGUSON AND C. AND R.&TYPE1=ALL&LOGIC1=AND&COLL=&SORT_TYPE=MTIC&item_count=5.

Western Engineering. Self-study notes - GAUSSIAN PLUMES. https://www.eng.uwo.ca/people/esavory/Gaussian plumes.pdf (accessed June 7, 2020).

Zhang N, Su B, Chan PT, Miao T, Wang P, Li Y. Infection spread and high-resolution detection of close contact behaviors. *Int J Environ Res Public Health* 2020; **17**. DOI:10.3390/ijerph17041445.

Farrell S, Leith DJ. Pairwise Handset Types and Orientations Are Sufficient to Blur Exposure Notification Thresholds. Trinity Coll. Dublin, Irel. 2020. https://down.dsg.cs.tcd.ie/tact/posorient.pdf (accessed Aug 6, 2020).

Jain S, Islam T, Chowdhury B, Chen Y, Son Y. Agent-based simulation to evaluate various entrance and exit policies and contact-caused risks in classrooms under a pandemic situation. Univ. Arizona. .

Guo S, Yu J, Shi X, *et al.* Droplet-transmitted infection risk ranking based on close proximity interaction. *Front Neurorobot* 2020; **13**: 1–11.

Setti L, Passarini F, De Gennaro G, *et al.* Airborne transmission route of covid-19: Why 2 meters/6 feet of inter-personal distance could not be enough. *Int J Environ Res Public Health* 2020; **17**. DOI:10.3390/ijerph17082932.

Chu DK, Akl EA, Duda S, *et al.* Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *Lancet* 2020; : 1973–87.

Rea E, Laflèche J, Stalker S, *et al.* Duration and distance of exposure are important predictors of transmission among community contacts of Ontario SARS cases. *Epidemiol Infect* 2007; **135**: 914–21.

Salathé M, Kazandjieva M, Lee JW, Levis P, Feldman MW, Jones JH. A high-resolution human contact network for infectious disease transmission. *Proc Natl Acad Sci U S A* 2010; **107**: 22020–5.

Ma J, Qi X, Chen H, *et al.* Exhaled breath is a significant source of SARS-CoV-2 emission. *medRxiv* 2020; : 2020.05.31.20115154.

Chen W, Zhang N, Wei J, Yen HL, Li Y. Short-range airborne route dominates exposure of respiratory infection during close contact. *Build Environ* 2020; **176**: 106859.

Brusca S, Famoso F, Lanzafame R, Mauro S, Garrano AMC, Monforte P. Theoretical and Experimental

476   Study of Gaussian Plume Model in Small Scale System. *Energy Procedia* 2016; **101**: 58–65.

477   24   Jang S, Han SH, Rhee J-Y. Cluster of Coronavirus Disease Associated with Fitness Dance Classes, South
478   Korea. *Emerg Infect Dis* 2020; **26**. DOI:10.3201/eid2608.200633.

479   25   Yang W, Elankumaran S, Marr LC. Concentrations and size distributions of airborne influenza A viruses
480   measured indoors at a health centre, a day-care centre and on aeroplanes. *J R Soc Interface* 2011; **8**: 1176–
481   84.

482   26   Feng Y, Marchal T, Sperry T, Yi H. Influence of wind and relative humidity on the social distancing
483   effectiveness to prevent COVID-19 airborne transmission : A numerical study. *J Aerosol Sci* 2020; **147**:
484   105585.

485   27   Qureshi Z, Jones N, Temple R, Larwood JP, Greenhalgh T, Bourouiba L. What is the evidence to support
486   the 2-metre social distancing rule to reduce COVID-19 transmission? CEBM. 2020.
487   https://www.cebm.net/covid-19/what-is-the-evidence-to-support-the-2-metre-social-distancing-rule-to-
488   reduce-covid-19-transmission/ (accessed July 15, 2020).

489   28   Prather KA, Wang CC, Schooley RT. Reducing transmission of SARS-CoV-2. *Science (80-)* 2020; **368**.

490   29   Lednicky JA, Lauzardo M, Fan ZH, *et al.* Viable SARS-CoV-2 in the air of a hospital room with COVID-19
491   patients. *medRxiv* 2020. DOI:10.1101/2020.08.03.20167395.

492   30   Fennelly KP. Particle sizes of infectious aerosols: implications for infection control. *Lancet Respir Med*
493   2020; **2600**: 1–11.

494   31   Jones NR, Qureshi ZU, Temple RJ, Larwood JPJ, Greenhalgh T, Bourouiba L. Two metres or one: what is
495   the evidence for physical distancing in covid-19? *BMJ* 2020; **370**: m3223.

496   32   Wei Y, Wei L, Liu Y, Huang L, Shen S, Zhang R. A systematic review and meta-analysis reveals long and
497   dispersive incubation period of COVID-19. *medRxiv* 2020.

498   33   Davies NG, Klepac P, Liu Y, Prem K, Jit M, Eggo RM. Age-dependent effects in the transmission and
499   control of COVID-19 epidemics. *medRxiv* 2020; : 2020.03.24.20043018.

500   34   Chen X, Zhang Y, Zhu B, *et al.* Associations of Clinical Characteristics and Antiviral Drugs with Viral
501   RNA Clearance in Patients with COVID-19 in Guangzhou, China. *medRxiv* 2020.
502   DOI:10.2139/ssrn.3566234.

503   35   Chau NVV, Thanh Lam V, Thanh Dung N, *et al.* The natural history and transmission potential of
504   asymptomatic SARS-CoV-2 infection. *Clin Infect Dis* 2020; **0813**: 1–30.

505   36   Yang R, Gui X, Xiong Y. Comparison of Clinical Characteristics of Patients with Asymptomatic vs
506   Symptomatic Coronavirus Disease 2019 in Wuhan, China. *JAMA Netw open* 2020; **3**: e2010182.

507   37   Xiao T, Wang Y, Yuan J, *et al.* Early viral clearance and antibody kinetics of COVID-19 among
508   asymptomatic carriers. *medRxiv* 2020. DOI:10.1101/2020.04.28.20083139.

509   38   Hu Z, Song C, Xu C, *et al.* Clinical characteristics of 24 asymptomatic infections with COVID-19 screened
510   among close contacts in Nanjing, China. *Sci China Life Sci* 2020; **63**: 706–11.

511   39   Ottino-Loffler B, Scott JG, Strogatz SH. Evolutionary dynamics of incubation periods. *Elife* 2017; **6**.
512   DOI:10.7554/eLife.30212.

513   40   Yang Y, Yang M, Shen C, *et al.* Evaluating the accuracy of different respiratory specimens in the laboratory
514   diagnosis and monitoring the viral shedding of 2019-nCoV infections. *medRxiv* 2020; :
515   2020.02.11.20021493.

516   41   Ai T, Yang Z, Hou H, *et al.* Correlation of Chest CT and RT-PCR Testing in Coronavirus Disease 2019
517   (COVID-19) in China: A Report of 1014 Cases. *Radiology* 2020; **2019**: 200642.

518   42   Kucirka LM, Lauer SA, Laeyendecker O, Boon D, Lessler J. Variation in False-Negative Rate of Reverse
519   Transcriptase Polymerase Chain Reaction–Based SARS-CoV-2 Tests by Time Since Exposure. *Ann Intern*
520   *Med* 2020. DOI:10.7326/m20-1495.

521   43   He X, Lau EHY, Wu P, *et al.* Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat*
522   *Med* 2020; **26**: 672–5.

523   44   Liu W Da, Chang SY, Wang JT, *et al.* Prolonged virus shedding even after seroconversion in a patient with
524   COVID-19. *J Infect* 2020. DOI:10.1016/j.jinf.2020.03.063.

525   45   Ferretti L, Ledda A, Wymant C, *et al.* The timing of COVID-19 transmission. *medRxiv* 2020.
526   DOI:10.1101/2020.09.04.20188516.

527   46   Cheng HY, Jian SW, Liu DP, Ng TC, Huang WT, Lin HH. Contact Tracing Assessment of COVID-19
528   Transmission Dynamics in Taiwan and Risk at Different Exposure Periods before and after Symptom Onset.
529   *JAMA Intern Med* 2020. DOI:10.1001/jamainternmed.2020.2020.

47   Böhmer MM, Buchholz U, Corman VM, *et al.* Investigation of a COVID-19 outbreak in Germany resulting

from a single travel-associated primary case: a case series. *Lancet Infect Dis* 2020; **3099**.
DOI:10.1016/S1473-3099(20)30314-5.
48   McDonald J. Unpacking WHO's Asymptomatic COVID-19 Transmission Comments. FactCheck.org. 2020.
https://www.factcheck.org/2020/06/unpacking-whos-asymptomatic-covid-19-transmission-comments/
(accessed June 16, 2020).
49   Lavezzo E, Franchin E, Ciavarella C, *et al.* Suppression of a SARS-CoV-2 outbreak in the Italian
municipality of Vo'. *Nature* 2020; published online June 30. DOI:10.1038/s41586-020-2488-1.
538