

Title

High Rate of SARS-CoV-2 Transmission due to Choir Practice in France at the Beginning of the COVID-19 Pandemic

Nathalie CHARLOTTE, MD, **SSR Val-Rosay**, 41 chemin Ferrand 69370 Saint-Didier-au-Mont-d'Or, France : *email* : nathalie.charlotte@ugecam.assurance-maladie.fr

Abstract

Background: There has been very little focus on the individual risk of acquiring COVID-19 related to choir practice. Methods: We report the case of a high transmission rate of SARS-CoV-2 linked to an indoor choir rehearsal in France in March 2020 at the beginning of the COVID-19 pandemic. Results: A total of 27 participants, including 25 male singers, a conductor and an accompanist attended a choir practice on March 12, 2020. The practice was indoor and took place in a non-ventilated space of 45 m². The mean age of the participants was 66.9 (range 35-86) years. 70% of the participants (19 of 27) were diagnosed with COVID-19 from 1 to 12 days after the rehearsal with a median of 5.1 days. 36% of the cases needed a hospitalization (7/19), and 21% (4/19) were admitted to an ICU. The index cases were possibly multiple. Discussion: The choir practice was planned in March 2020 at a period when the number of new cases of COVID-19 began to grow exponentially in France because SARS-CoV-2 was actively circulating. The secondary rate attack (70%) was much higher than it is described within households (10-20%) and among close contacts made outside households (0-5%). Singing might have contributed to enhance SARS-CoV-2 person-to-person transmission through emission of droplets and aerosolization in a closed non ventilated space with a relative high number of individuals. Conclusions: Indoor choir practice should be suspended during SARS-CoV-2 outbreaks. Further studies are necessary to test the spread of the virus by the act of singing. As the benefits of the barrier measures and social distancing are known to be effective in terms of a reduction in the incidence of the COVID-19, experts' recommendations concerning the resuming of choir practice are necessary.

Introduction

World Health Organization (WHO) declared the COVID-19 outbreak a pandemic on March 11, 2020¹. In response, all countries have implemented large scale public health and social measures in an attempt to reduce the transmission of SARS-CoV-2 and minimize the impact of the epidemic. Some social activities as fitness practice², night-clubbing³, Zumba practice², or working in open space offices⁴ have been described as super spreader events for COVID-19. Approximately 4.5% of the European population actively participate in collective singing activities⁵. There has been very little focus on the individual risk of acquiring COVID-19 related to choir practice. We report the case of a high transmission rate of SARS-CoV-2 linked to an indoor choir rehearsal in France at the beginning of the pandemic.

Methods – case analysis

The data were obtained by the collection of anonymous answers to a questionnaire. The participants were the persons who attended the choir practice. The president and the conductor of the choir were interviewed by telephone. The president of the choir collected all the questionnaires filled by the participants. The COVID-19 diagnosis was established by the general practitioners of the attendees or by hospital physicians for those who were hospitalized.

On March 12, 2020, 27 men (25 singers, 1 accompanist, 1 conductor) attended a choir 2-hours singing practice in France. The mean age of the participants was 66.9 (range 35-86) years. It took place inside a 45 m² and 3 m high room (mean surface available per person of 1.6 m²). The room wasn't ventilated. The members were sitting less close to one another than usual, but at a distance of less than 6 feet. Socialization acts were avoided during the practice (no shaking hands, no food sharing and no pauses). The male choir had a previous rehearsal on March 2. No choir member reported having been symptomatic for COVID-19 between March 2 and March 12. The individuals were split into 2 groups of 11 tenors and 14 basses. The mean age of the tenors was 66.9 (range 35-86) years. The mean age of the basses was 69.1 (range 48-84) years. The conductor was standing in front of the choir, the accompanist was near the tenors. (Figure 1).

Nineteen COVID-19 cases were identified among the members of the ensemble (Figure 2). A confirmed case was defined as an individual who has tested positive for SARS-CoV-2 by RT-PCR test performed on a nasopharyngeal sample and/or a case presenting a severe COVID-19 requiring hospitalization. A probable case was defined as a person presenting an acute respiratory illness relating to COVID-19 by a general practitioner 1 to 14 days following the rehearsal without having been evaluated with an RT-PCR test. Seven confirmed cases and 12 probable cases were identified. The dates of onset of symptoms among the cases ranged from 13/03 to 22/03 i.e. 1 to 10 days after the practice (median = 5 days, range 1-13 days). The secondary attack rate (SAR) was 33% among the confirmed cases and the overall attack rate was 70%. The SAR was 72% (8/11) among the tenors, of 64% (9/14) among the basses, and of 100% (2/2) among the non-choristers (conductor and accompanist) who were standing in front of the singers. The index cases were possibly multiple. Two attendees had a close contact with a COVID-19 case within 5 days before the practice. One chorister (tenor) who tested positive for COVID-19 has had contacts with the members of another choir

during 2 choir practices in a nearby area on March 9 and March 11. Several of these choristers reported COVID-19 symptoms 7 days later.

The mean age of the individuals who developed a severe COVID-19 defined by the need for a hospitalization, was 70.5 years (range 53-78). Seven of the 19 persons who became ill were hospitalized (36%), whose 57% (4/7) which represents 21% (4/19) of all patients were treated in an intensive care unit. The mean age of those who developed a mild COVID-19 (64%) was 60.9 years, (range 35-77). The fatality rate was 0%. Thirteen tertiary cases were later reported, mostly household contacts.

Discussion

From end February to end March 2020, SARS-CoV-2 was actively circulating in Europe. Several clusters of COVID-19 cases linked to choir practice were reported by the general media. In Whir au Val (Haut-Rhin, France), after a rehearsal of the Chorale de Saint Martin on February 28 with 20 choristers, 69 secondary cases were later identified and 9 persons died⁶. The Amsterdam Gemengd Koor (130 choristers) gave a 3-hour concert on March 8 at the Concertgebouw of Amsterdam (The Netherlands). The occurrence of 102 secondary COVID-19 cases was reported, including 4 fatal cases⁷. In Germany, within 2 weeks after gathering for a rehearsal on March 9, thirty members of the Berlin Cathedral Choir's 80 singers tested positive for COVID-19, and a further 30 reported COVID-19 symptoms⁸.

In the USA, Skagit County Public Health (Washington) investigated a cluster of 53 persons among 61 choristers after a choir practice on March 10. Two of them died of COVID-19⁹.

In France, the number of new cases of COVID-19 began to grow exponentially in March¹⁰. In the absence of effective treatment for the disease, efforts to control COVID-19 pandemic have relied on personal preventive actions, physical distancing, stay-at-home orders, school closures, and workplace restrictions adopted at the national level. On March 9 the gathering of more than 1 000 people was forbidden. On March 12 the closure of all schools and universities was ordered. On March 13 those restrictions were extended to the gathering of more than 100 people. The stay-at-home order was effective on March 17 after more than a thousand new confirmed cases were reported within 24 hours¹⁰. On the date the rehearsal took place, people in France were not yet familiar with the barrier measures.

The incubation of SARS-CoV-2 is known to be 1 to 14 days with a median of 5.1 days¹¹. The March 12 practice is the likely point-source exposure event because the surge of COVID-19 cases occurred 5 days later. The individual whose onset of symptoms was March 13 (within 24 hours after the practice), might have been a pre-symptomatic or asymptomatic carrier of SARS-CoV-2 at the time of the rehearsal, which could explain the shortness of the delay between the date of the event and the onset of the illness^{12,13,14}. The basic reproduction number of SARS-CoV-2 is approximately of 2.2 to 2.5¹⁵. The estimated secondary attack rate of the virus (SAR) within households is 10–20%, with a much smaller SAR among close contacts made outside households, (0–5%)¹⁶. In comparison, we found a much higher SAR which shows that SARS-CoV-2 has had the potential to spread widely during the choir rehearsal under its peculiar environmental circumstances: indoor setting, narrow and non-ventilated space, shortened inter-individual distances. A recent study shows that the transmission of

SARS-CoV-2 is likely at much higher risk in enclosed spaces than outside¹⁷. In addition, some other high human density environments like workplaces have been reported as important risk sites for the spread of SARS-CoV-2 and potential sources of further transmission⁴. For that reason, at the on-going re-opening phase of the pandemic, a minimum surface area of 4 m² available per person is recommended by the French Work Ministry for people working in the same room.¹⁸

SARS-CoV-2 is transmitted between people by infectious droplets emitted in close proximity to the nose, mouth or eyes of a susceptible person¹⁹. A recent study emphasizes the role of airborne transmission for the spread of the virus²⁰. Smaller droplets are able to be propelled further than 3 to 6 feet and to remain airborne longer after certain respiratory emissions^{21,22}, particularly in poorly ventilated spaces²³. Singing might have contributed to enhance SARS-CoV-2 person-to-person transmission through emission of droplets and aerosolization in a closed space with a relative high number of individuals. WHO recently acknowledged that transmission of the virus by aerosols may have been responsible for reported outbreaks of COVID-19 in some closed settings, such as places of worship or places of work where people may be shouting, talking, or singing²⁴. Despite avoiding social relations during the practice, it might have been occasions of facilitating the transmission of the virus like speaking face to face with each other shortly but recurrently.

This investigation has a number of limitations. The choristers were not interviewed by the author. The clinical features of the COVID-19 presentation of the cases were not available, prohibiting a detailed description of symptoms. The index case-patients were not identified. A majority of patients were not tested for SARS-CoV-2, which might have led to under-ascertainment of infections, particularly for those who were pre-symptomatic or asymptomatic. At mid-March 2020, France was at the third stage of the epidemic with the SARS-CoV-2 virus actively circulating countrywide. Testing patients reporting mild symptoms wasn't recommended in this context by the French High Council for Public Health (Haut Conseil de la Santé Publique, HCSP) who prioritized the access to diagnostic tests to patients at risk of severe forms of Covid-19 and to healthcare workers²⁵.

Conclusion

At the end of winter 2020 in Europe, choir practice indoors was a super spreader event because of the active circulation of the virus in the population at this moment, the characteristics of the viral transmission via close human contacts probably enhanced by the act of singing, and the social gathering. Indoor choir practice should be suspended during SARS-CoV-2 outbreaks. Further studies are necessary to test the spread of the virus by the act of singing. As the benefits of the barrier measures and social distancing are known to be effective in terms of a reduction in the incidence of the COVID-19, the question of the resuming of choir practice is being raised: experts' recommendations are necessary and must be set out to provide operational guidelines to singers at an amateur as well as a professional level.

Acknowledgements

We sincerely thank the president of the choir and its conductor for providing all the data and agreeing to participate actively in this investigation.

References:

1 -WHO announces COVID-19 outbreak a pandemic

<https://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/news/news/2020/3/who-announces-covid-19-outbreak-a-pandemic>

2- Cluster of Coronavirus Disease Associated with Fitness Dance Classes, South Korea
Sukbin Jang, Si Hyun Han, and Ji-Young Rhee.

Author affiliations: Dankook University Hospital, Dankook University College of Medicine, Cheonan, South Korea.

https://wwwnc.cdc.gov/eid/article/26/8/20-0633_article

3-New virus cases spike on steady rise in club-linked infections, another looming cluster.
Yonhap, in The Korea Herald Published : May 27, 2020.

http://www.koreaherald.com/view.php?ud=20200527000242&ACE_SEARCH=1

4 - Park SY, Kim YM, Yi S, et al. Coronavirus disease outbreak in call center, South Korea.
Emerg Infect Dis 2020 April 23 (Epub ahead of print). 11. Lee SH, Son H, Peck KR. Can po

5- A pilot research project coordinated by the European Choral Association – Europa Cantat
in the frame of the multi-annual Cultural Cooperation Project “VOICE – Vision on Innovation
for Choral Music in Europe” with the support of the EU.

https://europeanchoralassociation.org/wp-content/uploads/2019/01/singingeurope_report.pdf

6 – La croix 24 avril 2020 : Wihr-au-Val, le village meurtri par le coronavirus

<https://www.la-croix.com/France/Wihr-Val-village-meurtri-coronavirus-2020-04-22-1201090593>

7- <https://www.trouw.nl/verdieping/die-ene-passion-die-wel-doorging-met-rampzalige-gevolgen~b4ced33e/>

8- German choirs silenced as singing branded virus risk

<http://www.rfi.fr/en/wires/20200527-german-choirs-silenced-singing-branded-virus-risk>

9- High SARS-CoV-2 Attack Rate Following Exposure at a Choir Practice —
Skagit County, Washington, March 2020

Lea Hamner Polly Dubbel; Ian Capron Andy Ross Amber Jordan Jaxon Lee Joanne Lynn
Amelia Ball Simranjit Narwal Sam Russell Dale Patrick Howard Leibrand. MMWR / May 15,
2020 / Vol. 69 / No. 19.

10- Santé Publique France.

<https://www.santepubliquefrance.fr/maladies-et-traumatismes/maladies-et-infections-respiratoires/infection-a-coronavirus/articles/infection-au-nouveau-coronavirus-sars-cov-2-covid-19-france-et-monde#block-244210>

11- The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported
Confirmed Cases: Estimation and Application. Stephen A. Lauer, MS, Kyra H. Grantz,
Qifang Bi, Forrest K. Jones, Qulu Zheng, MHS; Hannah R. Meredith, PhD; Andrew S.
Azman, PhD; Nicholas G. Reich, and Justin Lessler. Ann Intern Med. 2020;172:577-582.
doi:10.7326/M20-0504

12- Temporal dynamics in viral shedding and transmissibility of COVID-19 Xi He et al. Nature medicine. <https://doi.org/10.1038/s41591-020-0869-5>

13 – Presymptomatic SARS-CoV-2 Infections and Transmission in a Skilled Nursing Facility. M.M. Arons et al. April 24,2020, NEJM.org.
DOI: <https://doi.org/10.1056/NEJMoa2008457>

14- The contribution of asymptomatic SARS-CoV-2 infections to transmission - a model-based analysis of the Diamond Princess outbreak.

Authors: Jon C Emery, Timothy W Russell, Yang Liu, Joel Hellewell, Carl AB Pearson, CMMID 2019-nCoV working group, Gwenan M Knight, Rosalind M Eggo, Adam J Kucharski, Sebastian Funk, Stefan Flasche, Rein M G J Houben. medRxiv preprint doi: <https://doi.org/10.1101/2020.05.07.20093849>.

15 - Riou J, Althaus CL. Pattern of early human-to-human transmission of Wuhan 2019 novel coronavirus (2019-nCoV), December 2019 to January 2020. Euro Surveill 2020; 25: 2000058.

16 - Effectiveness of isolation, testing, contact tracing and physical distancing on reducing transmission of SARS-CoV-2 in different settings: a mathematical modelling study

Status: Paper accepted at journal | First online: 23-04-2020 | Last update: 26-05-2020.

Authors: Adam J Kucharski, Petra Klepac, Andrew JK Conlan, Stephen M Kissler, Maria Tang, Hannah Fry, Julia R Gog, W John Edmunds & CMMID nCov working group.
https://cmmid.github.io/topics/covid19/reports/bbc_contact_tracing.pdf

17- Closed environments facilitate secondary transmission of coronavirus disease 2019 (COVID-19). Hiroshi Nishiura, Hitoshi Oshitani, Tetsuro Kobayashi, Tomoya Saito, Tomimasa Sunagawa, M.D., Ph.D. Tamano Matsui, M.D., Ph.D., Takaji Wakita MHLW COVID-19.

medRxiv preprint doi: <https://doi.org/10.1101/2020.02.28.20029272>

18 - Ministère du travail. République française.
<https://travail-emploi.gouv.fr/IMG/pdf/protocole-national-de-deconfinement.pdf>

19- World Health Organization, “Novel Coronavirus – China” (2020). URL <https://www.who.int/csr/don/12-january-2020-novel-coronavirus-china/en/>.

20 - Identifying airborne transmission as the dominant route for the spread of COVID-19 Renyi Zhang, Yixin Li, Annie L. Zhang, Yuan Wang, and Mario J. Molina, Department of Atmospheric Sciences, Texas A&M University,
www.pnas.org/cgi/doi/10.1073/pnas.2009637117

21- The airborne lifetime of small speech droplets and their potential importance in SARS-CoV-2 transmission

Valentyn Stadnytskyia, Christina E. Baxb, Adriaan Baxa,, and Philip Anfinrud, Laboratory of Chemical Physics, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD 20892-0520; and bPerelman School of Medicine, University of Pennsylvania, Philadelphia.
www.pnas.org/cgi/doi/10.1073/pnas.2006874117
<https://doi.org/10.5281/zenodo.3770559>

22- It is Time to Address Airborne Transmission of COVID-19. Lidia Morawska, Donald K Milton. Published: 06 July 2020.

Clinical Infectious Diseases, ciaa939, <https://doi.org/10.1093/cid/ciaa939>

<https://academic.oup.com/cid/article/doi/10.1093/cid/ciaa939/5867798>

23- Small droplet aerosols in poorly ventilated spaces and SARS-CoV-2 transmission. Lancet Respir Med 2020 Published Online May 27, 2020

[https://doi.org/10.1016/S2213-2600\(20\)30245-9](https://doi.org/10.1016/S2213-2600(20)30245-9)

24- WHO- What we know about aerosol transmission ? July 9, 2020.

<https://www.who.int/news-room/q-a-detail/q-a-how-is-covid-19-transmitted?referringSource=articleShare>

<https://www.who.int/news-room/q-a-detail/q-a-how-is-covid-19-transmitted?referringSource=articleShare>

25- Avis provisoire HCSP : Patients à risque de formes sévères de COVID-19 et priorisation du recours aux tests de diagnostic virologique.10/03/2020

<https://www.hcsp.fr/Explore.cgi/avisrapportsdomaine?clefr=780>




Figure 1: Sketch showing the position of the attendees in the practice room on May 12, 2020.




Figure 2 : Incidence of secondary COVID-19 cases among the choir attendees