Learning the Mental Health Impact of COVID-19 in the United States with Explainable Artificial Intelligence

Indra Prakash Jha MCA[1,*], Raghav Awasthi M.Sc.[1,*], Ajit Kumar MBA[2,3], Vibhor Kumar Ph.D.[1,#], Tavpritesh Sethi Ph.D.[1,#]

1 Indraprastha Institute of Information Technology, Delhi (IIIT-D)
2 XLRI - Xavier School of Management - Jamshedpur
3 Adobe
corresponding authors
* authors contributed equally to this work

Manuscript word count: 2902

Corresponding author Address:

Tavpritesh Sethi -
Phone: 011 26907 533
Email: tavpriteshsethi@iiitd.ac.in
Office: A-309, (R&D Block) IIIT Delhi, Delhi India, 110020

Vibhor Kumar -
Phone: 011 26907440
Email: vibhor@iiitd.ac.in
Office: A-304 (R&D Block) IIT Delhi, Delhi India, 110020

Key Points: Mental Health, Covid-19, Bayesian Network, Machine Learning

Question: What factors could affect mental health, and how could this level be predicted using the same factors during the of COVID-19?

Finding: In this survey-data based study, multiple factors such as Social isolation, digital communication, working, and schooling from home, were identified as crucial factors of mental illness during Covid-19. Interestingly, behavioral changes such as wearing a mask and avoiding restaurants and public places were not found to be associated with mental health.

Meaning: Regular non-virtual communication with friends and family, healthy social life and social security are key factors and especially taking care of people with mental disease history is even more important.

Abstract:

Importance: COVID-19 pandemic has deeply affected the health, economic, and social fabric of nations. Identification of individual-level susceptibility factors may help people in identifying and managing their emotional, psychological, and social well-being.

Objective: This work is focused on learning a ranked list of factors that could indicate a predisposition to a mental disorder during the COVID pandemic.

Data Sources and Study design: In this study, We have used a survey of 17764 adults in the USA at different age groups, genders, and socioeconomic statuses.

Methods: Through initial statistical analysis followed by Bayesian Network inference, we have identified key factors affecting Mental health during the COVID pandemic. Integrating Bayesian networks with classical machine learning approaches lead to effective modeling of the level of mental health.

Results: Overall, females are more stressed than males, and people of age-group 18-29 are more vulnerable to anxiety than other age groups. Using the Bayesian Network Model, we found that people with chronic medical condition of mental illness are more prone to mental disorders during the COVID age. The new realities of working from home, home-schooling, and lack of communication with family/friends/neighbors induces mental pressure. Financial assistance from social security helps in reducing mental stress during COVID generated economic crises. Finally, using supervised ML models, we predicted the most mentally vulnerable people with ~80% accuracy.

Introduction:

After 7 months of initial reporting, the coronavirus pandemic continues to rage across the world. The mental health consequences of COVID-19 pandemic are profound. More than half a million lives and more than 400 million jobs have been lost[1] causing a considerable degree of fear, worry, and concern. These effects are seen in the population at large and may be pronounced among certain groups in particular, such as youth, frontline workers, caregivers, and people with chronic medical conditions. The new world order has introduced unprecedented interventions of country-wide lockdowns that are necessary to control the spread but have led to increased social isolation. Loneliness, depression, harmful alcohol, drug use, and self-harm or suicidal behavior are also expected to rise.

The Lancet Psychiatry[2] recently highlighted the needs of vulnerable groups during this time, including those with severe mental illness, learning difficulties, and neurodevelopmental disorders, as well as socially excluded groups such as prisoners, the homeless, and refugees. Calls to action highlighting the need for engaging more early-career psychiatrists [3,4] , technology such as Telepsychiatry, and highlighting the high susceptibility of frontline medical workers themselves[5] highlights the magnitude of the problem. Further, interventions are expected to have a gender-specific impact with women more likely to be exposed to additional stressors related to informal care, already existing economic disparity, and school closures. Similarly, age and comorbidity status may have a direct impact on susceptibility to mental health challenges due to their relationship with COVID-19 morbidity and mortality. Indeed, it has been established that emotional distress is ubiquitous in affected populations — a finding certain to be echoed in populations affected by the Covid-19 pandemic[6]. Finally, the role of social media is complex with some research indicating the association of social media exposure with a higher prevalence of mental health problems[7].

However, most of these effects have been studied in isolation with a lack of modeling the collective impact of such factors. This study addresses this gap through the use of Bayesian networks, an explainable artificial intelligence approach that captures the joint multivariate distribution underlying a large survey data collected across the United States. We also address the gap of prediction of vulnerability to mental health events such as anxiety attacks using supervised machine learning models.

Methods:

Datasets

We extracted data of 17764 adults (https://www.covid-impact.org/)[8] from weekly surveys of the U.S. adult household population nationwide for 18 regional areas including 10 states (CA, CO, FL, LA, MN, MO, MT1, NY, OR, TX) and 8 Metropolitan Statistical Areas (Atlanta, Baltimore, Birmingham, Chicago, Cleveland, Columbus, Phoenix, Pittsburgh). To summarize, the dataset comprised variables on physical health, mental health, Insurance related Policy, economic security, and social dynamics. Table-1 shows the socio-demographic characteristics of respondents participating in the survey. Further details are available in the original source[8].

Analysis

Figure 1(a) shows the flow diagram for the analyses conducted. The survey questions were classified into several types of indicators such as mental-health, work-from-home, communication, COVID-symptoms, chronic-medical-conditions, behavioral-aspects, insurance-assistance, and many others *(see . Table-2)*. We constructed a model for mental-health indicators, *soc5a* (Felt nervous, anxious*, or on edge), *soc5b* (Felt depressed), *soc5c* (Felt lonely), *soc5d* (Felt hopeless about the future), and *soc5e*(sweating, trouble breathing, pounding heart, etc, in the last seven days) as outcome variables. Hence we first evaluated the consistency in answers to mental-health questions using Item Reliability Analysis**.** A scale for reliability measure of internal consistency, **Cronbach's alpha**, was calculated using the *Psych* package in R[9]. Thereafter, a pairwise chi-square test of independence was performed to examine associations of *mental health indicators* and other variables and a p-value of less than 0.05 was taken as the cut-off for significance.

Since mental health variables may have complex dependencies with potential confounding factors, mediation, and inter-causal dependency, we extended our association analysis with data-driven Bayesian network (BN) structure learning. The structure of the learned Bayesian network was made robust through bootstrapping and ensemble averaging of edge directions. Hill climbing optimizer [10] with Akaike Information Criterion(AIC)[11] based score was used to select the best PGM that explained the data. Bootstrapped learning and majority voting over 101 BNs were done. Exact Inference using the belief propagation algorithm[12] was learned in order to quantify the strength of learned associations. The Analysis was performed in the R using the package *wiseR*[13] Next, the *Markov-blanket*[14] of *mental-health-indicators* was extracted to select features that may predict responses to the *mental-health-indicators*. Data were partitioned into training (80%) and testing (20%) sets and the class imbalance was corrected using the Synthetic

Minority Oversampling Technique(SMOTE)[15]. Different supervised machine learning models - Random-Forest(RF), Support vector machine (SVM), logistic, naive-Bayes, were learned for predicting the response to mental health indicators using the Scikit-learn library [16]in Python.

Results:

Gender and age-related variation in mental-health indicators. *soc5a* (Felt nervous, anxious*,* or on edge), *soc5b* (Felt depressed), *soc5c* (Felt lonely), *soc5d* (Felt hopeless about the future) achieved a Cronbach's alpha approximating 0.8 [Figure 1(b)], thus confirming their internal consistency and suitability for modeling. Gender and age-specific difference was observed in *soc5a*, with females having a higher incidence than males (two proportion z-test, p-value < 2.2e-16) (Figure 2a), and young adults in the 18-29 age group (p-value < 2.2e-16) (Figure 2b). Age group 18 - 29 in both genders was most vulnerable to mental stress > 5 days in a week, thus indicating that COVID-19 may have disproportionately affected the mental health of youth due to a variety of factors.

Associations of anxiety in the United States. A Chi-square test revealed many significant associations of the mental health variables(Supplementary Figure 1). However, this analysis does not account for potential confounding or "explaining away" effects. Hence a data-driven Bayesian network structure-learning exercise was carried out and revealed interesting findings. From the learned structure, *soc5a* (Felt nervous, anxious, or on edge in the last seven days) was found to be the parent variable for other mental health indicators in almost 100% of the bootstrapped networks, represented as the strength of the edges [Figure 3(b)]. Being a driver variable in the structure, *soc5a* was taken as the primary dependent variable for downstream modeling analysis.

Impact of social life and work-related stressors. Our analysis using network inference via the Exact Inference algorithm showed a clear impact of in-person social communication on the reduction of anxiety level. A strong (>5% with CI ~1% at both sides) and (>6.5% with CI ~1.5% at both sides) monotonic increase between control of anxiety and frequency of speaking with neighbors *(soc2a, soc2b)* were observed. This effect was weaker (~1.5% with a wide confidence interval) with digital communication with friends and family conducted over phone, text, email, or other internet media (*soc3a, soc3b*). This finding underscores the importance of social communication while maintaining the appropriate measures such as masks and social distancing in order to maintain mental health during such isolating times. We also observed that the presence of kids in the house reduces the probability of depression by >11% with CI ~2% on both sides. Further, Exact Inference upon the network revealed an increase in the conditional

probability of anxiety (*soc5a*) arising from canceled or postponed work (>4% with CI ~1.4%), canceled, or postponed school(7% with CI ~1.5%), worked from home(>5% with CI ~1.3) and studied from home(>7% with CI ~1.8). Interestingly, although 83% of all volunteers chose to wear the mask, 77% avoided restaurants, 83% avoided public and crowded places these measures were not found to be associated with a significant change in anxiety levels as inferred from our model. These inferences are summarized in figure-4.

Impact of symptoms and comorbidities. We also investigated the relationship between mental stress and COVID symptoms indicators. WHO recommends contacting health service providers if any COVID symptoms (*phys1a* to *phys1q*) are experienced within the last seven days. Our network did not indicate any significant impact of these responses on mental health (*soc5a*), the conditional probability of which remained unchanged (62.2%) across the responses. Surprisingly, although medical conditions *(phys3a* to *phys3m)* are known to increase the risk of serious illness from COVID-19, our model showed that suffering from cancer (*phys3k*) and hypertension (*phys3b*) had a reverse impact on anxiety levels. Those suffering from cancer had approximately 8.3% (with ~2% CI) higher conditional probability of having less than one anxiety-ridden day in a week (>7% effect for hypertension with CI ~1.5%). Additionally, Cystic-fibrosis (phys3i) and Liver-disease (phys3j) have wide confidence intervals with non-significant differences in mean values. (Figure-4).

Impact of economic factors. Receiving income assistance through Social Security improved the conditional probability of less than one day of anxiety in a week by 10.4% (with CI ~1.5%) as compared with the segment of people who did not apply or receive it. Just applying for income assistance led to a 4% improvement (Figure-4). Similarly, Supplemental Social Security (~5.5 with CI ~4%) and Health insurance (~5% with CI ~2%) also led to similar results.
In addition to this, Old age people found health insurance more relaxing than young age people. COVID has also severely affected the financial condition of individuals. That may also lead to mental stress.

Predictive modeling for susceptibility to anxiety attacks. Our supervised modeling approach used the Markov blanket of the *soc5a* variable, i.e. age (*age4),* physical symptoms in the last seven days (*phys7_4),* stay at home (*phys2_18*), and prior clinical diagnosis of any mental health condition (*phys3h*) as predictors.

The following three prediction scenarios were considered-

1. Mental issues *less than one day* in a week (class 1) vs. Mental issues *more than **one** day* in a week (class 0)
2. Mental issues *less than one day* in a week (class 1) vs. Mental issues *more than **three** days* in a week (class 0)
3. Mental issues *less than one day* in a week (class 1) vs. Mental issues *more than **five** days* in a week (class 0)

We observed a decay (Accuracy 0.80 to 0.64, also confidence intervals mentioned in the table-2) in model predictability as we moved from high risk of depression (case-3) to low risk of depression (case-1) [figure4]. Such a trend was visible with all 4 machine learning techniques we used. Our ML properties (accuracy, sensitivity, specificity, AUROC) are summarized in table-2.

Discussion:

Mental Health is a serious public health concern. Mood disorders and suicide-related outcomes have increased significantly over the last decade among all age groups and genders[17] [18]. The rapid spread of coronavirus infection forced governments across the world to close public gathering places, schools, colleges, restaurants, and industries. Social isolation, digital communication, working, and schooling from home have become the new normal and many jobs have been lost. Collectively, this has triggered a high level of anxiety, stress, and depression, globally. We did not find studies that have used models to not just predict but also to explain the subtle effects of life-situations on mental health. An explainable probabilistic graphical modeling approach with bootstraps and exact inference allowed us to capture many of these effects in a robust manner. Our study revealed that individuals having a prior diagnosis of any mental illness are the most vulnerable for mental illness during the COVID-19 phase, which recommends building national-level policies to regularly track their mental status and treat them accordingly. Most importantly, our results re-iterate the economic underpinnings of collective mental health response. Income assistance via Social Security or Supplemental Social Security had a demonstrable effect on the alleviation of anxiety as inferred from our model, which provides the first scientific evidence, to the best of our knowledge, proving the utility of such efforts. The effect of the extent of such measures may be captured in such modeling studies conducted in various parts of the globe, with widely varying assistance structures during this time.

Our findings from the United States can also stimulate further cultural and social research in other geographies with similar or different social structures. For example, the effects of in-person communication, as opposed to digital connectedness, may be different in countries where

community living and joint families are still commonplace, e.g. India. Digital connectedness was not as effective as talking to a neighbor, at least in the United States highlighting that these are fundamentally different influences on mental health and need to be further explored in systematic studies. We conjecture that such differences may arise from the evolutionary mechanisms that have shaped human societies to live and share in close physical connectedness. Such an effect has been previously shown in primates kept in isolation who display depressive symptoms [19] [20]. Similarly, parenting and its association with neuropeptide hormones may partially explain [21] our results that the presence of kids reduces anxiety levels. Interestingly, the COVID-19 pandemic has created a unique natural experiment on the collective mental health response of individuals to a health emergency.

The life-cycle of such a response may need to be further studied as the world goes through various phases of the pandemic to its resolution. However, our study indicates that the mental health impact is observable within a span of a few months, especially on young individuals. Further research will be needed, ideally in a longitudinal setting, where the same individuals can be surveyed again to understand the dynamics of the collective mental health response.

Our results also highlight that modern technological development in virtual communication is not able to replace natural socializing. Hence it becomes imperative to design better and empathetic technological tools that may shape a society that prevents isolation and alienation even while maintaining physical distancing and preventive measures for limiting spread. Personalization and contextualization of such measures will also be important as our results indicate that persons with previous mental health conditions may be disproportionately affected.

Finally, our results indicate that it may be possible to identify people at the highest risk of developing mental health disturbances. Our model achieved its best performance for those who were most vulnerable (having mental stress more than five days in a week) vs. least vulnerable people (having no stress or less than one-day stress in a week). This can help in the segmentation of vulnerable populations such as front-line healthcare workers and who are facing disproportionately higher levels of stress during this time.

We could not explain why anxiety levels may be lower in persons with pre-existing cancer or hypertension. This may be a result of reduced work-environment related stress or more contact with family members at home. However, the current dataset is not suited to address this at a finer level of explainability. Our study has a few limitations. The dataset used for training the model is cross-sectional and we could not comment upon the temporality and persistence of the discovered effects. Secondly, our results were currently limited to only one geography, i.e. the United States. However, the relatively large sample size and multi-ethnic involvement in the survey allowed us to construct a robust network with ensemble averaging and majority voting from 101 bootstrapped networks. Therefore, the discovered influences are likely to hold true in the United States. Further, we believe that this is the first attempt to quantify the impact of social factors on mental health through an explainable AI model and many of our findings are intuitive. The ranking of features and quantification of this impact couldn't have been intuitively achieved

without modeling and this study will provide a basis for many further studies and design of effective social interventions to mitigate the mental health impact of natural health disasters.

Author Contribution:

Study Design : VK,TP,IJ , Dataset: IJ,AK , Data Analysis: IJ,RA , Paper writing: IJ,RA,AK,TP
Paper Review: VK,TP

Conflict of Interest:

None

Funding Statement:

The authors did not have specific funding for this work.

References:

1. COVID-19: Stimulating the economy and employment: As jobs crisis deepens, ILO warns of uncertain and incomplete labour market recovery. Accessed August 6, 2020.

2. Psychiatry TL. Mental health and COVID-19: change the conversation. *Lancet Psychiatry*. 2020;7(6):463. doi:10.1016/S2215-0366(20)30194-2

3. Pereira-Sanchez V, Adiukwu F, Hayek SE, et al. COVID-19 effect on mental health: patients and workforce. *Lancet Psychiatry*. 2020;7(6):e29-e30. doi:10.1016/S2215-0366(20)30153-X

4. Chen Q, Liang M, Li Y, et al. Mental health care for medical staff in China during the COVID-19 outbreak. *Lancet Psychiatry*. 2020;7(4):e15-e16. doi:10.1016/S2215-0366(20)30078-X

5. Mahase E. Covid-19: Mental health consequences of pandemic need urgent research, paper advises. *BMJ*. 2020;369:m1515. doi:10.1136/bmj.m1515

6. Pfefferbaum B, North CS. Mental Health and the Covid-19 Pandemic. *N Engl J Med*. 2020;0(0):null. doi:10.1056/NEJMp2008017

7. Gao J, Zheng P, Jia Y, et al. Mental health problems and social media exposure during COVID-19 outbreak. *PLOS ONE*. 2020;15(4):e0231924. doi:10.1371/journal.pone.0231924

8. Homepage | COVID Impact Survey. Untitled. Accessed June 26, 2020. https://www.covid-impact.org

9. Revelle W. *Psych: Procedures for Psychological, Psychometric, and Personality Research*.; 2020. Accessed June 26, 2020. https://CRAN.R-project.org/package=psych

10. Gámez JA, Mateo JL, Puerta JM. Learning Bayesian networks by hill climbing: efficient methods based on progressive restriction of the neighborhood. *Data Min Knowl Discov*. 2011;22(1):106-148. doi:10.1007/s10618-010-0178-6

11. Bozdogan H. Model selection and Akaike's Information Criterion (AIC): The general theory and its analytical extensions. *Psychometrika*. 1987;52(3):345-370. doi:10.1007/BF02294361

12. Yedidia JS, Freeman WT, Weiss Y. Generalized Belief Propagation. In: Leen TK, Dietterich TG, Tresp V, eds. *Advances in Neural Information Processing Systems 13*. MIT Press; 2001:689–695. Accessed June 28, 2020.

13. Sethi T, Maheshwari S. *WiseR: A Shiny Application for End-to-End Bayesian Decision Network Analysis and Web-Deployment*.; 2018. Accessed May 8, 2020. https://CRAN.R-project.org/package=wiseR

14. Probabilistic Graphical Models: Principles and Techniques - Daphne Koller, Nir Friedman - Google Books.

15. Chawla NV, Bowyer KW, Hall LO, Kegelmeyer WP. SMOTE: Synthetic Minority Over-sampling Technique. *J Artif Intell Res*. 2002;16:321-357. doi:10.1613/jair.953

16. scikit-learn: machine learning in Python — scikit-learn 0.23.1 documentation. Accessed June 26, 2020. https://scikit-learn.org/stable/

17. Bilsen J. Suicide and Youth: Risk Factors. *Front Psychiatry*. 2018;9. doi:10.3389/fpsyt.2018.00540

18. Brådvik L. Suicide Risk and Mental Disorders. *Int J Environ Res Public Health*. 2018;15(9). doi:10.3390/ijerph15092028

19. Hennessy MB, McCowan B, Jiang J, Capitanio JP. Depressive-like behavioral response of adult male rhesus monkeys during routine animal husbandry procedure. *Front Behav*

Neurosci. 2014;8. doi:10.3389/fnbeh.2014.00309

20. McKinney WT. Primate Social Isolation: Psychiatric Implications. *Arch Gen Psychiatry*. 1974;31(3):422-426. doi:10.1001/archpsyc.1974.01760150122018

21. Kohl J, Autry AE, Dulac C. The neurobiology of parenting: A neural circuit perspective. *BioEssays News Rev Mol Cell Dev Biol*. 2017;39(1):1-11. doi:10.1002/bies.201600159

Table 1: Socio-Demographics of Respondents Participated in the Survey

Factors	GENDER	MALE	FEMALE	REFUSED	SKIPPED	TOTAL
AGE	All Age Group	7,886	9,868	1	9	17,764
	18-24	438	559	0	1	998
	25-34	1,224	1,662	0	4	2,890
	35-44	1,130	1,471	0	2	2,603
	45-54	1,096	1,467	0	0	2,563
	55-64	1,578	2,110	1	2	3,691
	65-74	1,654	1,885	0	0	3,539
	75+	765	714	0	0	1,479
	Not Provided	1	0	0	0	1
EDUCATION	All education Group	7,886	9,868	1	9	17,764
	BA or above	4,439	4,922	0	4	9,365
	Some college	2,317	3,192	1	1	5,511
	HS graduate or equivalent	913	1,375	0	0	2,288
	No HS diploma	210	362	0	0	572
	Not sure	1	2	0	0	3
	REFUSED	2	4	0	0	6
	SKIPPED on web	4	11	0	4	19
GEO-LOCATION	All Geo-Locations	7,066	9,134	1	9	16,210
	Atlanta-Sandy Springs-Al..	317	458	0	0	775
	Baltimore-Columbia-Tow..	288	439	0	0	727
	Birmingham-Hoover, Ala..	264	409	0	0	673
	California	608	681	0	0	1,289
	Chicago-perville-Elgin, Ill..	376	473	0	0	849
	Cleveland-Elyria, Ohio	350	475	1	1	827
	Colorado	384	458	0	1	843
	Columbus, Ohio	335	429	0	0	764
	Florida	461	514	0	0	975
	Louisia	249	423	0	0	672
	Minnesota	363	438	0	2	803
	Missouri	288	457	0	0	745
	Monta	358	453	0	1	812
	New York	425	484	0	1	910
	Oregon	355	495	0	1	851
	Phoenix-Mesa-Chandler, ..	267	317	0	0	584
	Pittsburgh, Pennsylvania	336	457	0	1	794
	Texas	326	467	0	1	794
	Not Mentioned/Skipped	716	807	0	0	1,523
	Missing	820	734	0	0	1,554
RACE	All RACE Group	7,886	9,868	1	9	17,764
	Hispanic	622	899	0	1	1,522
	Black, non-Hispanic	521	1,057	0	0	1,578
	White, non-Hispanic	5,240	6,204	0	3	11,447
	Other, non-Hispanic	620	589	0	1	1,210
	REFUSED	2	5	1	0	8
	SKIPPED on web	8	2	0	2	12
	Removed for disclosure ri..	871	1,112	0	2	1,985
	DON'T KNOW	2	0	0	0	2
	Missing	0	0	0	0	0
LANGUAGE	All language Group	7,886	9,868	1	9	17,764
	English	7,815	9,757	1	9	17,582
	Spanish	71	111	0	0	182

Axis: 0K 2K 4K 6K 8K 0K 5K 10K -1 0 1 2 0 5 10 0K 10K

MALE | FEMALE | REFUSED | SKIPPED | TOTAL

Figure1: a) Outline of the analytical pipeline. b) Item Reliability Analysis of Mental Health Indicators revealed a high degree of internal consistency (Cronbach's alpha value >0.70) for most of the psychological variables, thus indicating suitability for the modeling exercise.

(a)   (b)

Figure2 : (a) Gender-wise and (b) age-wise distribution of Mental Issues (soc5a) variable. Significance was tested using two proportion z-test and chi-square test respectively, showing a higher prevalence of mental issues among youth and in women.

Figure 3: (a) Consensus structure learned through 101 bootstrapped samples. Hill-climbing search along with Bayesian Information Criterion were used to learn the structures and connections having edge strength and direction strength more than 90% are shown; **(b)** soc5a was found to be the parent Node of all other Mental Health Variables, therefore, leading to our choice of this variable as the Primary dependent variable.

Table 2: Model Performance Indicators of the Supervised Model for Prediction of Stress.

Data Set	Model Name	Accuracy (+- CI)	Sensitivity (+- CI)	Specificity (+- CI)	AU Roc (+- CI)
Mental issues less than one day in a week => class 1 Vs Mental issues more than five days in a week => class 0	RF	0.804 (+-0.016)	0.59(+-.063)	0.82(+-0.016)	0.71(+-0.026)
	SVM	0.802 (+-0.016)	0.56(+-.063)	0.82(+-0.016)	0.69(+-0.026)
	Naive Bayes	0.77 (+-0.017)	0.59(+-.063)	0.79(+-0.017)	0.69(+-0.025)
	Logistic	0.77 (+- 0.017)	0.59(+-.063)	0.78(+-0.017)	0.68(+-0.025)
Mental issues less than one day in a week => class 1 Vs Mental issues more than three days in a week => class 0	RF	0.72 (+- 0.018)	0.6(+-0.041)	0.75(+-0.018)	0.68(+-0.022)
	SVM	0.72 (+- 0.018)	0.6(+-0.041)	0.75(+-0.018)	0.67(+-0.022)
	Naive Bayes	0.74 (+-0.017)	0.56(+-0.041)	0.78(+-0.017)	0.67(+-0.022)
	Logistic	0.73(+-0.018)	0.57(+-0.041)	0.76(+-0.018)	0.67(+-.022)
Mental issues less than one day in a week => class 1 Vs Mental issues more than one day in a week => class 0	RF	0.66(+- 0.019)	0.48(+-0.027)	0.77(+-0.018)	0.62(+-0.019)
	SVM	0.66(+-0.019)	0.49(+-0.027)	0.76(+-0.018)	0.62(+-0.019)
	Naive Bayes	0.65 (+-0.019)	0.45(+-0.026)	0.77(+-0.018)	0.61(+-0.020)
	Logistic	0.62 (+- 0.019)	0.61(+-0.026)	0.64(+-0.020)	0.62(+-0.018)

Figure-4: Inferences from the Bayesian network: The difference in inferred probability was calculated after conditioning on the independent variables. A positive association implies a mental-stress-inducer whereas a negative association implies mental-stress-reducer. The red circle shows the mean value with green and blue showing confidence intervals.

Inferred Probability Difference

Supplementary Table 1: variable groups as Indicators

Code	Attributes Description
mental-health-indicators	
(questions related to mental stress in the last seven days)	
soc5a	Felt nervous, anxious, or on edge
soc5b	Felt depressed
soc5c	Felt lonely
soc5d	Felt hopeless about the future
soc5e	sweating, trouble breathing, pounding heart, etc
work-from-home-indicators	
(questions related to work and school from home)	
phys2_9	work from home
phys2_10	study from home
phys2_4	postponed work activities
phys2_5	postponed school activities
communication-indicators	
(questions related to communication with friends/neighbor/family)	
soc2a	in-person communication during COVID
soc2b	in-person communication before COVID
soc3a	digital communication during COVID
soc3b	digital communication before COVID
covid-symptoms-indicators	
(questions on covid symptoms Prescribed by WHO)	
phys1a	Fever
phys1b	Chills
phys1c	Runny or stuffy nose
phys1d	Chest congestion
phys1e	Skin rash
phys1f	Cough
phys1g	Sore throat
phys1h	Sneezing
phys1i	Muscle or body aches

phys1j	Headaches
phys1k	Fatigue or tiredness
phys1l	Shortness of breath
phys1m	Abdominal discomfort
phys1n	Nausea or vomiting
phys1o	Diarrhea
phys1p	Changed or lost sense of taste or smell
phys1q	Loss of appetite

chronic-medical-conditions-indicators
(questions on the chronic medical condition or prior clinical history of diseases)

phys3a	Diabetes
phys3b	High blood pressure or hypertension
phys3c	Heart disease, attack, stroke
phys3d	Asthma
phys3e	Chronic lung disease or COPD
phys3f	Bronchitis or emphysema
phys3g	Allergies
phys3h	A mental health condition
phys3i	Cystic fibrosis
phys3j	Liver disease
phys3k	Cancer
phys3l	compromised immune system
phys3m	obesity

behavioural-indicators
(questions on behavioral changes proposed/enforced by the government)

phys2_1	Canceled a doctor appointment
phys2_2	Worn a face mask
phys2_3	Visited a doctor or hospital
phys2_7	Canceled outside housekeepers
phys2_8	Avoided restaurants
phys2_11	postponed pleasure, social activities
phys2_12	Stockpiled food or water
phys2_13	Avoided public or crowded places

phys2_16	Washed or sanitized hands

insurance-assistance-indicators
(questions on insurance and different assistance)

econ6a	Unemployment insurance
econ6b	SNAP
econ6c	TANF
econ6d	Social Security
econ6e	Supplemental Social Security
econ6f	Health insurance
econ6g	aid from the government

other-useful-Indicators

phys7_4	did not feel hot, chilly, sweating
phys2_18	Stayed home because I felt unwell

Supplementary Figure 1: Heatmap for negative log p values for the significant association.

	SOC5A	SOC5B	SOC5C	SOC5D	SOC5E
PHYS2_1	3.929	5.110	4.370	3.732	5.999
PHYS2_2	5.834	3.518	5.701	5.602	5.999
PHYS2_3	4.527	4.797	3.146	5.810	6.000
PHYS2_6	5.999	6.000	6.000	6.000	5.978
PHYS2_7	0.986	3.797	2.125	2.636	1.698
PHYS2_9	6.000	6.000	6.000	6.000	5.420
PHYS2_14	1.086	1.349	3.591	1.484	0.768
PHYS2_15	6.000	6.000	6.000	6.000	6.000
PHYS2_16	2.699	3.970	2.889	3.165	2.324
PHYS2_17	5.772	1.669	2.388	2.563	0.668
PHYS2_18	6.000	6.000	6.000	6.000	6.000
PHYS2_19	4.722	3.945	5.556	3.338	6.000
ECON6C	5.973	6.000	5.999	6.000	6.000
PHYS3A	6.000	6.000	6.000	5.951	6.000
PHYS3B	6.000	6.000	6.000	6.000	5.993
PHYS3C	6.000	6.000	6.000	5.979	5.999
PHYS3E	6.000	6.000	6.000	4.895	6.000
PHYS3I	6.000	6.000	4.516	2.490	6.000
PHYS3J	6.000	6.000	5.777	5.756	6.000
PHYS3K	6.000	6.000	6.000	5.999	6.000
SOC2A	6.000	6.000	6.000	6.000	5.671
SOC2B	6.000	6.000	6.000	6.000	5.972
SOC3A	5.847	6.000	6.000	3.953	2.732
SOC3B	5.955	5.835	6.000	4.156	1.039

F3

0.668   6.000

Supplementary Figure 2: Bayesian Network topology