**The association between stigmatizing attitudes towards depression and help seeking attitudes in college students: Baseline results of a randomized controlled trial**

Virgínia da Conceição[1,2*], Inês Rothes[3,4], Ricardo Gusmão[1,2,5]

[1] EPIUnit - Institute of Public Health of the University of Porto (ISPUP), Portugal

[2] Laboratory for Integrative and Translational Research in Population Health (ITR), Porto, Portugal

[3] Faculty of Psychology and Education Science, University of Porto, Portugal

[4] Center for Psychology at University of Porto, Portugal

[5] Department of Public Health and Forensic Sciences, and Medical Education, Faculty of Medicine, University of Porto, Portugal

*Corresponding author

Virgínia da Conceição

virginia.mendes.conceicao@gmail.com

**Abstract**

College years represent a critical phase in young people's lives and concomitant with an increased risk of mental illness. Depression stigma has been considered a significant barrier to treatment. In this study, we aimed to understand the effects of age, gender, previous health care use, and symptomatology on depression stigma and analyze the impact of depression stigma on help-seeking attitudes. A total of 969 students with a mean age of 18.87 (SD=1.49) ranging between 16 and 25 were included in this study and completed the Depression Stigma Scale, the Attitude Toward Seeking Professional Psychological Help, the Patient Health Questionnaire-4 questionnaire, and a socio-demographic questionnaire. Data were analyzed using SPSS 24.0. Participants came from 14 University schools, and 64.6% were women. Stigma and help-seeking attitudes are positively affected by gender and previous access to mental healthcare services. Higher personal stigma weakened help-seeking attitudes. Depressive and anxiety symptoms influenced personal depression stigma and perceived stigma; however, we detected no direct symptomatology effect on help-seeking attitudes. Whereas personal depression stigma has an essential effect on help-seeking attitudes, depressive and anxiety symptoms do not. The promotion of literacy may decrease personal depression stigma and increase professional help-seeking intentions.

**Keywords:** Depression, social stigma, help-seeking behaviour, students

## 1. Introduction

Mental and behavioural disorders are estimated to account for 12% of the global burden of disease (World Health Organization, 2001). The number of people diagnosed with depressive disorders amounts to 4.4% of the world population, and depression ranks as the first contributor to non-fatal health burden, with 7.5% of all Years Lived with Disability (YLD) (World Health Organization, 2017).

College years are a critical developmental phase that typically coincides with emerging adulthood (Arnett, 2000), growingly conceptualized as 'late adolescence' (Sawyer et al., 2018; Twenge and Park, 2019). Entering a college or university mainly presents a high-risk transition due to several situational changes and upturns in life stressors, including academic and interpersonal difficulties (Uehara et al., 2010). Mental illness impacts academic performance negatively, increasing the odds of school attrition (Alonso et al., 2018; Auerbach et al., 2016). First-year college students have been identified as a risk group for developing depression, with reports of higher prevalence among university students than non-students for the same age group (Auerbach et al., 2018; Ibrahim et al., 2013).

People affected by depressive disorders face peer rejection (Moritz and Roberts, 2016), social and institutional stigmatizing responses similar to psychosis or chronic mental illness (Corrigan, 2006) with non-recognition of mental disorders as general health illness both by health professionals and society in general (McNair et al., 2002), healthcare access constraints (Knaak et al., 2017; Thornicroft et al., 2017), and increased difficulties in finding and maintaining a paid job (Khalema and Shankar, 2014; Rice et al., 2009).

Thus, reducing the stigma affecting people who have a mental illness is one of the main objectives of the Mental Health Action Plan 2013–2020 of the World Health Organization (World Health Organization, 2013). Identifying predictors of stigma is an important starting point to address stigmatizing attitudes (Griffiths et al., 2008).

Much of the recent research on depression stigma has focused on two types of social stigma: personal stigma and perceived stigma (Kosyluk et al., 2016). Personal stigma refers to one's own beliefs about depression, and perceived stigma refers to one's beliefs about the attitudes of others (Griffiths et al., 2008). Research on these two types of

stigma have reported lower levels of personal stigma comparing to higher perceived stigma (Boerema et al., 2016; Calear et al., 2011).

There has been a growing interest in mental health stigma interventions and research in Portugal in the last decade. A European study centred on depression stigma and attitudes towards professional help in the general adult population (Coppens et al., 2013) identified Portugal as the second country (in four) with the highest level of personal and perceived depression stigma, with 31.2% of the participants agreeing with personal stigmatizing affirmations, and 53.8% agreeing with perceived stigma affirmations, measured with the Depression Stigma Scale (Griffiths et al., 2004). In the attitudes towards professional help scale, Portugal ranked in the first place for openness to professional help; however, the value of professional service was not as good, with only 48.2% agreeability. Lower schooling levels, higher age, being a male, and not having previous contact with people with mental illness were associated with increased stigma scores (Dietrich et al., 2014; Oliveira and Azevedo, 2014).

Particular attention has been made to mental health stigma in health students as health professionals may act as literacy promoters (Xavier et al., 2013). In the past years, research about stigma in health students in Portugal has demonstrated the need for stigma reduction intervention in this group of students (Querido et al., 2016). Anti-stigma specific education (Gil, 2015; Telles-Correia et al., 2015).

Help-seeking has been a particular problem for first-year university students. Most of the WHO World Mental Health International College Student Initiative report participants reported they would hesitate to seek help in case of need, and attitudinal barriers were identified as the most important (Ebert et al., 2019).

Despite the increasing number of studies addressing mental health stigma and help-seeking  (Eisenberg et al., 2009; Jennings et al., 2015; Musa et al., 2020; Pedersen and Paves, 2014; Schnyder et al., 2017), most studies neglect to integrate the symptomatology of participants in the analysis.

Also, sampling in young adults has been a problem; usually, participants are health students or professionals (Wei et al., 2015), resulting in a lack of studies contemplating all study areas of the universities.

Lastly, the effects of previous mental healthcare utilization have not been well explored and can be helpful in the study of the relationship between stigma and help-seeking (Eisenberg et al., 2009).

*1.1 Aims*

With this study, we aim to (1) examine whether socio-demographic characteristics are related to personal and perceived stigma in students, (2) to understand the effects of the depressive and anxiety symptoms of depression stigma, and (3) examine whether stigmatizing attitudes towards depression is associated with help-seeking behaviour in students.

## 2. Methods

This study is part of a more extensive longitudinal research on university students' mental health, including an experimental single-blind randomized control trial (ISRCTN970936) to evaluate the effectiveness of a video intervention in depression stigma reduction and help-seeking promotion.

*2.1 Participants*

Using the SurveyMonkey platform, in February 2019, we contacted all 4493 first-year students from all 14 schools of the University of Porto through institutional email. We invited them to participate in the study. A reminder email was sent a week after the first email.

At the end of the questionnaire, a list of available healthcare responses was available, so students could seek help if they felt it was needed.

*2.2 Measures*

We asked participants to answer a socio-demographic questionnaire assessing gender, age, place of birth, university course, the Attitudes Toward Seeking Professional Psychological Help (ATSPPH) (Fischer and Turner, 1970), the Patient Health Questionnaire-4 (PHQ-4) (Khubchandani et al., 2016), and the Depression Stigma Scale (DSS) (Griffiths et al., 2004).

The ATSPPH is a ten-item four-point Likert scale, with scores varying from 0 to 30. The scale ranges from "disagree" (score 0) to "agree" (score 3) and comprises two subscales: the first one evaluates the openness to seeking treatment for emotional problems (items 1, 3, 5, 6 and 7), and the second evaluates the value and need on seeking treatment (items 2, 4, 8, 9 and 10, reversed before summed). The maximum score in each subscale is 15, and higher scores indicate better attitudes toward seeking treatment. ATSPPH is the most used scale in mental health help-seeking attitudes, showing excellent psychometric properties in its original form. One of the authors was responsible for the process of translation and adaptation of ATSPPH to the Portuguese population in the context of the OSPI program - Optimizing suicide prevention programs and their implementation in Europe (Coppens et al., 2013; Hegerl et al., 2009; Kohls et al., 2017).

PHQ-4 is a 4-item questionnaire with two items measuring depressive symptoms and two items for anxiety symptoms. Its score varies from 0 to 12; scores 0–4 represent absent symptomatology, 5–8 mild symptomatology, and 9–12 severe symptomatology (Löwe et al., 2010). The PHQ-4 used in the study resulted from the extraction of the corresponding items from the Portuguese version of the PHQ-9 (Ferreira et al., 2018; Monteiro et al., 2013) and GAD-7 (Sousa et al., 2015),

DSS is an 18 five-point Likert scale, with the first nine items measuring personal stigma (one's own beliefs about depression) and the last nine items measuring perceived stigma (one's beliefs about the attitudes of others). We converted the total score of both scales as a percentage according to the original score calculation (Griffiths et al., 2004), and the higher the percentage, the higher the stigma. The scale has been translated and validated to the Portuguese population using the OSPI program data (Coppens et al., 2013; Hegerl et al., 2009; Kohls et al., 2017).

*2.3 Data analysis*

We performed the analysis using SPSS 24 at a 95% confidence level.

We used percentages to describe participation rates, gender distribution, moving away from home, living situation, presence of mental illness in the family and previous access to mental healthcare. We used counts to describe participant distribution per faculties. Symptomatology stemming from the PHQ-4 scores resulted in three possible categories

of severity (absent, mild and severe), and we presented the distribution per category in percentages.

We divided previous access to mental healthcare in Group A, with no previous mental healthcare access, and Group B, with previous access to mental healthcare.

First, we calculated descriptive statistics: frequencies, means and standard deviations. Then, we performed variance analyses to examine gender differences, experience with previous healthcare, symptoms, depression stigma and help-seeking attitudes using One-Way ANOVA with Tukey HSD's multiple comparisons test among groups. To test the possible interaction effect between gender and previous mental healthcare groups on depression stigma and help-seeking attitudes, we performed a Two-Way ANOVA.

For normally distributed continuous variables (personal and perceived depression stigma and help-seeking attitudes) and performed T-tests to study differences between the gender, symptomatology groups and previous access to mental healthcare.

We carried out a Pearson correlation to examine associations between depression stigma and attitudes toward help-seeking.

Lastly, we executed hierarchical linear regressions to statistically control variables to see whether adding variables improves a model's ability to predict the Depression Stigma and the Help-seeking attitudes scores. Variables included in the models were gender, previous contact with mental illness, mental illness in the family and symptomatology group.

*2.4 Ethical considerations*

The Institute of Public Health of the University of Porto ethics committee approved the research on 26/12/2018 with I.D. reference CE18096.

## 3. Results

*3.1 Participants characteristics*

Of the 4493 students invited to participate in this study, a total of 1046 (23.3%) accepted to participate. Respondents came from all 14 schools: 224 from Engineering (22.6% of the 993 students invited), 179 from Humanities (23.6% of the 760 invited),

116 from Sciences (16.6% of the 698 invited), 86 from Biomedical Sciences (22.3% of the 386 invited), 82 from Economics (54.3% of the 151 invited), 81 from Law (33.1% of the 245 invited), 71 from Psychology and Educational Sciences (34.1% of the 208 invited), 53 from Arts (32.5% of the 163 invited), 49 from Pharmacy (26.5% of the 185 invited), 40 from Medicine (16.9% of the 236 invited), 27 from Nutrition (33.8% of the 80 students invited, 17 from Architecture (9.4% of the 139 invited), 13 from Sports (7.6% of the 171 invited), and eight from Dental Medicine (10.3% of the 78 invited).

To work with a sample within the late adolescent age limits, we excluded respondents aged 26 years and older, resulting in 969 participants (21.5%).

About two-thirds of the students were women (64.6%, 626), and 35.4% (343) were men, with a mean age of 18.87 (SD=1.49) ranging between 16 and 25. Approximately 43.9% of the students were away from home, and most of those (46.1%) were living with roommates.

Most of the participants had never previously sought mental health help care (59.6%), and 35.4% had a family member with a mental illness diagnosis. Of those who had previously sought mental healthcare, anxiety was the main reason (46.4%), followed by depression (25%).

In the PHQ-4, participants showed a mean of 5.65 (SD=3.35), and most of the participants had no symptomatology (51.8%). However, 25.5% showed mild symptomatology and 22.7% severe symptomatology. Only 47.3% of the participants with severe symptomatology had previously sought professional help.

### 3.2 Depression stigma
### 3.2.1   Personal Depression Stigma

As shown in Table 1, women had a lower personal stigma level than males. Participants in Group A obtained means of personal depression stigma statistically higher than those in Group B, but these differences turned out to be only significant among women. There was a significant interaction between gender and help-seeking group ($F_{(968,1)}=4.85$, $p<0.01$).

There was also a significant effect of symptomatology on personal depression stigma, and the mild symptoms groups presented the higher depression stigma score. Tukey

HSD's post-hoc analysis revealed that only the mild symptomatology group differed from the other severity symptoms groups: $p<0.01$ between absent and mild symptoms, and $p<0.001$ between mild and severe. When comparing depression stigma score differences among help-seeking groups, we verify significance in Group B; however, there was no significant interaction between symptomatology and help-seeking groups ($F_{(968,2)}=0.92$, $p=0.40$).

Table 1: Personal depression stigma mean differences according to gender, help-seeking and symptomatology groups.

| | Total | Group A (No) | Group B (Yes) | T-test$_{(df)}$ |
|---|---|---|---|---|
| Mean (SD) | 23.76 (12.37) | 25.72 (12.65) | 20.88 (11.53) | **t$_{(968)}$=-6.08, p<0.001** |
| Men: Mean (SD) | 27.37 (12.90) | 28.00 (13.22) | 26.11 (12.20) | t$_{(341)}$=-1.31, p=0.19 |
| Women: Mean (SD) | 21.79 (11.63) | 24.20 (12.08) | 18.78 (10.92) | **t$_{(625)}$= -5.98, p<0.001** |
| t-test$_{(df)}$ | **t$_{(968)}$=6.88, p<0.001** | **t$_{(577)}$= 3.56, p<0.001** | **t$_{(392)}$= 6.05, p<0.001** | |
| Absent symptoms (n=502) | 23.51 (11.89) | 25.11 (12.11) | 19.97 (9.98) | **t$_{(500)}$= -4.92, p<0.001** |
| Mild symptoms (n=247) | 25.91 (13.19) | 27.06 (13.44) | 23.95 (12.77) | t$_{(245)}$= -1.81, p=0.07 |
| Severe symptoms (n=158) | 21.72 (12.20) | 24.68 (13.19) | 18.36 (10.31) | **t$_{(218)}$= -3.93, p<0.001** |
| ANOVA$_{(df)}$ | **F$_{(966,2)}$=6.86, p<0.01** | F$_{(575,2)}$=1.51, p=0.22 | **F$_{(388,2)}$=4.20, p<0.01** | |

M=mean, SD=standard deviation; df=degrees of freedom. Significant results are in bold.

Participants with a family member with a mental illness diagnosis had a mean personal stigma score of 22.32 (SD=11.26), a significantly lower result ($t_{(966)}=-2.24$, $p<0.05$) than those with no family member with mental illness (M=24.17, SD=12.78).

Both gender and the help-seeking group significantly affected personal depression stigma, unlike having a family member with mental illness. Symptomatology was a significant predictor for those with mild symptoms compared with those with severe symptomatology (Table 2).

The hierarchical multiple regression revealed that at Stage one, gender contributed significantly to the regression model, $F_{(966,1)}= 48.62$, $p<0.001$ and accounted for 4.7% of the variation in Personal Depression Stigma. Introducing the previous care group variable explained an additional 2.9% variation, and this change in $R^2$ was significant, $F_{(966,1)}= 30.60$, $p <0.001$. Adding family mental illness and symptoms group did not produce an $R^2$ significant change. The most important predictor of personal stigma was gender which uniquely explained 4.8% of the variation.

Table 2: Effects of gender, previous help group, family mental illness, and symptomatology group on personal depression stigma

| | β | 95% CI | t | p |
|---|---|---|---|---|
| **Model 1** | | | | |
| Women | Ref. | | | |
| Men | **5.65** | **4.07, 7.25** | **48.61** | **<0.001** |
| **Model 2** | | | | |
| Women | Ref. | | | |
| Men | **5.16** | **3.58, 6.73** | **41.18** | **<0.001** |
| Previous help – Group A | Ref. | | | |
| Previous help – Group B | **-4.35** | **-5.89, -2.82** | **30.88** | **<0.001** |
| **Model 3** | | | | |
| Women | Ref. | | | |
| Men | **5.12** | **3.54, 6.69** | **40.40** | **<0.001** |
| Previous help – Group A | Ref. | | | |
| Previous help – Group B | **-4.20** | **-5.74, -2.65** | **20.20** | **<0.001** |
| Family with mental illness - No | Ref. | | | |
| Family with mental illness - Yes | -1.22 | -2.80, 0.36 | 2.30 | 0.13 |
| **Model 4** | | | | |
| Women | Ref. | | | |
| Men | **5.11** | **3.54, 6.68** | **40.74** | **<0.001** |
| Previous help – Group A | Ref. | | | |
| Previous help – Group B | **-4.12** | **-5.66, -2.59** | **27.43** | **<0.001** |
| Family with mental illness - No | Ref. | | | |
| Family with mental illness - Yes | -1.16 | -2.73, 0.41 | 2.09 | 0.15 |
| No symptoms | 1.02 | -0.86, 2.91 | 1.14 | 0.27 |
| Mild symptoms | **3.65** | **1.50, 5.79** | **11.04** | **<0.01** |
| Severe symptoms | Ref. | | | |

β=beta regression coefficients, Ref.=Reference category
* Model 1= gender; Model 2: Model 1 plus help-seeking group; Model 3: Model 2 plus family mental illness; Model 4: Model 3 plus symptomatology group.
Significant results are in bold.

### 3.2.2  *Perceived depression stigma*

Gender and help-seeking group differences were statistically significant in the total scores of the perceived depression stigma, with women and help-seeking group B

obtaining the highest scores. However, the interaction between gender and the help-seeking group was not significant ($F_{(968,1)}=0.01$, $p<0.94$).

The severe symptoms group presented the highest perceived stigma score. Tukey HSD's post-hoc analysis revealed a significant difference between the two groups: no symptoms and severe symptoms ($p<0.05$). The interaction between the symptoms and help-seeking groups was not significant ($F_{(968,2)}=0.49$, $p=0.62$).

Table 3: Perceived depression stigma as a function of previous experiences with mental health professionals, gender, and PHQ-4 scores.

| | Total | Group A (No) | Group B (Yes) | t-test$_{(df)}$ |
|---|---|---|---|---|
| Mean (SD) | 61.66 (17.60) | 60.58 (17.34) | 62.26 (18.24) | **$t_{(968)}=2.32$, $p<0.05$** |
| Men: Mean (SD) | 59.79 (16.17) | 58.92 (16.06) | 61.57 (16.03) | $t_{(341)}=1.42$, $p=0.15$ |
| Women: Mean (SD) | 62.68 (18.48) | 61.67 (18.07) | 63.94 (18.94) | $t_{(625)}=1.54$, $p=0.13$ |
| t-test$_{(df)}$ | **$t_{(968)}=-2.43$, $p<0.05$** | $t_{(577)}=-1.87$, $p=0.06$ | $t_{(392)}=-1.69$, $p=0.24$ | |
| Absent symptoms (n=502) | 60.89 (16.80) | 60.27 (16.50) | 61.91 (17.28) | $t_{(500)}=1.06$, $p=0.29$ |
| Mild symptoms (n=247) | 60.99 (18.31) | 59.96 (17.90) | 62.58 (18.91) | $t_{(245)}=1.10$, $p=0.27$ |
| Severe symptoms (n=158) | 64.52 (18.87) | 62.40 (18.71) | 66.90 (18.86) | $t_{(218)}=1.77$, $p=0.08$ |
| ANOVA$_{(df)}$ | **$F_{(966,2)}=3.53$, $p<0.05$** | $F_{(575,2)}=0.78$, $p=0.46$ | $F_{(388,2)}=2.68$, $p=0.07$ | |

M=mean, SD=standard deviation; df=degrees of freedom. Significant results are in bold.

We obtained non-significant ($t_{(968)}=1.77$, $p=0.08$) lower scores of perceived stigma in participants with no family member with mental illness (M=60.91, SD=17.31) compared with those with a family member bearing a mental disorder diagnosis (M=63.01, SD=18.49).

Since there were no differences between participants with and without mental illness in the family, we did not include this variable in the hierarchical linear regression.

As can be observed in Table 4, gender and symptomatology scores had significant effect on perceived depression stigma. Despite the significant effects verified, the variance

explained in each model is very small (0.6% in Model 1, 0,8% in Model 2, and 1.1% in model 3). Only Model 1 and Model 3 produced a $R^2$ significant change: Model 1 $F_{(967,1)}= 6.41$, p <0.05; Model 2 $F_{(967,1)}= 3.01$, p=0.08; Model 3 $F_{(967,1)}= 4.27$, p <0.05.

The most relevant predictor of perceived stigma was gender which explained merely 0.6% of the variation.

Table 4: Effects of gender, previous help group, family mental illness, and symptomatology group on perceived depression stigma.

| | β | 95% CI | t | p |
|---|---|---|---|---|
| **Model 1** | | | | |
| Women | Ref. | | | |
| Men | **-3.05** | **-5.38, -0.72** | **6.59** | **<0.05** |
| **Model 2** | | | | |
| Women | Ref. | | | |
| Men | **-2.76** | **-5.09, -0.42** | **5.36** | **<0.05** |
| Previous help – Group A | Ref. | | | |
| Previous help – Group B | **2.47** | **0.19, 4.74** | **4.50** | **<0.05** |
| **Model 3** | | | | |
| Women | Ref. | | | |
| Men | **-2.67** | **-5.00, -0.34** | **5.04** | **<0.05** |
| Previous help – Group A | Ref. | | | |
| Previous help – Group B | 2.26 | -0.02, 4.54 | 3.78 | 0.05 |
| No symptoms | **-3.29** | **-6.09, -0.49** | **5.32** | **<0.05** |
| Mild symptoms | **-3.28** | **-6.48, -0.08** | **4.04** | **<0.05** |
| Severe symptoms | Ref. | | | |

β=beta regression coefficients, Ref.=Reference category
* Model 1= gender; Model 2: Model 1 plus help-seeking group; Model 3: Model 2 plus symptomatology effect.
Significant results are in bold.

### 3.3 Help-seeking attitudes

Help-seeking attitudes presented a mean of 18.62 (SD=4.76). Women showed significantly higher means (19.20, SD=4.90) than men (17.56, SD=4.32) ($t_{(966)}$ =-5.20, p<0.001). Participants in the help-seeking Group B presented better help-seeking attitudes (20.00, SD=4.54) than those in Group A (17.69, SD=4.69, $t_{(966)}$ =7.66, p<0.001). Differences according to symptomatology were not significant: No symptoms M=18.77, SD=4.93; Mild symptoms M=18.15, SD=4.17; and Severe symptoms M=18.79, SD=4.96 with an ANOVA test of $F_{(966,2)}= 1.61$, p=0.20. We did not find a

significant interaction between the global score of the help-seeking attitudes scale and gender, help-seeking group, nor symptomatology group.

Participants with a family member with mental illness presented better help-seeking scores (M=19.22, SD=4.80) than those with no mental illness in the family (M=18.28, SD=4.80), $t_{(966)}$ =2.95, p<0.01.

Table 5: Effects of gender, previous help group, family mental illness, personal depression stigma and perceived depression stigma on help-seeking attitudes

|  | β | 95% CI | t | p |
|---|---|---|---|---|
| **Model 1** | | | | |
| Women | Ref. | | | |
| Men | **-1.64** | **-2.26, -1.02** | **5.20** | **<0.001** |
| **Model 2** | | | | |
| Women | Ref. | | | |
| Men | **-1.39** | **-1.20, -0.79** | **4.96** | **<0.001** |
| Previous help – Group A | Ref. | | | |
| Previous help – Group B | **2.17** | **1.58, 2.76** | **-7.18** | **<0.001** |
| **Model 3** | | | | |
| Women | Ref. | | | |
| Men | **-1.14** | **-1.99, -0.78** | **4.47** | **<0.001** |
| Previous help – Group A | Ref. | | | |
| Previous help – Group B | **2.08** | **1.48, 2.68** | **-6.84** | **<0.001** |
| Family with mental illness - No | Ref. | | | |
| Family with mental illness - Yes | 0.58 | -0.02, 1.19 | -1.87 | 0.06 |
| **Model 4** | | | | |
| Women | Ref. | | | |
| Men | **-0.64** | **-1.21, -0.07** | **2.19** | **<0.05** |
| Previous help – Group A | Ref. | | | |
| Previous help – Group B | **1.47** | **0.91, 1,67** | **-5.15** | **<0.001** |
| Family with mental illness – No | Ref. | | | |
| Family with mental illness – Yes | 0.40 | -0.16, 0.97 | -1.41 | 0.15 |
| Personal Depression Stigma | **-0.15** | **-0.17, -0.12** | **-12.64** | **<0.001** |
| **Model 5** | | | | |
| Women | Ref. | | | |
| Men | **-0.59** | **-1.16, -0.01** | **1.99** | **<0.05** |
| Previous help – Group A | Ref. | | | |
| Previous help – Group B | **1.44** | **0.88, 1.99** | **-5.02** | **<0.001** |
| Family with mental illness - No | Ref. | | | |
| Family with mental illness - Yes | 0.37 | -0.19, 0.93 | -1.293 | 0.20 |
| Personal Depression Stigma | **-0.15** | **-0.17, -0.12** | **-12.76** | **<0.001** |
| Perceived Depression Stigma | **0.02** | **0.01, 0.04** | **2.53** | **<0.05** |

β=beta regression coefficients, Ref.=Reference category
* Model 1= gender; Model 2: Model 1 plus help-seeking group; Model 3: Model 2 plus family mental illness; Model 4: Model 3 plus Personal Depression Stigma; Model 5: Model 4 plus Perceived Depression Stigma.
Significant results are in bold.

Personal depression stigma showed a moderate correlation with help-seeking attitudes (r=-0.42, p<0.01). The perceived stigma showed a weaker Pearson correlation with help-seeking attitudes (r=0.10, p<0.01).

The hierarchical multiple regression (table 5) revealed that at Stage one, gender contributed significantly to the regression model, $F_{(966,1)}$= 27.62, p<0.001 and accounted for 2.8% of the variation in the general score of the help-seeking attitudes scale. Introducing the previous care group variable explained an additional 4.9% variation, and this change in $R^2$ was significant, $F_{(966,1)}$= 50.80, p <0.001. Adding family mental illness did not produce an $R^2$ significant change. Personal depression stigma explained an additional 13.1% of the variance with a significant change in $R^2$ ($F_{(966,1)}$= 159.93, p <0.001), and perceived stigma explained an additional 0.01% of the variance. Model 5 total variance explained was 21.6%.

### 3.3.1    *Openness to seeking treatment for emotional problems*

The mean for openness to seeking treatment was 8.82 (SD=2.95), and the mean was higher among women (8.98, SD=3.10) than men (8.53, SD=2.64) with a significant difference: $t_{(966)}$ =-2.67, p<0.05. Participants in help-seeking Group B presented higher mean scores (9.71, SD=2.92) than those in Group A (8.21, SD=2.82), and the differences were statistically significant: $t_{(966)}$ =7.95, p<0.001.

Having a family member with mental illness beneficiated the openness to seeking treatment as the participants in this group obtained a higher mean (9.19, SD=2.88) than those with no mental illness in the family (8.61, SD=2.68), $t_{(966)}$ =2.92, p<0.01.

There were no differences according to symptomatology ($F_{(966,2)}$= 1.29, p=0.28).

Personal depression stigma showed a moderate correlation with openness to seeking treatment (r=-0.30, p<0.01), and the correlation with perceived stigma was not significant.

The hierarchical multiple regression revealed that gender contributed significantly to the regression model at model one, $F_{(966,1)}$= 5.30, p<0.05, but accounted only for 0.05% of the variation. Previous care explained an additional 5.8% variation, and this change in $R^2$ was significant, $F_{(966,1)}$= 32.87, p <0.001. Adding family mental illness group did not

produce an $R^2$ significant change ($F_{(966,1)}= 3.67$, p=0.06). Personal depression stigma explained an additional 12.7% of the variance with a significant change in $R^2$ ($F_{(966,1)}= 65.81$, p <0.001). Introducing perceived stigma in the model also did not result in an $R^2$ significant change: $F_{(966,1)}= 2.07$, p=1.51. Model 5 total variance explained was 12.4%. However, how we can see in table 1 of the supplementary file, in model 5, only previous care group and personal stigma had a significant beta regression coefficient.

### 3.3.2  Value and need of seeking treatment

Participants showed a mean of 9.81 (SD=2.56) in the value and need of seeking treatment. Woman presented significant higher means (10.23, SD=2.50) than men (9.03, SD=2.49), $t_{(966)} =-7.01$, p<0.001. Participants who have previously sought help (Group B) obtained better mean scores (10.29, SD=2.38) than participants in Group A (9.47, SD=2.63), with differences statistically significant: $t_{(966)} =4.97$, p<0.001. Participants with mental illness in the family presented better mean scores (10.03, SD=2.48) than participants with no family member with mental illness (9.67, SD=2.59), and these differences were statistically significant: $t_{(966)} =2.10$, p<0.05.

Symptomatology did not impact mean scores for value and need of seeking treatment ($F_{(966,2)}= 1.83$, p=0.16).

Personal depression stigma showed a moderate correlation with openness to seeking treatment (r=-0.45, p<0.001), and the correlation with perceived also showed a significant correlation, yet weaker (r=0.10, p<0.01).

Gender accounted for 5% of the variation as the first variable included in the hierarchical multiple regression, with a significant contribution for the $R^2$ change: $F_{(966,1)}= 51.37$, p<0.001. The previous care group variable explained an additional 1.8% variation, and the change produced in $R^2$ was also significant, $F_{(966,1)}= 17.79$, p <0.001. Family mental illness group did not change the $R^2$ ($F_{(966,1)}= 1.58$, p=0.21. Personal depression stigma explained an additional 15.3% of the variance with a significant change in $R^2$ ($F_{(966,1)}= 189.38$, p <0.001), and perceived stigma explained an additional lower amount of variance (0.1% ) yet significant in the $R^2$ change: $F_{(966,1)}= 8.94$, p<0.01. Model 5 explained a total variance of 22.9%. How we can see in table 2 of the

supplementary file, the previous help group and family with mental illness group did not have a significant beta regression coefficient.

## 4. Discussion

We contacted students to participate in this study via their institutional email addresses. Since they do not access their institutional email address very frequently, response rates were relatively low. Thus, the response rate was lower than the mean in similar studies in the general population (Burgard et al., 2019). Another explanation may be the recent increase in online surveys. Nevertheless, there was a comparable response rate within each school, except for Economics, with a much higher response rate than the mean, and Sports and Architecture with a response rate lower than 10%.

The gender distribution is comparable to the population: in the school year of 2018-2019, the University of Porto's records show that 59% of the students were women and 41% men.

Our sample presented lower stigma scores than the previous study in the general population (Coppens et al., 2013), which may be because our sample presents a much lower mean age, and depression stigma is associated with older age (Coppens et al., 2013; Dietrich et al., 2014).

As expected, personal stigma showed significantly lower means than perceived stigma, and women showed lower levels of personal stigma and higher levels of perceived stigma (Boerema et al., 2016; Calear et al., 2011). Some authors argue that may be due to masculine norms (Chatmon, 2020), a set of social norms and expected behaviour associated with men (Milner et al., 2018).

The previous use of mental health care had a significant effect on the personal stigma of women, but not men, which may also be because men have a more stigmatized perspective on mental illness. Thus, access to mental health care may not have the same stigma-reducing effect on men and women.

For perceived stigma, the difference becomes insignificant when analyzing the results by gender. This result agrees with previous studies, where the effects of the previous contact with depression are more evident for personal stigma than for perceived stigma (Busby Grant et al., 2016; Griffiths et al., 2008).

Depressive symptoms have a mitigating effect on personal stigma for those who have previously sought help, which may be due to the personal stigma reduction effects of mental health interventions.

Participants with a family member with mental illness, and thus having indirect experience with mental illness, showed lower personal depression stigma means; however, its effect on personal stigma was not significant.

Participants with milder symptoms showed more personal stigma and lower perceived stigma than those with severe symptomatology, resulting from a process of self-stigmatization and self-denial of a possible need for help, especially in the case of personal stigma. This particularity may be an essential element to consider when addressing personal stigma.

Symptomatology did not impact help-seeking attitudes, suggesting the importance of mental health literacy promotion to improve the recognized value of health care when needed.

The effect of gender is more substantial in value and need of seeking treatment than on the openness to seek treatment. While women see more value and need in the treatment-seeking, there are no significant differences when other effects are considered, such as personal stigma.

Past help-seeking had a more significant effect on the openness to seek treatment than on the value and need of seeking treatment, which may be interesting to research further as previous help-care experiences are an essential tool for literacy and help-seeking behaviours promotion.

Demonstration of a more substantial correlation of depression personal stigma with help-seeking attitudes than gender or previous contact with mental healthcare services confirms the importance of stigma interventions in promoting help-seeking behaviour (Gulliver et al., 2010; Thornicroft et al., 2017). Personal stigma concerning depression was the variable with the most significant magnitude effect on help-seeking attitudes, in line with the strong effect of attitudinal barriers in help-seeking (Ebert et al., 2019).

This result reinforces the importance of reducing stigma as a strategy to promote help-seeking behaviours.

One of the main limitations of this study is the sample's representativeness since the high rate of previous mental healthcare service usage, and the distributions of the PHQ-4 scores suggest some degree of selection bias. The second limitation is the lack of data on essential elements of stigma, namely the time and duration of mental health access care. Further studies should consider these elements to understand better the effects of previous mental healthcare usage on stigma and help-seeking attitudes, as well as the effect perfectionism on depression stigma (Shannon et al., 2018).

Despite the possible and inevitable sample bias in online studies, our sample compares well to the University of Porto population, both in distribution by the school as distribution by gender, strengthening the generalizability of the results.

Due to the low age variance in our sample – we only considered first-year students below 25 years old – age was not included as a study variable. Future research directed to the understanding of the relationship between age and depression stigma will address this gap, and as this particular sample may be followed, also the possible effect of stigma reduction as a result of the universities experiences and learnings, as reported on Portuguese health students (Querido et al., 2016)

Intensifying mental health promotion and easing access to mental healthcare will increase the probability of successful early intervention on college student's depression and anxiety disorders. More research will be needed to understand the different effects of mental health services on stigma reduction among men and women.

These findings also reinforce the importance of stigma reduction interventions as a tool to promote help-seeking behaviours.  Future research on the actual effect of stigma reduction on healthcare use is therefore essential.

**Funding:** National funds finance this work through FCT - Foundation for Science and Technology, I.P., under the project UIDB / 04750/2020

**Acknowledgements**: None

**Declaration of interest:** None

**References**

Alonso, J., Mortier, P., Auerbach, R.P., Bruffaerts, R., Vilagut, G., Cuijpers, P., Demyttenaere, K., Ebert, D.D., Ennis, E., Gutierrez-Garcia, R.A., Green, J.G., Hasking, P., Lochner, C., Nock, M.K., Pinder-Amaker, S., Sampson, N.A., Zaslavsky, A.M., Kessler, R.C., Collaborators, W.W.-I., 2018. Severe role impairment associated with mental disorders: Results of the WHO World Mental Health Surveys International College Student Project. Depress Anxiety 35 (9), 802-814.

Arnett, J.J., 2000. Emerging adulthood: A theory of development from the late teens through the twenties. American Psychologist 55 (5), 469-480.

Auerbach, R.P., Alonso, J., Axinn, W.G., Cuijpers, P., Ebert, D.D., Green, J.G., Hwang, I., Kessler, R.C., Liu, H., Mortier, P., Nock, M.K., Pinder-Amaker, S., Sampson, N.A., Aguilar-Gaxiola, S., Al-Hamzawi, A., Andrade, L.H., Benjet, C., Caldas-de-Almeida, J.M., Demyttenaere, K., Florescu, S., de Girolamo, G., Gureje, O., Haro, J.M., Karam, E.G., Kiejna, A., Kovess-Masfety, V., Lee, S., McGrath, J.J., O'Neill, S., Pennell, B.E., Scott, K., Ten Have, M., Torres, Y., Zaslavsky, A.M., Zarkov, Z., Bruffaerts, R., 2016. Mental disorders among college students in the World Health Organization World Mental Health Surveys. Psychol Med 46 (14), 2955-2970.

Auerbach, R.P., Mortier, P., Bruffaerts, R., Alonso, J., Benjet, C., Cuijpers, P., Demyttenaere, K., Ebert, D.D., Green, J.G., Hasking, P., Murray, E., Nock, M.K., Pinder-Amaker, S., Sampson, N.A., Stein, D.J., Vilagut, G., Zaslavsky, A.M., Kessler, R.C., Collaborators, W.W.-I., 2018. WHO World Mental Health Surveys International College Student Project: Prevalence and distribution of mental disorders. J Abnorm Psychol 127 (7), 623-638.

Boerema, A.M., Zoonen, K., Cuijpers, P., Holtmaat, C.J., Mokkink, L.B., Griffiths, K.M., Kleiboer, A.M., 2016. Psychometric Properties of the Dutch Depression Stigma Scale (DSS) and Associations with Personal and Perceived Stigma in a Depressed and Community Sample. PLoS One 11 (8), e0160740.

Burgard, T., Kasten, N., Bosnjak, M., 2019. Participation in online surveys in psychology. A meta-analysis., Research Synthesis 2019 conference. Leibniz Institute for Psychology Information, Dubrovnik, Croatia.

Busby Grant, J., Bruce, C.P., Batterham, P.J., 2016. Predictors of personal, perceived and self-stigma towards anxiety and depression. Epidemiol Psychiatr Sci 25 (3), 247-254.

Calear, A.L., Griffiths, K.M., Christensen, H., 2011. Personal and perceived depression stigma in Australian adolescents: magnitude and predictors. J Affect Disord 129 (1-3), 104-108.

Chatmon, B.N., 2020. Males and Mental Health Stigma. Am J Mens Health 14 (4), 1557988320949322.

Coppens, E., Van Audenhove, C., Scheerder, G., Arensman, E., Coffey, C., Costa, S., Koburger, N., Gottlebe, K., Gusmao, R., O'Connor, R., Postuvan, V., Sarchiapone, M., Sisask, M., Szekely, A., van der Feltz-Cornelis, C., Hegerl, U., 2013. Public attitudes toward depression and help-seeking in four European countries baseline survey prior to the OSPI-Europe intervention. J Affect Disord 150 (2), 320-329.

Corrigan, P.W., 2006. Mental Health Stigma as Social Attribution: Implications for Research Methods and Attitude Change. Clinical Psychology: Science & Practice 7 (1), 48-67.

Dietrich, S., Mergl, R., Rummel-Kluge, C., 2014. Personal and perceived stigmatization of depression: a comparison of data from the general population, participants of a depression congress and job placement officers in Germany. Psychiatry Res 220 (1-2), 598-603.

Ebert, D.D., Mortier, P., Kaehlke, F., Bruffaerts, R., Baumeister, H., Auerbach, R.P., Alonso, J., Vilagut, G., Martínez, K.I., Lochner, C., Cuijpers, P., Kuechler, A.M., Green, J., Hasking, P., Lapsley, C., Sampson, N.A., Kessler, R.C., 2019. Barriers of mental health treatment utilization among first-year college students: First cross-national results from the WHO World Mental Health International College Student Initiative. Int J Methods Psychiatr Res 28 (2), e1782.

Eisenberg, D., Downs, M.F., Golberstein, E., Zivin, K., 2009. Stigma and help seeking for mental health among college students. Med Care Res Rev 66 (5), 522-541.

Ferreira, T., Sousa, M., Salgado, J., 2018. Brief assessment of depression: Psychometric properties of the Portuguese version of the Patient Health Questionnaire (PHQ-9). The Psychologist: Practice & Research Journal 1 (2), 1-12.

Fischer, E.H., Turner, J.L., 1970. Development and Research Utility of an Attitude Scale. Journal of Consulting and Clinical Psychology 35 (1), 70-90.

Gil, I., 2015. Estigma e doença mental, Cadernos de Psiquiatria social e cultural, pp. 59-76.

Griffiths, K.M., Christensen, H., Jorm, A.F., 2008. Predictors of depression stigma. BMC Psychiatry 8, 25.

Griffiths, K.M., Christensen, H., Jorm, A.F., Evans, K., Groves, C., 2004. Effect of Web-Based Depression Literacy and Cognitive-Behavioural Therapy Interventions on Stigmatising Attitudes to Depression: Randomised Controlled Trial. The British Journal of Psychiatry(185), 342-349.

Gulliver, A., Griffiths, K.M., Christensen, H., 2010. Perceived barriers and facilitators to mental health help-seeking in young people: a systematic review. BMC Psychiatry 10 (113).

Hegerl, U., Wittenburg, L., Arensman, E., Van Audenhove, C., Coyne, J.C., McDaid, D., van der Feltz-Cornelis, C.M., Gusmao, R., Kopp, M., Maxwell, M., Meise, U., Roskar, S., Sarchiapone, M., Schmidtke, A., Varnik, A., Bramesfeld, A., 2009. Optimizing suicide prevention programs and their implementation in Europe (OSPI Europe): an evidence-based multi-level approach. BMC Public Health 9, 428.

Ibrahim, A.K., Kelly, S.J., Adams, C.E., Glazebrook, C., 2013. A systematic review of studies of depression prevalence in university students. J Psychiatr Res 47 (3), 391-400.

Jennings, K.S., Cheung, J.H., Britt, T.W., Goguen, K.N., Jeffirs, S.M., Peasley, A.L., Lee, A.C., 2015. How are perceived stigma, self-stigma, and self-reliance related to treatment-seeking? A three-path model. Psychiatr Rehabil J 38 (2), 109-116.

Khalema, N.E., Shankar, J., 2014. Perspectives on Employment Integration, Mental Illness and Disability, and Workplace Health. Advances in Public Health 2014.

Khubchandani, J., Brey, R., Kotecki, J., Kleinfelder, J., Anderson, J., 2016. The Psychometric Properties of PHQ-4 Depression and Anxiety Screening Scale Among College Students. Archives of Psychiatric Nursing 30 (4), 457-462.

Knaak, S., Mantler, E., Szeto, A., 2017. Mental illness-related stigma in healthcare: Barriers to access and care and evidence-based solutions. Healthcare Management Forum 30 (2), 111-116.

Kohls, E., Coppens, E., Hug, J., Wittevrongel, E., Van Audenhove, C., Koburger, N., Arensman, E., Szekely, A., Gusmao, R., Hegerl, U., 2017. Public attitudes toward depression and help-seeking: Impact of the OSPI-Europe depression awareness campaign in four European regions. J Affect Disord 217, 252-259.

Kosyluk, K.A., Al-Khouja, M., Bink, A., Buchholz, B., Ellefson, S., Fokuo, K., Goldberg, D., Kraus, D., Leon, A., Michaels, P., Powell, K., Schmidt, A., Corrigan, P.W., 2016. Challenging the Stigma of Mental Illness Among College Students. J Adolesc Health 59 (3), 325-331.

Löwe, B., Wahl, I., Rose, M., Spitzer, C., Glaesmer, H., Wingenfeld, K., Schneider, A., Brähler, E., 2010. A 4-item measure of depression and anxiety: Validation and standardization of the Patient Health Questionnaire-4 (PHQ-4) in the general population. Journal of Affective Disorders 122, 86-95.

McNair, B.G., Highet, N.J., Hickie, I.B., Davenport, T.A., 2002. Exploring the Perspectives of People Whose Lives Have Been Affected by Depression. The Medical Journal of Australia 196 (10), 69-76.

Milner, A., Kavanagh, A., King, T., Currier, D., 2018. The Influence of Masculine Norms and Occupational Factors on Mental Health: Evidence From the Baseline of the Australian Longitudinal Study on Male Health. Am J Mens Health 12 (4), 696-705.

Monteiro, S., Torres, A., Pereira, A., Albuquerque, E., Morgadinho, R., 2013. Preliminary validation study of a portuguese version of the patient health questionnaire (PHQ-9). European Psychiatry 28.

Moritz, D., Roberts, J.E., 2016. Individual Differences in Propensity to Socially Reject Depressed Others. Personality and Individual Differences 101, 500.

Musa, A., Ashraf, J., Tsai, F.J., Abolmagd, S., Liu, C., Hussain, H., Voslarova, E., Khalil, M.A., Wolitzky-Taylor, K.B., Lee, D., Sugar, J., Pendi, K., Lee, J., Abdelmaksoud, R., Adel, N., Baron, D., 2020. Depression Severity and Depression Stigma Among Students: A Survey of Universities in Five Countries. J Nerv Ment Dis 208 (11), 884-889.

Oliveira, A.R.F., Azevedo, S.M., 2014. Estigma na doença mental: estudo observacional. Revista Portuguesa de Medicina Geral e Familiar 30 (4), 227-234.

Pedersen, E.R., Paves, A.P., 2014. Comparing perceived public stigma and personal stigma of mental health treatment seeking in a young adult sample. Psychiatry Res 219 (1), 143-150.

Querido, A., Tomás, C., Carvalho, D., 2016. O estigma face à doença mental nos estudantes de saúde. Revista Portuguesa de Enfermagem de Saúde Mental(Spe. 3).

Rice, C.D., Howard, L., Leese, M., Jarrett, M., Thornicroft, G., 2009. Determinants of wanting to seek full versus part-time paid employment among people with severe mental illness. Journal of Mental Health 18:5, 424-432.

Sawyer, S.M., Azzopardi, P.S., Wickremarathne, D., Patton, G.C., 2018. The age of adolescence. The Lancet Child & Adolescent Health 2 (3), 223-228.

Schnyder, N., Panczak, R., Groth, N., Schultze-Lutter, F., 2017. Association between mental health-related stigma and active help-seeking: systematic review and meta-analysis. Br J Psychiatry 210 (4), 261-268.

Shannon, A., Goldberg, J.O., Flett, G.L., Hewitt, P.L., 2018. The relationship between perfectionism and mental illness stigma. Personality and Individual Differences 126, 66-70.

Sousa, T.V., Viveiros, V., Chai, M.V., Vicente, F.L., Jesus, G., Carnot, M.J., Gordo, A.C., Ferreira, P.L., 2015. Reliability and validity of the Portuguese version of the Generalized Anxiety Disorder (GAD-7) scale. Health Qual Life Outcomes 13, 50.

Telles-Correia, D., Marques, J.G., Gramaça, J., Sampaio, D., 2015. Stigma and Attitudes towards Psychiatric Patients in Portuguese Medical Students. Acta Médica Portuguesa 28 (6), 715-719.

Thornicroft, G., Chatterji, S., Evans-Lacko, S., Gruber, M., Sampson, N., Aguilar-Gaxiola, S., Al-Hamzawi, A., Alonso, J., Andrade, L., Borges, G., Bruffaerts, R., Bunting, B., de Almeida, J.M., Florescu, S., de Girolamo, G., Gureje, O., Haro, J.M., He, Y., Hinkov, H., Karam, E., Kawakami, N., Lee, S., Navarro-Mateu, F., Piazza, M., Posada-Villa, J., de Galvis, Y.T., Kessler, R.C., 2017. Undertreatment of people with major depressive disorder in 21 countries. Br J Psychiatry 210 (2), 119-124.

Twenge, J.M., Park, H., 2019. The Decline in Adult Activities Among U.S. Adolescents, 1976-2016. Child Dev 90 (2), 638-654.

Uehara, T., Takeuchi, K., Kubota, F., Oshima, K., Ishikawa, O., 2010. Annual transition of major depressive episode in university students using a structured self-rating questionnaire. Asia-Pacific Psychiatry 2 (2), 99-104.

Wei, Y., McGrath, P.J., Hayden, J., Kutcher, S., 2015. Mental health literacy measures evaluating knowledge, attitudes and help-seeking: a scoping review. BMC Psychiatry 15, 291.

World Health Organization, 2001. The WHO World Health Report 2001: New understanding - New hope. World Health Organization, Geneve.

World Health Organization, 2013. Mental Health Action Plan 2013-2020, pp. 1-44.

World Health Organization, 2017. Depression and Other Common Mental Disorders: Global Health Estimates, Geneva.

Xavier, S., Klut, C., Neto, A., Ponte, G.d., Melo, J.C., 2013. O Estigma da Doença Mental: Que Caminho Percorremos? Revista do Serviço de Psiquiatria do Hospital Prof. Doutor Fernando Fonseca, EPE 11 (2), 10-21.