Manuscript Number: NA

Article Summary Line: The loosening of the social distancing measures imposed by the Korean

government in May 2020 has resulted in the second wave of COVID-19 in the greater Seoul area

in the first two weeks of June, yielding reproduction numbers exceeding 3.0.

Running Title: Transmission potential of COVID-19 in South Korea

Keywords: Coronavirus; COVID-19; Republic of Korea; Seoul; severe acute respiratory

syndrome coronavirus 2; Coronavirus Infections; Reproduction number; Government

Title: Spatial and temporal variability in the transmission potential of COVID-19 in South Korea

including the second wave in the greater Seoul area, February to July 2020

Authors: Eunha Shim, Gerardo Chowell

Affiliations:

Department of Mathematics, Soongsil University, Seoul, Republic of Korea (E. Shim)

Department of Population Health Sciences, School of Public Health, Georgia State University,

Atlanta, USA (G. Chowell)

Abstract

In South Korea, 13,745 cases of COVID-19 have been reported as of 19 July, 2020. We used *EpiEstim R* package to investigate the time-varying reproduction numbers of the COVID-19 in the four most affected regions in South Korea: Seoul, Gyeonggi Province, Gyeongbuk Province, and Daegu. At the regional level, Seoul and Gyeonggi Province have experienced two waves with the first major peak of COVID-19 in early March, followed by the second wave in the first two weeks of June, with reproduction numbers in early May greater than 3.0. Gyeongbuk Province and Daegu are yet to experience a second wave of the disease, where the mean reproduction number reached values as high as 3.5-4.4. Our findings indicate that the loosening of the restrictions imposed by the government in May 2020 facilitated a second wave in the greater Seoul area.

Since the first COVID-19 cases reported in Wuhan, Hubei Province, China, in December

2019, more than 12 million cases of coronavirus disease (COVID-19), including 551,046 related

deaths, have been reported worldwide (1). In South Korea, the novel coronavirus was first

diagnosed in a 36-year-old Chinese woman who entered the country on 20 January 2020. South

Korea has since experienced two heterogenous waves of the disease with a total of 13,745 cases

including 295 deaths as of 19 July (2).

In the early phase of the COVID-19 outbreak in South Korea, public health authorities

primarily conducted strict contact tracing and isolation of confirmed cases as well as put under

quarantine orders those suspected with the novel coronavirus (3). As the number of COVID-19

cases increased, Korean public health authorities set the alert to the highest level (Level 4) on 23

February and required the population to report any symptoms related to COVID-19 for further

screening and testing. In addition, the country rapidly adopted a "test, trace, isolate, and treat"

strategy that has been deemed effective in stamping out localized outbreaks of the novel

coronavirus (2). However, the total number of confirmed cases in South Korea spiked from 31

cases on 18 February to 433 on 22 February. According to the Korea Centers for Disease Control

and Prevention (KCDC), this sudden jump was mainly attributed to a super-spreader (the 31[st]

case) who had participated in a religious gathering of attendees of the Shincheonji Church of

Jesus in Daegu (2). Superspreading events occurred in the Daegu and Gyeongbuk provincial

regions, leading to more than 5,210 secondary COVID-19 cases in Korea (2, 4). These events

facilitated sustained transmission chains, with 38% of the cases in the country being associated

with the church cluster in Daegu (5).

On 8 March, the KCDC announced that 79% of the confirmed cases were tied to clusters of

infections. Case clusters started to accumulate from churches in the Seoul Capital Area, and on

17 March, 79 church attendees developed COVID-19 after attending a service at the River of
Grace Community Church. Notwithstanding social distancing orders put forward by the
government, some churches held services, which led to new clusters of infections. For instance,
the Manmin Central Church in Seoul was involved in one of the clusters, with 41 infections
linked to a gathering in early March; other church clusters including one with 50 cases involving
the SaengMyeongSu Church in Gyeonggi Province (6).

As infection rates rose outside Korea, the number of imported cases increased, resulting in
476 imported out of 9,661 total cases as of 30 March. Consequently, as of 1 April, the KCDC
implemented self-quarantine measures for travellers from Europe or the U.S. (2). In addition,
incoming travellers that showed symptoms but tested negative, as well as asymptomatic short-
term visitors, were ordered to follow 2-week quarantines in government facilities (2).

After a sustained low incidence period with fewer than 20 cases per day, the government
eased its strict nationwide social distancing guidelines on 6 May, with phased reopening of
schools starting mid-May, 2020. However, a new cluster tied to nightclubs in Itaewon emerged in
central Seoul in early May, resulting in a resurgence of cases that led to a second wave in the
greater areas of Seoul. As of 29 May, the number of cases that were linked to this cluster had
reached 266 (2). Accordingly, the Seoul city government ordered all clubs, bars, and other
nightlife establishments in the city to close indefinitely (2). Moreover, one more cluster emerged
from an e-commerce warehouse in the Gyeonggi Province, resulting in 108 cases as of 30 May.

In the last week of May, the number of daily new cases increased and varied between 40 and
80 (2). Following this highest spike in the number of new COVID-19 infections in nearly 2
months, public health authorities reimplemented strict lockdown measures in Seoul and ordered
schools closed one more time across the nation. In June, it was announced that the strict social

distancing campaign would be indefinitely extended as a preventive measure in Seoul, Incheon,

and Gyeonggi Province; however, phased reopening of schools was initiated on May 20. It was

reported by the KCDC that a holiday weekend in early May triggered a new wave of infections

focused in the greater Seoul area, the so-called second wave of COVID-19 in South Korea (7). In

Seoul, the average number of new daily cases from 4 June to 17 June was 43 cases (2). In July,

sporadic clusters of infections across the country continued to occur, most of them related to

religious facilities and door-to-door salespeople, especially in the densely populated Seoul region

and adjacent areas. Therefore, since 10 July, the government banned churches from organising

small gatherings other than regular worship services (2).

To estimate the regional and temporal variability in the reproduction number of COVID-19

in South Korea, including the second wave concentrated in the greater Seoul areas, we analysed

the spatial-temporal progression of the epidemic in the country from mid-February to mid-July

2020. Here our focus is on estimating and interpreting the effective reproduction number R_t, a

metric that quantifies the time-dependent transmission potential, incorporating the effect of

control measures, susceptible depletion, and behavioural changes. This key epidemiological

parameter, R_t, represents the average number of secondary cases per case if conditions remain as

they were at time t. In this report, we estimated the effective reproduction number involving two

epidemic waves of the COVID-19 epidemic in South Korea by employing the time series of

cases by date of symptoms onset for the four most affected Korean regions: Seoul, Gyeonggi

Province, Gyeongbuk Province, and Daegu. We also discuss the spatial-temporal variability of

the reproduction number in terms of the public health policies that were put in place by the

Korean government.

Methods

Data

We collected the daily series of confirmed local COVID-19 cases in South Korea from 20

January to 19 July, which were published by national and local public health authorities,

including city or provincial departments of public health in South Korea (8). We focused our

analysis on the regions with the highest disease burden including Seoul, Gyeonggi Province,

Gyeongbuk Province, and Daegu (Figure 1).

Imputing the date of onset

To estimate epidemic growth rates with more accuracy, the epidemic curve should be

analysed according to the date of symptom onset rather than the date of reporting because

reporting delays can fluctuate substantially over the course of the epidemic. For the COVID-19

data in Korea, however, the symptom-onset date is only available for the 732 cases reported in

Gyeonggi Province. Therefore, we imputed the onset date for the cases with missing data using

the empirical distribution of the reporting delay from onset to diagnosis as in a previous study (9).

Specifically, we reconstructed 300 epidemic curves according to symptom-onset date, from

which we derived the mean incidence curve of local case incidence (9, 10). For the calculation of

$R_t(t)$, the mean incidence curve estimated based on symptom-onset date was used for the regions

of our interest (i.e., Seoul, Gyeonggi Province, Gyeongbuk Province, and Daegu) (Figure 2).

Calculation of R_t

To estimate the temporal variation in the transmission potential, we estimated $R_t(t)$ using the

distribution of the serial interval (SI) for COVID-19 and calculated the transmission ability (11).

The SI is defined as the period between the time of symptom onset in two successive cases in a

chain of transmission and its distribution is denoted by a probability distribution w_s, the

infectivity profile of the infected individual. The distribution of SI, w_s, is dependent on the time

since infection (s) but independent of calendar time (t). For example, an individual would be the

most infectious at time s when w_s is the largest. In addition, the infectiousness of a patient is a

function of the time since infection and is proportional to w_s if we set the timing of infection in

the primary case as the time zero of w_s and assume that the generation interval equals the SI.

Individual biological factors such as pathogen shedding or symptom severity can affect the

distribution w_s. The SI was assumed to follow a gamma distribution with a mean of 4.8 days and

a standard deviation of 2.3 days (12).

$R_t(t)$ can be estimated by the ratio of the number of new infections generated at time step t

(I_t) to the total infectiousness of infected individuals at time t, given by $\sum_{s=1}^{t} I_{t-s} w_s$ (13, 14).

Here, $\sum_{s=1}^{t} I_{t-s} w_s$ indicates the sum of infection incidence up to time step $t-1$, weighted by the

infectivity function w_s. Steady values of R_t above one indicate sustained disease transmission,

whereas values less than one indicate that the number of new cases is expected to follow a

declining trend. Analytical estimates of R_t were obtained within a Bayesian framework using

EpiEstim R package in *R* language version 3.6.3 (*R* Foundation for Statistical Computing, Vienna,

Austria) (12). R_t was estimated at 7-day intervals, and we reported the median and 95% credible

interval (CrI).

Results

As of 19 July, Seoul has reported a total of 1,474 cases (10.7% of the total reported in South

Korea), including 323 imported cases and 10 deaths, yielding an incidence rate estimated at 151

cases per million. In Seoul, the first peak based on the estimated dates of symptom onset

occurred during the second week of March (8^{th}–14^{th} March), with 18 new cases reported each

day as the number of new cases linked to a Guro-gu call centre was rising. Based on the

estimated dates of symptom onset, the 7-day moving average of daily cases reached 19 on 9

March (Figure 3) whereas R_t reached its highest value at 2.9 (95% CrI: 1.6-4.7) on 19 February

and stayed above one until 6 March (Figure 3).

After its first peak in February, the number of daily new cases by date of symptoms onset in

Seoul gradually declined, dropping below five on 1 April and staying under five new cases per

day for about a month (Figure 3). However, in early May, despite a steady decline in imported

cases, locally transmitted infections surged throughout the Seoul metropolitan area with case

clusters traced to clubs, churches, and sports facilities. Therefore, R_t increased, reaching 3.0 (95%

CrI: 1.6-5.0) on 4 May 2020.

The number of cases continued to increase, and in the first week of June, the average daily

number of confirmed COVID-19 cases in the capital surpassed the previous high point reached

in the middle of March. The major clusters in Seoul were linked to nightclubs (139 cases), the

Guro-gu call centre (99 cases), Manmin Central Church (41 cases), Richway (97 cases),

Yangcheon-gu table tennis club (41 cases), and Newly Planted Church in the Seoul Metropolitan

Region (37 cases) as of 18 June. On 14 June, the average R_t in the capital, which reflects the

average number of people infected by a patient, dropped below one (95% CrI: 0.8-1.2), implying

that the spread of the virus had slowed down substantially in the city (Figure 3). As of 15 July

2020, the R_t in Seoul was estimated at 0.9 (95% CrI: 0.7-1.2), straddling the epidemic threshold

of 1.0, and suggesting potential for further transmission of the virus.

Gyeonggi Province

Gyeonggi Province (literally meaning the "province surrounding Seoul") is located in the

western central region of Korea and is the most populous province in South Korea with a

population of 13.5 million people. In Gyeonggi Province, the daily number of new cases by date

of symptoms onset during the last weeks of February was 6.3 on average (Figure 4). Accordingly,

the first peak of R_t occurred on 22 February, reaching 8.9 (95% CrI: 4.8-14.2). In the second

week of March, South Korea recorded continuous drops in the number of daily new infections as

massive testing of the followers of a religious sect in the south-eastern city of Daegu, the

epicentre of COVID-19, was nearing its end; thereafter, the number of cases in Gyeonggi

Province gradually decreased.

However, clusters of infections in Gyeonggi Province raised concerns about further

community spread, with a resurgence of cases in the province occurring in late May and resulting

in the highest peak in early June. Between 1 June and 13 June, an average of 14 new cases were

reported each day in Gyeonggi Province. The second peak of R_t in the region occurred on 12

May, with an estimated value at 4.8 (95% CrI: 3.0-7.0). Since its second peak, R_t gradually

decreased (Figure 3); however, a series of sporadic clusters have continued to occur. The major

clusters in Gyeonggi Province included Grace River Church (67), Coupang warehouse (67),

nightclubs (59), Richway (58), Uijeongbu St. Mary's Hospital (50), Guro-gu call centre/Bucheon

SaengMyeongSu Church (50), door-to-door sales in the Seoul Metropolitan Region (32), and

Yangcheon-gu sports facility (28). As of 19 July, the number of local cases in Gyeonggi Province

was 1,027 (10.4% of the total reported cases in South Korea), including 29 deaths, with R_t

estimated at 0.8 (Figure 4). The incidence rate in the province was estimated at 108 per million.

Gyeongbuk Province

The first case in the Sincheonji cult cluster (the largest COVID-19 cluster in South Korea)

appeared on 18 February, resulting in sustained transmission chains, with 39% of the cases

associated with the church cluster in Gyeongbuk Province. Therefore, the virus alert level was

raised to "red" (the highest level) on 23 February, and health authorities focused on halting the

spread of the virus in Daegu and Gyeongbuk Provinces. Figure 5 shows that the peak of the

epidemic occurred in the first week of March (with a reproduction number greater than one until

9 March) (Figure 5). As of 18 July, the number of cases in Gyeongbuk Province was 1,393,

including 54 deaths. Among these cases, 566 were related to the Shincheonji cluster. The

incidence rate in Gyeongbuk Province was 523 per million, accounting for 10.2% of all

confirmed cases in South Korea (2). The major clusters in Gyeongbuk Province were linked to

Cheongdo Daenam Hospital (119 cases), Bonghwa Pureun Nursing Home (68 cases), Gyeongsan

Seo Convalescent Hospital (66 cases), pilgrimage to Israel (41 cases), Yecheon-gun (40 cases),

and Gumi Elim Church (11 cases).

City of Daegu

The epicentre of the South Korean COVID-19 outbreak has been identified in Daegu, a city

of 2.5 million people, approximately 150 miles south-east of Seoul. The rapid spread of COVID-

19 in Daegu was attributed to a superspreading event in a religious group called Shincheonji,

resulting in an explosive outbreak with 4,511 infections in the city of Daegu (Figure 6). Other

major clusters in Daegu included the second Mi-Ju Hospital (196 cases), Hansarang

Convalescent Hospital (124 cases), Daesil Convalescent Hospital (101 cases), and Fatima

Hospital (39 cases). Daegu was the most severely affected area in South Korea with 6,932

cumulative cases as of 19 July, accounting for 51.0% of all confirmed cases. According to our

model, the number of new cases based on the onset of symptoms was estimated to be the highest

on 27 February; the number gradually decreased thereafter. Accordingly, the estimated R_t was

above two until 27 February and dropped below one on 5 March, although recent sporadic

infections caused the R_t to fluctuate around one (Figure 6).

Discussion

The estimates of the transmission potential of COVID-19 in Korea displayed substantial

spatiotemporal variation. Indeed, several factors influence the value of the reproduction number,

including the transmissibility of an infectious agent, individual susceptibility, individual contact

rates, and control measures. Our results indicate that the effective reproduction number for

COVID-19 declined to low levels after the first wave and straddled the epidemic threshold of 1.0

in March and April suggesting that social distancing measures had a significant effect on

mitigating the spread of the novel coronavirus. In prior studies, the early national R_t for South

Korea has been estimated at 2.9 (95% CrI 2.0-3.9) in February (15) or 2.6 (95% CI: 2.3-2.9) in

March (16). These values are in good agreement with our estimates.

Our results suggest that South Korea has experienced two spatially heterogenous waves of

the novel coronavirus. At the regional level, Seoul and Gyeonggi Province have experienced two

waves whereas Daegu and Gyeongbuk Provinces are yet to experience the second wave of the

disease. The highest epidemic peak occurred in Daegu and Gyeongbuk Province in late February

and early March, with R_t being estimated at 4.4 (95% CrI: 2.6-6.6) and 3.5 (95% CrI: 0.9-7.3),

respectively. Similarly, in Gyeonggi Province and Seoul, the first wave was observed in late

February and early March, respectively. However, sporadic clusters of infections appeared in

Seoul and near Gyeonggi Province, immediately after the government eased its strict nationwide

social distancing guidelines on May 6. This resurgence of infections in Seoul and Gyeonggi

Province (i.e., the province surrounding Seoul) after a sustained period with fewer than 5 cases

per day in each region, led to the second epidemic wave with sub-exponential growth dynamics.

Accordingly, our findings revealed sustained local transmission in Seoul and Gyeonggi Province,

with the estimated reproduction number above one until the end of May. In late May, the country

implemented two weeks of tougher virus prevention guidelines for the metropolitan area, with

measures including shutting down public facilities and regulating bars and karaoke rooms. In the

second week of June, South Korea decided to indefinitely extend a period of tougher social

distancing measures, as nearly all locally transmitted cases were in the metropolitan area.

Although Korea has a relatively low number of reported cases compared with other

countries including the U.S. and China, it is believed that South Korea is currently experiencing

yet another resurgence of the virus. Originally, South Korean authorities predicted a resurgence

of the virus in the fall or winter; however, this possible second wave started in and around Seoul,

which, with 51.6 million inhabitants, accounts for about half of the entire population of the

country. Secondary waves of the disease can result from multiple factors, including easing of

travel restrictions and resuming social activities especially in the high population density areas of

Seoul and Gyeonggi Province. Furthermore, a substantial proportion of COVID-19 cases are

asymptomatic (17); thus, they are not detected by surveillance systems, resulting in the

underestimation of the epidemic growth curve. It was also recently reported that individuals aged

20-39 years in South Korea drove the COVID-19 epidemic throughout society with multiple

rebounds, and an increase in infection among the elderly was significantly associated with an

elevated transmission risk among young adults (18).

Our study is not exempted of limitations including the lack of dates of symptoms onset for

all of the cases, relying on a statistical reconstruction of the epidemic curve by dates of

symptoms onset as in a previous study (9). Overall, using most up-to-date epidemiological data

from South Korea, our study highlights the effectiveness of strong control interventions in South

Korea and emphasizes the need to maintain firm social distancing and contact tracing efforts to

mitigate the risk of additional waves of the disease.

Acknowledgments

This work was supported by the National Research Foundation of Korea (NRF) grant

funded by the Korea government (MSIT) [No. 2018R1C1B6001723] to E.S. For G.C., this work

was supported by RAPID NSF No. 2026797.

Author Bio

Dr. Shim is an Associate Professor of Mathematics at Soongsil University, Seoul, South

Korea. Her research interests include using mathematics and quantitative tools to understand

infectious disease dynamics and to evaluate the effectiveness of public health interventions.

Dr. Chowell is a professor of epidemiology and biostatistics and chair of the department of population health sciences at Georgia State University School of Public Health, Atlanta, Georgia. As a mathematical epidemiologist, he studies the transmission dynamics of emerging infectious diseases, such as Ebola, MERS, and SARS.

References

1.   WHO. Coronavirus disease (COVID-2019) situation reports.

2.   KCDC. The Updates of COVID-19 in Republic of Korea. Centers for Disease Control and

Prevention Korea; 2020.

3.   Covid-19 National Emergency Response Center E, Case Management Team KCfDC, Prevention.

Contact Transmission of COVID-19 in South Korea: Novel Investigation Techniques for Tracing

Contacts. Osong Public Health Res Perspect. 2020 Feb;11(1):60-3.

4.   Ryall J. Coronavirus: Surge in South Korea virus cases linked to church 'super-spreader'. The

Telegraph. 2020 Feb 20, 2020.

5.   Shim E, Tariq A, Choi W, Lee Y, Gerardo C. Transmission potential and severity of COVID-19

in South Korea. International Journal of Infectious Diseases. 2020 Mar 10, 2020.

6.   Park CK. Coronavirus cluster emerges at another South Korean church, as others press ahead

with Sunday services. South China Morning Post. 2020 Mar 30, 2020.

7.   Coronavirus: South Korea confirms second wave of infections. BBC. 2020 June 20.

8.   KCDC. The updates on COVID-19 in Korea as of 25 February. Seoul, Korea: Korea Centers for

Disease Control and Prevention 2020.

9.   Shim E, Tariq A, Choi W, Lee Y, Chowell G. Transmission potential and severity of COVID-19

in South Korea. International journal of infectious diseases : IJID : official publication of the International

Society for Infectious Diseases. 2020 Apr;93:339-44.

10.   Tariq A, Roosa K, Mizumoto K, Chowell G. Assessing reporting delays and the effective

reproduction number: The Ebola epidemic in DRC, May 2018-January 2019. Epidemics. 2019

Mar;26:128-33.

11.   Wallinga J, Teunis P. Different epidemic curves for severe acute respiratory syndrome reveal

similar impacts of control measures. American journal of epidemiology. 2004 Sep 15;160(6):509-16.

12.   Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to estimate time-

varying reproduction numbers during epidemics. American journal of epidemiology. 2013 Nov

1;178(9):1505-12.

13.   Chong KC, Zee BCY, Wang MH. Approximate Bayesian algorithm to estimate the basic

reproduction number in an influenza pandemic using arrival times of imported cases. Travel Med Infect

Dis. 2018 May - Jun;23:80-6.

14.   Fraser C. Estimating individual and household reproduction numbers in an emerging epidemic.

PloS one. 2007 Aug 22;2(8):e758.

15.   Ryu S, Ali S, Jang C, Kim B, Cowling BJ. Transmission dynamics of coronavirus disease 2019

outside of Daegu-Gyeongsangbuk provincial region in South Korea. 2020.

16.   Zhuang Z, Zhao S, Lin Q, Cao P, Lou Y, Yang L, et al. Preliminary estimates of the reproduction

number of the coronavirus disease (COVID-19) outbreak in Republic of Korea and Italy by 5 March

2020. International journal of infectious diseases : IJID : official publication of the International Society

for Infectious Diseases. 2020 Jun;95:308-10.

17.   Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of

coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama,

Japan, 2020. Euro surveillance : bulletin europeen sur les maladies transmissibles = European

communicable disease bulletin. 2020 Mar;25(10).

18.   Yu X, Duan J, Jiang Y, Zhang H. Distinctive trajectories of the COVID-19 epidemic by age and

gender: A retrospective modeling of the epidemic in South Korea. International journal of infectious

diseases : IJID : official publication of the International Society for Infectious Diseases. 2020 Jul

2;98:200-5.

Address for correspondence: Eunha Shim, Department of Mathematics, Soongsil University,

369 Sasngdo-ro, Dongjak-gu, Seoul, 06978 Republic of Korea; email: alicia@ssu.ac.kr

Figure 1. Map depicting the location of Seoul, Gyeonggi Province, Gyeongbuk Province, and

Daegu.

Figure 2. Timeline of confirmed cases of COVID-19 in Seoul, Gyeonggi Province, Gyeongbuk

Province, and Daegu. The daily number of COVID-19 cases by date of reported and by date of

symptom onset are shown. The empirical distribution of reporting delays from the onset to

diagnosis for 732 cases were used to impute the missing dates of onset for the remainder of the

cases with missing data.

Figure 3. The epidemic trajectory of coronavirus disease 2019 in Seoul, as of 19 July 2020.

Upper panel: The epidemic curve shows the daily number of new cases by date of symptom

onset in Seoul. The onset date for cases with missing data were imputed based on the empirical

distribution of the reporting delay from the onset to diagnosis. Lower panel: Real-time estimates

of the time-varying reproduction number (R_t) in Seoul. The solid line indicates the daily

estimated R_t and the grey area indicates the 95% credible interval of R_t. The dotted line indicates

the epidemic threshold of $R_t =1$.

Figure 4. The epidemic trajectory of coronavirus disease 2019 in Gyeonggi Province, as of 19

July 2020. Upper panel: The epidemic curve shows the daily number of new cases by date of

symptom onset in Gyeonggi Province. The onset date for cases with missing data were imputed

based on the empirical distribution of the reporting delay from the onset to diagnosis. Lower

panel: Real-time estimates of the time-varying reproduction number (R_t) in Gyeonggi Province.

The solid line indicates the daily estimated R_t and the grey area indicates the 95% credible

interval of R_t. The dotted line indicates the epidemic threshold of $R_t = 1$.

Figure 5. The epidemic trajectory of coronavirus disease 2019 in Gyeongbuk Province, as of 19

July 2020. Upper panel: The epidemic curve shows the daily number of new cases by date of

symptom onset in Gyeongbuk Province. The onset date for cases with missing data were

imputed based on the empirical distribution of the reporting delay from the onset to diagnosis.

Lower panel: Real-time estimates of the time-varying reproduction number (R_t) in Gyeongbuk

Province. The solid line indicates the daily estimated R_t and the grey area indicates the 95%

credible interval of R_t. The dotted line indicates the epidemic threshold of $R_t = 1$.

Figure 6. The epidemic trajectory of coronavirus disease 2019 in Daegu, as of 19 July 2020.

Upper panel: The epidemic curve shows the daily number of new cases by date of symptom

onset in Daegu. The onset date for cases with missing data were imputed based on the empirical

distribution of the reporting delay from the onset to diagnosis. Lower panel: Real-time estimates

of the time-varying reproduction number (R_t) in Daegu. The solid line indicates the daily

estimated R_t and the grey area indicates the 95% credible interval of R_t. The dotted line indicates

the epidemic threshold of $R_t = 1$.

Seoul

Gyeongi
Province

Gyeongbuk
Province

Daegu

Seoul

Daegu