Title: Anatomy of digital contact tracing: role of age, transmission setting, adoption and case detection

Short title: Anatomy of digital contact tracing

Jesús A. Moreno López[1,2], Beatriz Arregui García[1,2], Piotr Bentkowski[1], Livio Bioglio[3], Francesco Pinotti[1], Pierre-Yves Boëlle[1], Alain Barrat[4,5], Vittoria Colizza[1], Chiara Poletto[1]*

[1]INSERM, Sorbonne Université, Pierre Louis Institute of Epidemiology and Public Health, Paris, France

[2]Instituto de Física Interdisciplinar y Sistemas Complejos IFISC (CSIC-UIB), Campus UIB, Palma de Mallorca, Spain

[3]Department of Computer Science, University of Turin, Turin, Italy

[4]Aix Marseille Univ, Université de Toulon, CNRS, CPT, Turing Center for Living Systems, Marseille, France

[5]Tokyo Tech World Research Hub Initiative (WRHI), Tokyo Institute of Technology, Tokyo, Japan

*chiara.poletto@inserm.fr

Abstract

The efficacy of digital contact tracing against COVID-19 epidemic is debated: smartphone

penetration is limited in many countries, non-uniform across age groups, with low coverage

among elderly, the most vulnerable to SARS-CoV-2. We developed an agent-based model to

precise the impact of digital contact tracing and household isolation on COVID-19

transmission. The model, calibrated on French population, integrates demographic, contact-

survey and epidemiological information to describe the risk factors for exposure and

transmission of COVID-19. We explored realistic levels of case detection, app adoption,

population immunity and transmissibility. Assuming a reproductive ratio $R = 2.6$ and 50%

detection of clinical cases, a ~20% app adoption reduces peak incidence by ~35%. With $R =$

1.7, >30% app adoption lowers the epidemic to manageable levels. Higher coverage among

adults, playing a central role in COVID-19 transmission, yields an indirect benefit for elderly.

These results may inform the inclusion of digital contact tracing within a COVID-19 response

plan.

Introduction

Intervention measures aiming at preventing transmission have been key to control the first

wave of the COVID-19 pandemic. Many countries have adopted lockdown and strong social

distancing during periods of intense epidemic activity to suppress the epidemic and reduce

hospital occupancy below saturation levels (*1*, *2*). Due to their huge economic and societal

costs these interventions can only be implemented for a limited amount of time. The building

of population immunity has been slow (*3–5*), so that new waves are possible after temporary

lockdowns and lifting of restrictions. Sustainable strategies are required to maintain the

epidemic under control while enabling the close-to-normal functioning of the society.

Widespread testing, case finding and isolation, contact-tracing, use of face masks and

enhanced hygiene are believed to be crucial components of these strategies.

Contact-tracing aims to avoid transmission by isolating at an early stage only those

individuals who are infectious or potentially infectious, in order to minimize the societal costs

associated to isolation. Considerable resources are therefore directed at improving

surveillance capacities to allow efficient and rapid investigation and isolation of cases and

their contacts. To enhance tracing capacities, the use of digital technologies has been

proposed, leveraging the wide-spread use of smartphones. Therefore, proximity-sensing

applications have been designed and made available – e.g. in Australia, France, Germany,

Iceland, Italy, Switzerland – to automatically trace contacts, notify users about potential

exposure to COVID-19 and invite them to isolate.

Empirical studies of the impact of these digital applications are however limited (*6–8*), and

the utility of this intervention is debated. Some built-in features make it more efficient than

manual contact tracing: it is automated, reducing the burden of manual contact tracing and

limiting recall bias; it is faster, as information can be transmitted in real time. However,

coverage is uneven. In particular, most children and elderly do not own a smartphone or are

less familiar with digital technologies. The overall adoption of the app among smartphone

owners will also be a limiting factor, as well as the fraction of cases actually triggering the

alert to the contacts and the adherence to isolation of the app adopters who receive an alert.

These variables must be gauged in light of the risk factors for exposure and transmission

driving the COVID-19 epidemic. First, individuals of different age contribute differently to the

transmission dynamics of COVID-19. Younger individuals tend to have more contacts than

adults or the elderly. On the other hand, a marked feature of COVID-19 is the strong age

imbalance among cases (*9–13*), that may be explained by both a reduced susceptibility (*9,*

10) and an increased rate of subclinical infections in children compared to adults (*10, 11, 13*).

As subclinical cases are harder to detect, this implies that identification of cases and of their

contacts may be dependent on age. Second, SARS-CoV-2 transmission risk varies

substantially by setting. Transmissions were registered predominantly in households, in

specific workplaces and in the community (linked to shopping centers, meals, parties, sport

classes, etc.) (*14*, *15*). This is due, at least in part, to the higher risk of contagion of crowded

and indoor environments (*14–16*). Notably, contacts occurring in the community are also the

ones more affected by recall biases, thus more difficult to trace with manual contact tracing.

Several modelling studies have quantified the impact of contact tracing (*17–26*), with some of

them addressing specific aspects of digital contact tracing (*18–23*). Still the interplay between

age and setting heterogeneity in determining the efficacy of this intervention is largely

unexplored. Here we provide a systematic exploration of the different variables at play. We

considered France as a case study and integrated different sources of data to realistically

describe the French population, in terms of its demography and social contact behavior. We

accounted for the dynamics of contacts according to age and setting, and for the setting-

specific risk of transmission. We used COVID-19 epidemiological characteristics for

parametrization. We then modelled case detection and quarantining, isolation of their

household contacts and digital contact tracing, under different hypotheses of potential

reduction in transmissibility due to other effects (e.g. face-masks and increased hygiene). We

quantified the impact of digital contact tracing on the whole population and on different

population groups and settings, as a function of several variables such as the rate of app

adoption, the probability of detection of clinical and subclinical cases, population immunity,

compliance to isolation and transmission potential. Our results provide quantitative

information regarding the impact of digital contact tracing within a broader response plan.

90   Results

91   Dynamic multi-setting contact network

We modelled the French population integrating available demographic and social-contact

data. We collected population statistics on age, household size and composition (Figure 1 A,

B), workplace and school size, smartphone penetration (Figure 1 E), and commuting fluxes.

Then, by following standard approaches in the literature (*27*, *28*) individuals were created in-

silico with a given age and assigned to a municipality, a household, and a workplace/school

according to the statistics. Smartphones were assigned to individuals depending on their age

according to available statistics on French users (Figure 1 E) (*29*). Overall smartphone

penetration was 64%, that represents the upper bound limit of app adoption in the population

– reached when 100% of individuals owning a smartphone download the app. This synthetic

population reproduced the location statistics of individuals in different settings, yielding the

basis of a multi-setting network of daily face-to-face contacts in household, school,

workplace, community and transport (Figure 1 H) (*30–32*). We parametrized the network

from a social contact survey providing information on contacts by age and setting (*33*)

(Figure 1 C, D). As contacts may occur repeatedly, we associated an activation rate to each

contact and sampled each day contacts based on their activation rate (Figure 1 G). We

imposed that 35% of the contacts registered during one day occur with daily frequency, as

found in (*33*). Figure 1 F and I show that the features of the resulting daily contact network

matched the data: the distribution of the number of contacts was right-skewed as the

empirical one reported in (*33*) and the contact matrix showed age assortativity and the

characteristic parent-children (off-diagonal) contact pattern. As a case study we restricted our

study to a municipality with a population size of ~100,000 individuals (see Material and

Methods and Supplementary Material for additional details).

COVID-19 epidemic dynamics

We modelled coronavirus transmission and outcome as shown in Figure 2 A, B. Individuals

could be susceptible, S, exposed, E, pre-symptomatic preceding subclinical infection, $I_{p,sc}$,

pre-symptomatic preceding clinical infection, $I_{p,c}$, subclinically infectious, I_{sc}, clinically

infectious, I_c, and recovered, R. Subclinical cases had symptoms that ranged from no

symptoms to mild and continued their normal activity throughout the infectious period.

Clinical cases had moderate to critical symptoms and stayed at home after the onset of

symptoms (*11*, *13*) – we did not consider hospitalization. Individuals in compartments $I_{p,sc}$,

$I_{p,c}$, I_{sc}, I_c transmitted the infection, with subclinical individuals characterized by a lower risk

of transmission than clinical ones (see Material and Methods). We accounted for the

heterogeneous susceptibility and clinical manifestation by age as parametrized from (*9, 13*)

(Table 1). In order to parametrize the infection's natural history, we combined evidence from

epidemiological and viral shedding studies. We used 5.2 days for the incubation period (*34*),

2.3 days for the average length of the pre-symptomatic phase (*35*), and 7 days on average

for the infectivity period after symptoms' onset (*35*).

We first simulated an uncontrolled epidemic assuming transmission levels corresponding to

$R_0 = 3.1$, within the range of values estimated for COVID-19 in France at the early stage of

the pandemic (*1, 36*). The generation time resulting from our model and parameters had

mean value of 6.0 days (95% CI [2,17]), in agreement with epidemiological estimates (*11, 35,*

37). Figure 2 C and D show the repartition of cases among age groups and settings at the

early stage and during the whole course of the epidemic. Age-specific infection probability

was higher among young adults, while clinical infections were shifted towards older

population with respect to the overall (clinical and subclinical) cases, as noted in previous

observational and modelling works (*10*). The age profile changed in time with children

infected later as the epidemic unfolded (*10, 38*). Transmissions occurred predominantly in

household and workplaces followed by the community setting (*14*).

Contact tracing

We quantified the impact of combined household isolation and digital contact tracing

considering the possible scenario of a new epidemic wave emerging after the release of strict

lockdown measures in the country. We thus assumed some level of immunity to the virus –

exploring a range from 0 to 15% of the population. We considered interventions based on the

use of digital contact tracing, coupled with testing and isolation of clinical cases and

households. 50% of individuals with clinical symptoms were assumed to get tested after

consulting a doctor and to isolate if positive. Higher and lower percentages were also

considered.

Case tracing was assumed to start when a case with clinical symptoms tested positive and

was isolated, with an average delay of ~1 day. Household members were also invited to

isolate – we assumed that 90% of them accepted to isolate and that their isolation occurred

at the same time as the detected case. If the index case had the app installed, the contacts

he/she registered in the previous $D = 7$ days were notified and could decide to isolate with a

compliance probability of 90% – lower values of compliance were also explored. Note that

only contacts occurring between individuals who both use the app can be registered, so only

app adopters can be notified. We explored several levels of app adoption in the population.

In addition to the detection of clinical cases, we assumed that a proportion of subclinical

cases was also identified. These may be cases with very mild, unspecific symptoms who

decided to get tested as part of vulnerable groups (i.e. co-morbidity) or because highly

exposed to the infection (health care professionals). We hypothesized this proportion to be

small in the baseline scenario (5%), and we later varied it up to 45%. Isolated individuals

resumed normal daily life if infection was not confirmed. We took 7 days as the time needed

for being confirmed negative because multiple tests and some delay since the exposure are

needed for a negative result to be reliable. Infected individuals got out of quarantine after 14

days unless they still have clinical symptoms after the time is passed. They may, however,

decide to drop out from isolation each day with a probability of 2% if they don't have

symptoms (*21, 26*).

Figure 3 summarizes the effect of the interventions. We compared the uncontrolled scenario

($R = R_0 = 3.1$) with scenarios where the transmissibility is reduced due to the adoption of

barrier measures (R down to 1.5). We also assumed 10% of the population to be immune to

the infection (*36*). Panels A-C shows the results for $R = 2.6$ and $R = 1.7$. With $R = 2.6$ (Figure

3 A, C), the relative reduction of peak incidence due to household isolation only would be

27%. The inclusion of digital contact tracing would increase the relative reduction to 35% with

~20% app adoption, and to 66% with ~60% app adoption – i.e., 90% of individuals owning a

smartphone use the app. This corresponds to an additional mitigation effect ranging from

30% to 144% provided by contact tracing compared to household isolation only. With $R = 1.7$

(Figure 3 B, C), we find that ~20% app adoption would reduce the peak incidence by 45%

(additional mitigation effect of 25%), while the reduction would reach 89% in a scenario of

~60% app adoption (additional mitigation effect of 147%). According to the projections in

Ref.(*1*), intensive care units occupation would remain below the saturation level with

incidence below 0.4 /1000 hab. In the scenario with $R = 1.7$, this would be reached with app

adoption greater than ~30% (grey dashed line in Figure 3 B). Stronger reductions could be

obtained with more efficient detection of clinical cases (Figure 3 E, H, obtained with $R = 2.6$)

and of subclinical ones (Figure 3 L, $R = 2.6$). The relative reduction in peak incidence

produced by ~20% app adoption would be 47% with an 80% detection rate of clinical cases,

compared to the 35% relative reduction obtained with 50% detection rate. Results show

similar trends across different levels of population immunity, with higher relative impacts

predicted for low immunity (Figure 3 F, I, $R = 2.6$). Compliance to isolation of household

contacts had an appreciable effect at low app adoption (Figure 3 K, $R = 2.6$). 50%

compliance would reduce peak incidence of 19%, compared to 27% reduction for 90%

compliance, in the case of household isolation only. Compliance of notified contacts to

isolation, instead, has a larger effect on peak incidence only when app adoption is high, as

expected. For example, if the app was adopted by 60% of individuals the reduction in peak

incidence would pass from 55% to 66% if compliance changed from 50% to 90% (Figure 3 J,

$R = 2.6$).

We analyzed the simulation outputs to characterize index cases and their contacts and relate

this to the reduction in number of cases by age and setting. We found that adults

represented the majority of index cases (Figure 4 D), while their household contacts were

mostly children. The app registered mostly contacts with adults, and the tracked contacts

were occurring predominantly in workplaces and in the community (Figure 4 A). This results

in a heterogenous reduction in transmission (TRR) by setting and age group. Household

isolation reduced transmission in all settings, with the smallest effect in workplaces (Figure 4

B). Digital contact tracing has instead a high TRR at work, in the community and in transports

(Figure 4 C). Household isolation reached mostly children (< 15 years old) and the elderly

(especially the 75+ group) with the smallest effect in the 15-59 years old (Figure 4 E).

Adopting digital contact tracing led to an increased TRR with age, even among the oldest

age range (Figure 4 F). This result shows the indirect effect of digital tracing: due to the

central role of adults in the transmission of SARS-CoV-2 towards all age-groups, avoiding

adult infections led to less transmission to the elderly. We also tested the case in which

elderly people (70+) owning a smartphone did not install the app at all, assuming they are

less familiar with digital technologies, and we found no appreciable effect. These results and

additional details are provided in the Supplementary Material.

Traced and Isolated individuals

Feasibility of contact tracing depends on the number of traced contacts who require

assistance and virological tests. In a scenario with high detection rate (80%), we found that

for each detected case 1.5 contacts were identified on average through household isolation

but up to 7.5 with app adoption at 57%, for $R = 2.6$ and 10 for $R = 1.7$ (Figure 5 D). This

number was however subject to fluctuations (Figure 5 A). Overall, the maximal fraction of the

population quarantined at any given time was ~50 per 1000 habitants in a scenario with $R =$

2.6, and was between ~1 and ~4.5 per 1000 habitants when $R = 1.7$ (Figure 5 B and E). The

latter case corresponded to the situation in which high levels of app adoption were able to

strongly reduce spreading, thus the proportion of isolated individuals declined in time,

signaling the success of quarantining in preventing the propagation of the infection. A total of

30 per 1000 habitants were isolated in a scenario with $R = 1.7$, assuming high app adoption.

At $R = 2.6$, 1030 per 1000 habitants were isolated at the end of the epidemic meaning that

certain individuals were isolated more than once. In all scenarios, the increase of app

adoption inevitably determined an increase in the proportion of people that were

unnecessarily isolated, i.e. of individuals that were not infected but still isolated (Figure 5C,

F): this proportion increased from 61% to 84% with the increase of app adoption from 0% to

57% (note that the case of 0% app adoption implies that 61% of individuals who were

isolated through household isolation were not infected). These numbers were similar for the

two tested values of $R = 1.7$ and 2.6 (Figure 5 C and F).

Discussion

Quantifying the impact of digital contact tracing is essential to envision this strategy within a

wider response plan against the COVID-19 epidemic. We modelled this intervention together

with household isolation assuming a 50% detection of clinical cases. In a scenario of high

transmissibility ($R = 2.6$), we found that household isolation by itself would produce a

reduction in peak incidence of 27%, while the inclusion of digital contact tracing could

increase this effect by 30% for a reasonably achievable app adoption (~20% of the

population), and by 144% for a large-scale app adoption (~60%). At a moderate

transmissibility level ($R = 1.7$), the app would substantially damp transmission (36% to 89%

peak incidence reduction for increasing app adoption), bringing the epidemic to manageable

levels if adopted by 32% of the population or more. Importantly, the app-based tracing and

household isolation have different effects across settings, the first intervention efficiently

preventing transmissions at work that are not well targeted by the second. Moreover, app-

based contact tracing also yields a protection for the elderly despite the lower penetration of

smartphones in this age category.

Lockdown and social distancing have been effective in reducing transmission in the first

epidemic wave in many countries. However, their huge societal and economic costs made

their prolonged implementation impossible. Phasing out lockdown occurred at the beginning

of the summer in Europe, with high temperatures, increased ventilation and outdoor activities

helping reducing the risk of contagion (16). The relaxation of almost all restrictive measures,

the start of activities in the fall and the cold season accelerated transmission, reaching a

point in which strict non-pharmaceutical interventions were again necessary to curb the

epidemic increase. At the time of writing, national or local lockdowns were restored in several

countries in Europe (*39*). This highlights the need for exit strategies based on sustainable

non-pharmaceutical interventions, able to suppress COVID-19 spread while having limited

impact on the economy and on individuals' daily life (*1*).

Many countries have increased their capacity to detect cases and track their contacts. In

France, thousands of transmission clusters have been identified and controlled since the end

of the first lockdown period (*40*). Under-detection of cases was however estimated over

summer, and the system was predicted to deteriorate rapidly for increasing epidemic activity

(*41*). The automated tracking of contacts could then provide an important complementary

tool. Here we found that digital contact tracing could reduce attack rate and peak incidence,

in agreement with previous works (*18*, *19*, *26*). The impact of the measure would depend on

population immunity, thus geographical heterogeneities should be expected, as regions were

differentially hit by the first wave of the epidemic (*4*). On the other hand, app adoption as well

may be higher in these areas because of risk aversion behavior (*42*). Also, higher

participation rates may be expected in dense urban areas to protect from exposure from

random encounters (e.g. in public transports).

Under realistic hypotheses, the intervention would not be able alone to bring the epidemic

under control in a scenario where transmission is high (*18*, *19*, *26*), mainly due to the strong

role of asymptomatic transmission in fueling the epidemic (*11*, *12*, *43*). We explored different

values of the reproductive number R, to effectively account for non-pharmaceutical measures

mitigating the epidemic and for the adoption of preventive measures substantially hindering

SARS-COV-2 transmission. We found that a reduction of the epidemic to a manageable level

would be possible with a moderate R (e.g. R =1.7 explored here).

Improved case finding is the first step towards a successful contact-tracing intervention. We

found that the increase in detection of clinical cases substantially reduced peak incidence

and improved the efficacy of contact tracing. Many countries progressively increased testing

capacity (*41*) and lifted restrictions on access to testing (*44*). Easy access to testing is

essential to detect cases, because of the substantial fraction of subclinical cases and the

similarity of COVID-19 clinical presentation to the one of other respiratory infections. In the

period from September to November 2020, the French network of virological surveillance run

by general practitioners reported that only 22.7% of Acute Respiratory Infections were

caused by SARS-CoV2, against 46.5% attributed to rhinovirus (*45*). Given that the majority of

cases do not require hospitalization, case detection effectiveness is also influenced by the

consultation rate. This has been estimated to be around ~30% with peaks at ~45% by the

participatory surveillance platform covidnet.fr (*40, 41*). Increased population awareness is

thus essential for the efficient monitoring of the epidemic and its containment through contact

tracing.

Little information is available on compliance to isolation. Low compliance to isolation was

reported in the UK and in a university campus in the US (*46, 47*). However, this may vary

greatly according to cultural, socioeconomic and demographic context. Due to a self-

selection bias, individuals who decided to download and use the mobile application may be

more akin to follow the recommendation and isolate if they receive a notification. Step-by-

step recommendations provided by the app can further help in increasing compliance.

Strengthened communication and compensations (such as paid work leave, loss-of-income

payments for self-employed professionals, medical school-absence certificates) should be

implemented to increase the acceptability of isolation (*48*).

App adoption remains the key factor determining the efficacy of digital contact tracing.

Adoption levels were initially low (<5%) in many countries (e.g. Italy, France), increasing later

as the second wave was rising, likely due to increased concerns of the population. As of

November 2020, 17% and 13% of the population had downloaded the app in Italy and

France, respectively (*49, 50*). Higher levels were observed, e.g., in Australia (6 millions

download, 25% of the population) (*51*) and Iceland (~150 thousand, 38%) (*52*). Importantly,

official figures may overestimate real adoption levels, since many individuals may download

the application without using it. In France this proportion was 60% among university students

(*53*). Individuals may be more inclined to use the app if they perceive a direct and immediate

benefit from its use. This may be implemented through, e.g., easy access to testing in case

they are notified as contacts and assistance by public health professionals. Moreover, even if

the application preserves users' privacy and can be downloaded on a voluntary basis in

many countries, increased transparency and ethical debate remain essential to reassure the

population about data treatment (*53–55*).

The results presented here are based on an agent-based model that describes age-specific

risk factors for exposure and transmission: contact rates, contacts by location, susceptibility

to the virus, probability of being detected and rate of app adoption. The interplay between

these features has a profound impact on COVID-19 spread and affects the efficacy of

household isolation and digital contact tracing. To account for contact heterogeneities we

used statistics on population demography, combined with social contact surveys to build a

multi-setting contact network, similarly to previous works (*17*, *21*, *26*, *30–32*). The network is

also dynamic in time as it captures the repetition of a certain number of contacts (e.g.

relationships) and the occurrence of random encounters. Social contact data provide an

invaluable information source to study the current COVID-19 outbreak (*1*, *36*). Previous

projections on the impact of contact tracing rely on a similar approach in some cases (*17*, *24*,

26). Other works make use of high resolution data (*18*, *19*, *22*), that are more reliable than

contact surveys, but are restricted to specific settings or population groups. Despite the

difference in the data source and approach, the results of these studies are consistent and in

agreement with our work on the overall impact of the intervention.

We modelled age-specific epidemiological characteristics based on available knowledge in

the literature. Children are less impacted by the COVID-19 epidemic (*9–13*). This may be

explained by reduced susceptibility and severity, with accumulating evidence that both

effects are acting simultaneously (*10*). The strength of these effects is still debated and the

infection risk for children should not be minimized. However, these differences have

implications for digital contact tracing. Indeed, it is precisely in the group that plays a central

role in transmission and where cases are more likely symptomatic (i.e., adults) that the app

coverage is already the highest. Our model shows that taken together, these characteristics

reinforce the impact of digital tracing and provides indirect protection in the elderly

population. This occurs even if no adoption is registered in the elderly population.

Our study is affected by limitations. First, we analyzed the effect of digital contact tracing on

COVID-19 incidence in the general population. Crucial information for public health

authorities would be to quantify the effect in time of these measures on hospitalizations. This

would require to couple our model for COVID-19 transmission in the general population with

a model describing disease severity and within-hospital patient trajectories (*17*, *21*, *26*).

Second, the model does not account for transmission in nursing homes. This setting is where

the majority of transmissions among elderly occurred. At the same time, however, the

response to the COVID-19 epidemic in this setting relies mostly on routine screening of

symptoms and frequent testing of residents, together with face masks and strict hygiene

rules for visitors. Third, results may be conservative as clustering effects and large

fluctuations in the number of contacts per person (*56*) are only partially captured by the

model thanks to the repetition of contacts, but effects may be larger in real contact patterns.

This also includes crowding events playing an important role in the transmission dynamics

(*15*). Overall behaviors obtained with our synthetic network of contacts are however

compatible with findings obtained with real contact data (*18*). In a future work the description

of temporal and topological properties of contacts in workplaces, schools and community

could be improved by using modeling frameworks informed by detailed contact data, that has

become available for specific settings (*57–59*). For this purpose, frameworks such as hidden

variable models or other recent dynamical models for social networks could be employed

(*60–62*). Fourth, other assumptions may be instead optimistic, regarding the probability of

detection of index cases, and compliance to isolation, for example. Few data are available to

inform these parameters that may also vary over time (depending on the epidemic context

and increased population fatigue) and across countries (depending on cultural aspects and

regulations in place). While we explored a range of parameter values, more detailed

information will be needed to contextualize our approach to a specific epidemic situation in a

given country.

Material and Methods

Synthetic population

The model simulates the population of Metropolitan France representing individual

inhabitants. This approach is similar to studies done previously e.g. for Italy[27] and for

USA[28]. The French synthetic population is based on the National Institute of Statistics and

Economic Studies (INSEE) censuses. Individuals were assigned to municipalities according

to the administrative borders. The number of households and the age structure of their

inhabitants, sizes of schools and workplaces, fluxes of commuters between municipalities

also followed the distribution of these statistics found in the INSEE data. Population size was

kept constant through a simulation as we aimed at simulating one season of the epidemic.

To generate the population, we defined several statistics derived from INSEE publicly

available data:

• The list of municipalities ("les communes de France") of Metropolitan France (2015) with

each municipality described by its INSEE code, population size, number of schools of six

different levels (from kindergarten to university), number of workplaces in given size

categories (0-9, 10-49, 50-99, 100-499, 500-999 and over 1000 employees) (Populations

légales 2017, INSEE, https://www.insee.fr/fr/statistiques/4265429?sommaire=4265511).

• Statistics regarding the percentage of people in given age groups enrolled in each of six

school levels, employed and unemployed (Bilan démographique 2010, INSEE,

https://www.insee.fr/fr/statistiques/1280950).

• The age pyramid for France as the population fractions of individuals of a given age

(Bilan démographique 2010, INSEE, https://www.insee.fr/fr/statistiques/1280950).

• The number of people commuting to work between each pair of municipalities (Mobilités

professionnelles en 2016: déplacements domicile - lieu de travail, INSEE,

https://www.insee.fr/fr/statistiques/4171554).

• The number of people commuting to school between each pair of municipalities

(Mobilités scolaires en 2015: déplacements domicile - lieu d'études, INSEE,

https://www.insee.fr/fr/statistiques/3566470).

• The probability distributions of sizes of households in France (Couples - Familles -

Ménages en 2010. INSEE, https://www.insee.fr/fr/statistiques/2044286/?geo=COM-

34150.)

• The probability of individuals belonging to a particular age class, given their role in the

household: child of a couple, child of a single adult, adult in a couple without children,

adult in a couple with children (Couples - Familles - Ménages en 2010. INSEE,

https://www.insee.fr/fr/statistiques/2044286/?geo=COM-34150).

With the above statistics, the synthetic population was generated in the following steps:

1. Initialization of all the municipalities with an appropriate number of schools of each type

and workplaces of given sizes.

2. Creation of schools in each municipality according to given statistics.

3. Creation of workplaces in each municipality according to given statistics.

4. Definition of the commuter fluxes between municipalities.

Each municipality has a defined number of inhabitants and individuals are created (one by

one) until this number is reached. Each individual was assigned an age, a school or a

workplace (or is assigned to stay at home) according to probability distributions derived from

the data mentioned above.

The numbers of households within each municipality were not defined explicitly, but

depended on the number of individuals. The municipal population size and statistics

regarding family demographics constrain the number of households. Additional details on the

algorithm for the population reconstruction are provided in the Supplementary Material.

415   Face-to-face contact network

The synthetic population encodes information on the school, workplace, household and

community each individual belongs to. We used this information to extract a dynamic network

representing daily face-to-face contacts. We parametrized this network based on contacts'

statistics for the French population (*33*).

First, we generated a time aggregated network representing all contacts that can potentially

occur – we will call this the *acquaintance network,* with some abuse of language since it

includes also sporadic contacts. Second, to each contact we assigned a daily rate of

activation. Then, in the course of the simulation we sampled contacts each day based on

their rate.

The acquaintance network has five distinct layers representing contacts in household (layer

H), workplace (layer W), school (layer S), community (layer C) and transports (layer T). The

household layer is formed by a collection of complete networks linking individuals in the

same household. The W, S, C, and T layers are formed by collections of Erdős–

Rényi networks generated in each location i, with average degree χ_i. A location can be a

workplace (W layer), a school (S layer) and a municipality (C and T layers). χ_i is extracted at

random for each place and depends on the type and size of the location. In particular, when

the size of a location is small we assume that each individual enters in contact with all the

others frequenting the same place. As the size increases the number of contacts saturates.

Once the acquaintance network was built a daily activation rate x was assigned to each link

according to a cumulative distribution that depends on the layer s. For simplicity we assumed

this distribution to be the same for $s = W, S, C$, while we allowed it to be different in household

(where contacts are more frequent) and in transports (where contacts are sporadic).

Parameters were tuned based on average daily number of contacts, proportion of contacts

by setting, and contact frequency as provided in (*33*) (Figure 1 C D). Additional details on the

network reconstruction and parametrization are provided in the Supplementary Material.

Transmission model

We defined a minimal model of COVID-19 spread in the general population that accounts for

two levels of symptoms: none to mild (subclinical cases, I_{sc}), and moderate to severe (clinical

cases, I_c). We assumed that clinical cases stay at home after developing symptoms.

Susceptible individuals, if in contact with infectious ones, may get infected and enter the

exposed compartment (E). After an average latency period ϵ^{-1} they become infectious,

developing a subclinical infection with probability p_{sc}^A and a clinical infection otherwise. From

E, before entering in either I_{sc} or I_c, individuals enter first a prodromal phase (either $I_{p,sc}$ or

$I_{p,c}$), that lasts on average μ_p^{-1} days and where individuals do not show any sign of illness,

despite being already infectious. Contact-tracing, population-screening and modelling studies

provide evidence that infectivity is related to the level of symptoms, with less severely hit

individuals being also less infectious (*11*, *43*). Therefore, we assumed that subclinical cases,

$I_{p,sc}$ and I_{sc} have a reduced transmissibility compared to $I_{p,c}$ and I_c. This is modulated by the

scaling factor β_I. We neglected hospitalization and death and assumed that with rate μ

infected individuals become recovered.

The impact of COVID-19 is heterogeneous across age groups (*9–13*). This may be driven by

differences in susceptibility (*9*), differences in clinical manifestation (*11*, *13*) or both (*10*). We

considered here both effects in agreement with recent modelling estimates (*10*).

Susceptibility by age, σ_A, was parametrized from (*9*), while clinical manifestation, p_{sc}^A, was

parametrized from a large-scale descriptive study of the COVID19 outbreak in Italy (*13*).

Transition rates are summarized in Figure 2 B, and parameters and their values are listed in

Table 1. The incubation period was estimated to be around 5.2 days from an early analysis

of 425 patients in Wuhan (*34*). COVID-19 transmission potential varies across settings,

populations and social contexts (*14–16*). In particular, indoor places were found to increase

the odds of contagion 18.7 times compared to an open-air environment (*16*). In our model we

assumed that all contacts at work, school and transport occur indoor and have the same

transmission risk (ω_S). In the contact survey of Béraud et al. (*33*), 46% of contacts in the

community were occurring outdoors. Combining this information with the 18.7 indoor vs.

outdoor risk ratio leads to a 60% relative risk of community contacts with respect to

workplace/school/transport contacts. Contacts within households are generally associated to

a higher risk with respect to other settings, because they last longer and there is a higher risk

of environmental transmission. We assumed that the transmission risk associated with them

is twice the one in workplace/school/transport. For the basic reproductive ratio of COVID-19

we took $R_0 \sim 3.1$ (*1, 36*). We also explored lower levels of transmission potential, i.e.

reproductive ratios R down to 1.5, to effectively account for behavioral changes and adoption

of barrier measures. Our definition of R does not integrate population immunity. We explicitly

indicate the initial level of population immunity to disentangle the relative role of the two

quantities.

Modelling contact tracing

Self-isolation and isolation of household contacts

Self-isolation and isolation of household contacts was modelled according to following rules:

• As an individual shows clinical symptoms, s/he is detected with probability

$p_{d,c}$ (baseline value 50%, additional explored values 30% and 80%). If detected, case

confirmation, isolation and contacts' isolation occur with rate $r_{d,c} = 0.9$ upon symptoms

onset.

• Subclinical individuals are also detected with probability $p_{d,sc}$ (baseline value 5%,

additional explored values 25% and 45%), and rate $r_{d,sc} = 0.5$.

• The individual's family members are isolated with probability $p_{c,h} = 0.9$ (0.5 and 0.7 were

also explored).

• We assume that contacts are tested and the follow up guarantees that all individuals who

got infected prior to isolation are detected. Thus, contacts that are negative (either

susceptible or recovered at the time of isolation) terminate their isolation after 7 days.

The index-case and the positive contacts are isolated for 14 days. Contacts with no

clinical symptoms have a daily probability $p_{drop} = 0.02$ to drop-out from isolation.

• For both the case and the contacts, isolation is implemented by assuming no contacts

outside the household and contacts within a household having an associated

transmission risk (i.e. the weight ω_H) reduced by a factor $\iota = 0.5$.

Digital contact tracing

We assumed that contact tracing is adopted in combination with self-isolation and isolation of

household members. Therefore, we added the following rules to the ones outlined above:

• At the beginning of the simulation, a smartphone is assigned to individuals with

probability p_{sm}^A, based on the statistics of smartphone penetration (0% for [0,11], 86% for

[12,17], 98% for [18,24], 95% for [25,39], 80% for [40,59], 62% for [60,69], 44% for 70+)

(*29*).

• Each individual with a smartphone has a probability p_a to download the app (we explored

values between 0 and 0.9).

• Only contacts occurring between individuals with a smartphone and the app are traced.

• If the individual owns a smartphone and downloaded the app the contacts that s/he has

traced in the period since $D = 7$ days before his/her detection are isolated with probability

$p_{c,a} = 0.9$ (0.5 and 0.7 were also tested).

• We assume contacts are tested and the follow up guarantees that all individuals who got

infected prior to isolation are detected. Thus, contacts that are negative (either

susceptible or recovered at the time of isolation) terminate their isolation after 7 days.

The index-case and the positive contacts are isolated for 14 days. Contacts with no

clinical symptoms have a daily probability $p_{drop} = 0.02$ to drop out from isolation.

- For both the case and the contacts, isolation is implemented by assuming no contacts

  outside the household and contacts within a household having an associated

  transmission risk (i.e. the weight ω_H) reduced by a factor $\iota = 0.5$.

Reference

1. L. Di Domenico, G. Pullano, C. E. Sabbatini, P.-Y. Boëlle, V. Colizza, Impact of lockdown on COVID-19 epidemic in Île-de-France and possible exit strategies. *BMC Medicine*. **18**, 240 (2020).

2. S. Flaxman, S. Mishra, A. Gandy, H. J. T. Unwin, T. A. Mellan, H. Coupland, C. Whittaker, H. Zhu, T. Berah, J. W. Eaton, M. Monod, A. C. Ghani, C. A. Donnelly, S. M. Riley, M. A. C. Vollmer, N. M. Ferguson, L. C. Okell, S. Bhatt, Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe. *Nature*, 1–8 (2020).

3. M. Pollán, B. Pérez-Gómez, R. Pastor-Barriuso, J. Oteo, M. A. Hernán, M. Pérez-Olmeda, J. L. Sanmartín, A. Fernández-García, I. Cruz, N. F. de Larrea, M. Molina, F. Rodríguez-Cabrera, M. Martín, P. Merino-Amador, J. L. Paniagua, J. F. Muñoz-Montalvo, F. Blanco, R. Yotti, F. Blanco, R. G. Fernández, M. Martín, S. M. Navarro, M. Molina, J. F. Muñoz-Montalvo, M. S. Hernández, J. L. Sanmartín, M. Cuenca-Estrella, R. Yotti, J. L. Paniagua, N. F. de Larrea, P. Fernández-Navarro, R. Pastor-Barriuso, B. Pérez-Gómez, M. Pollán, A. Avellón, G. Fedele, A. Fernández-García, J. O. Iglesias, M. T. P. Olmeda, I. Cruz, M. E. F. Martinez, F. D. Rodríguez-Cabrera, M. A. Hernán, S. P. Fernández, J. M. R. Aguirre, J. M. N. Marí, B. P. Borrás, A. B. P. Jiménez, M. Rodríguez-Iglesias, A. M. C. Gascón, M. L. L. Alcaine, I. D. Suárez, O. S. Álvarez, M. R. Pérez, M. C. Sanchís, C. J. V. Gomila, L. C. Saladrigas, A. H. Fernández, A. Oliver, E. C. Feliciano, M. N. G. Quintana, J. M. B. Fernández, M. A. H. Betancor, M. H. Febles, L. M. Martín, L.-M. L. López, T. U. Miota, I. D. B. Población, M. S. C. Pérez, M. N. V. Fernández, T. M. Enríquez, M. V. Arranz, M. D.-G. González, I. Fernández-Natal, G. M. Lobón, J. L. M. Bellido, P. Ciruela, A. M. i Casals, M. D. Botías, M. A. M. Maeso, D. P. del Campo, A. F. de Castro, R. L. Ramírez, M. F. E. Retamosa, M. R. González, M. S. B. Lobeiras, A. F. Losada, A. Aguilera, G. Bou, Y. Caro, N. Marauri, L. M. S. Blanco, I. del C. González, M. H. Pascual, R. A. Fernández, P. Merino-Amador, N. C. Castro, A. T. Lizcano, C. R. Almagro, M. S. Hernández, N. A. Elizaga, M. E. Sanz, C. E. Baquedano, A. B. Bascaran, S. I. Tamayo, L. E. Otazua, R. B. Benarroch, J. L. Flores, A. V. de la Villa, Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological study. *The Lancet*. **0** (2020), doi:10.1016/S0140-6736(20)31483-5.

4. F. Carrat, X. de Lamballerie, D. Rahib, H. Blanche, N. Lapidus, F. Artaud, S. Kab, A. Renuy, F. S. de Edelenyi, L. Meyer, N. Lydie, M.-A. Charles, P.-Y. Ancel, F. Jusot, A. Rouquette, S. Priet, P. M. S. Villaroel, T. Fourie, C. Lusivika-Nzinga, J. Nicol, S. Legot, N. Druesne-Pecollo, Y. Essedik, C. Lai, J.-M. Gagliolo, J.-F. Deleuze, N. Bajos, G. Severi, M. Touvier, M. Zins, Seroprevalence of SARS-CoV-2 among adults in three regions of France following the lockdown and associated risk factors: a multicohort study. *medRxiv* (2020), doi:10.1101/2020.09.16.20195693.

5. S. L. Vu, G. Jones, F. Anna, T. Rose, J.-B. Richard, S. Bernard-Stoecklin, S. Goyard, C. Demeret, O. Helynck, C. Robin, V. Monnet, L. P. de Facci, M.-N. Ungeheuer, L. Léon,

Y. Guillois, L. Filleul, P. Charneau, D. Lévy-Bruhl, S. van der Werf, H. Noel, Prevalence of SARS-CoV-2 antibodies in France: results from nationwide serological surveillance. *medRxiv* (2020), doi:10.1101/2020.10.20.20213116.

6. M. Kendall, L. Milsom, L. Abeler-Dörner, C. Wymant, L. Ferretti, M. Briers, C. Holmes, D. Bonsall, J. Abeler, C. Fraser, Epidemiological changes on the Isle of Wight after the launch of the NHS Test and Trace programme: a preliminary analysis. *The Lancet Digital Health*. **0** (2020), doi:10.1016/S2589-7500(20)30241-7.

7. I. Braithwaite, T. Callender, M. Bullock, R. W. Aldridge, Automated and partly automated contact tracing: a systematic review to inform the control of COVID-19. *The Lancet Digital Health*. **2**, e607–e621 (2020).

8. M. Salathé, C. Althaus, N. Anderegg, D. Antonioli, T. Ballouz, E. Bugnon, S. Čapkun, D. Jackson, S.-I. Kim, J. Larus, N. Low, W. Lueks, D. Menges, C. Moullet, M. Payer, J. Riou, T. Stadler, C. Troncoso, E. Vayena, V. von Wyl, Early evidence of effectiveness of digital contact tracing for SARS-CoV-2 in Switzerland. *Swiss Medical Weekly*. **150** (2020), doi:10.4414/smw.2020.20457.

9. J. Zhang, M. Litvinova, Y. Liang, Y. Wang, W. Wang, S. Zhao, Q. Wu, S. Merler, C. Viboud, A. Vespignani, M. Ajelli, H. Yu, Changes in contact patterns shape the dynamics of the COVID-19 outbreak in China. *Science* (2020), doi:10.1126/science.abb8001.

10. N. G. Davies, P. Klepac, Y. Liu, K. Prem, M. Jit, R. M. Eggo, Age-dependent effects in the transmission and control of COVID-19 epidemics. *Nature Medicine*, 1–7 (2020).

11. Q. Bi, Y. Wu, S. Mei, C. Ye, X. Zou, Z. Zhang, X. Liu, L. Wei, S. A. Truelove, T. Zhang, W. Gao, C. Cheng, X. Tang, X. Wu, Y. Wu, B. Sun, S. Huang, Y. Sun, J. Zhang, T. Ma, J. Lessler, T. Feng, Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study. *The Lancet Infectious Diseases*. **0** (2020), doi:10.1016/S1473-3099(20)30287-5.

12. E. Lavezzo, E. Franchin, C. Ciavarella, G. Cuomo-Dannenburg, L. Barzon, C. Del Vecchio, L. Rossi, R. Manganelli, A. Loregian, N. Navarin, D. Abate, M. Sciro, S. Merigliano, E. De Canale, M. C. Vanuzzo, V. Besutti, F. Saluzzo, F. Onelia, M. Pacenti, S. G. Parisi, G. Carretta, D. Donato, L. Flor, S. Cocchio, G. Masi, A. Sperduti, L. Cattarino, R. Salvador, M. Nicoletti, F. Caldart, G. Castelli, E. Nieddu, B. Labella, L. Fava, M. Drigo, K. A. M. Gaythorpe, A. R. Brazzale, S. Toppo, M. Trevisan, V. Baldo, C. A. Donnelly, N. M. Ferguson, I. Dorigatti, A. Crisanti, Suppression of a SARS-CoV-2 outbreak in the Italian municipality of Vo'. *Nature*. **584**, 425–429 (2020).

13. F. Riccardo, M. Ajelli, X. D. Andrianou, A. Bella, M. D. Manso, M. Fabiani, S. Bellino, S. Boros, A. M. Urdiales, V. Marziano, M. C. Rota, A. Filia, F. D'Ancona, A. Siddu, O. Punzo, F. Trentini, G. Guzzetta, P. Poletti, P. Stefanelli, M. R. Castrucci, A. Ciervo, C. D. Benedetto, M. Tallon, A. Piccioli, S. Brusaferro, G. Rezza, S. Merler, P. Pezzotti, the C.-19 working Group, Epidemiological characteristics of COVID-19 cases and estimates of the reproductive numbers 1 month into the epidemic, Italy, 28 January to 31 March 2020. *Eurosurveillance*. **25**, 2000790 (2020).

14. Q. J. Leclerc, N. M. Fuller, L. E. Knight, CMMID COVID-19 Working Group, S. Funk, G. M. Knight, What settings have been linked to SARS-CoV-2 transmission clusters? *Wellcome Open Research*. **5**, 83 (2020).

15. B. M. Althouse, E. A. Wenger, J. C. Miller, S. V. Scarpino, A. Allard, L. Hébert-Dufresne,
H. Hu, Superspreading events in the transmission dynamics of SARS-CoV-2:
Opportunities for interventions and control. *PLOS Biology*. **18**, e3000897 (2020).

16. H. Nishiura, H. Oshitani, T. Kobayashi, T. Saito, T. Sunagawa, T. Matsui, T. Wakita, M.
C.-19 R. Team, M. Suzuki, Closed environments facilitate secondary transmission of
coronavirus disease 2019 (COVID-19). *medRxiv* (2020),
doi:10.1101/2020.02.28.20029272.

17. A. Aleta, D. Martín-Corral, A. Pastore y Piontti, M. Ajelli, M. Litvinova, M. Chinazzi, N. E.
Dean, M. E. Halloran, I. M. Longini Jr, S. Merler, A. Pentland, A. Vespignani, E. Moro,
Y. Moreno, Modelling the impact of testing, contact tracing and household quarantine
on second waves of COVID-19. *Nature Human Behaviour*. **4**, 964–971 (2020).

18. G. Cencetti, G. Santin, A. Longa, E. Pigani, A. Barrat, C. Cattuto, S. Lehmann, B. Lepri,
Using real-world contact networks to quantify the effectiveness of digital contact tracing
and isolation strategies for Covid-19 pandemic. *medRxiv* (2020),
doi:10.1101/2020.05.29.20115915.

19. A. J. Kucharski, P. Klepac, A. J. K. Conlan, S. M. Kissler, M. L. Tang, H. Fry, J. R. Gog,
W. J. Edmunds, J. C. Emery, G. Medley, J. D. Munday, T. W. Russell, Q. J. Leclerc, C.
Diamond, S. R. Procter, A. Gimma, F. Y. Sun, H. P. Gibbs, A. Rosello, K. van
Zandvoort, S. Hué, S. R. Meakin, A. K. Deol, G. Knight, T. Jombart, A. M. Foss, N. I.
Bosse, K. E. Atkins, B. J. Quilty, R. Lowe, K. Prem, S. Flasche, C. A. B. Pearson, R. M.
G. J. Houben, E. S. Nightingale, A. Endo, D. C. Tully, Y. Liu, J. Villabona-Arenas, K.
O'Reilly, S. Funk, R. M. Eggo, M. Jit, E. M. Rees, J. Hellewell, S. Clifford, C. I. Jarvis, S.
Abbott, M. Auzenbergs, N. G. Davies, D. Simons, Effectiveness of isolation, testing,
contact tracing, and physical distancing on reducing transmission of SARS-CoV-2 in
different settings: a mathematical modelling study. *The Lancet Infectious Diseases*. **0**
(2020), doi:10.1016/S1473-3099(20)30457-6.

20. L. Ferretti, C. Wymant, M. Kendall, L. Zhao, A. Nurtay, L. Abeler-Dörner, M. Parker, D.
Bonsall, C. Fraser, Quantifying SARS-CoV-2 transmission suggests epidemic control
with digital contact tracing. *Science* (2020), doi:10.1126/science.abb6936.

21. R. Hinch, W. J. M. Probert, A. Nurtay, M. Kendall, C. Wymatt, M. Hall, K. Lythgoe, A. B.
Cruz, L. Zhao, A. Stewart, L. Ferritti, D. Montero, J. Warren, N. Mather, M. Abueg, N.
Wu, A. Finkelstein, D. G. Bonsall, L. Abeler-Dorner, C. Fraser, OpenABM-Covid19 - an
agent-based model for non-pharmaceutical interventions against COVID-19 including
contact tracing. *medRxiv* (2020), doi:10.1101/2020.09.16.20195925.

22. A. Barrat, C. Cattuto, M. Kivelä, S. Lehmann, J. Saramäki, Effect of manual and digital
contact tracing on COVID-19 outbreaks: a study on empirical contact data. *medRxiv*
(2020), doi:10.1101/2020.07.24.20159947.

23. M. Mancastroppa, C. Castellano, A. Vezzani, R. Burioni, Stochastic sampling effects
favor manual over digital contact tracing. *arXiv:2010.03399 [cond-mat, physics:physics,*
q-bio] (2020) (available at http://arxiv.org/abs/2010.03399).

24. L. Lorch, H. Kremer, W. Trouleau, S. Tsirtsis, A. Szanto, B. Schölkopf, M. Gomez-
Rodriguez, Quantifying the Effects of Contact Tracing, Testing, and Containment
Measures in the Presence of Infection Hotspots. *arXiv:2004.07641 [physics, q-bio, stat]*
(2020) (available at http://arxiv.org/abs/2004.07641).

25. M. E. Kretzschmar, G. Rozhnova, M. C. J. Bootsma, M. van Boven, J. H. H. M. van de
Wijgert, M. J. M. Bonten, Impact of delays on effectiveness of contact tracing strategies
for COVID-19: a modelling study. *The Lancet Public Health*. **5**, e452–e459 (2020).

26. M. Abueg, R. Hinch, N. Wu, L. Liu, W. J. M. Probert, A. Wu, P. Eastham, Y. Shafi, M.
Rosencrantz, M. Dikovsky, Z. Cheng, A. Nurtay, L. Abeler-Dörner, D. G. Bonsall, M. V.
McConnell, S. O'Banion, C. Fraser, Modeling the combined effect of digital exposure
notification and non-pharmaceutical interventions on the COVID-19 epidemic in
Washington state. *medRxiv* (2020), doi:10.1101/2020.08.29.20184135.

27. M. Ajelli, B. Gonçalves, D. Balcan, V. Colizza, H. Hu, J. J. Ramasco, S. Merler, A.
Vespignani, Comparing large-scale computational approaches to epidemic modeling:
Agent-based versus structured metapopulation models. *BMC Infectious Diseases*. **10**,
190 (2010).

28. D. L. Chao, M. E. Halloran, V. J. Obenchain, I. M. L. Jr, FluTE, a Publicly Available
Stochastic Influenza Epidemic Simulation Model. *PLOS Comput Biol*. **6**, e1000656
(2010).

29. Baromètre du numérique 2019 (2019), (available at
https://www.credoc.fr/publications/barometre-du-numerique-2019).

30. L. Fumanelli, M. Ajelli, P. Manfredi, A. Vespignani, S. Merler, Inferring the Structure of
Social Contacts from Demographic Data in the Analysis of Infectious Diseases Spread.
PLOS Computational Biology. **8**, e1002673 (2012).

31. Q.-H. Liu, M. Ajelli, A. Aleta, S. Merler, Y. Moreno, A. Vespignani, Measurability of the
epidemic reproduction number in data-driven contact networks. *PNAS*. **115**, 12680–
12685 (2018).

32. D. Mistry, M. Litvinova, A. Pastore y Piontti, M. Chinazzi, L. Fumanelli, M. F. C. Gomes,
S. A. Haque, Q.-H. Liu, K. Mu, X. Xiong, M. E. Halloran, I. M. Longini, S. Merler, M.
Ajelli, A. Vespignani, Inferring high-resolution human mixing patterns for disease
modeling. *Nature Communications*. **12**, 323 (2021).

33. G. Béraud, S. Kazmercziak, P. Beutels, D. Levy-Bruhl, X. Lenne, N. Mielcarek, Y.
Yazdanpanah, P.-Y. Boëlle, N. Hens, B. Dervaux, The French Connection: The First
Large Population-Based Contact Survey in France Relevant for the Spread of Infectious
Diseases. *PLOS ONE*. **10**, e0133203 (2015).

34. Q. Li, X. Guan, P. Wu, X. Wang, L. Zhou, Y. Tong, R. Ren, K. S. M. Leung, E. H. Y. Lau,
J. Y. Wong, X. Xing, N. Xiang, Y. Wu, C. Li, Q. Chen, D. Li, T. Liu, J. Zhao, M. Liu, W.
Tu, C. Chen, L. Jin, R. Yang, Q. Wang, S. Zhou, R. Wang, H. Liu, Y. Luo, Y. Liu, G.
Shao, H. Li, Z. Tao, Y. Yang, Z. Deng, B. Liu, Z. Ma, Y. Zhang, G. Shi, T. T. Y. Lam, J.
T. Wu, G. F. Gao, B. J. Cowling, B. Yang, G. M. Leung, Z. Feng, Early Transmission
Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia. *New England
Journal of Medicine*. **382**, 1199–1207 (2020).

35. X. He, E. H. Y. Lau, P. Wu, X. Deng, J. Wang, X. Hao, Y. C. Lau, J. Y. Wong, Y. Guan,
X. Tan, X. Mo, Y. Chen, B. Liao, W. Chen, F. Hu, Q. Zhang, M. Zhong, Y. Wu, L. Zhao,
F. Zhang, B. J. Cowling, F. Li, G. M. Leung, Temporal dynamics in viral shedding and
transmissibility of COVID-19. *Nature Medicine*, 1–4 (2020).

36. H. Salje, C. T. Kiem, N. Lefrancq, N. Courtejoie, P. Bosetti, J. Paireau, A. Andronico, N.
Hozé, J. Richet, C.-L. Dubost, Y. L. Strat, J. Lessler, D. Levy-Bruhl, A. Fontanet, L.

Opatowski, P.-Y. Boelle, S. Cauchemez, Estimating the burden of SARS-CoV-2 in
France. *Science* (2020), doi:10.1126/science.abc3517.

37. D. Cereda, M. Tirani, F. Rovida, V. Demicheli, M. Ajelli, P. Poletti, F. Trentini, G.
Guzzetta, V. Marziano, A. Barone, M. Magoni, S. Deandrea, G. Diurno, M. Lombardo,
M. Faccini, A. Pan, R. Bruno, E. Pariani, G. Grasselli, A. Piatti, M. Gramegna, F.
Baldanti, A. Melegaro, S. Merler, The early phase of the COVID-19 outbreak in
Lombardy, Italy. *arXiv:2003.09320 [q-bio]* (2020) (available at
http://arxiv.org/abs/2003.09320).

38. mbevand, *mbevand/florida-covid19-line-list-data* (2020;
https://github.com/mbevand/florida-covid19-line-list-data).

39. Stay-at-home requirements during the COVID-19 pandemic. *Our World in Data*,
(available at https://ourworldindata.org/grapher/stay-at-home-covid).

40. SPF, COVID-19 : point épidémiologique du 19 novembre 2020, (available at /maladies-
et-traumatismes/maladies-et-infections-respiratoires/infection-a-
coronavirus/documents/bulletin-national/covid-19-point-epidemiologique-du-19-
novembre-2020).

41. G. Pullano, L. Di Domenico, C. E. Sabbatini, E. Valdano, C. Turbelin, M. Debin, C.
Guerrisi, C. Kengne-Kuetche, C. Souty, T. Hanslik, T. Blanchon, P.-Y. Boëlle, J. Figoni,
S. Vaux, C. Campèse, S. Bernard-Stoecklin, V. Colizza, Underdetection of COVID-19
cases in France threatens epidemic control. *Nature*, 1–9 (2020).

42. G. Pullano, E. Valdano, N. Scarpa, S. Rubrichi, V. Colizza, Evaluating the effect of
demographic factors, socioeconomic factors, and risk aversion on mobility during the
COVID-19 epidemic in France under lockdown: a population-based study. *The Lancet
Digital Health*. **2**, e638–e649 (2020).

43. R. Li, S. Pei, B. Chen, Y. Song, T. Zhang, W. Yang, J. Shaman, Substantial
undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-
CoV-2). *Science*. **368**, 489–493 (2020).

44. Coronavirus (COVID-19) Testing - Statistics and Research. *Our World in Data*, (available
at https://ourworldindata.org/coronavirus-testing).

45. Réseau Sentinelles > France > Accueil, (available at
https://websenti.u707.jussieu.fr/sentiweb/).

46. L. E. Smith, H. W. W. Potts, R. Amlot, N. T. Fear, S. Michie, J. Rubin, Adherence to the
test, trace and isolate system: results from a time series of 21 nationally representative
surveys in the UK (the COVID-19 Rapid Survey of Adherence to Interventions and
Responses [CORSAIR] study). *medRxiv* (2020), doi:10.1101/2020.09.15.20191957.

47. K. Chang, A University Had a Great Coronavirus Plan, but Students Partied On. *The
New York Times* (2020), (available at
https://www.nytimes.com/2020/09/10/health/university-illinois-covid.html).

48. L. Atlani-Duault, B. Lina, D. Malvy, Y. Yazdanpanah, F. Chauvin, J.-F. Delfraissy,
COVID-19: France grapples with the pragmatics of isolation. *The Lancet Public Health*.
5, e573–e574 (2020).

49. TousAntiCovid sur Twitter. *Twitter*, (available at
https://twitter.com/TousAntiCovid/status/1328272121177452551).

50. Immuni - I numeri di Immuni, (available at www.immuni.italia.it).

51. The curve of people downloading COVIDSafe has flattened. Has the app done enough?
(2020), (available at https://www.abc.net.au/news/2020-06-02/coronavirus-covid19-
covidsafe-app-how-many-downloads-greg-hunt/12295130).

52. Nearly 40% of Icelanders are using a covid app—and it hasn't helped much. *MIT*
Technology Review, (available at
https://www.technologyreview.com/2020/05/11/1001541/iceland-rakning-c19-covid-
contact-tracing/).

53. I. Montagni, N. Roussel, R. Thiébaut, C. Tzourio, The French Covid-19 contact tracing
app: knowledge, attitudes, beliefs and practices of students in the health domain. *J Med*
Internet Res (2021), doi:10.2196/26399.

54. *DP-3T/documents* (DP^3T, 2020; https://github.com/DP-3T/documents).

55. M. Nanni, G. Andrienko, A.-L. Barabási, C. Boldrini, F. Bonchi, C. Cattuto, F.
Chiaromonte, G. Comandé, M. Conti, M. Coté, F. Dignum, V. Dignum, J. Domingo-
Ferrer, P. Ferragina, F. Giannotti, R. Guidotti, D. Helbing, K. Kaski, J. Kertesz, S.
Lehmann, B. Lepri, P. Lukowicz, S. Matwin, D. M. Jiménez, A. Monreale, K. Morik, N.
Oliver, A. Passarella, A. Passerini, D. Pedreschi, A. Pentland, F. Pianesi, F. Pratesi, S.
Rinzivillo, S. Ruggieri, A. Siebes, V. Torra, R. Trasarti, J. van den Hoven, A.
Vespignani, Give more data, awareness and control to individual citizens, and they will
help COVID-19 containment. *Ethics Inf Technol* (2021), doi:10.1007/s10676-020-
09572-w.

56. I. Z. Kiss, D. M. Green, R. R. Kao, Infectious disease control using contact tracing in
random and scale-free networks. *Journal of The Royal Society Interface*. **3**, 55–62
(2006).

57. J. Stehlé, N. Voirin, A. Barrat, C. Cattuto, L. Isella, J.-F. Pinton, M. Quaggiotto, W. V. den
Broeck, C. Régis, B. Lina, P. Vanhems, High-Resolution Measurements of Face-to-
Face Contact Patterns in a Primary School. *PLOS ONE*. **6**, e23176 (2011).

58. M. Génois, A. Barrat, Can co-location be used as a proxy for face-to-face contacts? *EPJ*
Data Sci. **7**, 1–18 (2018).

59. D. J. A. Toth, M. Leecaster, W. B. P. Pettey, A. V. Gundlapalli, H. Gao, J. J. Rainey, A.
Uzicanin, M. H. Samore, The role of heterogeneity in contact timing and duration in
network models of influenza spread in schools. *Journal of The Royal Society Interface*.
12, 20150279 (2015).

60. H. Hartle, F. Papadopoulos, D. Krioukov, Dynamic Hidden-Variable Network Models.
arXiv:2101.00414 [physics] (2021) (available at http://arxiv.org/abs/2101.00414).

61. M. Boguñá, R. Pastor-Satorras, Class of correlated random networks with hidden
variables. *Phys. Rev. E*. **68**, 036112 (2003).

62. G. Laurent, J. Saramäki, M. Karsai, From calls to communities: a model for time-varying
social networks. *Eur. Phys. J. B*. **88**, 301 (2015).

Acknowledgements

Funding: This study was partially supported by the ANR project DATAREDUX (ANR-19-CE46-0008-01) to AB, VC and PYB; EU H2020 grants MOOD (H2020-874850) to VC, CP, and PYB, and RECOVER (H2020-101003589) to VC; the Municipality of Paris (https://www.paris.fr/) through the programme Emergence(s) to JAML, BAG, PB, FP and CP; the ANR and Fondation de France through the project NoCOV (00105995) to PYB, CP; the Spanish Ministry of Science and Innovation to JAML and BAG, the AEI and FEDER (EU) under the grant PACSS (RTI2018-093732-B-C22) JAML and BAG; the Maria de Maeztu program for Units of Excellence in R&D (MDM-2017-0711) to JAML and BAG.

Author contributions: CP conceived the study; PYB, VC, AB, CP designed the study and interpreted the results; CP, wrote the original draft; PB, LB, FP implemented the synthetic population model; JAML, BAG, implemented the epidemic model; All authors edited the manuscript and approved its final version. **Competing interests:** All authors declare that they have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper or available from public sources cited on the paper.

Figures and Tables

Figure 1 Synthetic population. A-E Key statistics used as input for the synthetic population reconstruction. **A**
Age pyramid for France (source INSEE). **B** Household size (source INSEE). **C** Ratio of contacts by setting with
respect to household contacts (*33*). **D** Fraction of contacts occurring each day or less frequently (*33*). **E**
Smartphone penetration by age. The overall average adoption in the population is 64% (*29*). **F** Distribution of the
number of daily contacts in the model. **G** Cumulative distribution of the activation rate associated to the contacts
in the model, calibrated in order to be consistent with the information of panel D. **H** Sketch of the construction of
the contact network: contacts among individuals were represented as a multi-layer dynamical network, where
each layer includes contacts occurring in a specific setting. **I** Age contact matrix computed from the contact
network model.

A

B

transition	transition rate
$S \to E$	$\beta\,\sigma_A\,\beta_I\,\omega_s$
$E \to I_{p,sc}$	$\epsilon\,p_{sc}^A$
$E \to I_{p,c}$	$\epsilon\,(1 - p_{sc}^A)$
$I_{p,sc} \to I_c,\ I_{p,c} \to I_c$	μ_p
$I_{sc},\ I_c \to R$	μ

C   **D**

Figure 2. Modelling COVID-19 epidemic. **A, B** Compartmental model summarizing the epidemic states and

transitions between states. Parameters and their values are reported in Table 1. **C** Cases by age for an

uncontrolled epidemic. We show all cases (clinical and subclinical) in red and clinical cases in black. The grey line

shows the clinical cases in the early stage of the epidemic (here defined as the first 30 days), with less cases

among children than in later stages. **D** Transmission by setting (H, W, S, C, T stand respectively for household,

workplace, school, community, transport). The simulations were done with $\beta = 0.25$ corresponding to $R_0 = 3.1$.

Additional aspects of the outbreak are reported in the Supplementary Material.

Figure 3 Impact of digital contact tracing and household isolation on the epidemic. A, B Incidence (clinical
cases) according to app adoption for $R = 2.6$ and $R = 1.7$, respectively. The black curve shows the scenario with
no intervention (NI). Other curves correspond to app adoption levels ranging from 0% (household isolation only)
to 57% (90% of smartphone users). Incidence threshold level corresponding to ICU saturation is showed as a

dashed grey line in panel B. **C** Relative reduction (RR) in attack rate (AR) and peak incidence (PI) as a function of

the app adoption for the scenarii shown in A and B. RR is computed as $\frac{x_{ref}-x}{x_{ref}}$, where x is either PI or AR and x_{ref}

is the value of the quantity with no intervention. Attack rate is computed as cumulative incidence discounting initial

immunity (10%). **D, G** Peak incidence and attack rate according to reproduction ratio R and app adoption. **E, H**

Peak incidence and attack rate according to app adoption and percentage of clinical cases detected. **F, I** Peak

incidence and attack rate according to app adoption and initial immunity. **J** Peak incidence according to app

adoption and compliance to isolation of contacts notified by the app. **K** Peak incidence according to app adoption

and compliance to isolation of household contacts. **L** Peak incidence according to app adoption and percentage

of subclinical cases detected. Except as otherwise indicated, parameters values were: initial immunity 10%,

clinical case detection 50%, subclinical case detection 5%, compliance to isolation of contacts notified by the app

90%, compliance to isolation of household contacts 90%, $R = 2.6$.

Figure 4 Effect of digital contact tracing and household isolation by age and setting. A Repartition among

the different settings of the contacts detected by contact tracing (57% app adoption). **B** Relative reduction in

transmission (TRR) by setting obtained with household isolation. The relative reduction in transmission is here

defined as $TRR(s) = \frac{I_{ref}^{s}-I^{s}}{I_{ref}^{s}}$, where I^{s} is the total number of clinical and subclinical cases infected in setting s, in

Figure 5 impact of combined digital contact tracing and household isolation on the isolation of

individuals. **A** Distribution of the number of isolated individuals per detected case (DC) for 57% of

app adoption. **D** Average number of isolated individuals per detected case as a function of app

adoption. **B, E** Percentage of the population isolated as a function of time for $R = 1.7$ (B) $R = 2.6$ (E).

C, F: Fraction of unnecessary isolated, i.e. fraction of contacts isolated without being positive.

Table 1. Compartmental model parameters and their values

Parameter	Description	Values	Source
IP	Incubation period	5.2 days	(*34*)
μ_p	Rate of developing symptoms for pre-symptomatic individuals	$(2.3 \text{ days})^{-1}$	(*35*)
ϵ	Rate of becoming infectious for exposed individuals	$(2.9 \text{ days})^{-1}$	$IP - \mu_p^{-1}$
μ	Recovery rate	$(7 \text{ days})^{-1}$	(*35*)
β_I	transmissibility rescaling according to the infectious stage	0.51 for $I_{p,sc}$, I_{sc} 1 for $I_{p,c}$, I_c	(*1*)
ω_s	Transmission risk by layer	1 for H layer 0.3 for C layer 0.5 otherwise	(*16, 33*)
β	Transmission rate	Explored between 0.1 to 0.25	
σ_A	susceptibility	0.23 for A in [0,14] 0.68 for A in [15,64] 1 for A in 65 +	(*9*)
p_{sc}^A	Proportion of subclinical cases	0.27 for A in [0,1] 0.48 for A in [2,6] 0.57 for A in [7,19] 0.43 for A in [20,29] 0.38 for A in [30,39] 0.30 for A in [40,49] 0.24 for A in [50,59] 0.15 for A in [60,69] 0.11 for A in [70,79] 0.12 for A in [80,89] 0.26 for A in 90+	(*13*)

Supplementary Material of

Anatomy of digital contact tracing: role of age,
transmission setting, adoption and case detection

Additional methods

Algorithms for the generation of the synthetic population

The generation of the synthetic population is a stochastic process resulting in a contact

network being slightly different each time it is generated. Thanks to this mechanism, we can

account for some of the uncertainty concerning the input data. It also allows for population

scaling to reduce the population by respecting its composition and spatial distribution, thus

increasing computational efficiency. Specifically, the scaling decreases the number of

individuals in municipalities and the fluxes of commuters between them, but it does not

impact the number of municipalities nor the number of schools and workplaces. The smaller

population has a smaller number of households, but it maintains the statistics regarding

family size and age structure given by the INSEE data.

Households

Census data on age structure and household type and size are used to randomly assign age

and locate individuals in households. Five different types of household are considered: single

person, single with children, couple without children, couple with children, other household

groups; some of the household types may also contain an additional adult member (usually

an elderly person or a relative: if the number of additional adults is greater than one, the

household falls in the "other" category). For each municipality m with population size pop_m,

we generate new households until the size of the virtual population of the municipality vir_m

reaches the real size of population pop_m. For each household, we determine its type, its size

and the presence of an additional member (if the household type is not single-person or

couple without children, and in the case of a household with children if the size is greater

than the number of adults plus one): then, according to the role of each individual (adult,

child or other) we randomly extract his/her age, with some additional conditions:

C1: the age of any child is between 15 and 45 years less than that of the youngest parent;

C2: spouses' ages differ by no more than 15 years.

The detailed procedure is summarized in Algorithm 1.

Algorithm S1: Creation of households in municipalities

for *each municipality* **do**

     $pop_m \leftarrow$ number of individuals in *m*;

     $vir_m \leftarrow 0$;

     while $vir_m < pop_m$ **do**

         determine household type $t \sim M(p_T)$, where $M(p_T)$ is a multinomial distribution with probabilities p_T given in the first column of Table 1;

         determine age class of household head $c \sim M(p_C(t))$, where $M(p_C(t))$ is a multinomial distribution with probabilities $p_C(t)$ given in Table 2;

         determine if the household contains an additional member according to probabilities $p_E(t)$ given in the second column of Table 1;

         determine age of household head $a \sim M(p_A(c))$, where $M(p_A(c))$ is the multinomial distribution of the French age structure in the interval *c*;

         for *each other member of household* **do**

             determine the role *r* of the member (other adult in couple, children of couple; children of single, other);

             determine age class $c_m \sim M(p_{Cm}(r))$, where $M(p_{Cm}(r))$ is a multinomial distribution with probabilities $p_{Cm}(r)$ given in Table 4;

             determine the exact age $a_m \sim M(p_A(c_m))$ with the additional constraints C1 and C2;

end

$vir_m \leftarrow vir_m + s$

end

End

889 Table S1 For each household type, the fist column shows its frequency; the second column the probability that

890 the household type contains an additional member; the third column the frequency of individuals in each

891 household type

Type of household	Frequency	Fr. of add. member	Fr. of individuals
Single without children	0.338	0.0	0.149
Couple without children	0.268	0.0204	0.243
Couple with children	0.277	0.0096	0.475
Single with children	0.09	0.0123	0.106
Other	0.027	0.0	0.027

893 Table S2 For each household type, the frequency of the age class of the household head

Type of household	0 - 14	15 - 19	20 - 24	25 - 39	40 - 54	55 - 64	65 - 79	80 - 100
Single without children	0.0	0.019	0.075	0.191	0.186	0.162	0.212	0.155

Couple without children	0.0	0.029	0.132	0.263	0.264	0.133	0.094	0.085
Couple with children	0.0	0.002	0.023	0.271	0.47	0.116	0.073	0.045
Single with children	0.0	0.001	0.021	0.251	0.313	0.197	0.17	0.047
Other	0.0	0.008	0.042	0.233	0.283	0.176	0.173	0.085

895 Table S3 Probability distributions of the size of the household (except singles and couples without children,

896 having size 1 and 2, respectively) for each age class of the household head. Rows are the age class of the

897 household head, while columns are the size of the household

Age class	size: 1	size: 2	size: 3	size: 4	size: 5	size: 6
0 - 19	0.0	0.3386	0.4432	0.1169	0.0479	0.0534
20 - 24	0.0	0.1247	0.6325	0.1732	0.044	0.0256
25 - 29	0.0	0.169	0.5217	0.2291	0.0585	0.0217
30 - 34	0.0	0.1327	0.3919	0.343	0.1008	0.0316
35 - 39	0.0	0.106	0.273	0.411	0.1594	0.0506
40 - 44	0.0	0.1087	0.2546	0.3984	0.1767	0.0616
45 - 49	0.0	0.1123	0.3323	0.3568	0.1434	0.0552
50 - 54	0.0	0.1021	0.4621	0.2887	0.1005	0.0466
55 - 59	0.0	0.0997	0.5513	0.2269	0.0763	0.0458
60 - 64	0.0	0.1119	0.5888	0.1884	0.0657	0.0452
65 - 69	0.0	0.2312	0.5307	0.1469	0.0526	0.0386
70 - 74	0.0	0.3896	0.4509	0.1026	0.0349	0.022
75 - 79	0.0	0.5679	0.3392	0.0643	0.0183	0.0103
80 - 100	0.0	0.7572	0.1956	0.0335	0.0087	0.005

899   Table S4 Probability of age class of individuals depending on his role in the household, excluding the household

900   head

Age class	Child of couple	Child of a single	Adult of couple with children	Adult of couple without children	Other
0 - 4	0.2384	0.1288	0.0	0.0	0.0161
5 - 9	0.2264	0.181	0.0	0.0	0.0176
10 - 14	0.212	0.2125	0.0	0.0	0.0226
15 - 19	0.1777	0.2131	0.0035	0.0008	0.0825
20 - 24	0.0886	0.1144	0.0445	0.0164	0.1587
25 - 29	0.0305	0.0448	0.071	0.0718	0.1051
30 - 34	0.0105	0.0204	0.0414	0.1392	0.066
35 - 39	0.0064	0.0182	0.0244	0.1848	0.0576
40 - 44	0.0045	0.0181	0.0224	0.1873	0.0577
45 - 49	0.0028	0.0172	0.0434	0.1642	0.0578
50 - 54	0.0014	0.0139	0.0902	0.111	0.057
55 - 59	0.0006	0.0101	0.1379	0.0621	0.057
60 - 64	0.0002	0.0053	0.1537	0.0306	0.0506
65 - 69	0.0	0.0016	0.1126	0.0135	0.0364
70 - 74	0.0	0.0005	0.0993	0.0086	0.0356
75 - 79	0.0	0.0001	0.0812	0.0056	0.0382
80 - 100	0.0	0.0	0.0745	0.0041	0.0835

Employment

School and industry census data are used to randomly assign an employment category to

each individual on the basis of age: the probabilities are reported in Table S5. The table

assumes that all the children attend school from elementary to high school, while the

attendance of universities is taken from census data. Census data contains the number of

individuals commuting from one municipality to another one, specifying if they are students or

workers, and the same for the number of individuals that study or work in the same

municipality where they live. Such information is used to randomly assign students and

workers to a municipality of work, which may be different from the one where they live, and

then to a random school building or workplace inside this municipality. Students are assigned

to a random school building, while workers are assigned to a random workplace type (5

types, depending on the workplace size, i.e., the number of employees) and then to a

random workplace building. Students are not grouped in classes.

Table S5 Probability of workplace kind by age: individuals in the last column stay at home

Age class	creche	maternel	elementaire	college	lycee	universite	work	neet
0 - 3	0.6	-	-	-	-	-	-	0.4
4 - 6	-	1.0	-	-	-	-	-	-
7 - 11	-	-	1.0	-	-	-	-	-
12 - 15	-	-	-	1.0	-	-	-	-
16 - 18	-	-	-	-	1.0	-	-	-
19 - 21	-	-	-	-	-	0.365	0.448	0.187
22 - 23	-	-	-	-	-	0.332	0.471	0.197
24 - 26	-	-	-	-	-	0.026	0.687	0.287
27 - 62	-	-	-	-	-	-	0.705	0.295
63 - 100	-	-	-	-	-	-	-	1.0

Face-to-face contact network

Acquaintance network

For each place, i – a workplace (W layer), a school (S layer) and a municipality (C and T

layers) – we build an Erdős–Rényi network, with average degree χ_i. The latter is a

stochastic variable and depends on place layer, s_i, and size, n_i. We draw it from a gamma

distribution with average $\overline{\chi(s,n)}$ and coefficient of variation CV.

We expect that when the size of a place is small each individual enters in contact with

everybody. As the size increases the number of contacts saturates. We model this by

assuming $\overline{\chi(s,n)} = \frac{w_s\,(n-1)}{w_s+(n-1)}$. The function approaches $(n-1)$ for small n and saturates to w_s

as n increases (Figure S1).

For each setting the parameter w_s is tuned to reproduce the overall proportion of contacts

occurring in the layer. CV rules the level of heterogeneity among places of the same kind and

size. For simplicity we assume it to be the same for all settings. Additional details on the

parametrization are provided below.

Daily contact network

Once the acquaintance network is built a daily activation rate x is assigned to each link

according to a cumulative distribution $F_s(x)$ that depends on the layer s. We model $F_s(x)$ with

a sigmoid function with two parameters, A_s and B_s. For simplicity we assume $F_s(x)$ to be the

same for $s = W, S, C$, while we allow it to be different in households (where contacts are more

frequent) and in transports (where contacts are sporadic). On average, a fraction $\langle x \rangle_s$ of the

links of the acquaintance network is active each day. By indicating with K_s and k_s the

average degree of a layer in the acquaintance and daily networks, respectively, we have

$\langle x \rangle_s = \dfrac{k_s}{K_s}.$

Parametrization

We tuned parameters w_s, A_s, B_s and CV to reproduce the contact statistics in (*33*), namely:

• the average daily number of contacts is 10;

• the contact distribution is skewed with mode 3;

• being k_s the average daily number of contacts in setting s and $r_s = \dfrac{k_s}{k_H}$, the survey reports

$r_W = 3.17$, $r_S = 1.55$, $r_C = 1.86$, and $r_T = 0.23$.

• 35% of the registered contacts were with people met every day, while the rest with

people met less frequently.

Specifically, these data provide the following constraints:

• Combining point 1) and 2) above we get $k_H(1 + r_W + r_S + r_C) = 10$, meaning $k_H = 1.28$.

The household statistics used for our synthetic population reconstruction yields $K_H =$

1.97. This implies $\langle x \rangle_H = 0.65$.

•   We assume that the daily contact network has 35% of links with activity rate >0.95. In

order to do so we must account for the fact that being $f_s(x)$ the distribution of activation

rate values assigned to links of the acquaintance network (i.e. the probability density

function associated to $F_s(x)$), the distribution of links sampled in the daily network is

biased toward higher rates, i.e. it is given by $\frac{x}{\langle x \rangle} f_s(x)$.

In addition to these, we assume uncommon but still possible to meet more than once people

in transports within a time frame of one or few months. Thus, we assume an activation rate

for links in the transport layer as high as a few percent or lower. Based on these constraints

we design the frequency distribution as in Figure S2.

Once $F_s(x)$ is parametrized we tune the parameters w_s of the acquaintance network to

reproduce the proportion r_s of daily contacts in different settings. We then fix $CV = 0.2$ to

reproduce the mode of the distribution. The main properties of the network (contact

distribution, link activation frequency, and age contact matrix) are shown in Figure 1 of the

main paper. Other features are summaries in Table S1.

Figure S2 Cumulative distribution of the daily activation rate of contacts. We model it with the function $F(x) =$

$A \tanh^{-1}(1 - 2x) + B$. Parameters are the following: $A = 0.25$ for all settings; $B_H = 0.65, B_T = -0.40, B_s = 0.56$

$(s = S, W, C)$.

Table S6 Main network features

Parameter	Description	Value
$\langle x \rangle_s, s = S, W, C$	Average activation rate of contacts for the school, workplace, community layer	0.56
$\langle x \rangle_H$	Average activation rate of contacts for the household layer	0.65
$\langle x \rangle_T$	Average frequency of contacts for the transport layer	0.007
k_H (K_H)	Average degree of the daily network (acquaintance network) in a household	1.79 (2.31)
k_W (K_W)	Average degree of the daily network (acquaintance network) in a workplace	12.9 (22)
k_S (K_S)	Average degree of the daily network (acquaintance network) in a school	7.9 (14)
k_C (K_C)	Average degree of the daily network (acquaintance network) in a community	2.6 (4.3)
k_T (K_T)	Average degree of the daily network (acquaintance network) in transport	1.1 (21)

Details of the numerical simulations

977   Simulations are discrete-time and stochastic. At each time step, corresponding to one day,

978   three processes occur: *(i)* the contact network is sampled according to the activation rate of

979   each link; *(ii)* for each node, the infectious status is updated; and *(iii)* cases and contacts are

980   isolated, or get out from isolation.

981 The transmission process is modelled through the links of the multi-layer network as follows.

982 At each time step, a susceptible node i gets infected with the following probability

983 $\Lambda_i = 1 - \left(\prod_s \prod_{j \in v_s} (1 - \sigma_{A,i} \beta \, \beta_{I,j} \, \omega_s \, \delta_j) \right),$

984 where j is a node belonging to the neighborhood v_s of i on layer s, and δ_j is 1 if j belongs to

985 any infectious stage $(I_{p,sc}, I_{p,c}, I_{sc}, I_c)$ and 0 otherwise. Links of layer s have weight ω_s, that

986 represent the average level of risk associated to contacts occurring in the setting s. We

987 assume that individuals in the I_c state stay at home due to illness and can therefore transmit

988 the disease only through the links of the household layer.

989 Simulations are run on the synthetic municipality of Strasbourg. The population is reduced by

990 a factor 3 to shorten simulation time to feasible levels, yielding a population of 92,423

991 individuals.

992 A single-run simulation is executed with no modelled intervention, until the desired immunity

993 level is reached. This guarantees that immune individuals are realistically clustered in space.

994 Then the simulations of the epidemic with contact tracing are started, considering 15

995 individuals initially infected randomly assigned in the population.

996 Quantities shown in Figures 2, 3, 4, and 5 of the main paper are computed by averaging

997 results over different stochastic realizations. We run 100 stochastic simulations. Increased

998 statistics (300 runs) was necessary to accurately compare the relative reduction in incidence

999 obtained with low app adoption levels – i.e. household isolation only, app adoption 6% and

1000 app adoption 19%.

1001 We vary COVID-19 transmission potential by tuning the daily transmission rate per contact β.

1002 The reproductive number R is computed numerically as the average number of infections

1003 each infected individual generates throughout its infectious period. To do so, population

1004 immunity at the beginning of the simulation is set to 0 and R is computed considering the

infections generated by individuals who get infected the first two time-steps of the simulations

to guarantee that the whole population is susceptible. We find that $\beta = 0.1, 0.125, 0.15,$

$\ldots, 0.25$ corresponds to the following R values: 1.47 95% CI [0.0, 4.86], 1.75 95% CI [0.0,

5.78], 2.05 95% CI [0.0, 5.31], 2.25 95% CI [0.0, 5.31], 2.61 95% CI [0.0, 6.13], 2.95 95% CI

[0.06, 7.29], 3.09 95% CI [0.68, 8.11]. We also computed numerically the generation time

from the infector-infected pairs.

1011

Additional results

Uncontrolled epidemic

1015   Figure S3 Epidemic in an uncontrolled scenario. **A** Incidence of clinical cases. **B** Incidence of all cases. **C** Attack

1016   rate. The bundle of curves shows 70 stochastic realizations. The epidemic is obtained with transmission rate $\beta =$

1017 0.25 corresponding to R_0 = 3.1.

App adoption variable by age

The model accounted for age-varying smartphone penetration. However, we assumed that

the probability of downloading the app, provided an individual owns a smartphone, is uniform

and independent on age – uniform app adoption scenario, U. This hypothesis is simplified.

Indeed, elderly people may be less inclined to use the app even when they own a

smartphone. We also tested the extreme case scenario in which no individual in the 70+ age

cohort adopts the app – non-uniform app adoption scenario, NU. We consider a 32% app

coverage over the whole population, and we compared U and NU scenarios, assuming the

same number of apps are downloaded in the two cases. Figure S4 shows the attack rate

relative reduction $ARR = \frac{AR_U^A - AR_{NU}^A}{AR_U^A}$ by age group, where AR_{sce}^A is the attack rate of the

epidemic for the age group, A, and the scenario, $sce = U, NU$. We found that ARR is close to

zero, meaning that $AR_{NU}^A \simeq AR_U^A$, for all age groups (Figure S4). This means that distributing

the app only to individuals younger than 70 years would not reduce the protection in the 70+

age group.

1032

1033

1034   Figure S4 Comparison between the U and the NU contact tracing scenario. Here $R = 2.6$, Immunity is 10%,

1035   detection probability is 50% and app penetration is 32%. The line shows the average attack rate relative reduction

1036   and the shaded area is standard deviation.

1037

1038