

A reference for mortality in Spain from 2001 to 2019 records with an accurate estimate of excess deaths during the 2020 spring covid-19 outbreak

José María Martín-Olalla*

Universidad de Sevilla. Facultad de Física. Departamento de Física de la Materia Condensada. ES41012 Sevilla. Spain

(Dated: 20200722.1327+0200)

Spanish official records of mortality and population in the 21st century are analyzed to determine the reference value of all cause, all age, all sex per capita death rate in the country.

This reference is used to analyze the mortality in 2020, largely influenced by a massive anomaly in spring due to covid-19.

The most probable mortality excess from Monday March 2 to Sunday May 24, 2020 — W10 to W21 — is 1055(17) deaths per one million population or 49 940(730) total deaths. Standard deviation is 67 deaths per one million population. The excess amounts to 53 % relative to the expected number of deaths from 21st century records and yields a z -score $z = 16$ —extremely high excess, according to EuroMoMo classification scheme.

Taking into account nationwide seroprevalence(Pollán *et al.*, 2020) that is an infection fatality rate of 2.1 %.

Keywords: mortality excess; z -score; EuroMoMo; INE; Instituto de Salud Carlos III

CONTENTS

1. Introduction	2
2. Datasets	2
3. A reference for mortality in Spain from the series 2001 to 2019	3
4. Excess death rate in 2020	4
5. Spring 2020 covid-19 excess deaths	6
6. Conclusions	6
7. Conflict of interest statement	8
8. Data availability	8
References	8
Acknowledgments	10

* olalla@us.es; Twitter: [@MartinOlalla_JM](https://twitter.com/MartinOlalla_JM);



1. INTRODUCTION

The illness designated covid-19 caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) caught worldwide attention since its identification in late 2019 (Zhu *et al.*, 2020). Spain is one of the European countries most impacted by the disease during the spring of 2020. Confirmed covid-19 cases climbed up to several hundred thousands—some few thousands per one million population—and confirmed covid-19 deaths to some 28 000 or six hundred deaths per one million population.¹ Nonetheless these numbers are only a fraction of the excess death recorded by mortality monitors (ISC3, 2020)² and a fraction of infections as detected by seroprevalence studies (Pollán *et al.*, 2020).

Total excess deaths is a key quantity to understand the acute impact of a pandemic (Aaron *et al.*, 2020). Its determination sensibly requires a reference mortality and a time interval over which the excess is determined. This manuscript takes official records of all cause, all age, all sex weekly mortality in Spain in the 21st century to ascertain a sensible reference value of weekly per capita mortality as a function of the week of the year. When this reference is used, 2020 mortality from Monday March 2 to Sunday May 24 2020 yields an excess of 1055 deaths per one million population or 49 940 excess deaths. In the classification scheme of EuroMoMo the z -score is 16, an extremely high deviation for an elapsed time interval equal to eleven weeks.

2. DATASETS

The Spanish Instituto Nacional de Estadística (INE) usually releases at the end of a calendar year mortality reports corresponding to the preceding year. INE has access to every single death certificate in Spain.

On June 3th 2020 the INE announced preliminary reports (INE, 2020a) on weekly all cause, all age number of deaths in Spain for 2020. On their first press release INE announced³ 44 000 excess deaths in Spain until week 21 of the year 2020, with the reference set to mortality in 2019 until week 21.

A csv dataset is available at the link provided in the reference. This report will analyze the July 15th, 2020 update which collects reports until Sunday July 5th, 2020.

The dataset contains consolidated all cause, all age weekly mortality deaths from 2000 to 2018 and the preliminary results from 2019, alongside with 2020 fresh data. The dataset is organized by the ISO-8601 week date which considers regular years with $w = 52$ weeks (364 days) and extraordinary (“leap”) years with $w = 53$ weeks (371 days). In the dataset 2004, 2009, 2015 and 2020 are “leap” years.

The dataset follows the European convention: (1) a week begins on Monday and ends on Sunday; (2) a year starts on the Monday following the first Thursday of a calendar year and ends on the Sunday following the last Thursday of the calendar year. Every single calendar date belongs to only one week of the ISO-8601 week date; there is no gap and no overlap. Therefore the first week (W01) and the last week of a year (W52 or W53) may contain dates from adjacent calendar years: dates from December 29st to January 3rd can fall on the first week of a year or in the last week of a year.

Any analysis of mortality over a range of 20 years must of necessity account for population changes during the period. INE (INE, 2020b) publishes (table 31 304) the official numbers of people living in Spain on a regular basis. The dataset starts on 1996 and it is updated twice a year with the population for January 1st and for July 1st. The last available figures are those of January 1st, 2020.

In this work this dataset will be translated in week dates and linear interpolation with splines will produce an array N_{ij} of estimated Spanish population from 2000-W01 to 2020-W01. Every week of the year 2020 will equally collect the last available data.

Since the start of the covid-19 worldwide outbreak the Centro de Coordinación de Alertas y Emergencias Sanitarias (CCAES)—a Spanish equivalent for the Centers for Disease Control and Prevention CDC, and enrolled in the Instituto de Salud Carlos III (ISC3)—released situation reports and, later, started to release the number of covid-19 cases detected in Spain and the number of deaths within this subset of population. This dataset has been largely criticized during in the spring of 2020. It was often refreshed and it usually contained inconsistent figures. Since mid June the dataset—showing total number of cases—gained stability and accuracy reporting by date of symptom start or, if unknown, by the sixth day prior to the date of diagnostic positive test. The dataset is available at the link provided in footnote 1. The database on covid-19 deaths was also cleaned up but it is no longer available within

¹ Instituto de Salud Carlos III covid-19 in Spain <https://cnecovid.isciii.es/covid19>

² https://momo.isciii.es/public/momo/dashboard/momo_dashboard.html

³ https://www.ine.es/prensa/experimental_edes.pdf




FIG. 1 All-cause all-age weekly per capita death rate in Spain during the 21st century. Wavy black line is observed death rate while orange line is the 52-week moving average or the yearly per capita all-cause all-age death toll (right axis). Left and right scales are conformal. They have only been displaced relative to each other to improve the readability of both lines. Blueish straight line shows the overall trend, an increase of 1% for the full range or 500 parts per million and per year. The null hypothesis “per capita deaths does not depend on week date” sustains ($p = 0.43$).

official repositories. Yet journalists like Datadista maintain accessible csv files on GitHub where information can be retrieved.⁴⁵

3. A REFERENCE FOR MORTALITY IN SPAIN FROM THE SERIES 2001 TO 2019

From 2001 to 2019 weekly number of deaths D_{ij} and weekly population N_{ij} were used to collect the weekly all-cause, all-age, all sex per capita⁶ death rate in Spain $d_{ij} = D_{ij}/N_{ij}$ (see Figure 1). Save for the “high frequency” seasonal variability the relative change in d_{ij} is on average 1% for the full range. If bivariate descriptive analysis where the predictor x is week date and the response y is d_{ij} are performed then the null hypothesis “per capita weekly death rate is unrelated to x ” sustains ($p = 0.43$) at the standard level of confidence. Contrastingly, D_{ij} increased on average 16% and its bivariate analysis does not sustain ($p = 3 \times 10^{-16}$) the null hypothesis.

Therefore a sensitive conclusion is d_{ij} remained still over the elapsed 19 years or 991 weeks, save for “high frequency” seasonal variability. In practical terms all cause, all age, all sex weekly deaths in Spain can be expressed as $D = N \times f(x_i)$, where $f(x_i)$ is some function of social inputs—including sex and age—and environmental inputs—like subsolar latitude—which remained unaltered over the span of twenty years. This shows the stability of this period.

Figure 2 collapses the full set of weekly death data into a $w = 52$ week x -axis corresponding to every of the first w weeks in a week year. The multicolored step lines display the consolidated values in the range 2001 to 2018 and the provisional 2019 value.

Every week i exhibits a distribution d_{ij} of $n = 19$ values from 2001 to 2019. A Kolmogorov-Smirnov test of normality sustains the null hypothesis “sample comes from a normal distribution” at the standard level of confidence for every week except for W33 ($p = 0.043$). Univariate central tendency and dispersion will describe the sample through the

⁴ <https://github.com/datadista/datasets/tree/master/COVID%2019>

⁵ https://raw.githubusercontent.com/datadista/datasets/master/COVID%2019/ccaa_covid19_fallecidos_por_fecha_defuncion_nueva_serie_long.csv

⁶ This manuscript will translate per capita values into “per one million population”. The symbol M^{-1} will be used for this purpose. Notice, however, that the use of isolated SI prefixes is not recommended. In this case M^{-1} means a fraction 1×10^{-6} of 1. For that purpose μ could have been equally used. Sometimes rates will be computed per capita and per some unit of time, like $M^{-1}a^{-1}$ which means per one million population and per year and not per one million years, symbol Ma^{-1} . Notice again that $M^{-1}a^{-1}$ is not a recommended notation.




FIG. 2 Spanish all cause, all age per capita death rate in the 21st century as a function of the week calendar. Multicolored thin step lines display values for 2001 to 2019. Black thick step line shows average value r_i with standard deviation displayed by the light blue colored strip band ($\pm 2.33s_i$). Vertical axis display values per one million population and per week (left) or per year (right).

sample average r_i , sample standard deviation s_i and sample error of the mean (SEM) $u_i = s_i/\sqrt{n}$. Figure 2 displays the sample average r_i (thick black step line) and a strip band located at $r_i \pm 2.33s_i$ which under the null hypothesis of normality should contain 99% of the observations. Therefore 978 out of the $w \times n = 988$ observations should lie within the colored strip band in Figure 2.

There are 18 observations lying outside the colored strip band. However 4 of this outliers are located the 2003 heat wave while 3 of them are located on 2012 and can be traced to the impact of a strong influenza wave (León-Gómez *et al.*, 2015), also annotated in Figure 1. The remaining 11 outliers are already compatible with the number of outliers under normality even though they might be also linked to the impact of some disease or to environmental conditions.

Figure 1 and Figure 2 also display the seasonality of death rates. Generally speaking death rates are higher from W48 —early November— to W10 —early March— in the following year. This is associated to winter mortality, the impact of influenza and cold waves. Mortality is lower from W10 to W48 but variability still remains in the summer weeks as a result of the impact of variable heat waves. Specifically 2003–W32 and 2003–W33 are easily spotted on Figure 2. They are contemporaneous with the strong 2003 heat wave that increased mortality across Europe and which is annotated in Figure 1.

Table 1 displays the average r_i —a reference for the 2001 to 2019 death rate in Spain— and the sample error of the mean (SEM) in brackets. The average value r_i is the most probable value for the death rate in 2020 under the null-hypothesis of normality following the preceding nineteen years.

The table also shows the accumulated value in a 52 week year which is $8607(71) \text{ M}^{-1} \text{ a}^{-1}$ (death rate per one million population and per year) or $165.5(14) \text{ M}^{-1} \text{ w}^{-1}$ (death rate per one million population and per week) or $23.64(19) \text{ M}^{-1} \text{ d}^{-1}$ (death rate per one million population and per day).

4. EXCESS DEATH RATE IN 2020

Table 1 also lists observed all cause, all age death rates in 2020 o_i per week and per one million population, alongside with the accumulated value until W27. It is assumed that the observed 2020 value has zero uncertainty, albeit it is a provisional figure. The hypothesis just means o_i is not a source of uncertainty, which in practical terms implies excess death uncertainty is given by the sample error of the mean for the 2001 to 2019 series.

In the same way Table 1 displays three statistics related to the 2020 excess rate of death.

	Series 2001 – 2019		2020 values		2020 Excess			Breakdown	
	Average (SEM)	Cumulative (SEM)	Observed	Cumulative	Absolute	Relative	z -score	Covid	Unknown
	$r_i(u_i)$	$\sum_{k=1}^i r_k$	o_i	$\sum_{k=1}^i o_k$	$e_i = o_i - r_i$	e_i/r_i	e_i/s_i	c_i	$k_i = e_i - c_i$
	$M^{-1} w^{-1}$	M^{-1}	$M^{-1} w^{-1}$	M^{-1}	$M^{-1} w^{-1}$	%		$M^{-1} w^{-1}$	$M^{-1} w^{-1}$
W01	200.3(49)	200.0(49)	188.00	188.00	-12.3	-6.2	-0.6		-12.3
W02	203.2(55)	404(10)	203.06	391.06	-0.2	-0.1	0.0		-0.2
W03	203.7(55)	607(16)	213.50	604.56	9.8	4.8	0.4		9.8
W04	198.7(48)	806(20)	202.94	807.50	4.3	2.2	0.2		4.3
W05	197.7(47)	1004(24)	203.66	1011.16	6.0	3.0	0.3		6.0
W06	195.2(40)	1199(28)	198.54	1209.70	3.3	1.7	0.2		3.3
W07	192.6(43)	1391(31)	189.75	1399.45	-2.8	-1.5	-0.2	0.02	-2.9
W08	189.1(42)	1580(33)	179.63	1579.08	-9.4	-5.0	-0.5		-9.4
W09	185.7(37)	1766(36)	168.92	1748.00	-16.8	-9.0	-1.0		-16.8
W10	182.6(29)	1949(38)	182.74	1930.74	0.2	0.1	0.0	0.51	-0.3
W11	176.5(24)	2125(40)	196.09	2126.83	19.6	11.1	1.9	9.36	10.2
W12	171.7(16)	2297(41)	267.59	2394.42	95.9	56	13.3	49.65	46.3
W13	169.2(17)	2466(42)	405.43	2799.85	236.2	140	31	115.57	120.6
W14	167.4(15)	2634(43)	435.73	3235.58	268.3	160	42	118.97	149.4
W15	164.9(15)	2798(44)	363.74	3599.33	198.9	121	31	96.30	102.6
W16	162.0(16)	2960(45)	277.18	3876.51	115.2	71	16.5	71.58	43.6
W17	160.5(14)	3121(46)	218.25	4094.76	57.7	36	9.4	47.01	10.7
W18	156.8(13)	3278(47)	186.56	4281.32	29.7	19.0	5.3	31.12	-1.4
W19	157.7(12)	3435(47)	181.30	4462.63	23.6	15.0	4.4	21.64	2.0
W20	156.4(14)	3592(48)	159.94	4622.57	3.5	2.2	0.6	12.21	-8.7
W21	155.2(14)	3747(48)	161.50	4784.07	6.3	4.0	1.1	8.77	-2.5
W22	152.9(19)	3900(48)	152.08	4936.15	-0.8	-0.6	-0.1	5.79	-6.6
W23	152.1(16)	4052(49)	151.98	5088.13	-0.1	-0.1	0.0	3.44	-3.5
W24	153.5(23)	4206(50)	145.68	5233.81	-7.9	-5.1	-0.8	2.05	-9.9
W25	155.7(24)	4361(51)	149.84	5383.65	-5.8	-3.7	-0.6	0.68	-6.5
W26	155.7(19)	4517(51)	155.00	5538.65	-0.7	-0.5	-0.1	0.42	-1.1
W27	155.4(21)	4672(52)	142.17	5680.82	-13.3	-8.5	-1.5	0.21	-13.5
W28	154.5(23)	4827(52)							
W29	152.4(21)	4979(53)							
W30	153.3(19)	5133(54)							
W31	155.8(26)	5289(55)							
W32	154.2(33)	5443(56)							
W33	150.9(34)	5594(58)							
W34	148.5(16)	5742(58)							
W35	146.5(12)	5889(59)							
W36	144.3(13)	6033(59)							
W37	142.35(88)	6175(59)							
W38	143.8(14)	6319(60)							
W39	146.0(15)	6465(61)							
W40	148.5(10)	6613(62)							
W41	150.9(11)	6764(62)							
W42	153.21(91)	6918(63)							
W43	155.6(11)	7073(63)							
W44	157.3(12)	7230(64)							
W45	160.2(15)	7391(65)							
W46	164.2(18)	7555(66)							
W47	165.6(17)	7720(66)							
W48	169.3(17)	7890(67)							
W49	171.7(17)	8061(68)							
W50	175.7(20)	8237(69)							
W51	181.2(21)	8418(70)							
W52	188.2(32)	8607(71)							

TABLE 1 Sample average value of weekly per capita death rate from 2001 to 2019 and sample error of the mean (SEM) in brackets alongside to accumulated values. Also observed death rate in 2020 until W27 and accumulated values. The last set displays the 2020 per capita excess: absolute value, relative value, z -score and accumulated excess death rate. Total numbers for 2020 can be computed considering the Spanish population $N = 47\,329\,981$.

Absolute excess: The difference $e_i = o_i - r_i$ from observed death rate to predicted death rate under the normal hypothesis for the series. Absolute excess shows an anomaly around W14 —March 30 to April 5—, where the largest deviation occurs. The anomaly starts on W10 and ends on W21. Thereafter it will be referred as the 2020 spring anomaly.

Relative excess: The ratio e_i/r_i , shown as a percentage. Largest value is 160% on W14. Relative to r_i observed death rate more than doubled during three weeks.

z -score: or normalized excess. It is the ratio e_i/s_i , which tests the observed rates of deaths against a normal distribution with zero mean and unit standard deviation. The largest z -score is $z = 42$. Under the null hypothesis of normality such a deviation would have likely been observed in a sample consisting of $\sim 10^{350}$ years.

The European Monitor of Mortality uses the z -score to compare mortality across countries with different sizes. They bin the scores according to: $2 < z \leq 4$ is a low excess; $4 < z \leq 7$ is moderate; $7 < z \leq 10$ is a high excess; $10 < z \leq 15$ is very high; and $z > 15$ is extremely high. For four weeks in a row the z -score was extremely high in Spain.

Finally, Table 1 lists the breakdown of the 2020 excess in Spain with the numbers of confirmed covid-19 deaths c_i and the numbers of excess deaths without attributed cause —unknown cause— $k_i = e_i - c_i$. Both of them show an anomaly around W14. Notably, $k_i > c_i$ for W13 to W15.

Table 2 provides a better insight on how W01 to W27 accumulated mortality changed in Spain during the past twenty years. The table shows population, deaths and per capita deaths alongside with deviation from the average value for the 2001–2019 years and the expected values for 2020. Notice that population is only listed for W01, while interpolated values of population were used to obtain o . As of W27 death excess in Spain amounts to 1008(52) deaths per one million population or 47 700(2400) deaths, a 22% relative to the reference. The z -score is $z = 4.5$, meaning a moderate excess. Notice, however, this is a score computed on an elapsed time interval equal to 27 weeks.

5. SPRING 2020 COVID-19 EXCESS DEATHS

From W10 to W21 —Monday March 2 to Sunday May 24— excess death rate is anomalously high in Spain —see Table 1—. Figure 3 displays the observed death rates (o_i , black step line) in Spain for the year 2020; the reference r_i (greenish step line) and the number of deaths with unknown attributed cause (magenta step line). Succinctly Table 1 is displayed graphically in Figure 3. Moreover the dashed line displays the weekly rate confirmed covid-19 cases. It is advanced some two weeks relative to the mortality anomaly.

The darkest shaded area highlights the total number of expected deaths from W10 to W21. Intermediate shaded area is the number of deaths with unknown cause in the same period of time and the lightest shaded area is the number of confirmed covid-19 deaths in the time window. Table 3 lists accumulated values for e_i, c_i and k_i . In the elapsed time from W10 to W21 excess death rate amounts to 1055(15) per million population or 49 940(730) deaths; 55% of them (27 555 deaths) were confirmed covid-19 cases and the remaining 45% (22 383 still lacks an attributed cause). Total excess is 55% relative to expected deaths in the same period of time and the z -score for the W10–W21 period of time is 15.8, extremely high according to the EuroMoMo classification. Note however that this is the z -score for the twelve-week period of time. Weekly z -scores are listed in Table 1.

Table 4 lists the variability of the W10 to W21 accumulated number of deaths in the past twenty years and population numbers for W01. Notice however that interpolated values of population were used to compute per capita death rates. Bivariate descriptive statistics where the calendar year plays the role of a predictor and the observed per capita mortality o is the response sustains ($p = 0.48$) the null hypothesis of no relationship at the standard level of confidence. The confidence interval for the slope is $[-4, +8]M^{-1} a^{-1}$, in accordance with Figure 1. In sharp contrast if the response is set to the observed mortality O then the null hypothesis does not sustain ($p = 7 \times 10^{-8}$) at the standard level of significance. The confidence interval for the slope is $[572, 923]a^{-1}$.

6. CONCLUSIONS

A reference value for weekly per capita mortality in Spain obtained from public records which includes data in the 21st century is reported (see Table 1 and Figure 2).

W01 to W27 analysis								
Year	Observed			Deviation			2020 values	
	Population N_i	Absolute O_i	per capita $o_i = O_i/N_i$ M^{-1}	absolute $e_i = o_i - r$ M^{-1}	relative e_i/r %	z -score $z_i = e_i/s$	absolute $o_i \times N_{20}$	deviation $(o_i - r) \times N_{20}$
2001	40 665 545	187 272	4599.4	-73(52)	-1.6	-0.3	217 688	-3459(2400)
2002	41 035 271	201 227	4882.3	210(52)	4.5	0.9	231 078	9931(2400)
2003	41 827 836	199 554	4750.6	78(52)	1.7	0.3	224 843	3697(2400)
2004	42 547 454	196 676	4606.4	-66(52)	-1.4	-0.3	218 023	-3124(2400)
2005	43 310 423	216 705	4984.8	312(52)	6.7	1.4	235 931	14 784(2400)
2006	44 023 452	195 191	4417.0	-255(52)	-5.5	-1.1	209 057	-12 090(2400)
2007	44 784 659	205 474	4565.6	-107(52)	-2.3	-0.5	216 088	-5059(2400)
2008	45 668 938	205 332	4481.5	-191(52)	-4.1	-0.8	212 109	-9038(2400)
2009	46 239 271	208 111	4494.9	-178(52)	-3.8	-0.8	212 744	-8403(2400)
2010	46 489 539	202 037	4342.5	-330(52)	-7.1	-1.5	205 528	-15 619(2400)
2011	46 669 832	205 416	4398.4	-274(52)	-5.9	-1.2	208 175	-12 972(2400)
2012	46 816 223	222 828	4762.1	90(52)	1.9	0.4	225 388	4241(2400)
2013	46 727 890	207 953	4456.4	-216(52)	-4.6	-1.0	210 923	-10 224(2400)
2014	46 512 199	211 829	4556.9	-116(52)	-2.5	-0.5	215 678	-5469(2400)
2015	46 449 565	234 821	5057.4	385(52)	8.2	1.7	239 365	18 218(2400)
2016	46 440 475	216 271	4656.5	-16(52)	-0.3	-0.1	220 391	-756(2400)
2017	46 527 263	228 130	4902.8	230(52)	4.9	1.0	232 052	10 905(2400)
2018	46 658 447	234 864	5030.1	358(52)	7.7	1.6	238 073	16 926(2400)
2019	46 937 060	227 139	4831.2	159(52)	3.4	0.7	228 659	7512(2400)
AVERAGE	45 070 071	210 886	$r = 4672 \pm 52$	0.0	0.0	0.0	221 147	0.0
2020	47 329 981	268 873	5680.8	1008(52)	21.6	4.5	268 873	47 726(2400)

TABLE 2 Death rate in Spain from W01 to W27 in a span of twenty years. First column, resident population in Spain. Second column, observed accumulated deaths from W01 to W27 followed by deaths per one million population. Columns four to six list deviation from the average 2001 to 2019 (standard error of the mean in brackets); relative deviation; and the z -score or normalized deviation. Finally column 7 shows expected values for 2020 population and column 8, the expected deviation. The standard deviation of the distribution is $s = SEM \times \sqrt{19} = 230 M^{-1}$ or $s = 11\,000$ deaths for Spanish population in 2020. Averages printed in blue ink are the only statistically meaningful. Linear regression for calendar year versus observed per capita death rate sustains ($p = 0.30$) the null hypothesis of no relationship. The null hypothesis does not sustain ($p = 1.1 \times 10^{-5}$) if, instead, absolute deaths are tested.

March 2 to May 24				
	Total (SEM)	Covid	Unknown	z -score
	$a_i = \sum_{k=10}^{i \leq 21} e_k$ M^{-1}	$p_i = \sum_{k=10}^{i \leq 21} c_k$ M^{-1}	$a_i - p_i$ M^{-1}	z_i
W10	0.2(29)	0.51	-0.35	0.0
W10-W11	19.7(52)	9.87	9.87	0.9
W10-W12	115.7(66)	59.52	56.14	4.0
W10-W13	351.8(80)	175.09	176.75	10.1
W10-W14	620.2(90)	294.06	326.13	15.8
W10-W15	819(10)	390.37	428.68	18.8
W10-W16	934(11)	461.95	472.27	19.0
W10-W17	992(12)	508.96	483.00	18.6
W10-W18	1022(13)	540.08	481.61	17.9
W10-W19	1045(14)	561.72	483.62	17.4
W10-W20	1049(15)	573.93	474.92	16.5
W10-W21	1055(15)	582.70	472.41	15.8

TABLE 3 Accumulated per capita values of total excess death rate (with SEM) a_i , confirmed covid-19 deaths c_i and deaths without attributed cause $k_i = e_i - c_i$ for 2020 W10 to W21 in Spain. The z -score is $a_i \sqrt{19}/u_i$ where u_i is the SEM. Total numbers can be computed considering the Spanish population in 2020 $N = 47\,329\,981$.




FIG. 3 Weekly all cause, all age per capita death rate in Spain in 2020 (black step line) compared to the reference value 2001 to 2019 (greenish step line). Magenta step line displays deaths with unattributed cause and the dashed step line the number of confirmed covid-19 cases. Blueish shaded area is the $\pm 2.33s_i$ where s_i is the standard deviation of the reference. Lightest shaded area in the peak is the confirmed covid-19 deaths from W10 to W21. The darkest shaded area is the expected number of deaths W10 to W21. Shaded area in intermediate color represent the number of unattributed deaths W10 to W21.

This reference value is used to compute the excess of mortality in the year 2020 from W10 to W21 when weekly mortality displays an extremely high anomaly (see Figure 3) due to the impact of covid-19. The per capita excess death is 1055(15) per one million population or 49 940(730) (see Table 4).

Considering a nationwide seroprevalence(Pollán *et al.*, 2020) of 5.0% this translates into an upper bound of the infection fatality rate (IFR) of 2.1% if every excess death is attributed to the disease. The lower bound for IFR is 1.2%, obtained after considering confirmed covid-19 deaths only.

7. CONFLICT OF INTEREST STATEMENT

The author declares no conflict of interest.

8. DATA AVAILABILITY

A tsv file containing the Spanish weekly deaths from 2000 to 2020 exported from INE table 35176 and the computed values of Spanish population used in this manuscript is available at <https://personal.us.es/olalla/covid/ESPDeathsPopulation.tsv>.

REFERENCES

- Aaron, Janine, John Mullbauer, Charlie Giattino, and Hannah Ritchie (2020), “A pandemic primer on excess mortality statistics and their comparability across countries,” .
- INE, (2020a), “Estimación del número de defunciones semanales durante el brote de Covid-19,” .
- INE, (2020b), “Población residente por fecha, sexo y edad,” .
- ISC3, (2020), “Monitor de Mortalidad en España,” .

W10 to W21 analysis

Year	Observed			Deviation			2020 values	
	Population N_i	Absolute O_i	per capita $o_i = O_i/N_i$ M^{-1}	absolute $e_i = o_i - r$ M^{-1}	relative e_i/r %	z -score $z_i = e_i/s$	absolute $o_i \times N_{20}$	deviation $(o_i - r) \times N_{20}$
2001	40 665 545	81 377	1998.3	17(15)	0.9	0.3	94 578	819(730)
2002	41 035 271	84 157	2040.2	59(15)	3.0	0.9	96 564	2805(730)
2003	41 827 836	84 756	2016.5	36(15)	1.8	0.5	95 440	1681(730)
2004	42 547 454	86 210	2018.1	37(15)	1.9	0.6	95 514	1756(730)
2005	43 310 423	87 876	2019.7	39(15)	2.0	0.6	95 591	1832(730)
2006	44 023 452	84 433	1909.5	-71(15)	-3.6	-1.1	90 378	-3381(730)
2007	44 784 659	87 823	1949.7	-31(15)	-1.6	-0.5	92 280	-1479(730)
2008	45 668 938	88 214	1924.3	-57(15)	-2.9	-0.8	91 077	-2681(730)
2009	46 239 271	87 668	1893.1	-88(15)	-4.4	-1.3	89 598	-4161(730)
2010	46 489 539	87 370	1877.6	-103(15)	-5.2	-1.5	88 869	-4890(730)
2011	46 669 832	87 598	1875.4	-106(15)	-5.3	-1.6	88 765	-4994(730)
2012	46 816 223	94 284	2015.1	34(15)	1.7	0.5	95 376	1618(730)
2013	46 727 890	90 097	1931.2	-50(15)	-2.5	-0.7	91 403	-2356(730)
2014	46 512 199	89 921	1934.6	-46(15)	-2.3	-0.7	91 564	-2195(730)
2015	46 449 565	96 390	2076.1	95(15)	4.8	1.4	98 263	4504(730)
2016	46 440 475	94 855	2042.3	61(15)	3.1	0.9	96 661	2902(730)
2017	46 527 263	92 972	1998.1	17(15)	0.9	0.3	94 569	811(730)
2018	46 658 447	97 897	2096.4	115(15)	5.8	1.7	99 221	5462(730)
2019	46 937 060	95 099	2022.1	41(15)	2.1	0.6	95 707	1948(730)
AVERAGE	45 070 071	89 421	$r = 1981 \pm 15$	0.0	0.0	0.0	93 759	0.0
2020	47 329 981	143 697	3036.1	1055(15)	53	15.8	143 697	49 938(730)

TABLE 4 Death rate in Spain from W10 to W21 in a span of twenty years. First column, resident population in Spain. Second column, observed accumulated deaths from W10 to W21 followed by deaths per one million population. Columns four to six list deviation from the average 2001 to 2019 (standard error of the mean in brackets); relative deviation; and the z -score or normalized deviation. Finally column 7 shows expected values for 2020 population and column 8, the expected deviation. The standard deviation of the distribution is $s = SEM \times \sqrt{19} = 67 M^{-1}$ or $s = 3200$ deaths for Spanish population in 2020. Averages printed in blue ink are the only statistically meaningful. Linear regression for calendar year versus observed per capita death rate sustains ($p = 0.48$) the null hypothesis of no relationship. The null hypothesis does not sustain ($p = 7 \times 10^{-8}$) if, instead, absolute deaths are tested.

León-Gómez, Inmaculada, Concepción Delgado-Sanz, Silvia Jiménez-Jorge, Víctor Flores, Fernando Simón, Diana Gómez-Barroso, Amparo Larrauri, Salvador De, and Mateo Ontañón (2015), “Exceso de mortalidad relacionado con la gripe en España en el invierno de 2012,” *Gac Sanit* **29** (4), 258–265.

Pollán, Marina, Beatriz Pérez-Gómez, Roberto Pastor-Barriuso, Jesús Oteo, Miguel A Hernán, Mayte Pérez-Olmeda, Jose L Sanmartín, Aurora Fernández-García, Israel Cruz, Nerea Fernández de Larrea, Marta Molina, Francisco Rodríguez-Cabrera, Mariano Martín, Paloma Merino-Amador, Jose León Paniagua, Juan F Muñoz-Montalvo, Faustino Blanco, Raquel Yotti, Faustino Blanco, Rodrigo Gutiérrez Fernández, Mariano Martín, Saturnino Mezcuca Navarro, Marta Molina, Juan F. Muñoz-Montalvo, Matías Salinero Hernández, Jose L. Sanmartín, Manuel Cuenca-Estrella, Raquel Yotti, José León Paniagua, Nerea Fernández de Larrea, Pablo Fernández-Navarro, Roberto Pastor-Barriuso, Beatriz Pérez-Gómez, Marina Pollán, Ana Avellón, Giovanni Fedele, Aurora Fernández-García, Jesús Oteo Iglesias, María Teresa Pérez Olmeda, Israel Cruz, Maria Elena Fernandez Martinez, Francisco D. Rodríguez-Cabrera, Miguel A. Hernán, Susana Padrones Fernández, José Manuel Rumbao Aguirre, José M. Navarro Marí, Begoña Palop Borrás, Ana Belén Pérez Jiménez, Manuel Rodríguez-Iglesias, Ana María Calvo Gascón, María Luz Lou Alcaine, Ignacio Donate Suárez, Oscar Suárez Álvarez, Mercedes Rodríguez Pérez, Margarita Cases Sanchís, Carlos Javier Villafáfila Gomila, Lluís Carbo Saladrigas, Adoración Hurtado Fernández, Antonio Oliver, Elías Castro Feliciano, María Noemí González Quintana, José María Barrasa Fernández, María Araceli Hernández Betancor, Melisa Hernández Febles, Leopoldo Martín Martín, Luis-Mariano López López, Teresa Ugarte Miota, Inés De Benito Población, María Sagrario Celada Pérez, María Natalia Vallés Fernández, Tomás Maté Enríquez, Miguel Villa Arranz, Marta Domínguez-Gil González, Isabel Fernández-Natal, Gregoria Megías Lobón, Juan Luis Muñoz Bellido, Pilar Ciruela, Ariadna Mas i Casals, Maria Doladé Botías, M. Angeles Marcos Maeso, Dúnia Pérez del Campo, Antonio Félix de Castro, Ramón Limón Ramírez, Maria Francisca Elías Retamosa, Manuela Rubio González, María Sinda Blanco Lobeiras, Alberto Fuentes Losada, Antonio Aguilera, German Bou, Yolanda Caro, Noemí Marauri, Luis Miguel Soria Blanco, Isabel del Cura González, Montserrat Hernández Pascual, Roberto Alonso Fernández, Paloma Merino-Amador, Natalia Cabrera Castro, Aurora Tomás Lizcano, Cristóbal Ramírez Almagro, Manuel Segovia Hernández, Nieves Ascunce Elizaga, María Ederra Sanz, Carmen Ezpeleta Baquedano, Ana Bustinduy Bascaran, Susana Iglesias Tamayo, Luis Elorduy Otazua, Rebeca Benarroch Benarroch, Jesús Lopera Flores, and Antonia Vázquez de la Villa (2020), “Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological study,” *The Lancet* **0** (0), 10.1016/S0140-6736(20)31483-5.

Zhu, Na, Dingyu Zhang, Wenling Wang, Xingwang Li, Bo Yang, Jingdong Song, Xiang Zhao, Baoying Huang, Weifeng Shi, Roujian Lu, Peihua Niu, Faxian Zhan, Xuejun Ma, Dayan Wang, Wenbo Xu, Guizhen Wu, George F. Gao, and Wenjie Tan (2020), “A novel coronavirus from patients with pneumonia in China, 2019,” [New England Journal of Medicine](#) **382** (8), 727–733.

ACKNOWLEDGMENTS

Counting and its how-to are the least things I can do in the loving memory of those who died during the spring of 2020.

JMMO heartfully thanks the Instituto Nacional de Estadística and the Instituto de Salud Carlos III for making publicly available the datasets. JMMO also thanks to Datadista (www.datadista.com) for realising public records of covid-19 confirmed deaths.

This work was performed using free software running on xubuntu 18.04.1LTS. Data bases have been imported into GNU octave-4.2.2 (<https://www.gnu.org/software/octave/>). Pictures were developed thanks to gnuplot-5.2.2 (<http://www.gnuplot.info/>). The manuscript was typeset in GNU emacs-25.2.2 (<https://www.gnu.org/software/emacs/>) assisted by AUCTeX(<https://www.gnu.org/software/auctex/>). Data in pictures and in tables have been exported directly from octave.

This project started on July 3, 2020.