Herd immunity thresholds for SARS-CoV-2 estimated from unfolding epidemics

Ricardo Aguas[1], Jessica G. King[2], Guilherme Gonçalves[3], Marcelo U. Ferreira[4], M. Gabriela M. Gomes[5]*

[1] *Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom.*

[2] *Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, United Kingdom.*

[3] *Unidade Multidisciplinar de Investigação Biomédica, Instituto de Ciências Biomédicas Abel Salazar, Universidade do Porto, Porto, Portugal.*

[4] *Instituto de Ciências Biomédicas, Universidade de São Paulo, São Paulo, Brazil.*

[5] *Department of Mathematics and Statistics, University of Strathclyde, Glasgow, United Kingdom.*

* Correspondence and requests for materials should be addressed to M.G.M.G. (email: gabriela.gomes@strath.ac.uk).

Variation in individual susceptibility or exposure to infection accelerates the rate

at which populations acquire immunity by natural infection. Individuals that are

more susceptible or more exposed tend to be infected earlier and hence

selectively removed from the susceptible pool, decelerating the incidence of new

infections. Eventually, susceptible numbers become low enough to prevent

epidemic growth or, in other words, the herd immunity threshold is reached.

Here we fit epidemiological models, with inbuilt distributions of susceptibility or

exposure, to SARS-CoV-2 epidemics in Spain and Portugal, to estimate basic

reproduction numbers (R_0) alongside coefficients of individual variation (CV)

and the effects of control measures. Herd immunity thresholds are then

calculated as $1 - (1/R_0)^{1/(1+CV^2)}$ or $1 - (1/R_0)^{1/(1+2CV^2)}$, depending on

whether variation is in susceptibility or exposure. Our inferences result in lower

herd immunity thresholds than what would be expected if population immunity

was to be induced by random infection or vaccination, $1 - 1/R_0$.

Introduction

Selection acting on unmeasured individual variation is a well-known source of bias in the analysis of populations. It has been shown to affect measured rates of mortality (Keyfitz and Littman; Vaupel et al 1979; Vaupel and Yashin 1985), the survival of endangered species (Kendall and Fox 2002; Jenouvrier et al 2018), the scope of neutral theories of biodiversity and molecular evolution (Steiner and Tuljapurkar 2012, Gomes et al 2019), the risk of diseases whether non-communicable (Aalen et al 2015; Stensrud and Valberg 2017) or infections (Anderson et al 1986; Dwyer et al 1997; Smith et al 2005; Bellan et al 2015; Gomes et al 2019; Corder et al 2020), and the efficacy of interventions such as vaccines (Halloran et al 1996; O´Hagan et al; Gomes et al 2014; Gomes et al 2016; Langwig et al 2017) or symbionts (Pessoa et al 2016; King et al 2018). Building on this knowledge, we addressed how selection on individual variation might affect the course of the coronavirus disease (COVID-19) pandemic (Gomes et al 2020).

COVID-19 is an infectious respiratory disease caused by a virus (severe acute respiratory syndrome coronavirus 2 [SARS-CoV-2]), which was first identified in China in late 2019 and has since spread worldwide leading to considerable human suffering and social disruption. European and American continents have been the most affected, with 0.16% and 0.20% of the respective total populations having died as of the 15 July 2021 (WHO 2021). Here we analyse series of daily deaths attributed to COVID-19 in Spain and Portugal (Iberian Peninsula) to study how individual variation in susceptibility and exposure to a respiratory virus affects its epidemic trajectory. Besides adding to the compendium of neglected effects of selection in population dynamics we hope to stimulate a new approach to study epidemic

dynamics.

The main idea developed here is that individual variation in susceptibility or exposure

(connectivity) accelerates the acquisition of immunity in populations. More

susceptible and more connected individuals have a higher propensity to be infected

and thus are likely to become immune earlier. Due to this selective immunization by

natural infection, heterogeneous populations acquire herd immunity by natural

infection more efficiently than suggested by models that do not fully account for these

types of variation. We integrate continuous distributions of susceptibility or

connectivity in otherwise basic epidemiological models for COVID-19, which

necessarily account for non-pharmaceutical intervention effects, and generate three

types of results. First, at national levels the herd immunity threshold by natural

infection declines from around 70% to 20-30%. This is newly reported here for Spain

and in agreement with recent estimates for England and Scotland (Gomes et al 2021).

Second, these inferences can be made relatively early in the pandemic, such as

between first and second waves, provided the first wave is sufficiently large and

spatially synchronous. Third, we include a selection of results for Portugal to illustrate

how the inferential procedure degenerates when national data do not meet certain

conditions.

Individual variation in SARS-CoV-2 transmission

SARS-CoV-2 is transmitted primarily by respiratory droplets and modelled as a

susceptible-exposed-infectious-recovered (SEIR) process.

Variation in susceptibility to infection

Individual variation in susceptibility is integrated as a continuously distributed factor

that multiplies the force of infection upon individuals (Diekmann et al 1990) in the

form of an infinite system of ordinary differential equations (ODEs):

$$\dot{S}(x) = -\lambda x S(x), \tag{1}$$

$$\dot{E}(x) = \lambda x [S(x) + \sigma R(x)] - \delta E(x), \tag{2}$$

$$\dot{I}(x) = \delta E(x) - \gamma I(x), \tag{3}$$

$$\dot{R}(x) = (1 - \phi)\gamma I(x) - \sigma \lambda x R(x), \tag{4}$$

where $S(x)$ is the density of individuals with susceptibility x, $E(x)$ and $I(x)$ are the

densities of individuals who originally had susceptibility x and became exposed and

infectious, while $R(x)$ represents those who have recovered and have their

susceptibility reduced to a reinfection factor σ due to acquired immunity. Parameter δ

is the rate of progression from exposed to a period of maximal infectiousness ($=$

$1/5.5$ per day [McAloon et al. 2020]), γ is the rate of recovery from maximal

infectiousness ($= 1/4$ per day [Nishiura et al. 2020; Lauer et al. 2020; Li et al.

2020]), ϕ is the proportion of individuals who die as a result of infection ($= 0.009$

[Ward et al. 2021]), and:

$$\lambda = \frac{\beta}{N} \int \rho E(y) + I(y) \, dy \tag{5}$$

is the average force of infection upon susceptible individuals in a population of

approximately constant size N and transmission coefficient β. Standardizing so that

susceptibility distributions have mean $\int x g(x) \, dx = 1$, given a probability density

function $g(x)$, the basic reproduction number, defined as the expected number of

secondary infections generated by an infected individual in a population that has no

specific immunity to the virus (Diekmann et al 1990), is:

$$\mathcal{R}_0 = \beta \left(\frac{\rho}{\delta} + \frac{1}{\gamma} \right), \tag{6}$$

where ρ is a factor measuring the infectiousness of individuals in compartment E in

relation to those in I (= 0.5 [Wei et al. 2020; To et al. 2020; Arons et al. 2020; He et

al. 2020]). The coefficient of variation in individual susceptibility $CV =$

$\sqrt{\int (x-1)^2 g(x)\, dx}$ is also treated as a parameter.

The basic reproduction number $\mathcal{R}_0$ is a theoretical framework. It cannot be exactly

measured directly but it can be approximated from the initial growth in case numbers.

However, as the virus spreads through the population, infected and immune

individuals accumulate, reducing the availability of susceptible hosts. As a result,

growth in case numbers deviates from being a direct indication of $\mathcal{R}_0$ but rather of a

so-called effective reproduction number $\mathcal{R}_{eff}$.

When susceptibility is given by a gamma distribution and acquired immunity is

totally protective, the effective reproduction number is:

$$\mathcal{R}_{eff}(t) = \mathcal{R}_0 \left(\frac{S(t)}{N} \right)^{1+CV^2}, \tag{7}$$

where $S(t) = \int S(x, t)\, dx$ is the total number of susceptible individuals at time t, and

(Equations 1-4) reduce to a finite system of ODEs:

$$\dot{S} = -\beta(\rho E + I)\left(\frac{S}{N} \right)^{1+CV^2}, \tag{8}$$

$$\dot{E} = \beta(\rho E + I)\left(\frac{S}{N} \right)^{1+CV^2} - \delta E, \tag{9}$$

$$\dot{I} = \delta E - \gamma I, \tag{10}$$

$$\dot{R} = (1 - \phi)\gamma I, \tag{11}$$

where S, E, I and R are the total numbers of susceptible, exposed, infectious and

recovered individuals, respectively (Novozhilov 2008; Montalbán et al. 2020).

Variation in connectivity

In a directly transmitted infectious disease, such as COVID-19, variation in exposure

to infection is primarily governed by patterns of connectivity among individuals. Here

we incorporate this in the system (Equations 1-4) under the assumption that

individuals mix at random (Pastor-Satorras and Vespignani 2001; Miller et al. 2012),

while in Supplementary Information we conduct some sensitivity analyses to this

assumption. Under random mixing and heterogeneous connectivity, the force of

infection is written as:

131
$$\lambda = \frac{\beta}{N} \frac{\int y[\rho E(y) + I(y)]\, dy}{\int y g(y)\, dy}, \tag{12}$$

132   and the basic reproduction number is:

133
$$\mathcal{R}_0 = (1 + CV^2)\beta \left(\frac{\rho}{\delta} + \frac{1}{\gamma}\right). \tag{13}$$

134   In this setup, when connectivity is given by a gamma distribution and acquired

135   immunity is totally protective, the effective reproduction number is:

136
$$\mathcal{R}_{eff}(t) = \mathcal{R}_0 \left(\frac{S(t)}{N}\right)^{1+2CV^2}, \tag{14}$$

137   and (Equations 1-4) reduce to the finite system of ODEs:

138
$$\dot{S} = -(1 + CV^2)\beta(\rho E + I)\left(\frac{S}{N}\right)^{1+2CV^2}, \tag{15}$$

$$\dot{E} = (1 + CV^2)\beta(\rho E + I)\left(\frac{S}{N}\right)^{1+2CV^2} - \delta E,$$ (16)

$$\dot{I} = \delta E - \gamma I,$$ (17)

$$\dot{R} = (1 - \phi)\gamma I,$$ (18)

where S, E, I and R are the total numbers of susceptible, exposed, infectious and

recovered individuals, respectively (Novozhilov 2008; Montalbán et al. 2020).

We have used the reduced systems (Equations 8-11) and (Equations 15-18) in

combination with non-pharmaceutical interventions (NPIs) to analyse series of daily

deaths in the study countries.

Non-pharmaceutical interventions and other transmissibility modifiers

NPIs designed to control transmission typically reduce β and hence $\mathcal{R}_0$. Denoting the

time-dependent reproduction number when control measures are in place by $\mathcal{R}_c(t)$,

the modified effective reproduction number is obtained by replacing $\mathcal{R}_0$ with $\mathcal{R}_c(t)$ in

(Equation 7) and (Equation 14) as appropriate. For the estimation of $\mathcal{R}_c(t)$ we

introduce flexible transmissibility profiles $c(t)$ as illustrated in Figure 1.

Figure 1: Transmissibility profile. Schematic illustration of factor $c(t)$ representing
the combined effects of NPIs, seasonality and viral evolution on the reproduction
number $\mathcal{R}_c(t)$. T_0 is the time when $\mathcal{R}_0$ begins to decrease due to behavioural change
or seasonality; T_1 is the time interval over which transmission reduces gradually until
the first lockdown; $c_1 \leq 1$ is the average $c(t)$ achieved during the first lockdown; L_1,
L_2 and L_3, denote the length in days of the successive periods of strictest NPI
measures (L_1 being the first lockdown). These profiles are adopted in fits until: 1 July
2020 (top); 1 March 2021 (bottom).

One-wave transmissibility profile

When the model is applied to the first pandemic wave only (until 1 July 2020) we use

the top profile in Figure 1. T_0 is the time when $\mathcal{R}_0$ begins to decrease due to

behavioural change or seasonality; T_1 is the time interval over which transmission

reduces gradually (assumed linearly, until the first lockdown; $c_1 \leq 1$ is the average

$c(t)$ from the beginning of the first lockdown onwards (14 March in Spain, 19 March

in Portugal). Mathematically this is constructed as:

169
$$
c(t) = \begin{cases} 1, & \text{if } 0 < t \leq T_0, \\ 1 - (1 - c_1) \cdot \dfrac{(1 - T_0)}{T_1}, & \text{if } T_0 < t \leq T_0 + T_1, \\ c_1, & \text{otherwise.} \end{cases}
\tag{19}
$$

Two-wave transmissibility profile

Applying the model over longer periods which capture multiple waves and multiple

lockdowns requires additional features on the transmissibility profile. Denoting by L_1

the duration of the first lockdown (29 days in Spain, 44 days in Portugal), we allow

restrictions to be progressively relaxed at the end of this period by letting transmission

begin a linear increase such that $c(t)$ reaches 1 in T_2 days, which may or may not be

within the range of the study. Changes in other factors that affect transmission (such

as seasonality or viral evolution) are inseparable from contact changes in this

framework and are also accounted for by $c(t)$. Mathematically this is constructed as:

$$179 \quad c_0(t) = \begin{cases} 1, & \text{if } 0 < t \leq T_0, \\ 1 - (1 - c_1) \cdot \dfrac{(1 - T_0)}{T_1}, & \text{if } T_0 < t \leq T_0 + T_1, \\ c_1, & \text{if } T_0 + T_1 < t \leq T_0 + T_1 + L_1, \\ 1 - (1 - c_1) \cdot \dfrac{(T_0 + T_1 + L_1 + T_2 - t)}{T_1}, & \text{otherwise.} \end{cases} \quad (20)$$

180   Second and third lockdowns in the autumn and winter season are implemented as a

181   further reduction in transmission (by factors c_2 and c_3, respectively) over the

182   stipulated time periods (L_2 and L_3 in the bottom panel of Figure 1):

$$183 \quad c_{\text{Spain}}(t) = \begin{cases} c_2 \cdot c_0(t), & \text{if } t \in [26 \text{ October } 2020, 22 \text{ December } 2020], \\ c_3 \cdot c_0(t), & \text{if } t \in [7 \text{ January } 2021, \text{ last data point}], \\ c_0(t), & \text{otherwise.} \end{cases} \quad (21)$$

$$184 \quad c_{\text{Portugal}}(t) = \begin{cases} c_2 \cdot c_0(t), & \text{if } t \in [9 \text{ November } 2020, 21 \text{ December } 2020], \\ c_3 \cdot c_0(t), & \text{if } t \in [14 \text{ January } 2021, \text{ last data point}], \\ c_0(t), & \text{otherwise.} \end{cases} \quad (22)$$

In Spain, second and third lockdowns were effectively a single intervention mildly interrupted

by a short relaxation over Christmas, and hence we assume $c_2 = c_3$ in this country. This is

contrasted by Portugal where the third lockdown was much stricter than the second.

Herd immunity thresholds

Individual variation in risk of acquiring infection is under selection by the force of

infection, whether individual differences are due to biological susceptibility,

exposure, or both. The most susceptible or exposed individuals are selectively

removed from the susceptible pool as they become infected and eventually recover

with immunity (some die), resulting in decelerated epidemic growth and accelerated

acquisition of immunity in the population. The herd immunity threshold (HIT) defines

the percentage of the population that needs to be immune to reverse epidemic growth

and prevent future waves. In the absence of NPIs or other transmissibility modifiers,

if individual susceptibility or connectivity is gamma-distributed and mixing is

random, basic HIT curves ($\mathcal{H}_0$) can be derived analytically (Montalbán et al 2020)

from the model systems (Equations 1-4, with the respective forces of infections). In

the case of variation in susceptibility to infection we obtain

$$201 \quad \mathcal{H}_0 = 1 - \left[\frac{1 - \sigma R_0}{(1 - \sigma)R_0}\right]^{\frac{1}{1 + CV^2}}, \tag{23}$$

202   while variable connectivity results in

$$203 \quad \mathcal{H}_0 = 1 - \left[\frac{1 - \sigma R_0}{(1 - \sigma)R_0}\right]^{\frac{1}{1 + 2CV^2}}. \tag{24}$$

In less straightforward cases, such as when the characteristics follow a distribution

other than gamma (Gomes et al. 2020), when mixing is not random (Supplementary

Information), when both distributions in susceptibility and connectivity are

considered, or when contact networks are rewired as result of NPIs or otherwise, $\mathcal{H}_0$

can be obtained numerically.

Both (Equation 23) and (Equation 24) convey substantial declines in HIT as

individual variation increases (Figure 2, Gomes et al. 2020), most strikingly over

relatively low CV (from 0 up to 1 or 2). For concreteness, when $\mathcal{R}_0 = 3$, $\mathcal{H}_0 = 67\%$

for $CV = 0$, while $CV = 1$ brings $\mathcal{H}_0$ down to 42% for heterogeneous susceptibility

and 30% for heterogeneous connectivity, and $CV = 2$ brings $\mathcal{H}_0$ further down to 20%

and 11%, respectively. Accounting for reinfection (i.e. $\sigma > 0$ in the formulas) might

moderate these reductions. In any case, such differences in HIT are an indication of

strong sensitivity of the epidemic dynamics to the parameter CV. Since early in the

COVID-19 pandemic we and others (Gomes et al. 2020, Britton et al 2020) have

advocated for attention to this issue. Given that the steeper decline in $\mathcal{H}_0$ appears to

occur over a range of realistic CVs (Gomes et al. 2020, Gomes et al 2021) it is

plausible that some aspects of the pandemic dynamics need to be reinterpreted in a

broader framework.

Figure 2: Herd immunity threshold and epidemic final size. Herd immunity
thresholds (solid curves) are calculated according to (Equation 23) for heterogeneous
susceptibility and (Equation 24) for heterogeneous connectivity. Final sizes of the
corresponding unmitigated epidemics are also shown (dashed). Curves are generated
for different values of the efficacy of immunity conferred by natural infection ($1 - \sigma$)
as displayed in the legend.

We emphasise, however, that $\mathcal{H}_0$ is a theoretical framework almost to the extent that

$\mathcal{R}_0$ is a theoretical framework. It cannot be measured directly when epidemic

trajectories are affected by interventions, but it can be inferred indirectly from

epidemiological data. By construction, $\mathcal{H}_0$ changes if the parameters that determine

its value change. Most notably, natural changes in $\mathcal{R}_0$ through time, which can

happen due to seasonal forces or viral evolution, transfer to $\mathcal{H}_0$ according to

(Equation 23), (Equation 24), or even their homogeneity equivalent $1 - 1/\mathcal{R}_0$. As a

result, the percentage of the population immune required to prevent sustained

epidemic growth may deviate from the initial $\mathcal{H}_0$. Notwithstanding, a model with

lower $\mathcal{H}_0$ outputs smaller epidemics than a model with higher $\mathcal{H}_0$, all non-basic

processes being the same.

The phenomenon of variation and selection which accounts for lower HIT was widely

explained around mid-2020 (Hartnett 2020) and generated broad public interest in the

context of COVID-19. By early 2021, vaccines had become available, and the

narrative changed to imply that the HIT might be unachievable for COVID-19

(Aschwanden 2021). This apparent contradiction stems from the adoption of less

fundamental meaning of HIT by the latter. While Hartnett (2020) writes about using

the basic $\mathcal{H}_0$ to assess pandemic potential, stressing how that is weakened by

variation and selection by natural infection, Aschwanden (2021) endorses a more

common sensical view that reinfection may become frequent enough to make herd

immunity unachievable. In the light of the theory presented here – specifically

(Equation 23) and (Equation 24) – these views are transversal and not contradictory.

Data

We fit compartmental models to publicly available epidemiological data from the

coronavirus dashboards for Spain [https://cnecovid.isciii.es/covid19] and Portugal

[https://covid19.min-saude.pt/ponto-de-situacao-atual-em-portugal] to fit the models

and estimate parameters of interest. Namely, we fit model reconstructed mortality

timeseries assuming a fixed infection fatality ratio to datasets containing daily deaths,

$\{(k, \tilde{y}_k)\}_{k=1}^n$, where $k = 1$ is the day when the cumulative moving average of death

numbers exceed $5 \cdot 10^{-7}$ of the population (7 March in Spain, 19 March 2020 in

Portugal).

Model fits were carried out to the raw series of daily deaths until the 1 July 2020 in

the first instance (to cover the first wave of the epidemic in the study countries), and

until the 1 March 2021 in an extended analysis (as a compromise between having a

series sufficiently long to capture much of the second wave and not so long that it

would be affected by vaccination and require the vaccine to be modelled). We defined

the initial conditions as:

$R(0) = 0,$ (25)

$I(0) = \dfrac{\tilde{y}_{1+\eta}}{1 - \exp{(-\phi\gamma)}},$ (26)

$E(0) = \dfrac{I(0)}{1 - \exp{(-\gamma)}},$ (27)

$S(0) = N - E(0) - I(0) - R(0),$ (28)

and the population sizes N were obtained from the most recent respective census

(approx. 46.94 million in Spain, 10.28 million in Portugal, 3.57 million in the North

Region of Portugal, 3.66 million in Lisbon and Tagus Valley Region of Portugal).

Model fitting and parameter estimating

We assumed that reinfection was negligible throughout the study period. A study

conducted in England (Hall et al. 2021), between June 2020 and January 2021,

concluded that previous SARS-CoV-2 infection induced 84% effective immunity to

future infections. In (Gomes et al. 2121) we fit models to daily COVID-19 deaths in

England and Scotland, assuming no reinfection (i.e. 100% effective immunity) or

90% effective immunity, and found this alteration to have no significant effect on

projected model trajectories. Basically, the fit readjusts the parameters when

reinfection is added to the model in such a way that the output solutions remain

similar.

Parameter estimation was performed with the software MATLAB using the PESTO

(Parameter EStimation Toolbox) package (Stapor et al. 2018). We assumed the

number of SARS-CoV-2 infections to be Poisson distributed.

We approximate the dynamics of COVID-19 deaths by estimating the set of

parameters θ that maximises the log-likelihood (LL) of observing the daily numbers

of reported deaths Y:

$$LL(\theta|Y) = -\sum_{k=1}^{n} y(k,\theta) + \sum_{k=1}^{n} \tilde{y}(k)\ln\big(y(k,\theta)\big) - \sum_{k=1}^{n} \ln(\tilde{y}(k)!), \qquad (29)$$

where $y(k,\theta)$ are the simulated model output numbers of COVID-19 deaths at day k

for the set of parameters θ, $Y = \{(k,\tilde{y}_k)\}_{k=1}^{n}$ are the numbers of daily reported

deaths, and n is the total number of days included in the analysis.

Fitting models to one pandemic wave

The models exploring heterogeneity in susceptibility (Equations 8-11) and

connectivity (Equations 15-18) both with transmissibility profile as in (Equation 19)

were fit to COVID-19 deaths recorded daily until 1 July 2020, in Spain and Portugal.

A homogeneous version obtained by setting $CV = 0$ in either model was also fitted.

Results for Spain are shown in Table 1 and Figure 3, and for Portugal in Table 2 and

Figure 4.

We estimate the basic reproduction number $\mathcal{R}_0$ with 95% credible intervals (CI)

around $3.5 - 3.8$ in Spain and $2.3 - 3.4$ in Portugal. For the minimal transmissibility

factor c_1 we estimate $0.21 - 0.27$ when individual variation is allowed and $0.18 -$

0.19 with the homogeneity constraint in Spain, while in Portugal we estimate the

wider intervals $0.26 - 0.39$. For coefficients of variation in Spain, we estimate CV in

the range $1.3 - 2.4$ under heterogeneous susceptibility and $1.0 - 1.7$ under

heterogeneous connectivity. In Portugal, we obtain the much wider and uninformative

ranges $0.0 - 3.2$.

Left plots in Figures 3 and 4 show the best fitting model solutions generated from the

median posterior estimates of each parameter in the respective countries as well as the

95% CI generated from 100,000 posterior samples. The basic herd immunity

thresholds $\mathcal{H}_0$, calculated from $\mathcal{R}_0$ and CV estimates, are $\mathcal{H}_0 = 19\%$ (95% CI, 13-

32%) under heterogeneous susceptibility, $\mathcal{H}_0 = 19\%$ (95% CI, 13-36%) under

heterogeneous connectivity, and $\mathcal{H}_0 = 72\%$ (95% CI, 71-73%) when homogeneity is

imposed, in Spain. In Portugal, we obtain $\mathcal{H}_0 = 19\%$ (95% CI, 5-69%) under

heterogeneous susceptibility, $\mathcal{H}_0 = 13\%$ (95% CI, 3-69%) under heterogeneous

connectivity, and $\mathcal{H}_0 = 67\%$ (95% CI, 66-69%) when homogeneity is imposed.

Credible intervals for $\mathcal{H}_0$ in Portugal are wide and uninformative when individual

variation is allowed which is expected given the wide ranges obtained for CV. NPIs in

Portugal were initiated very early in the epidemic which resulted in transmissibility

reductions blending with $\mathcal{R}_0$, making parameter identification a major challenge from

the data accessible to us.

In Spain, where the three models provide good fits to the data, model selection criteria

such as AIC (Akaike information criterion) support the heterogeneous

implementations. To better distinguish the various models, we run the respective

systems of equations forward, under a set of conventions, and compare the respective

projected outcomes. Right plots in Figures 3 were generated by taking the end

conditions of the left plots, moving all exposed and infectious individuals to

recovered (except a residual proportion to seed a new outbreak) and running each

model with the estimated $\mathcal{R}_0$ and CV until the susceptible pool has been effectively

depleted in all implementations. In this manner we can visualise how much more

infection appears to be ahead when models are constrained to be homogeneous (a

manifestation of their relatively higher $\mathcal{H}_0$. Roughly, epidemics peak one order of

magnitude higher when models are homogeneous. This must have broad implications,

which we believe remain largely unappreciated, for how a population will experience

an epidemic, irrespective of how this basic scenario is adapted to specific factors such

behavioural patterns, seasonality, viral evolution, or vaccination.

The same analysis applied to Portugal, selects in favour of the homogeneity

assumption and larger projected waves. We recall, however, the great uncertainty

associated with these specific results. We investigate this further through fittings to

longer data series, in the first instance.

Table 1: Model parameters for Spain (one wave). Estimated by Bayesian inference
based on daily deaths until 1 July 2020. Model selection based on maximum log-
likelihood (LL) and Akaike information criterion (AIC). Best fitting models have
lower AIC scores (best in red, second best in blue). Herd immunity threshold ($\mathcal{H}_0$)
derived from estimated $\mathcal{R}_0$ and CV.

	Heterogeneous susceptibility		Heterogeneous connectivity		Homogeneous	
	Median	95% CI	Median	95% CI	Median	95% CI
Estimated parameters						
T_0	5.44	[3.52, 6.07]	5.40	[3.45, 6.07]	5.48	[3.73, 6.04]
c_1	0.24	[0.21, 0.27]	0.24	[0.21, 0.28]	0.18	[0.18, 0.19]
η	16^*	[15.53, 16.47]	16^*	[15.52, 16.48]	16^*	[15.52, 16.47]
$\mathcal{R}_0$	3.59	[3.51, 3.79]	3.59	[3.51, 3.80]	3.53	[3.47, 3.71]

CV	1.98	[1.33, 2.44]	1.41	[0.97, 1.74]	0	-
Derived from estimated parameters						
$\mathcal{H}_0$	23%	[17%, 38%]	23%	[16%, 37%]	72%	[71%, 73%]
Model selection						
LL	-503.78		-503.50		-518.33	
AIC	1017.60		1017.00		1044.70	

346 * Rounded median resulting in integer number of days.

Table 2: Model parameters for Portugal (one wave). Estimated by Bayesian inference based on daily deaths until 1 July 2020. Model selection based on maximum log-likelihood (LL) and Akaike information criterion (AIC). Best fitting models have lower AIC scores (best in red, second best in blue). Herd immunity threshold ($\mathcal{H}_0$) derived from estimated $\mathcal{R}_0$ and CV.

	Heterogeneous susceptibility		Heterogeneous connectivity		Homogeneous	
	Median	95% CI	Median	95% CI	Median	95% CI
Estimated parameters						
T_0	0.05	[0.00, 0.10]	0.05	[0.00, 0.10]	0.05	[0.00, 0.10]
c_1	0.31	[0.26, 0.36]	0.32	[0.26, 0.39]	0.31	[0.26, 0.35]
η	20*	[14.75, 25.72]	20*	[13.27, 25.94]	21*	[15.56, 26.04]
$\mathcal{R}_0$	2.62	[2.29, 3.21]	2.66	[2.27, 3.39]	2.61	[2.28, 3.15]
CV	0.31	[0.01, 3.24]	0.38	[0.01, 3.04]	0	-
Derived from estimated parameters						
$\mathcal{H}_0$	58%	[7%, 69%]	53%	[4%, 70%]	62%	[56%, 68%]
Model selection						
LL	-287.10		-287.14		-287.34	
AIC	584.20		584.28		582.67	

352 * Rounded median resulting in integer number of days.

Figure 3: Estimating SARS-CoV-2 transmission in Spain by fitting one wave of COVID-19 deaths. Variation in susceptibility (top panels); variation in connectivity (middle panels); and homogeneous model (bottom panels). Susceptibility or connectivity factors implemented as gamma distributions. Controlled (R_c) and effective (R_{eff}) reproduction numbers are displayed on shallow panels underneath the main plots. Basic reproduction number, coefficients of variation and transmissibility

profile parameters estimated by Bayesian inference as described in Methods
(estimates in Table 1). Curves represent model reconstructions from the median
posterior parameter estimates. Shades represent 95% credible intervals from 100,000
posterior samples.

Figure 4: Estimating SARS-CoV-2 transmission in Portugal by fitting one wave of
COVID-19 deaths. Variation in susceptibility (top panels); variation in connectivity
(middle panels); and homogeneous model (bottom panels). Susceptibility or
connectivity factors implemented as gamma distributions. Controlled (R_c) and
effective (R_{eff}) reproduction numbers are displayed on shallow panels underneath the
main plots. Basic reproduction number, coefficients of variation and transmissibility
profile parameters estimated by Bayesian inference as described in Methods
(estimates in Table 2). Curves represent model reconstructions from the median
posterior parameter estimates. Shades represent 95% credible intervals from 100,000
posterior samples.

Fitting models to two serial pandemic waves

Here we take the models with heterogeneity in susceptibility (Equations 8-11) and

connectivity (Equations 15-18) and apply the transmissibility profile in (Equations

20-22) to both before fitting the model outputs to COVID-19 deaths recorded daily

until 1 March 2021, in Spain and Portugal. As before a homogeneous version

obtained by setting $CV = 0$ was also fitted. Results for Spain are shown in Table 3

and Figure 5, and for Portugal in Table 4 and Figure 6.

We estimate $\mathcal{R}_0$ with 95% CI around $3.9 - 4.4$ in Spain and $2.7 - 4.6$ in Portugal.

For the minimal transmissibility factor c_1 we estimate $0.20 - 0.21$ when individual

variation is allowed and 0.17 with the homogeneity constraint in Spain, while in

Portugal we obtain $0.23 - 032$ with individual variation and $0.16 - 018$ without. For

coefficients of variation in Spain, we estimate CV around 2.0 under heterogeneous

susceptibility and 1.3 under heterogeneous connectivity. In Portugal, the estimates are

around 0.5.

Left plots in Figures 5 and 6 show best fitting model solutions as well as the

respective 95% CI. In Spain, basic herd immunity thresholds calculated from best-

fitting $\mathcal{R}_0$ and CV are $\mathcal{H}_0 = 25\%$ under heterogeneous susceptibility, $\mathcal{H}_0 = 27\%$

under heterogeneous connectivity, and $\mathcal{H}_0 = 76\%$ when homogeneity is imposed. In

Portugal, we obtain again wider and higher ranges: $\mathcal{H}_0 = 56\%$ (95% CI, 52-57%)

under heterogeneous susceptibility, $\mathcal{H}_0 = 56\%$ (95% CI, 54-57%) under

heterogeneous connectivity, and $\mathcal{H}_0 = 77\%$ (95% CI, 76-78%) when homogeneity is

imposed.

Puzzling, as in the case of shorter data series, model selection supports the

homogeneous model for Portugal while still favouring the incorporation of individual

variation in Spain. Moreover, for each country, results are consistent whether we base

our estimates on one or two waves of the national epidemic. This consistency was

also verified in England and Scotland (Gomes et al 2021). This suggests that the

earlier initiation of NPI in Portugal may not in itself explain why results for Portugal

contrast with those for other countries studied.

Table 3: Model parameters for Spain (two waves). Estimated by Bayesian
inference based on daily deaths until 1 March 2021. Model selection based on
maximum log-likelihood (LL) and Akaike information criterion (AIC). Best fitting
models have lower AIC scores (best in red, second best in blue). Herd immunity
threshold ($\mathcal{H}_0$) derived from estimated $\mathcal{R}_0$ and CV.

	Heterogeneous susceptibility		Heterogeneous connectivity		Homogeneous	
	Median	95% CI	Median	95% CI	Median	95% CI
	Estimated parameters					
T_0	3.22	[3.18, 3.31]	6.06	[6.06, 6.07]	5.99	[5.98, 5.99]
T_2	448.42	[446.58, 452.53]	429.91	[427.04, 433.98]	783.88	[777.04, 784.12]
c_1	0.20	[0.20, 0.20]	0.21	[0.21, 0.21]	0.17	[0.17, 0.17]
c_2	0.71	[0.71, 0.71]	0.68	[0.68, 0.68]	0.50	[0.50, 0.50]
η	12*	[12.42, 12.50]	12*	[11.54, 12.45]	10*	[10.04, 10.07]
$\mathcal{R}_0$	4.36	[4.32, 4.37]	3.93	[3.92, 3.95]	4.11	[4.10, 4.11]
CV	2.02	[2.01, 2.02]	1.30	[1.28, 1.31]	0	-
	Derived from estimated parameters					
$\mathcal{H}_0$	25%	[25%, 25%]	27%	[27%, 27 %]	76%	[76%, 76%]
	Model selection					
LL	-3511.40		-3269.90		-5826.40	
AIC	7036.80		6553.90		11665.00	

* Rounded median resulting in integer number of days.

Table 4: Model parameters for Portugal (two waves). Estimated by Bayesian
inference based on daily deaths until 1 March 2021. Model selection based on
maximum log-likelihood (LL) and Akaike information criterion (AIC). Best fitting

models have lower AIC scores (best in red, second best in blue). Herd immunity
threshold ($\mathcal{H}_0$) derived from estimated $\mathcal{R}_0$ and CV.

	Heterogeneous susceptibility		Heterogeneous connectivity		Homogeneous	
	Median	95% CI	Median	95% CI	Median	95% CI
Estimated parameters						
T_0	0.03	[0.01, 0.05]	0.04	[0.00, 0.10]	0.05	[0.00, 0.10]
T_2	585.14	[573.66, 672.90]	644.26	[615.71, 695.38]	899.67	[872.69, 946.07]
c_1	0.27	[0.27, 0.32]	0.24	[0.23, 0.25]	0.17	[0.16, 0.18]
c_2	0.71	[0.71, 0.87]	0.71	[0.70, 0.77]	0.66	[0.64, 0.68]
c_3	0.34	[0.27, 0.48]	0.42	[0.39, 0.48]	0.35	[0.34, 0.36]
η	16^*	[15.52, 17.39]	14^*	[12.76, 15.34]	12^*	[11.54, 12.33]
$\mathcal{R}_0$	2.94	[2.66, 2.96]	3.25	[3.14, 3.32]	4.36	[4.20, 4.58]
CV	0.55	[0.54, 0.59]	0.47	[0.47, 0.49]	0	-
Derived from estimated parameters						
$\mathcal{H}_0$	56%	[52%, 57%]	56%	[54%, 57%]	77%	[76%, 78%]
Model selection						
LL	-1550.80		-1376.80		-1234.30	
AIC	3117.50		2769.50		2482.50	

415 * Rounded median resulting in integer number of days.

Figure 5: Estimating SARS-CoV-2 transmission in Spain by fitting two serial waves of COVID-19 deaths. Variation in susceptibility (top panels); variation in connectivity (middle panels); and homogeneous model (bottom panels). Susceptibility or connectivity factors implemented as gamma distributions. Controlled (R_c) and effective (R_{eff}) reproduction numbers are displayed on shallow panels underneath the main plots. Basic reproduction number, coefficients of variation and transmissibility

profile parameters estimated by Bayesian inference as described in Methods (estimates in Table 3). Curves represent model reconstructions from the median posterior parameter estimates. Shades represent 95% credible intervals from 100,000 posterior samples.

Figure 6: Estimating SARS-CoV-2 transmission in Portugal by fitting two serial
waves of COVID-19 deaths. Variation in susceptibility (top panels); variation in
connectivity (middle panels); and homogeneous model (bottom panels). Susceptibility
or connectivity factors implemented as gamma distributions. Controlled (R_c) and
effective (R_{eff}) reproduction numbers are displayed on shallow panels underneath the
main plots. Basic reproduction number, coefficients of variation and transmissibility
profile parameters estimated by Bayesian inference as described in Methods
(estimates in Table 4). Curves represent model reconstructions from the median
posterior parameter estimates. Shades represent 95% credible intervals from 100,000
posterior samples.

Fitting models to regional data in Portugal

Intrigued by the puzzling results for Portugal at country level, we gathered regional

data. We found that the epidemic was considerably different between the two largest

regions: North, home to roughly one third of the Portuguese population; and Lisbon

and Tagus Valley, home to another third. Asynchrony of epidemic dynamics between

regions of similar sizes may require disaggregated analyses. We then decided to fit

the mortality data for the two regions simultaneously, estimating common parameters

to describe country-wide lockdowns, and regions-specific parameters to describe the

basic transmission dynamics ($\mathcal{R}_0$ and CV). The results are provided in Tables 5, 6 and

Figures 7-9. According to these analyses, the best-fitting models include

heterogeneity.

Basic herd immunity thresholds inferred by fits to the longer series (until 1 March

2021) are around $25 - 34\%$ in the North (similar as Spain, England, and Scotland)

and $47 - 52\%$ in Lisbon and Tagus Valley. The higher $\mathcal{H}_0$ in the capital region

results from the estimation of a lower CV. This may be real and due to the more urban

character of Lisbon and Tagus Valley or, in contrast, it may be a spurious result of the

lack of a first wave in the region to inform the model. It would be interesting to

replicate the regional analysis in other countries to investigate to what extent more

urban regions have higher HIT.

Fits to the shorter series (until 1 July 2020) are again inconclusive. Not only the

uncertainty around parameter estimates is large but some estimates are implausible.

First, $\mathcal{R}_0$ around 2 or less is on the low end of consensus estimates (Flaxman et al.

2020; Keeling et al. 2020; Viana et al. 2021; Wood 2021). More strikingly, the

algorithm is incapable of estimating CV from these regional series, resulting in

convergence to the upper and lower limits of the prior distribution (uniform between

$0.01 - \sqrt{12}$) in the North and Lisbon regions, respectively.

In summary, the results reported in this section support two notions. First,

asynchronous dynamics may compromise parameter estimation based on model

fittings to aggregated data. This should depend on whether the asynchrony in question

is between similar-sized regions. Second, the early estimation of parameters for

heterogeneous models (especially CV) may require a larger first wave than that

required by models that are either homogeneous (Flaxman et al. 2020; Wood 2021) or

have their heterogeneity informed directly by specific data (Keeling et al. 2020). The

downside of these other approaches, however, is that heterogeneity is either absent or

possibly incomplete, resulting in reduced selection and biased estimates. Among the

studies we have completed, England, Scotland and Spain had sufficiently sized first

waves to inform the inference of CV while Portugal did not.

Table 5: Model parameters for the North and Lisbon regions of Portugal (one
wave). Estimated by Bayesian inference based on daily deaths until 1 July 2020.
Model selection based on maximum log-likelihood (LL) and Akaike information
criterion (AIC). Best fitting models have lower AIC scores (best in red, second best in
blue). Herd immunity threshold ($\mathcal{H}_0$) derived from estimated $\mathcal{R}_0$ and CV.

	Heterogeneous susceptibility		Heterogeneous connectivity		Homogeneous	
	Median	95% CI	Median	95% CI	Median	95% CI

	Median	95% CI	Median	95% CI	Median	95% CI
	Common parameters					
T_0	0.05	[0.00, 0.10]	0.05	[0.00, 0.10]	0.05	[0.00, 0.10]
c_1	0.48	[0.43, 0.52]	0.55	[0.47, 0.61]	0.43	[0.40, 0.47]
η	24[*]	[19.45, 28.99]	21[*]	[14.18, 27.37]	29[*]	[28.50, 28.50]
	North					
$\mathcal{R}_0$	2.17	[2.00, 2.45]	2.43	[2.17, 2.92]	1.91	[1.80, 2.09]
CV	3.44	[3.38, 3.46]	3.45	[3.39, 3.46]	0	-
$\mathcal{H}_0$	5.9%	[5.2%, 7.0%]	3.5%	[3.0%, 4.4%]	48%	[44%, 52%]
	Lisbon and Tagus Valley					
$\mathcal{R}_0$	1.78	[1.65, 2.00]	1.65	[1.49, 1.96]	1.85	[1.74, 2.02]
CV	0.07	[0.01, 0.56]	0.07	[0.01, 0.57]	0	-
$\mathcal{H}_0$	44%	[32%, 50%]	39%	[21%, 49%]	46%	[43%, 51%]
	Model selection					
LL	-520.56		-479.75		-578.50	
AIC	1055.10		973.49		1167.00	

[*] Rounded median resulting in integer number of days.

Table 6: Model parameters for the North and Lisbon regions of Portugal (two waves). Estimated by Bayesian inference based on daily deaths until 1 March 2021. Model selection based on maximum log-likelihood (LL) and Akaike information criterion (AIC). Best fitting models have lower AIC scores (best in red, second best in blue). Herd immunity threshold ($\mathcal{H}_0$) derived from estimated $\mathcal{R}_0$ and CV.

	Heterogeneous susceptibility		Heterogeneous connectivity		Homogeneous	
	Median	95% CI	Median	95% CI	Median	95% CI
	Common parameters					
T_0	0.06	[0.00, 0.10]	0.09	[0.06, 0.10]	0.05	[0.03, 0.05]
T_2	498.77	[489.85, 515.47]	499.41	[462.35, 547.99]	960.64	[960.28, 961.42]
c_1	0.27	[0.26, 0.29]	0.25	[0.25, 0.25]	0.17	[0.17, 0.17]
c_2	0.75	[0.73, 0.82]	0.71	[0.69, 0.75]	0.73	[0.73, 0.73]
c_3	0.44	[0.42, 0.45]	0.49	[0.45, 0.50]	0.36	[0.36, 0.36]
η	16[*]	[15.36, 16.46]	15[*]	[13.75, 15.16]	12[*]	[11.62, 11.64]
	North					
$\mathcal{R}_0$	2.99	[2.88, 3.03]	3.12	[3.05, 3.18]	4.34	[4.34, 4.35]
CV	1.40	[1.39, 1.41]	0.99	[0.93, 1.20]	0	-
$\mathcal{H}_0$	31%	[30%, 31%)	32%	[25%, 34%]	77%	[77%, 77%]
	Lisbon and Tagus Valley					
$\mathcal{R}_0$	2.81	[2.69, 2.86]	2.95	[2.87, 3.01]	4.39	[4.38, 4.40]
CV	0.67	[0.65, 0.70]	0.54	[0.53, 0.57]	0	-
$\mathcal{H}_0$	51%	[49%, 52%]	49%	[47%, 51%]	77%	[77%, 77%]
	Model selection					
LL	-2313.50		-2471.00		-2764.30	
AIC	4647.10		4961.90		5544.60	

[*] Rounded median resulting in integer number of days.

Figure 7: Estimating SARS-CoV-2 transmission in the two larger regions of Portugal. Heterogeneous susceptibility implemented as a gamma distribution. Controlled (R_c) and effective (R_{eff}) reproduction numbers are displayed on shallow panels underneath the main plots. Basic reproduction number, coefficients of variation and transmissibility profile parameters estimated by Bayesian inference as described in Methods (estimates in Table 5). Curves represent model reconstructions

Figure 8: Estimating SARS-CoV-2 transmission in the two larger regions of
Portugal. Heterogeneous connectivity implemented as a gamma distribution.
Controlled (R_c) and effective (R_{eff}) reproduction numbers are displayed on shallow
panels underneath the main plots. Basic reproduction number, coefficients of
variation and transmissibility profile parameters estimated by Bayesian inference as
described in Methods (estimates in Table 5). Curves represent model reconstructions

Figure 9: Estimating SARS-CoV-2 transmission in the two larger regions of Portugal. Homogeneous model. Controlled (R_c) and effective (R_{eff}) reproduction numbers are displayed on shallow panels underneath the main plots. Basic reproduction number, coefficients of variation and transmissibility profile parameters estimated by Bayesian inference as described in Methods (estimates in Table 5). Curves represent model reconstructions from the median posterior parameter estimates. Shades represent 95% credible intervals from 100,000 posterior samples.

Discussion

We fit SEIR models, with inbuilt distributions of individual susceptibility or exposure to infection, to daily series of COVID-19 deaths in Spain and Portugal. In similarity with other studies (Flaxman et al. 2020, Keeling et al. 2020, Bertuzzo et al. 2021, López & Rodó 2021, Viana et al. 2021), we estimated relevant transmission parameters, such as the basic reproduction number $\mathcal{R}_0$, and a time dependent

transmissibility profile $c(t)$ which multiplies $\mathcal{R}_0$ to account for effects of NPIs,

seasonality, viral evolution, or any unexplicit factor that modifies the ability of the

virus to infect new hosts. In contrast, our sets of estimated parameters include

coefficients of variation that characterise distributions of unmeasured individual

susceptibility or connectivity while more common practices are to impose measured

variation into the model and ignore the unmeasured.

The two approaches have different advantages and disadvantages, and which to adopt

depends on study objectives. Using directly measured variation to inform the model

allows phenomena to be represented on a micro scale as desired for designing targeted

interventions, for instance, but ignoring unmeasured variation may create biases on

the macro scale if relevant selection is occurring on omitted characteristics. This may

result in the model overpredicting the impact of uniform interventions, for example.

The approach of inferring the amount of variation that is under selection by fitting the

model on a macro scale is conceived to eliminate those biases, but not suitable for

informing targeted actions. In addition, the latter approach appears to require larger

infected numbers than the former. In a global crisis such as the current COVID-19

pandemic the two approaches can have a role, helping to overcome each other's

limitations.

The inference of selectable variation presented here is relatively new to infectious

disease modelling. Prior to attempting this in the context of the COVID-19 pandemic,

we and others have been accelerating progress in systems that were either

experimentally controlled (Dwyer et al. 1997; Ben-Ami et al. 2008; Zwart et al. 2011;

Pessoa et al. 2014; Langwig et al. 2017; King et al. 2018) or already endemic (Smith

et al. 2005; Bellan et al. 2015; Corder et al. 2020). Several aspects of the pandemic

made it more challenging. There was an urgency for early results from inherently

scarce data, a challenge amplified by non-pharmaceutical interventions designed to

suppress the epidemic. In countries where interventions began earlier, such as

Portugal, it has been impossible to reach conclusive results without refining data and

methods further. In Spain (this study) and England and Scotland (Gomes et al 2021),

on the other hand, interventions started later, and results were consistent internally

and with each other. Moreover, comparing our coefficients of variation for individual

connectivity to those measured directly by contact surveys, we found them to be less

than 40% higher. This suggests that there was more selectable variation than the

surveys had captured, as expected, but not enough to seem implausible.

In conclusion, we believe that the results presented here constitute a major step in the

establishment of a novel approach to improve the accuracy and predictability of

infectious disease modelling. It holds good promise for use in conjunction with other

tools to help devise future pandemic policies.

Acknowledgements

We thank Jan Hasenauer, Rodrigo Corder and Antonio Montalbán for technical

advice and contributions to related research. M.U.F. received funding from Conselho

Nacional de Desenvolvimento Científico e Tecnológio (CNPq), Brazil.

Author contributions

M.G.M.G. conceived the study. R.A. and M.G.M.G. performed the analyses. All

authors interpreted the data and wrote the paper.

Competing interests

The authors declare no competing interests.

References

1.   Keyfitz, N. & Littman, G. (1979) Mortality in a heterogeneous population. *Popul. Stud.*
33, 333-342.

2.   Vaupel, J., Manton, K. & Stallard, E. (1979) Impact of heterogeneity in individual frailty
on the dynamics of mortality. *Demography* **16**, 439-454.

3.   Vaupel, J., & Yashin, A. (1985) Heterogeneity ruses – some surprising effects of
selection on population dynamics. *Am. Stat.* **39**, 176-185.

4.   Kendall, B. E. & Fox, G. A. (2002) Variation among individuals and reduced
demographic stochasticity. *Conserv. Biol.* **16**, 109-116.

5.   Jenouvrier, S, Aubry, L. M., Barbraud, C, Weimerskirch, H & Caswell, H. (2018)
Interacting effects of unobserved heterogeneity and individual stochasticity in the life
history of the southern fulmar. *J. Anim. Ecol.* **87**, 212-222.

6.   Steiner, U. K. & Tuljapurkar, S. (2012) Neutral theory for life histories and individual
variability in fitness components. *Proc. Natl. Acad. Sci U. S. A.* **109**, 4684-4689.

7.   Gomes, M. G. M., King, J. G., Nunes, A., Colegrave, N. & Hoffmann, A. (2019) The
effects of individual nonheritable variation on fitness estimation and coexistence. *Ecol.*
Evol. **16**, 8995-9004.

8.   Aalen, O. O., Valberg, M., Grotmol, T. & Tretli, S. (2015) Understanding variation in
disease risk: the elusive concept of frailty. *Int. J. Epidemiol.* **4**, 1408-1421.

9.   Stensrud, M. J. & Valberg, M. (2017) Inequality in genetic cancer risk suggests bad
genes rather than bad luck. *Nat. Commun.* **8**, 1165.

10.   Anderson, R. M., Medley, G. F., May, R. M. & Johnson, A. M. (1986) A preliminary
study of the transmission dynamics of the human immunodeficiency virus (HIV), the
causative agent of AIDS. *IMA J. Math. Appl. Med. Biol.* **3**, 229-263.

11.   Dwyer, G., Elkinton, J. S. & Buonaccorsi, J. P. (1997) Host heterogeneity in
susceptibility and disease dynamics: Tests of a mathematical model. *Am. Nat.* **150**, 685-
707.

12. Smith, D. L., Dushoff, J., Snow, R. W. & Hay, S. I. (2005) The entomological

inoculation rate and *Plasmodium falciparum* infection in African children. *Nature* **438**,

492-495.

13. Bellan, S. E., Dushoff, J., Galvani, A. P. & Meyers, L. A. (2015) Reassessment of HIV-1

acute phase infectivity: accounting for heterogeneity and study design with simulated

cohorts. *PLOS Med.* **12**, e1001801.

14. Gomes, M. G. M., et al. (2019) Introducing risk inequality metrics in tuberculosis policy

development. *Nat. Commun.* **10**, 2480.

15. Corder, R. M., Ferreira, M. U. & Gomes, M. G. M. (2020) Modelling the epidemiology

of residual Plasmodium vivax in a heterogeneous host population: a case study in the

Amazon Basin. *PLOS Comput. Biol.* **16**, e1007377.

16. Halloran, M. E., Longini, I. M. Jr. & Struchiner, C. J. (1996) Estimability and

interpretability of vaccine efficacy using frailty mixing models. *Am. J. Epidemiol.* **144**,

83-97.

17. O'Hagan, J. J., Hernán, M. A., Walensky, R. P. & Lipsitch, M. (2012) Apparent

declining efficacy in randomized trials: Examples of the Thai RV144 HIV vaccine and

CAPRISA 004 microbicide trials. *AIDS* **26**, 123.

18. Gomes, M. G. M., et al. (2014) A missing dimension in measures of vaccination

impacts. *PLOS Pathog.* **10**, e1003849.

19. Gomes, M. G. M., Gordon, S. B. & Lalloo, D. G. (2016) Clinical trials: the mathematics

of falling vaccine efficacy with rising disease incidence. *Vaccine* **34**, 3007.

20. Langwig, K. E., et al. (2017) Vaccine effects on heterogeneity in susceptibility and

implications for population health management, *mBio* **8**, e00796-17.

21. Pessoa, D., et al. (2016) Unveiling time in dose-response models to infer host

susceptibility to pathogens. *PLOS Comput. Biol.* **10**, e1003773.

22. King, J. G., Souto-Maior, C., Sartori, L. M., Maciel-de-Freitas, R. & Gomes, M. G. M.

(2018) Variation in *Wolbachia* effects on *Aedes* mosquitoes as a determinant of

invasiveness and vectorial capacity. *Nat. Commun.* **9**, 1-8.

23. Gomes, M. G. M., et al. (2020) Individual variation in susceptibility or exposure to

SARS-CoV-2 lowers the herd immunity threshold. *medRvix*

10.1101/2020.04.27.20081893.

24. World Health Organization (2021) COVID-19 weekly epidemiological update:

https://www.who.int/publications/m/item/weekly-epidemiological-update-on-covid-19---

13-july-2021.

25. Gomes, M. G. M., Ferreira, M. U., Chikina, M., Pegden, W. & Aguas, R. (2021) Frailty

variation models for susceptibility and exposure to SARS-CoV-2. *medRvix*

10.1101/2021.05.25.21257766.

26. Diekmann, O., Heesterbeek, J. A. P. & Metz, J. A. J. (1990) On the definition and

computation of the basic reproduction ratio R_0 in models for infectious diseases in

heterogeneous populations. *J. Math. Biol.* **28**, 365-382.

27. McAloon, C., et al. (2020) Incubation period of COVID-19: a rapid systematic review

and meta-analysis of observational research. *BMJ Open* **10**, e039652.

28. Nishiura, H., Linton, N. M. & Akhmetzhanov, A. R. (2020) Serial interval of novel

coronavirus (COVID-19) infections. *Int. J. Infect. Dis.* **93**, 284-6.

29. Lauer, S. A., et al. (2020) The Incubation Period of Coronavirus Disease 2019 (COVID-

19) From Publicly Reported Confirmed Cases: Estimation and Application. *Ann. Intern.*

Med. **172**, 577-582.

30. Li, Q., et al. (2020) Early transmission dynamics in Wuhan, China, of novel coronavirus-

infected pneumonia. *N. Engl. J. Med.* **382**, 1199-1207.

31. Wei, W. E., et al. (2020) Presymptomatic Transmission of SARS-CoV-2 — Singapore,

January 23–March 16, 2020. *MMWR Morb. Mortal. Wkly. Rep.* **69**, 411-415.

32. To, K. K. W., et al. (2020) Temporal profiles of viral load in posterior oropharyngeal

saliva samples and serum antibody responses during infection by SARS-CoV-2: an

observational cohort study. *Lancet Infect. Dis.* **20**, 565-74.

33. Arons, M. M., et al. (2020) Presymptomatic SARS-CoV-2 Infections and Transmission

in a Skilled Nursing Facility. *N. Engl. J. Med.* **382**, 2081-2090.

34. He, X., et al. (2020) Temporal dynamics in viral shedding and transmissibility of

COVID-19. *Nat. Med.* **26**, 672-675.

35. Ward, H., et al. (2021) SARS-CoV-2 antibody prevalence in England following the first

peak of the pandemic. *Nat. Commun.* **12**, 905.

36. Novozhilov, A. S. (2008) On the spread of epidemics in a closed hoterogeneous

population. Math. Biosci. 215, 177-185.

37. Montalbán, A., Corder, R. M. & Gomes, M. G. M. (2020) Herd immunity under

individual variation and reinfection. *arXiv* 200800098v2.

38. Pastor-Satorras, R. & Vespignani, A. (2001) Epidemic dynamics and endemic states in

complex networks. *Phys. Rev. E* **63**, 066117.

39. Miller, J. C., Slim, A. C. & Volz, E. M. (2012) Edge-based compartmental modelling for

infectious disease spread. *J. R. Soc. Interface* **9**, 890-906.

40. Britton, T., Ball, F. & Trapman, P. (2020) A mathematical model reveals the influence

of population heterogeneity on herd immunity to SARS-CoV-2. *Science* **369**, 846-849.

41. Hartnett, K. (2020) The tricky math of herd immunity for COVID-19. *Quanta Magazine*.

https://www.quantamagazine.org/the-tricky-math-of-covid-19-herd-immunity-

20200630/

42. Aschwanden, C. (2021) Five reasons why COVID herd immunity is probably

impossible. *Nature* **591**, 520-522. https://www.nature.com/articles/d41586-021-00728-2

43. Hall, et al. (2021) SARS-CoV-2 infection rates of antibody-positive compared with

antibody-negative health-care workers in England: a large, multicentre, prospective

cohort study (SIREN). *Lancet* **397**, 1459-1469.

44. Flaxman, S., et al. (2020) Estimating the effects of non-pharmaceutical interventions on

COVID-19 in Europe. *Nature* **584**, 257-261.

45. Keeling, M. J., et al. (2020) Fitting to the UK COVID-19 outbreak, short-term forecasts

and estimating the reproductive number. *medRvix* 10.1101/2020.08.04.20163782.

46. Viana, J. et al. (2021) Controlling the pandemic during the SARS-CoV-2 vaccination

rollout. *Nat. Commun.* **12**, 3674.

47. Wood, S. N. (2021) Inferring UK COVID-19 fatal infection trajectories from daily

mortality data: Were infections already in decline before the UK lockdowns? *Biometrics*

10.1111/biom.13462

48. Bertuzzo, E. et al. (2020) The geography of COVID-19 spread in Italy and implications

for the relaxation of confinement measures. *Nat. Commun.* **11**, 4264.

49. López, L. & Rodó, X. (2021) A modified SEIR model to predict the COVID-19

outbreak in Spain and Italy: Simulating control scenarios and multi-scale epidemics.

Results Phys. **21**, 103746.