Herd immunity thresholds for SARS-CoV-2 estimated from unfolding epidemics

Ricardo Aguas[1†], Rodrigo M. Corder[2†], Jessica G. King[3], Guilherme Gonçalves[4],

Marcelo U. Ferreira[2], M. Gabriela M. Gomes[5,6*]

[1] *Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine,*

University of Oxford, Oxford, United Kingdom.

[2] *Instituto de Ciências Biomédicas, Universidade de São Paulo, São Paulo, Brazil.*

[3] *Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, United*

Kingdom.

[4] *Unidade Multidisciplinar de Investigação Biomédica, Instituto de Ciências*

Biomédicas Abel Salazar, Universidade do Porto, Porto, Portugal.

[5] *Department of Mathematics and Statistics, University of Strathclyde, Glasgow,*

United Kingdom.

[6] *Centro de Matemática da Universidade do Porto, Porto, Portugal.*

[†] These authors contributed equally to this work.

[*] Correspondence and requests for materials should be addressed to M.G.M.G. (email:

gabriela.gomes@strath.ac.uk).

As severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) spreads, the susceptible subpopulation declines causing the rate at which new infections occur to slow down. Variation in individual susceptibility or exposure to infection exacerbates this effect. Individuals that are more susceptible or more exposed tend to be infected and removed from the susceptible subpopulation earlier. This selective depletion of susceptibles intensifies the deceleration in incidence. Eventually, susceptible numbers become low enough to prevent epidemic growth or, in other words, the herd immunity threshold is reached. Here we fit epidemiological models with inbuilt distributions of susceptibility or exposure to SARS-CoV-2 outbreaks to estimate basic reproduction numbers (R_0) alongside coefficients of individual variation (CV) and the effects of containment strategies. Herd immunity thresholds are then calculated as $1 - (1/R_0)^{1/(1+CV^2)}$ or $1 - (1/R_0)^{1/(1+2CV^2)}$, depending on whether variation is on susceptibility or exposure. Our inferences result in herd immunity thresholds around 10-20%, considerably lower than the minimum coverage needed to interrupt transmission by random vaccination, which for R_0 higher than 2.5 is estimated above 60%. We emphasize that the classical formula, $1 - 1/R_0$, remains applicable to describe herd immunity thresholds for random vaccination, but not for immunity induced by infection which is naturally selective. These findings have profound consequences for the governance of the current pandemic given that some populations may be close to achieving herd immunity despite being under more or less strict social distancing measures.

Scientists throughout the world have engaged with governments, health agencies, and with each other, to address the ongoing pandemic of coronavirus disease (COVID-19). Mathematical models have been central to important decisions concerning

contact tracing, quarantine, and social distancing, to mitigate or suppress the initial

pandemic spread[1]. Successful suppression, however, may leave populations at risk to

resurgent waves due to insufficient acquisition of immunity. Models have thus also

addressed longer term SARS-CoV-2 transmission scenarios and the requirements for

continued adequate response[2]. This is especially timely as countries relax lockdown

measures that have been in place over recent months with varying levels of success in

tackling national outbreaks.

Here we demonstrate that individual variation in susceptibility or exposure

(connectivity) accelerates the acquisition of immunity in populations. More

susceptible and more connected individuals have a higher propensity to be infected

and thus are likely to become immune earlier. Due to this selective immunization by

natural infection, heterogeneous populations require less infections to cross their herd

immunity threshold (HIT) than suggested by models that do not fully account for

variation. We integrate continuous distributions of susceptibility or connectivity in

otherwise basic epidemic models for COVID-19 which account for realistic

intervention effects and show that as coefficients of variation (CV) increase from 0 to

5, HIT declines from over 60%[3,4] to less than 10%. We then fit these models to series

of daily new cases to estimate CV alongside basic reproduction numbers (R_0) and

derive the corresponding HITs.

Effects of individual variation on SARS-CoV-2 transmission

SARS-CoV-2 is transmitted primarily by respiratory droplets and modelled as a

susceptible-exposed-infectious-recovered (SEIR) process.

Variation in susceptibility to infection

Individual variation in susceptibility is integrated as a continuously distributed factor

that multiplies the force of infection upon individuals[5] as

$\dot{S}(x) = -\lambda(x)xS(x),$ (1)

$\dot{E}(x) = \lambda(x)x[S(x) + \sigma R(x)] - \delta E(x),$ (2)

$\dot{I}(x) = \delta E(x) - \gamma I(x),$ (3)

$\dot{R}(x) = (1 - \phi)\gamma I(x) - \sigma\lambda(x)xR(x),$ (4)

where $S(x)$ is the number of individuals with susceptibility x, $E(x)$ and $I(x)$ are the

numbers of individuals who originally had susceptibility x and became exposed and

infectious, while $R(x)$ counts those who have recovered and have their susceptibility

reduced to a reinfection factor σ due to acquired immunity. δ is the rate of

progression from exposed to infectious, γ is the rate of recovery or death, ϕ is the

proportion of individuals who die as a result of infection and $\lambda(x) =$

$(\beta/N) \int [\rho E(y) + I(y)] \, dy$ is the average force of infection upon susceptible

individuals in a population of size N and transmission coefficient β. Standardizing so

that susceptibility distributions have mean $\int xg(x) \, dx = 1$, given a probability

density function $g(x)$, the basic reproduction number is

$R_0 = \beta\left(\dfrac{\rho}{\delta} + \dfrac{1}{\gamma}\right),$ (5)

where ρ is a factor measuring the infectivity of individuals in compartment E in

relation to those in I. The coefficient of variation in individual susceptibility $CV =$

$\sqrt{\int (x - 1)^2 g(x) \, dx}$ is explored as a parameter. Non-pharmaceutical interventions

(NPIs) designed to control transmission typically reduce β and hence R_0. We denote

the resulting controlled reproduction number by R_c. The effective reproduction

number R_{eff} is another useful indicator obtained by multiplying R_c by the

susceptibility of the population, in this case written as $R_{eff}(t) =$

$R_c(t) \int xS(x,t)\, dx / N(t)$ to emphasize its time dependence.

Figure 1 depicts model trajectories fitted to suppressed epidemics (orange) in 4

European countries (Belgium, England, Portugal and Spain) assuming gamma

distributed susceptibility and no reinfection ($\sigma = 0$). We estimate: R_0 rounding 5

(Belgium), 2.9 (England), 4.3 (Portugal) and 4.1 (Spain); individual susceptibility CV

reaching 3.9 (Belgium), 1.9 (England), 4.3 (Portugal) and 3.2 (Spain); and overall

intervention efficacy at maximum (typically during lockdown) being 60% (Belgium),

48% (England), 69% (Portugal) and 63% (Spain). Another estimated parameter is the

day when NPIs begin to affect transmission, after which we assume a linear

intensification from baseline over 21 days, remaining at maximum intensity for 30

days and linearly lifting back to baseline over a period of 120 days (although we have

confirmed that the results do not change significantly if measures are lifted over

slightly longer time frames, such as 150 or 180 days). Denoting by $d(t)$ the

proportional reduction in average risk of infection due to interventions, in this case we

obtain $R_c(t) = [1 - d(t)]R_0$ which is depicted for each country, alongside $R_{eff}(t)$,

underneath the respective epidemic trajectories. Overlaid on the R_c plots are mobility

data from Google[6], showing excellent agreement with our independently chosen

framework and estimate for the time R_{eff} starts declining. To assess the potential for

case numbers to overshoot if NPIs had not been applied, we rerun the model with

$d(t) = 0$ and obtain the unmitigated epidemics (black). Further details and sensitivity

analyses are described in Methods.

Variation in connectivity

In a directly transmitted infectious disease, such as COVID-19, variation in exposure

to infection is primarily governed by patterns of connectivity among individuals. We

incorporate this in the system (Equations 1-4) assuming that individuals mix at

random (but see Methods for more general formulations that enable other mixing

patterns). Under random mixing and heterogeneous connectivity, the force of

infection[7] is written as $\lambda(x) = (\beta/N)(\int y[\rho E(y) + I(y)]\, dy / \int y g(y)\, dy)$, the basic

reproduction number is

$$R_0 = (1 + CV^2)\beta\left(\frac{\rho}{\delta} + \frac{1}{\gamma}\right), \qquad\qquad (6)$$

$R_c(t)$ is as above and $R_{eff}(t)$ is derived by a more general expression given in

Methods. Applying this model to the same epidemics as before we estimate: R_0

rounding 7.1 (Belgium), 3.8 (England), 7.9 (Portugal) and 6.6 (Spain); individual

susceptibility CV reaching 2.9 (Belgium), 1.6 (England), 4.0 (Portugal) and 2.7

(Spain); and intervention efficacy during lockdown being 73% (Belgium), 58%

(England), 80% (Portugal) and 72% (Spain).

Comparing the two models, variation in connectivity systematically leads to estimates

that are higher for R_0, lower for CV, and higher for the efficacy of non-

pharmaceutical interventions. Nevertheless, the percentage of the population required

to be immune to curb the epidemic and prevent future waves when interventions are

lifted appears remarkably conserved across models: 9.6 vs 11% (Belgium); 20 vs 21%

(England); 7.3 vs 6.0% (Portugal); and 12 vs 11% (Spain). This property is further

explored below.

Herd immunity thresholds and their conserveness across models

Individual variation in risk of acquiring infection is under selection by the force of

infection, whether individual differences are due to biological susceptibility,

exposure, or both. The most susceptible or exposed individuals are selectively

removed from the susceptible pool as they become infected and eventually recover

(some die), resulting in decelerated epidemic growth and accelerated induction of

immunity in the population. In essence, the *herd immunity threshold* defines the

percentage of the population that needs to be immune to reverse epidemic growth and

prevent future waves. When individual susceptibility or connectivity is gamma-

distributed and mixing is random, HIT curves can be derived analytically[8] from the

model systems (Equations 1-4, with the respective forces of infections). In the case of

variation in susceptibility to infection we obtain

$$HIT = 1 - \left[\frac{1 - \sigma R_0}{(1 - \sigma)R_0}\right]^{\frac{1}{1+CV^2}},$$ (7)

while variable connectivity results in

$$HIT = 1 - \left[\frac{1 - \sigma R_0}{(1 - \sigma)R_0}\right]^{\frac{1}{1+2CV^2}}.$$ (8)

In more complex cases HIT curves can be approximated numerically. Figure 3 shows

the expected downward trends in HIT and the sizes of the respective unmitigated

epidemics for SARS-CoV-2 without reinfection ($\sigma = 0$) as the coefficients of

variation are increased (gamma distribution shapes adopted here are illustrated in

Extended Data Figure 1; for robustness of the trends to other distributions see Gomes

et al[9]). Values of R_0 and CV estimated for our study countries are overlaid to mark

the respective HIT and final epidemic sizes. While herd immunity is expected to

require 60-80% of a homogeneous population to have been infected, at the cost of

infecting almost the entire population if left unmitigated, given an R_0 between 2.5 and

5, these percentages drop to the range 10-20% or lower when CV is roughly between

2 and 5.

When acquired immunity is not 100% effective ($\sigma > 0$) HITs are relatively higher

(Extended Data Figure 2). However, there is an upper bound for how much it is

reasonable to increase σ before the system enters a qualitatively different regime.

Above $\sigma = 1/R_0$ – the *reinfection threshold*[10,11]– infection becomes stably endemic

and the HIT concept no longer applies. Respiratory viruses are typically associated

with epidemic dynamics below the reinfection threshold, characterized by seasonal

epidemics intertwined with periods of undetection.

Individual variation in exposure, in contrast with susceptibility, accrues from complex

patterns of human behaviour which have been simplified in our model. To explore the

scope of our results we generalise our models (Methods) by relaxing some key

assumptions. First, we enable mixing to be assortative in the sense that individuals

contact predominantly with those of similar connectivity. Formally, an individual

with connectivity x, rather than being exposed uniformly to individuals of all

connectivities y, has contact preferences described by a normal distribution on the

difference $y - x$. We find this modification to have negligible effect on HIT

(Extended Data Figure 3). Second, we allow connectivity distributions to change in

shape (not only scale) when subject to social distancing. In particular we modify the

model so that CV reduces in proportion to the intensity of social distancing (Extended

Data Figure 4) and replicate the fittings to epidemics in our study countries (Extended

Data Figure 5). We find a general tendency for this model to estimate higher values

for R_0 and CV while HIT remains again remarkably robust to the change in model

assumptions.

Herd immunity thresholds and seroprevalence at sub-national levels

As countries are conducting immunological surveys to assess the extent of exposure

to SARS-CoV-2 in populations it is of practical importance to understand how HIT

may vary across regions. We have redesigned our analyses to address this question.

Series of daily new cases were stratified by region. Fitting the models simultaneously

to the multiple series enabled the estimation of local parameters (R_0 and CV) while

the effects of NPIs were estimated at country level. Extended Data Figures 6-9 show

how the modelled epidemics fit the regional data and include an additional metric to

describe the cumulative infected percentage. These model projections are comparable

to data from seroprevalence studies such as Spain[12]. We emphasise that

seroprevalence estimates generally lie slightly below our cumulative infection curves

(Extended Data Figure 9) consistently with recent findings that a substantial fraction

of infected individual does not exhibit detectable antibodies[13]. In addition to their

practical utility these results begin to unpack some of the variation in HIT within

countries: Belgium (9.4-11%), England (16-26%), Portugal (7.1-9.9%) and Spain

(7.5-21%).

Discussion

The concept of *herd immunity* was developed in the context of vaccination

programs[14,15]. Defining the percentage of the population that must be immune to

cause infection incidences to decline, HITs constitute useful targets for vaccination

coverage. In idealized scenarios of vaccines delivered at random and individuals

mixing at random, HITs are given by a simple formula $(1 - 1/R_0)$ which, in the case

of SARS-CoV-2, suggests that 60-80% of randomly chosen subjects of the population

would need be immunized to halt spread considering estimates of R_0 between 2.5 and

5. This formula does not apply to infection-induced immunity because natural

infection does not occur at random. Individuals who are more susceptible or more

exposed are more prone to be infected and become immune, providing greater

community protection than random vaccination[16]. In our model, the HIT declines

sharply when coefficients of variation increase from 0 to 2 and remains below 20%

for more variable populations. The magnitude of the decline depends on what

property is heterogeneous and how it is distributed among individuals, but the

downward trend is robust as long as susceptibility or exposure to infection are

variable (Figure 3 and Extended Data Figures 3) and acquired immunity is efficacious

enough to keep transmission below the reinfection threshold (Extended Data Figure

2).

Several candidate vaccines against SARS-CoV-2 are showing promising safety and

immunogenicity in early-phase clinical trials[17,18], although it is not yet known how

this will translate into effective protection. We note that the reinfection threshold[10,11]

informs not only the requirements on naturally acquired immunity but, similarly, it

sets a target for how efficacious a vaccine needs to be in order to effectively interrupt

transmission. Specifically, given an estimated value of R_0 we should aim for a

vaccine efficacy of $1 - 1/R_0$ (60% or 80% if R_0 is 2.5 or 5, respectively). A vaccine

whose efficacy is insufficient to bring the system below the reinfection threshold will

not interrupt transmission.

Heterogeneity in the transmission of respiratory infections has traditionally focused

on variation in exposure summarized into age-structured contact matrices. Besides

overlooking differences in susceptibility given exposure, the aggregation of

individuals into age groups reduces coefficients of variation. We calculated CV for

the landmark POLYMOD matrices[19,20] and obtained values between 0.3 and 0.5.

Recent studies of COVID-19 integrated contact matrices with age-specific

susceptibility to infection (structured in three levels)[21] or with social activity (three

levels also)[22] which, again, resulted in coefficients of variation less than unity. We

show that models with coefficients of variation of this magnitude would appear to

differ only moderately from homogeneous approximations when compared with our

estimates, which are consistently above 1 in England and above 2 in Belgium,

Portugal and Spain. In contrast with reductionistic procedures that aim to reconstruct

variation from correlate markers left on individuals (such as antibody or reactive T

cells for susceptibility, or contact frequencies for exposure), we have embarked on a

holistic approach designed to infer the whole extent of individual variation from the

imprint it leaves on epidemic trajectories. Our estimates are therefore expected to be

higher and should ultimately be confronted with more direct measurements as these

become available. Adam at et[23] conducted a contact tracing study in Hong Kong and

estimated a coefficient of variation of 2.5 for the number of secondary infections

caused by individuals, attributing 80% of transmission to 20% of cases. This

statistical dispersion has been interpreted as reflecting a common pattern of contact

heterogeneity which has been corroborated by studies that specifically measure

mobility[24]. According to our inferences, 20% of individuals may be responsible for

47-94% infections depending on model and country. In parallel, there is accumulating

evidence of individual variation in the immune system′s ability to control SARS-

CoV-2 infection following exposure[25,26]. While our inferences serve their purpose of

improving accuracy in model predictions, diverse studies such as these are necessary

for developing interventions targeting individuals who may be at higher risk of being

infected and propagating infection in the community.

Country-level estimates of R_0 reported here are in the range 3-5 when individual

variation in susceptibility is factored and 4-8 when accounting for variation in

connectivity. The homogeneous version of our models would have estimated R_0

between 2.4 and 3.3, in line with other studies[27]. Estimates for England suggest lower

baseline R_0 and lower CV in comparison with the other study countries (Belgium,

Portugal and Spain). The net effect is a slightly higher HIT in England which

nevertheless we estimate around 20%. The lowest HIT, at less than 10%, is estimated

in Portugal, with higher R_0 and higher CV. NPIs reveal less impact under variable

susceptibility (48-69%), followed by variable connectivity (58-80%), and finally

appear to inflate and agree with Flaxman et al[27] when homogeneity assumptions are

made (65-89%), although this does not affect the HIT which relates to pre-pandemic

societies.

More informative than reading these numbers, however, is to look at simulated

projections for daily new cases over future months (Figures 1 and 2). In all four

countries considered here we foresee HIT being achieved between July and October

and the COVID-19 epidemic being mostly resolved by the end of 2020. Looking

back, we conclude that NPIs had a crucial role in halting the growth of the initial

wave between February and April. Although the most extreme lockdown strategies

may not be sustainable for longer than a month or two, they proved effective at

preventing overshoot, keeping cases within health system capacities, and may have

done so without impairing the development of herd immunity.

1.   Ferguson, N. M., et al. Impact of non-pharmaceutical interventions (NPIs) to reduce

COVID-19 mortality and healthcare demand (Imperial College COVID-19 Response

Team, 2020). 10.25561/77482.

2.   Kissler, S. M., Tedijanto, C., Goldstein, E., Grad, Y. H. & Lipsitch, M. Projecting the

transmission dynamics of SARS-CoV-2 through the postpandemic period. *Science* **368**,

860-868 (2020).

3.   Wu, J. T., Leung, K. & Leung, G. M. Nowcasting are forecasting the potential domestic

and international spread of the 2019-nCov outbreak originating in Wuhan, China: a

modelling study. *Lancet* **395**, 689-697 (2020).

4.   Kwok, K. O., Lai, F., Wei, W. I., Wong, S. Y. S. & Tang, J. Herd immunity – estimating

the level required to halt the COVID-19 epidemics in affected countries. *J. Infect.* **80**,

e32-e33 (2020).

5.   Diekmann, O., Heesterbeek, J. A. P. & Metz, J. A. J. On the definition and computation

of the basic reproduction ratio R_0 in models for infectious diseases in heterogeneous

populations. *J. Math. Biol.* **28**, 365-382 (1990).

6.   Google, COVID-19 Community Mobility Reports (2020).

7.   Pastor-Satorras, R. & Vespignani, A. Epidemic dynamics and endemic states in

complex networks. *Phys. Rev. E* **63**, 066117 (2001).

8.   Gomes, M. G. M. & Montalbán, A. A SEIR model with variable susceptibility or

exposure to infection. *arXiv* (2020).

9.   Gomes, M. G. M., et al. Individual variation in susceptibility or exposure to SARS-CoV-

2 lowers the herd immunity threshold. *medRvix* 10.1101/2020.04.27.20081893 (2020).

10.   Gomes, M. G. M., White, L. J. & Medley, G. F. Infection, reinfection, and vaccination

under suboptimal immune protection: Epidemiological perspectives. *J. Theor. Biol.* **228**,

539-549 (2004).

11. Gomes, M. G. M., Gjini, E., Lopes, J. S., Souto-Maior, C. & Rebelo, C. A theoretical

framework to identify invariant thresholds in infectious disease epidemiology. *J. Theor.*

Biol. **395**, 97-102 (2016).

12. Pollán, M., et al. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide,

population-based seroepidemiological study. *Lancet* 10.1016/s0140-6736(20)31483-5

(2020).

13. Sekine, T., et al. Robust T cell immunity in convalescent individuals with asymptomatic

or mild COVID-19. *medRvix* 10.1101/2020.06.29.174888 (2020).

14. Gonçalves, G. Herd immunity: recent uses in vaccine assessment. *Expert Rev. Vaccines*

7, 1493-1506 (2008).

15. Fine, P., Eames, K. & Heymann, D. L. "Herd immunity": a rough guide, *Clin. Infect.*

Dis. **52**, 911-916 (2011).

16. Ferrari, M. J., Bansal, S., Meyers, L. A. & Bjornstad, O. N. Network frailty and the

geometry of herd immunity. *Proc. R. Soc. B* **273**, 2743-2748 (2006).

17. Folegatti, P. M., et al. Safety and immunogenicity of the ChAdOx1 nCoV-19 vaccine

against SARS-CoV-2: a preliminary report of a phase 1/2, single-blind, randomised

controlled trial. Lancet 10.1016/S0140-6736(20)31604-4 (2020).

18. Zhu, F.-C., et al. Immunogenicity and safety of a recombinant adenovirus type-5-

vectored COVID-19 vaccine in healthy adults aged 18 years or older: a randomised,

double-blind, placebo-controlled, phase 2 trial. Lancet 10.1016/S0140-6736(20)31611-1

(2020).

19. Mossong, J., et al. Social contacts and mixing patterns relevant to the spread of

infectious diseases. *PLOS Med*. **5**, e74 (2008).

20. Prem, K., Cook, A. R. & Jit, M. Projecting social contact matrices in 152 countries using

contact surveys and demographic data. *PLOS Comput. Biol.* **13**, e1005697 (2017).

21. Zhang, J., et al. Changes in contact patterns shape the dynamics of the COVID-19

outbreak in China. *Science* **368**, 1481-1486 (2020).

22. Britton, T., Ball, F. & Trapman, P. A mathematical model reveals the influence of

population heterogeneity on herd immunity to SARS-CoV-2. *Science*

10.1126/science.abc6810 (2020).

23. Adam, D., et al. Clustering and superspreading potential of severe acute respiratory

syndrome coronavirus 2 (SARS-CoV-2) infections in Hong Kong. 10.21203/rs.3.rs-

29548/v1

24. Eubank, S., et al. Modelling disease outbreaks in realistic urban social networks. *Nature*

429, 180-184 (2004).

25. Grifoni, A., et al. Targets of T cell responses to SARS-CoV-2 coronavirus in humans

with COVID-19 disease and unexposed individuals. *Cell* **181**, 1489-1501.e15 (2020).

26. Le Bert, N., et al. SARS-CoV-2-specific T cell immunity in cases of COVID-19 and

SARS, and uninfected controls. *Nature* 10.1038/s41586-020-2550-z (2020).

27. Flaxman, S., et al. Estimating the effects of non-pharmaceutical interventions on

COVID-19 in Europe. *Nature* 10.1038/s41586-020-2405-7 (2020).

Fig. 1| SARS-CoV-2 transmission with individual variation in susceptibility to infection. Suppressed wave and subsequent dynamics in Belgium, England, Portugal and Spain (orange). Estimated epidemic in the absence of interventions revealing overshoot (black). Blue bars are daily new cases. Basic (R_0) and effective ($R_{eff} = \{\int \lambda(x)x[S(x) + \sigma R(x)]\,dx / \int \rho E(x) + I(x)\,dx\}\{\rho/\delta + 1/\gamma\}$) reproduction numbers are displayed on shallow panels underneath the main plots. Blue shades represent social distancing (intensity reflected in R_0 trends and shade density). Susceptibility factors implemented as gamma distributions. Consensus parameter values (Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per day; and $\rho = 0.5$. Fraction of infected individuals identified as positive (reporting fraction): 0.06 (Belgium); 0.024 (England); 0.09 (Portugal); 0.06 (Spain). Basic reproduction number, coefficients of variation and social distancing parameters estimated by Bayesian inference as described in Methods (estimates in Extended Data Table 1). Curves represent mean model predictions from 10^4 posterior samples. Orange shades represent 95% credible intervals. Vertical lines represent the expected time when herd immunity threshold will be achieved. Circles depict independent mobility data Google[6] not used in our parameter estimation.

Fig. 2| SARS-CoV-2 transmission with individual variation in exposure to infection. Suppressed wave and subsequent dynamics in Belgium, England, Portugal and Spain (orange). Estimated epidemic in the absence of interventions revealing overshoot (black). Blue bars are daily new cases. Basic (R_0) and effective ($R_{eff} = \{\int \lambda(x)x[S(x) + \sigma R(x)]\,dx/\int \rho E(x) + I(x)\,dx\}\{\rho/\delta + 1/\gamma\}$) reproduction numbers are displayed on shallow panels underneath the main plots. Blue shades represent social distancing (intensity reflected in R_0 trends and shade density). Exposure factors implemented as gamma distributions. Consensus parameter values (Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per day; and $\rho = 0.5$. Fraction of infected individuals identified as positive (reporting fraction): 0.06 (Belgium); 0.024 (England); 0.09 (Portugal); 0.06 (Spain). Basic reproduction number, coefficients of variation and social distancing parameters estimated by Bayesian inference as described in Methods (estimates in Extended Data Table 2). Curves represent mean model predictions from 10^4 posterior samples. Orange shades represent 95% credible intervals. Vertical lines represent the expected time when herd immunity threshold will be achieved. Circles depict independent mobility data Google[6] not used in our parameter estimation.

385
386

Fig. 3| Herd immunity threshold with gamma-distributed susceptibility or exposure to infection. Curves generated with the SEIR model (Equation 1-4) assuming values of R_0 estimated for the study countries (Extended Data Tables 1 and 2) assuming gamma-distributed: susceptibility (top); connectivity (bottom). Herd immunity thresholds (solid curves) are calculated according to the formula $1 - (1/R_0)^{1/(1+CV^2)}$ for heterogeneous susceptibility and $1 - (1/R_0)^{1/(1+2CV^2)}$ for heterogeneous connectivity. Final sizes of the corresponding unmitigated epidemics are also shown (dashed).

METHODS

Model structure and underlying assumptions

The model presented here is a differential equation SEIR model, where susceptible individuals become exposed at a rate that depends on their susceptibility, the number of potentially infectious contacts they engage in, and the total number of infectious people in the population per time unit. Upon exposure, individuals enter an asymptomatic incubation phase, during which they slowly become infectious[29-32]. Thus, infectivity of exposed individuals is made to be $1/2$ of that of infectious ones ($\rho = 0.5$). After a few days, individuals develop symptoms – on average 4 days after the exposure to the virus ($\delta = 1/4$) – and thus become fully infectious[33-35]. They recover, i.e., they are no longer infectious 4 days after that ($\gamma = 1/4$), on average[36].

Efficacy of acquired immunity

We conducted the core of our analysis under the assumption that no reinfection occurs after recovery due to acquired immunity ($\sigma = 0$). To analyse the sensitivity of these results to leakage in immune response ($\sigma > 0$) we calculated herd immunity thresholds (HIT) as a function of coefficients of variation (CV) for different values of σ. The results displayed in Extended Data Figure 2 confirm the expectation that as the efficacy of acquired immunity decreases (σ increases) larger percentages of the population are infected before herd immunity is reached. Less intuitive is that there is an upper bound for how much it is reasonable to increase σ before the system enters a qualitatively different regime – the reinfection threshold[10-11] ($\sigma = 1/R_0$) – above which infection becomes stably endemic and the notion of herd immunity threshold

no longer applies. Respiratory viruses are typically associated with epidemics

dynamics below the reinfection threshold.

Effective reproduction number

The effective reproduction number (R_{eff}, also denoted by R_e or R_t by other authors)

is a time-dependent quantity which we calculate as the incidence of new infections

divided by the total number of active infections (affected by ρ for individuals in E)

multiplied by the average duration of infection (also affected by ρ for individuals in

E)

$$R_{eff}(t) = \frac{\int \lambda(x,t)x[S(x,t) + \sigma R(x,t)] \, dx}{\int \rho E(x,t) + I(x,t) \, dx} \left(\frac{\rho}{\delta} + \frac{1}{\gamma}\right).$$ (9)

Assortative mixing

In the main text we assumed random mixing among individuals, but human

connectivity patterns are assortative due societal structures and human behaviours. To

explore the sensitivity of our results to deviations from random mixing, we develop

an extended formalism that allows individuals to connect preferentially with those

with similar connectivity, formally $\lambda(x) =$

$(\beta/N)(\int y \, h(y-x)[\rho E(y) + I(y)] \, dy / \int y g(y) \, dy)$, where $h(y-x)$ is a normal

distribution on the difference between connectivity factors (Extended Data Figure 3).

Dynamic coefficients of variation

The formulation of the variable connectivity model in the main text assumes that

coefficients of variation are constant irrespective of interventions. Social distancing

has been assumed to reduce connectivity of every individual by the same factor (from

x to $[1-d]x$) leaving the coefficient of variation unchanged. The possibility that CV

might reduce with social distancing (d), causing a drop in the intensity of selection,

might affect our results. To study sensitivity to this type of CV dynamics, we

formulate an extended model where connectivity is reformulated as $(1-d)[1+$

$(1-d)(x-1)]$, and whose CV decreases with social distancing (Extended Data

Figure 4). This does not change the way the model is written but special care is

needed in analysis and interpretation to account for the new dynamics. The basic

reproduction number, in particular, depends explicitly on a CV which is now

dependent on social distancing

$$R_0 = [1 + CV(d)^2]\beta \left(\frac{\rho}{\delta} + \frac{1}{\gamma}\right), \tag{10}$$

which is noticeable in the curvilinear shape of the controlled R_0 (R_c) trajectories

(Extended Data Figure 5).

Non-pharmaceutical interventions

We implemented non-pharmaceutical interventions (NPI) as a gradual decrease in

viral transmissibility in the population and thus a lowering of the controlled and

effective reproduction numbers (R_c and R_{eff}). Once containment measures are put in

place in each country, we postulate it takes 21 days until the maximum effectiveness

of social distancing measures is reached. In the simulations presented throughout we

have held this condition (maximum "lockdown" efficacy) for 30 days, after which

period, social distancing measures are progressively relaxed, slowly returning to pre-

pandemic conditions. Both the implementation and relaxing of the social distancing

measures are imposed to be linear in this scheme.

Bayesian Inference

The model laid out above is amenable to theoretical exploration as presented in the main manuscript and provides a perfect framework for inference. Fundamentally, to be able to reproduce the inception of any epidemic, we would need to estimate when local transmission started to occur (t_0), and the pace at which individuals infected each other in the very early stages of the epidemic (R_0). All countries, to different extents and at different timepoints of the epidemic, enforced some combination of social distancing measures. To fully understand the interplay between herd immunity and the impact of NPIs, we then set out to estimate the time at which social distancing measures started to have an impact on daily incidence (t_0^d), what their maximum effectiveness (d_{max}) is, the basic reproduction number (R_0) and what the underlying variance in heterogeneity is for both susceptibility to infection and number of infectious contacts.

In order to preserve identifiability, we made two simplifying assumptions: (*i*) the fraction of infectious individuals reported as COVID-19 cases (reporting fraction) is constant throughout the study period and is comparable between countries proportionally to the number of tests performed per person; (*ii*) local transmission starts (t_0) when countries/regions report 1 case per 5 million population in one day. To calculate the reporting rates, we used the Spanish national serological survey[12] as a reference and divided the total number of reported cases up to May 11[th] by the estimated number of people that had been exposed to the virus. This gives us a reporting rate for Spain around 6%. Unfortunately, there are no other national serological surveys that could inform the proportion of the population infected in other countries, so we had to extrapolate the reporting rate for those. Assuming the

reporting rate is highly dependent on the testing effort employed in each country,

reflected in the number of tests per individual, we estimate the reporting rate by

scaling the reporting rate recorded in Spain according to the ratio of PCR tests per

person in other countries relative to the Spanish reference of 0.9 tests per thousand

people (https://ourworldindata.org/coronavirus-testing). This produced estimated case

reporting rates (ratio of reported cases to infections) of 9% for Portugal, 6% for

Belgium (and Spain) and 2.4% for England.

Whist national case and mortality data is easily available for most countries, more

spatially resolute data is difficult to find in the public domain. Thus, we restricted our

analysis to countries for which disaggregated regional case data was easily available.

We collected the data at two time points. First, we compiled all available data from

the day the countries started reporting COVID-19 cases to the initial collection date

(May 20th) and later collated available data from May 21st to July 10th.

Parameter estimation was performed with the software MATLAB, using PESTO

(Parameter EStimation Toolbox)[37], and assuming the reported case data can be

accurately described by a Poisson process. We first fixed the beginning of local

transmission (parameter t_0) in each data series as the day in which reported cases

surpassed 1 in 5 million individuals. Next, we optimized the model for the set of

parameters $\theta = \{R_0, CV, t_0^d, d_{max}\}$ by maximizing the logarithm of the likelihood

(*LL*) (Equation 11) of observing the daily reported number of cases in each country

$D = \{(k, \tilde{y}_k)\}_{k=0}^n$:

$$LL(\theta|D) = -\sum_{k=1}^n y(k,\theta) + \left(\sum_{k=1}^n \tilde{y}(k) \ln(y(k,\theta))\right) - ln\left(\prod_{k=1}^n \tilde{y}(k)!\right) \quad (11)$$

in which $y(k, \theta)$ is the simulated model output number of COVID-19 cases at day k

(with respect to t_0), and n is the total number of days included in the analysis for each

country.

When fitting the model to disaggregated data, we follow the procedure outlined above

and estimate region-specific R_0 and CV, with common t_0^d and d_{max}. To ensure that

the estimated maximum is a global maximum, we performed 50 multi-starts

optimizations, and selected the combination of parameters resulting in the maximal

Loglikelihood as a starting point for 10^4 Markov Chain Monte-Carlo iterations. From

the resulting posterior distributions, we extract the median estimates for each

parameter and the respective 95% credible intervals for the set of parameters $\theta =$

$\{R_0, CV, t_0^d, d_{max}\}$. We used uniformly distributed priors with ranges {1-9, 0.0025-

8,1-60, 0-0.7}.

This fitting procedure was applied to 4 countries (Belgium, England, Portugal and

Spain) for both the national and disaggregated case data series and repeated for each

of the 4 model variants considered here (homogeneous, heterogeneous susceptibility,

heterogeneous connectivity with constant CV, and heterogeneous connectivity with

CV reducing in proportion to social distancing). In the fitting procedures using sub-

national data, we assumed regions had the same start date for interventions that

mitigate transmission (t_0^d), and that these measures produced the same maximum

impact on transmission (d_{max}) everywhere. Thus, the only region-specific parameters

to be estimated are $\{R_{0_i}, CV_i\}$. Parameter estimates obtained from each of the model

variants are displayed in Extended Data Table 1 (heterogeneity in susceptibility),

Extended Data Table 2 (heterogeneity in connectivity with constant CV), Extended

Data Table 3 (heterogeneity in connectivity with dynamic CV) and Extended Data

Table 4 (homogeneous model), are comparable to those obtained in other studies[27,38-

43]. Finally, we apply the Akaike information criterion (AIC) for each estimation

procedure to inform on the quality of each model's fit to the datasets of reported cases

(Extended Data Table 5). In all cases, heterogeneous models are preferred over the

homogeneous approximation. Homogeneous models systematically fail to fit the

maintenance of low numbers of cases after the relaxation of social distancing

measures in many countries and regions (images not shown). The three heterogeneous

models are roughly equally well supported by the data used in this study. Further

research should complement this with discriminatory data types and hybrid models to

enable the integration of different forms of individual variation.

Data and code availability

Datasets are publicly available at the respective national ministry of health websites

(44-48). Core models implemented in MATLAB available from:

https://github.com/mgmgomes1/covid

28. Wei, W. E., et al. Presymptomatic Transmission of SARS-CoV-2 — Singapore, January

23–March 16, 2020. MMWR Morb Mortal Wkly Rep [Internet]. 2020 Apr 10 [cited

2020 May 4];69(14):411–5. Available from:

http://www.cdc.gov/mmwr/volumes/69/wr/mm6914e1.htm?s_cid=mm6914e1_w

29. To, K. K. W., et al. Temporal profiles of viral load in posterior oropharyngeal saliva

samples and serum antibody responses during infection by SARS-CoV-2: an

observational cohort study. *Lancet Infect. Dis.* **20**, 565-74 (2020).

30. Arons, M. M., et al. Presymptomatic SARS-CoV-2 Infections and Transmission in a

Skilled Nursing Facility. *N. Engl. J. Med.* **382**, 2081-2090 (2020).

31. He, X., et al. Temporal dynamics in viral shedding and transmissibility of COVID-19.

Nat. Med. **26**, 672-675 (2020).

32. Lauer, S. A., et al. The Incubation Period of Coronavirus Disease 2019 (COVID-19)

From Publicly Reported Confirmed Cases: Estimation and Application. *Ann. Intern.*

Med. **172**, 577-582 (2020).

33. Li, Q., et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-

infected pneumonia. *N. Engl. J. Med.* **382**, 1199-1207 (2020).

34. Zhang, J., et al. Evolving epidemiology and transmission dynamics of coronavirus

disease 2019 outside Hubei province, China: A descriptive and modelling study. *Lancet*

Infect. Dis. **20**, 793-802 (2020).

35. Nishiura, H., Linton, N. M. & Akhmetzhanov, A. R. Serial interval of novel coronavirus

(COVID-19) infections. *Int. J. Infect. Dis.* **93**, 284-6 (2020).

36. Stapor, P., et al. PESTO: Parameter EStimation TOolbox. *Bioinformatics* **34**, 705-707

(2018).

37. Jarvis, C. I., et al. Quantifying the impact of physical distance measures on the

transmission of COVID-19 in the UK. *BMC Medicine* **18**, 124 (2020).

38. Prem, K., et al. The effect of control strategies to reduce social mixing on outcomes of

COVID-19 epidemic in Wuhan, China: a modelling study. *Lancet Public Health* **5**, e261-

e270.

39. Tian, H., et al. An investigation of transmission control measures during the first 50 days

of the COVID-19 epidemic in China. *Science* **368**, 638-642.

40. Kucharski, A. J., et al. Effectiveness of isolation, testing, contact tracing and physical

distancing on reducing transmission of SARS-CoV-2 in different settings: a

mathematical modelling study. *Lancet Infect. Dis.* 10.1016/s1473-3099(20)30457-6

(2020).

41. Salje, H., et al. Estimating the burden of SARS-CoV-2 in France. *Science* **369**, 208-211

(2020).

42. Di Domenico, L., Pullano, G., Sabbatini, C. E., Boelle, P.-Y. & Colizza, V. Expected

impact of lockdown in Île-de-France and possible exit strategies. *medRxiv*

10.1101/2020.04.13.20063933 (2020).

43. https://ourworldindata.org/coronavirus-testing#source-information-country-by-

country. Accessed on July 10th 2020.

44. https://cnecovid.isciii.es/covid19.

45. https://covid-19.sciensano.be/nl/covid-19-epidemiologische-situatie.

46. https://covid19.min-saude.pt/ponto-de-situacao-atual-em-portugal.

47. https://coronavirus.data.gov.uk.

Acknowledgements

We thank Jan Hasenauer and Antonio Montalbán for helpful discussions concerning

statistical inference and mathematics, respectively. R.M.C. and M.U.F. receive

scholarships from the Conselho Nacional de Desenvolvimento Científico e

Tecnológio (CNPq), Brazil.

Author contributions

M.G.M.G. conceived the study. R.A. and R.M.C. and M.G.M.G. performed the

analyses. All authors interpreted the data and wrote the paper.

Competing interests

The authors declare no competing interests.

Extended Data Fig. 1| Distributions used for variable susceptibility and
connectivity. Gamma distribution probability density functions with mean 1 and
various coefficients of variation: $\left[x^{1/CV^2-1}e^{-x/CV^2}\right]/\left[\Gamma(1/CV^2)CV^{(2/CV^2)}\right]$, where Γ
is the Gamma function. For numerical implementations we discretized gamma
distributions into N bins, calculated the susceptibility or connectivity factor as well as
the fraction of the population in each bin, and derived the associated $4N$-dimensional
systems of ordinary differential equations.
613

Extended Data Fig. 2| Herd immunity threshold and epidemic final size with
reinfection. Curves in the main panels generated with the SEIR model (Equation 1-4)
assuming $R_0 = 3$ and gamma-distributed susceptibility (top) or connectivity (bottom).
Efficacy of acquired immunity is captured by a reinfection parameter σ, potentially
ranging between $\sigma = 0$ (100% efficacy) and $\sigma = 1$ (0 efficacy). This illustration
depicts final sizes of unmitigated epidemics and associated HIT curves for 6 values of
σ: $\sigma = 0$ (black); $\sigma = 0.1$ (green); $\sigma = 0.2$ (blue); $\sigma = 0.3$ (magenta); $\sigma = 1/3$ (red);
and $\sigma = 0.4$ (orange);. Above $\sigma = 1/R_0$ (reinfection threshold (Gomes et al 2004;
2016)) the infection becomes stably endemic and there is no herd immunity threshold.
Representative epidemics of the regime $\sigma \leq 1/R_0$ are shown on the right while the
regime $\sigma > 1/R_0$ is illustrated on top. All depicted dynamics are based on the
rightmost CVs represented on the main panel.
629

Extended Data Fig. 3| Herd immunity threshold and epidemic final size with gamma-distributed exposure to infection and assortative mixing. Curves in central panel generated with the SEIR model (Equation 1-4) assuming $R_0 = 3$ and gamma-distributed connectivity. Assortative mixing is implemented by imposing a normal distribution for contact preferences such that individuals contact preferentially with those with the similar contact degree (left). This illustration used normal distributions with standard deviation $SD = 50$ (green); $SD = 10$ (blue); and $SD = 2$ (magenta). More assortative mixing leads to more skewed epidemics. Herd immunity thresholds were calculated numerically as the percentage of the population no longer susceptible when new outbreaks are effectively prevented (approximately when the exposed fraction crosses the peak in the absence of mitigation). Final sizes of the corresponding unmitigated epidemics are also shown. Representative epidemics are depicted on the right based on the rightmost CVs represented on the main panel (with vertical lines marking the point when herd immunity is achieved).

Extended Data Fig. 4| Connectivity distributions with reducing coefficient of
variation in proportion to social distancing. Individual variation in connectivity is
originally implemented as a gamma distribution of mean 1 parameterised by the
coefficient of variation (CV) (black). Social distancing is initially implemented as a
reduction in connectivity by the same factor to every individual, from x to $(1 - d)x$
(top panels). Sensitivity of the results to the possibility that CV might reduce with
social distancing with replicated the analyses with a model connectivity is
reformulated as $(1 - d)[1 + (1 - d)(x - 1)]$ (bottom panels).
657

Extended Data Fig. 5| SARS-CoV-2 transmission with individual variation in exposure reduced by social distancing. Suppressed wave and subsequent dynamics in Belgium, England, Portugal and Spain. Blue bars are daily new cases. Basic (R_0) and effective ($R_{eff} = \{\int \lambda(x)x[S(x) + \sigma R(x)] \, dx / \int \rho E(x) + I(x) \, dx\}\{\rho/\delta + 1/\gamma\}$) reproduction numbers are displayed on shallow panels underneath the main plots. Blue shades represent social distancing (intensity reflected in R_0 trends and shade density). Exposure factors implemented as gamma distributions. Consensus parameter values (Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per day; and $\rho = 0.5$. Fraction of infected individuals identified as positive (reporting fraction): 0.06 (Belgium); 0.024 (England); 0.09 (Portugal); 0.06 (Spain). Basic reproduction number, coefficients of variation and social distancing parameters estimated by Bayesian inference as described in Methods (estimates in Extended Data Table 3). Curves represent mean model predictions from 10^4 posterior samples. Orange shades represent 95% credible intervals. Vertical lines represent the expected time when herd immunity threshold will be achieved. Circles depict independent mobility data (Google 2020) not used in our parameter estimation.

Extended Data Fig. 6| SARS-CoV-2 transmission at subnational levels in Belgium.
Suppressed wave and subsequent dynamics in Flanders and the rest of Belgium, with
individual variation in susceptibility (left) or exposure (right). Blue bars are daily new
cases. Shades represent social distancing (intensity reflected in shade density).
Susceptibility or exposure factors implemented as gamma distributions. Consensus
parameter values (Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per day; and $\rho = 0.5$.
Fraction of infected individuals identified as positive (reporting fraction): 0.06. Basic
reproduction number, coefficients of variation and social distancing parameters
estimated by Bayesian inference as described in Methods (estimates in Extended Data
Table 1 and 2). Curves represent mean model predictions from 10^4 posterior samples.
Orange shades represent 95% credible intervals. Red curves represent cumulative
infected percentages.
690

Extended Data Fig. 7| SARS-CoV-2 transmission at subnational levels in England.
Suppressed wave and subsequent dynamics in London, Northwest, Southeast and the
rest of England, with individual variation in susceptibility (left) or exposure (right).
Blue bars are daily new cases. Shades represent social distancing (intensity reflected
in shade density). Susceptibility or exposure factors implemented as gamma
distributions. Consensus parameter values (Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per
day; and $\rho = 0.5$. Fraction of infected individuals identified as positive (reporting
fraction): 0.024. Basic reproduction number, coefficients of variation and social
distancing parameters estimated by Bayesian inference as described in
Methods(estimates in Extended Data Table 1 and 2). Curves represent mean model
predictions from 10^4 posterior samples. Orange shades represent 95% credible
intervals. Red curves represent cumulative infected percentages.

Extended Data Fig. 8| SARS-CoV-2 transmission at subnational levels in Portugal.
Suppressed wave and subsequent dynamics in the North and Centre regions versus the
rest of Portugal, with individual variation in susceptibility (left) or exposure (right).
Blue bars are daily new cases. Shades represent social distancing (intensity reflected
in shade density). Susceptibility or exposure factors implemented as gamma
distributions. Consensus parameter values (Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per
day; and $\rho = 0.5$. Fraction of infected individuals identified as positive (reporting
fraction): 0.09. Basic reproduction number, coefficients of variation and social
distancing parameters estimated by Bayesian inference as described in Methods
(estimates in Extended Data Table 1 and 2). Curves represent mean model predictions
from 10^4 posterior samples. Orange shades represent 95% credible intervals. Red
curves represent cumulative infected percentages.

Extended Data Fig. 9| SARS-CoV-2 transmission at subnational levels in Spain.
Suppressed wave and subsequent dynamics in Madrid, Catalunya and the rest of
Spain, with individual variation in susceptibility (left) or exposure (right). Blue bars
are daily new cases. Shades represent social distancing (intensity reflected in shade
density). Susceptibility or exposure factors implemented as gamma distributions.
Consensus parameter values (Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per day; and $\rho =$
0.5. Fraction of infected individuals identified as positive (reporting fraction): 0.06.
Basic reproduction number, coefficients of variation and social distancing parameters
estimated by Bayesian inference as described in Methods (estimates in Extended Data
Table 1 and 2). Curves represent mean model predictions from 10^4 posterior samples.
Orange shades represent 95% credible intervals. Red curves represent cumulative
infected percentages and vertical red segments mark seroprevalences (95% CI)
according to a recent study[12].

 Extended Data Table 1| Estimated parameters for heterogeneous susceptibility
model. Estimates generated from model fit to the national datasets are in the grey
shaded rows. The remaining rows provide the region-specific estimates. Best
parameter estimates are presented as a bold median bounded by the lower and upper
ends for the 95% credible interval. Model runs are initiated on the day (t_0) when
reported cases surpassed 1 in 5 million individuals: Belgium (day 1); England (day
29); Portugal (day 3); Spain (day 8).

Country/Region	R_0			CV			$1 - d_{max}$			t_0^d		
Belgium	4.99	**5.03**	5.06	3.87	**3.88**	3.91	0.40	**0.40**	0.41	1.00	**1.01**	1.12
Flanders	4.96	**5.00**	5.02	3.89	**3.91**	3.93	0.41	**0.41**	0.41	1.00	**1.02**	1.15
Rest	4.97	**5.01**	5.03	3.87	**3.89**	3.91						
Portugal	4.23	**4.26**	4.31	4.22	**4.26**	4.30	0.30	**0.31**	0.31	7.37	**7.71**	7.91
North/Centre	3.54	**3.58**	3.61	3.72	**3.76**	3.79	0.34	**0.35**	0.35	7.51	**7.73**	8.00
Rest	4.27	**4.32**	4.36	3.69	**3.72**	3.74						
Spain	4.08	**4.10**	4.11	3.20	**3.21**	3.22	0.37	**0.37**	0.37	16.02	**16.13**	16.23
Madrid	4.38	**4.39**	4.39	2.37	**2.37**	2.38	0.37	**0.37**	0.37	16.40	**16.41**	16.44
Catalunya	4.20	**4.21**	4.21	2.49	**2.50**	2.50						
Rest	3.96	**3.97**	3.97	3.80	**3.81**	3.82						
England	2.93	**2.94**	2.95	1.94	**1.94**	1.95	0.52	**0.52**	0.52	40.56	**40.69**	40.80
London	2.95	**2.96**	2.96	2.24	**2.24**	2.24	0.52	**0.53**	0.53	41.35	**41.51**	41.64
NorthWest	3.03	**3.04**	3.05	1.66	**1.67**	1.68						
SouthEast	2.80	**2.81**	2.82	2.07	**2.07**	2.07						
Rest	2.88	**2.88**	2.89	1.64	**1.64**	1.65						

Extended Data Table 2| Estimated parameters for heterogeneous connectivity
model (constant CV). Estimates generated from model fit to the national datasets are
in the grey shaded rows. The remaining rows provide the region-specific estimates.
Best parameter estimates are presented as a bold median bounded by the lower and
upper ends for the 95% credible interval. Model runs are initiated on the day (t_0)
when reported cases surpassed 1 in 5 million individuals: Belgium (day 1); England
(day 29); Portugal (day 3); Spain (day 8).

Country/Region	R_0			CV			$1 - d_{max}$			t_0^d		
Belgium	7.12	**7.14**	7.17	2.86	**2.87**	2.88	0.27	**0.27**	0.27	1.00	**1.01**	1.04
Flanders	7.09	**7.11**	7.13	2.86	**2.87**	2.89	0.27	**0.27**	0.28	1.00	**1.01**	1.03
Rest	7.11	**7.13**	7.15	2.86	**2.87**	2.89						
Portugal	7.76	**7.94**	8.14	4.01	**4.04**	4.09	0.19	**0.20**	0.20	2.67	**2.98**	3.27
North/Centre	5.06	**5.08**	5.09	3.24	**3.24**	3.24	0.25	**0.25**	0.25	7.21	**7.22**	7.24
Rest	5.68	**5.69**	5.69	2.79	**2.81**	2.83						
Spain	6.59	**6.60**	6.60	2.73	**2.73**	2.73	0.28	**0.28**	0.28	10.00	**10.01**	10.02
Madrid	7.81	**7.83**	8.82	1.98	**1.99**	2.06	0.24	**0.26**	0.26	5.38	**7.02**	7.06
Catalunya	8.00	**8.02**	9.00	2.33	**2.34**	2.43						
Rest	7.97	**7.99**	8.96	3.58	**3.59**	3.72						
England	3.81	**3.82**	3.83	1.55	**1.55**	1.55	0.42	**0.42**	0.42	36.45	**36.52**	36.61
London	3.70	**3.70**	3.71	1.69	**1.69**	1.70	0.43	**0.43**	0.43	37.50	**37.52**	37.55
NorthWest	3.83	**3.83**	3.84	1.32	**1.32**	1.32						
SouthEast	3.58	**3.59**	3.59	1.66	**1.67**	1.68						
Rest	3.60	**3.60**	3.61	1.30	**1.31**	1.31						

753
754
755
756

Extended Data Table 3| Estimated parameters for heterogeneous connectivity
model (dynamic CV). Estimates generated from model fit to the national datasets are
in the grey shaded rows. The remaining rows provide the region-specific estimates.
Best parameter estimates are presented as a bold median bounded by the lower and
upper ends for the 95% credible interval. Model runs are initiated on the day (t_0)
when reported cases surpassed 1 in 5 million individuals: Belgium (day 1); England
(day 29); Portugal (day 3); Spain (day 8).

Country/Region	R_0			CV			$1 - d_{max}$			t_0^d		
Belgium	8.79	**8.84**	8.88	3.98	**4.00**	4.02	0.56	**0.56**	0.56	1.00	**1.01**	1.05
Flanders	8.78	**8.82**	8.86	3.98	**4.00**	4.02	0.56	**0.56**	0.56	1.00	**1.01**	1.04
Rest	8.82	**8.86**	8.89	3.98	**4.00**	4.02						
Portugal	9.86	**9.92**	9.95	5.73	**5.77**	5.80	0.50	**0.50**	0.50	3.00	**3.01**	3.07
North/Centre	6.65	**6.72**	6.80	3.75	**3.78**	3.81	0.56	**0.57**	0.57	5.84	**6.02**	6.19
Rest	5.98	**6.05**	6.13	4.09	**4.14**	4.19						
Spain	5.97	**6.09**	6.10	3.04	**3.09**	3.10	0.56	**0.56**	0.56	14.46	**14.50**	14.89
Madrid	6.19	**6.20**	6.21	2.43	**2.43**	2.44	0.57	**0.57**	0.57	13.80	**13.81**	13.83
Catalunya	6.30	**6.32**	6.33	2.61	**2.62**	2.62						
Rest	6.34	**6.35**	6.36	3.80	**3.81**	3.82						
England	4.01	**4.05**	4.09	1.90	**1.92**	1.93	0.66	**0.66**	0.66	36.79	**37.03**	37.29
London	3.78	**3.79**	3.80	1.99	**2.00**	2.01	0.67	**0.67**	0.67	39.06	**39.20**	39.28
NorthWest	3.91	**3.92**	3.94	1.64	**1.65**	1.66						
SouthEast	3.67	**3.69**	3.70	1.89	**1.89**	1.90						
Rest	3.64	**3.65**	3.66	1.58	**1.58**	1.59						

765
766
767

Extended Data Table 4| Estimated parameters for the homogenous model.
Estimates generated from model fit to the national datasets are in the grey shaded
rows. The remaining rows provide the region-specific estimates. Best parameter
estimates are presented as a bold median bounded by the lower and upper ends for the
95% credible interval. Model runs are initiated on the day (t_0) when reported cases
surpassed 1 in 5 million individuals: Belgium (day 1); England (day 29); Portugal
(day 3); Spain (day 8).

Country/Region	R_0			$1 - d_{max}$			t_0^d		
Belgium	3.298	**3.301**	3.310	0.208	**0.209**	0.210	16.299	**16.354**	16.357
Flanders	3.235	**3.239**	3.308	0.208	**0.212**	0.213	16.324	**17.039**	17.064
Rest	3.235	**3.238**	3.307						
Portugal	2.900	**2.910**	2.914	0.242	**0.243**	0.244	20.551	**20.622**	20.693
North/Centre	3.542	**3.578**	3.608	0.343	**0.345**	0.348	7.514	**7.725**	7.999
Rest	4.274	**4.321**	4.361						
Spain	3.028	**3.031**	3.034	0.149	**0.150**	0.150	28.329	**28.360**	28.393
Madrid	4.113	**4.116**	4.120	0.111	**0.111**	0.112	20.000	**20.000**	20.000
Catalunya	4.208	**4.214**	4.218						
Rest	3.735	**3.751**	3.752						
England	2.434	**2.435**	2.437	0.355	**0.355**	0.356	55.047	**55.070**	55.074
London	2.307	**2.308**	2.310	0.359	**0.360**	0.360	54.577	**54.578**	54.578
NorthWest	2.602	**2.603**	2.604						
SouthEast	2.368	**2.370**	2.371						
Rest	2.502	**2.503**	2.504						

776
777
778

Extended Data Table 5| Model selection criteria. Displays the maximum
Loglikelihood obtained for each combination of model and data partitioning for each
country, as well as the Akaike information criterion. Models are labelled by a sort
name as follows: homog (homogenous); hetsus (heterogeneity in susceptibility);
hetcon (heterogeneity in connectivity with constant CV); hetdyn (heterogeneity in
connectivity with dynamic CV).

Country	Model	LL	AIC
Aggregate Data			
Portugal	homog	2.25E+05	-4.50E+05
	hetsus	2.30E+05	-4.59E+05
	hetcon	2.30E+05	-4.60E+05
	hetdyn	2.30E+05	-4.59E+05
Spain	homog	1.77E+06	-3.54E+06
	hetsus	1.79E+06	-3.59E+06
	hetcon	1.79E+06	-3.58E+06
	hetdyn	1.79E+06	-3.58E+06
England	homog	1.67E+06	-3.33E+06
	hetsus	1.68E+06	-3.36E+06
	hetcon	1.68E+06	-3.36E+06
	hetdyn	1.68E+06	-3.36E+06
Belgium	homog	3.30E+05	-6.60E+05
	hetsus	3.33E+05	-6.66E+05
	hetcon	3.33E+05	-6.65E+05
	hetdyn	3.33E+05	-6.66E+05
Regional Data			
Portugal	homog	3.36E+05	-6.73E+05
	hetsus	3.55E+05	-7.11E+05
	hetcon	3.55E+05	-7.10E+05
	hetdyn	3.55E+05	-7.11E+05
Spain	homog	1.87E+06	-3.75E+06
	hetsus	1.96E+06	-3.91E+06
	hetcon	1.96E+06	-3.91E+06
	hetdyn	1.47E+06	-2.95E+06
England	homog	2.59E+06	-5.18E+06
	hetsus	2.63E+06	-5.25E+06
	hetcon	2.62E+06	-5.25E+06
	hetdyn	2.62E+06	-5.25E+06
Belgium	homog	3.74E+05	-7.48E+05
	hetsus	3.78E+05	-7.56E+05
	hetcon	3.78E+05	-7.55E+05
	hetdyn	3.78E+05	-7.55E+05