Herd immunity thresholds for SARS-CoV-2 estimated from unfolding epidemics

Ricardo Aguas[1†], Rodrigo M. Corder[2†], Jessica G. King[3], Guilherme Gonçalves[4], Marcelo U. Ferreira[2], M. Gabriela M. Gomes[5,6]*

[1] *Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom.*

[2] *Instituto de Ciências Biomédicas, Universidade de São Paulo, São Paulo, Brazil.*

[3] *Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, United Kingdom.*

[4] *Unidade Multidisciplinar de Investigação Biomédica, Instituto de Ciências Biomédicas Abel Salazar, Universidade do Porto, Porto, Portugal.*

[5] *Department of Mathematics and Statistics, University of Strathclyde, Glasgow, United Kingdom.*

[6] *Centro de Matemática da Universidade do Porto, Porto, Portugal.*

[†] These authors contributed equally to this work.

[*] Correspondence and requests for materials should be addressed to M.G.M.G. (email: gabriela.gomes@strath.ac.uk).

As severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) spreads, the susceptible subpopulation declines causing the rate at which new infections occur to slow down. Variation in individual susceptibility or exposure to infection exacerbates this effect. Individuals that are more susceptible or more exposed tend to be infected and removed from the susceptible subpopulation earlier. This selective depletion of susceptibles intensifies the deceleration in incidence. Eventually, susceptible numbers become low enough to prevent epidemic growth or, in other words, the herd immunity threshold is reached. Here we fit epidemiological models with inbuilt distributions of susceptibility or exposure to SARS-CoV-2 outbreaks to estimate basic reproduction numbers (R_0) alongside coefficients of individual variation (CV) and the effects of containment strategies. Herd immunity thresholds are then calculated as $1 - (1/R_0)^{1/(1+CV^2)}$ or $1 - (1/R_0)^{1/(1+2CV^2)}$, depending on whether variation is on susceptibility or exposure. Our inferences result in herd immunity thresholds around 10-20%, considerably lower than the minimum coverage needed to interrupt transmission by random vaccination, which for R_0 higher than 2.5 is estimated above 60%. We emphasize that the classical formula, $1 - 1/R_0$, remains applicable to describe herd immunity thresholds for random vaccination, but not for immunity induced by infection which is naturally selective. These findings have profound consequences for the governance of the current pandemic given that some populations may be close to achieving herd immunity despite being under more or less strict social distancing measures.

Scientists throughout the world have engaged with governments, health agencies, and with each other, to address the ongoing pandemic of coronavirus disease (COVID-19). Mathematical models have been central to important decisions concerning

contact tracing, quarantine, and social distancing, to mitigate or suppress the initial

pandemic spread[1]. Successful suppression, however, may leave populations at risk to

resurgent waves due to insufficient acquisition of immunity. Models have thus also

addressed longer term SARS-CoV-2 transmission scenarios and the requirements for

continued adequate response[2]. This is especially timely as countries apply, relax and

reapply lockdown measures with varying levels of success in tackling national

outbreaks.

Here we demonstrate that individual variation in susceptibility or exposure

(connectivity) accelerates the acquisition of immunity in populations. More

susceptible and more connected individuals have a higher propensity to be infected

and thus are likely to become immune earlier. Due to this selective immunization by

natural infection, heterogeneous populations require less infections to cross their herd

immunity threshold (HIT) than suggested by models that do not fully account for

variation. We integrate continuous distributions of susceptibility or connectivity in

otherwise basic epidemic models for COVID-19 which account for realistic

intervention effects and show that as coefficients of variation (CV) increase from 0 to

5, HIT declines from over 60%[3,4] to less than 10%. We then fit these models to series

of daily new cases to estimate CV alongside basic reproduction numbers (R_0) and

derive the corresponding HITs.

Effects of individual variation on SARS-CoV-2 transmission

64   SARS-CoV-2 is transmitted primarily by respiratory droplets and modelled as a

65   susceptible-exposed-infectious-recovered (SEIR) process.

66   *Variation in susceptibility to infection*

67 Individual variation in susceptibility is integrated as a continuously distributed factor

68 that multiplies the force of infection upon individuals[5] as

69
$$\dot{S}(x) = -\lambda(x)xS(x), \tag{1}$$

70
$$\dot{E}(x) = \lambda(x)x[S(x) + \sigma R(x)] - \delta E(x), \tag{2}$$

71
$$\dot{I}(x) = \delta E(x) - \gamma I(x), \tag{3}$$

72
$$\dot{R}(x) = (1 - \phi)\gamma I(x) - \sigma\lambda(x)xR(x), \tag{4}$$

where $S(x)$ is the number of individuals with susceptibility x, $E(x)$ and $I(x)$ are the

numbers of individuals who originally had susceptibility x and became exposed and

infectious, while $R(x)$ counts those who have recovered and have their susceptibility

reduced to a reinfection factor σ due to acquired immunity. δ is the rate of

progression from exposed to infectious, γ is the rate of recovery or death, ϕ is the

proportion of individuals who die as a result of infection and $\lambda(x) =$

$(\beta/N) \int [\rho E(y) + I(y)]\, dy$ is the average force of infection upon susceptible

individuals in a population of size N and transmission coefficient β. Standardizing so

that susceptibility distributions have mean $\int xg(x)\, dx = 1$, given a probability

density function $g(x)$, the basic reproduction number is

83
$$R_0 = \beta\left(\frac{\rho}{\delta} + \frac{1}{\gamma}\right), \tag{5}$$

where ρ is a factor measuring the infectivity of individuals in compartment E in

relation to those in I. The coefficient of variation in individual susceptibility $CV =$

$\sqrt{\int (x-1)^2 g(x)\, dx}$ is explored as a parameter. Non-pharmaceutical interventions

(NPIs) designed to control transmission typically reduce β and hence R_0. We denote

the resulting controlled reproduction number by R_c. The effective reproduction

number R_{eff} is another useful indicator obtained by multiplying R_c by the

susceptibility of the population, in this case written as $R_{eff}(t) =$

$R_c(t) \int x S(x, t)\, dx / N(t)$ to emphasize its time dependence.

Top panels of Figure 1 depict model trajectories fitted to suppressed epidemics

(orange) in 2 European countries (Belgium and England) assuming gamma distributed

susceptibility and no reinfection ($\sigma = 0$). We estimate: R_0 rounding 5 (Belgium) and

2.9 (England); individual susceptibility CV reaching 3.9 (Belgium) and 1.5 (England);

and overall intervention efficacy at maximum (typically during lockdown) being 60%

(Belgium) and 54% (England). Other estimated parameters are the day when NPIs

begin to affect transmission, after which we assume a linear intensification from

baseline over 21 days, remaining at maximum intensity for T_{max} days and linearly

lifting back to baseline over a period of T_{lift} days (both T_{max} and T_{lift} are estimated).

Denoting by $d(t)$ the proportional reduction in average risk of infection due to

interventions, in this case we obtain $R_c(t) = [1 - d(t)]R_0$ which is depicted for each

country, alongside $R_{eff}(t)$, underneath the respective epidemic trajectories. To assess

the potential for case numbers to overshoot if NPIs had not been applied, we rerun the

model with $d(t) = 0$ and obtain the unmitigated epidemics (black). Further details

are described in Methods.

Variation in connectivity

In a directly transmitted infectious disease, such as COVID-19, variation in exposure

to infection is primarily governed by patterns of connectivity among individuals. We

incorporate this in the system (Equations 1-4) assuming that individuals mix at

random (but see Methods for more general formulations that enable other mixing

patterns). Under random mixing and heterogeneous connectivity[6], the force of

infection is written as $\lambda(x) = (\beta/N)(\int y[\rho E(y) + I(y)]\, dy / \int y g(y)\, dy)$, the basic

reproduction number is

$$R_0 = (1 + CV^2)\beta\left(\frac{\rho}{\delta} + \frac{1}{\gamma}\right), \tag{6}$$

$R_c(t)$ is as above and $R_{eff}(t)$ is derived by a more general expression given in

Methods. The results from this basic variable connectivity model are shown in

Extended Data Figure 1). To allow for the possibility that social distancing (d) may

change not only the scale but also the shape of connectivity distributions, we consider

an extended model where connectivity is reformulated as $(1 - d)[1 +$

$(1 - d)(x - 1)]$ (Extended Data Figure 2). This does not change the way the model

is written but special care is needed in analysis and interpretation to account for the

dynamic contact patterns. The basic reproduction number, in particular, depends

explicitly on a CV which is now dependent on social distancing.

Applying this model to the same epidemics as before we obtain the bottom panels of

Figure 1. The estimated epidemiological parameters are: R_0 rounding 8.9 (Belgium)

and 3.7 (England); individual connectivity CV reaching 3.9 (Belgium) and 1.6

(England); and intervention efficacy during lockdown being 45% (Belgium) and 37%

(England). The reported CVs correspond to baseline contact patterns.

Comparing the two models, variation in connectivity systematically leads to higher R_0

estimates. The effect attributed to NPIs is lower in England than in Belgium. In both

countries, contacts appear to reach minimal levels around lockdown in April (R_c

plots), beginning to re-intensify thereafter but remaining below about half of pre-

lockdown levels by the end of the data series used for model fittings (solid portions of

the R_c line), consistently with the CoMix contact survey[7]. To illustrate how the

epidemic may unfold beyond the data used in this study we simulate a hypothetical

scenario whereby contacts continue to reactivate linearly until eventually reaching the

pre-pandemic baseline. This leads to low epidemic activity over a period that

coincides with the Summer months, followed by a resurge as contacts intensify. This

resurgence is likely to be aggravated by seasonality as the Winter approaches,

although seasonality has not been included in our models. The percentage of the

population required to be immune to curb the epidemic and prevent future waves

when interventions are lifted appears remarkably conserved across models: 10 vs 11%

(Belgium); and 27 vs 25% (England).

We then fit to the same data the model constrained by $CV = 0$, i.e. assuming no

relevant heterogeneity in susceptibility or exposure to infection. The results are shown

in Figure 2. In this case epidemic potential is considerably larger and hence the effect

attributed to NPIs to fit the data must also be larger. The percentage of the population

required to be immune to prevent epidemic growth when interventions are lifted rises

to 73% (Belgium) and 63% (England).

Herd immunity thresholds

Individual variation in risk of acquiring infection is under selection by the force of

infection, whether individual differences are due to biological susceptibility,

exposure, or both. The most susceptible or exposed individuals are selectively

removed from the susceptible pool as they become infected and eventually recover

(some die), resulting in decelerated epidemic growth and accelerated induction of

immunity in the population. In essence, the *herd immunity threshold* defines the

percentage of the population that needs to be immune to reverse epidemic growth and

prevent future waves. When individual susceptibility or connectivity is gamma-

distributed and mixing is random, HIT curves can be derived analytically[8] from the

model systems (Equations 1-4, with the respective forces of infections). In the case of

variation in susceptibility to infection we obtain

$$HIT = 1 - \left[\frac{1 - \sigma R_0}{(1 - \sigma)R_0}\right]^{\frac{1}{1+CV^2}},\tag{7}$$

while variable connectivity results in

$$HIT = 1 - \left[\frac{1 - \sigma R_0}{(1 - \sigma)R_0}\right]^{\frac{1}{1+2CV^2}}.\tag{8}$$

In more complex cases (such as the variable connectivity with dynamic CV or

assortative mixing) HIT curves can be approximated numerically. Figure 3 shows the

expected downward trends in HIT and the sizes of the respective unmitigated

epidemics for SARS-CoV-2 without reinfection ($\sigma = 0$) as the coefficients of

variation are increased (here we adopt gamma distributions; for robustness of the

trends to other distributions see Gomes et al[9]). Values of R_0 and CV estimated for our

study countries are overlaid to mark the respective HIT and final epidemic sizes.

While herd immunity is expected to require 60-80% of a homogeneous population to

have been infected, at the cost of infecting almost the entire population if left

unmitigated, given an R_0 between 2.5 and 5, these percentages drop to the range 10-

20% when CV is roughly between 1.5 and 4.

When acquired immunity is not 100% effective ($\sigma > 0$) HITs are relatively higher

(Extended Data Figure 3). However, there is an upper bound for how much it is

reasonable to increase σ before the system enters a qualitatively different regime.

Above $\sigma = 1/R_0$ – the *reinfection threshold*[10,11]– infection becomes stably endemic

and the HIT concept no longer applies. Respiratory viruses are typically associated

with epidemic dynamics below the reinfection threshold, characterized by seasonal

epidemics intertwined with periods of low detection.

Individual variation in exposure, in contrast with susceptibility, accrues from complex

patterns of human behaviour which have been simplified in our basic model. To

explore the scope of our results we generalised the models by relaxing some key

assumptions. First, we allowed connectivity distributions to change in shape (not only

scale) when subject to social distancing (Figure 1, bottom panels). Second, we enable

mixing to be assortative in the sense that individuals contact predominantly with those

of similar connectivity (Methods). Formally, an individual with connectivity x, rather

than being exposed uniformly to individuals of all connectivities y, has contact

preferences described by a normal distribution on the difference $y - x$. We find this

modification to have negligible effect on HIT (Extended Data Figure 4).

Herd immunity thresholds and seroprevalence at sub-national levels

As countries conduct immunological surveys to assess the extent of exposure to

SARS-CoV-2 in populations it is of practical importance to understand how HIT may

vary across regions. We have redesigned our analyses to address this question. Series

of daily new cases in 4 European countries (Belgium, England, Portugal and Spain)

were stratified by region. Fitting the models simultaneously to the multiple series

enabled the estimation of local parameters (R_0 and CV) while the effects of NPIs

were estimated at country level. Extended Data Figures 5-8 show how the modelled

epidemics fit the regional data and include an additional metric to describe the

cumulative infected percentage. These model projections are comparable to data from

seroprevalence studies such as in Spain[12]. In addition to their practical utility these

results begin to unpack some of the variation in HIT within countries: Belgium (9.4-

11%), England (16-26%), Portugal (7.1-9.9%) and Spain (7.5-21%). The sub-national

stratification has also enabled the application of the models to countries where the

epidemic was sufficiently asynchronous across regions to compromise the ability of

the models to fit the aggregate data (Portugal and Spain).

Discussion

The concept of *herd immunity* was developed in the context of vaccination

programs[13,14]. Defining the percentage of the population that must be immune to

cause infection incidences to decline, HITs constitute useful targets for vaccination

coverage. In idealized scenarios of vaccines delivered at random and individuals

mixing at random, HITs are given by a simple formula $(1 - 1/R_0)$ which, in the case

of SARS-CoV-2, suggests that 60-80% of randomly chosen subjects of the population

would need be immunized to halt spread considering estimates of R_0 between 2.5 and

5. This formula does not apply to infection-induced immunity because natural

infection does not occur at random. Individuals who are more susceptible or more

exposed are more prone to be infected and become immune, providing greater

community protection than random vaccination[15]. In our model, the HIT declines

sharply when coefficients of variation increase from 0 to 2 and remains below roughly

20% for more variable populations. The magnitude of the decline depends on what

property is heterogeneous and how it is distributed among individuals, but the

downward trend is robust as long as susceptibility or exposure to infection are

variable (Figure 3 and Extended Data Figures 4) and acquired immunity is efficacious

enough to keep transmission below the reinfection threshold (Extended Data Figure

3).

Several candidate vaccines against SARS-CoV-2 are showing promising safety and

immunogenicity in early-phase clinical trials[16,17], although it is not yet known how

this will translate into effective protection. We note that the reinfection threshold[10,11]

informs not only the requirements on naturally acquired immunity but, similarly, it

sets a target for how efficacious a vaccine needs to be in order to effectively interrupt

transmission. Specifically, given an estimated value of R_0 we should aim for a

vaccine efficacy of $1 - 1/R_0$ (60% or 80% if R_0 is 2.5 or 5, respectively), which

seems to be materialising according to preliminary results from phase 3 trials.

Heterogeneity in the transmission of respiratory infections has traditionally focused

on variation in exposure summarized into age-structured contact matrices. Besides

overlooking differences in susceptibility given exposure, the aggregation of

individuals into age groups reduces coefficients of variation. We calculated CV for

the landmark POLYMOD matrices[18,19] and obtained values between 0.3 and 0.5.

Recent studies of COVID-19 integrated contact matrices with age-specific

susceptibility to infection (structured in three levels)[20] or with social activity (three

levels also)[21] which, again, resulted in coefficients of variation less than unity. We

show that models with coefficients of variation of this magnitude would appear to

differ only moderately from homogeneous approximations when compared with our

estimates, which are consistently above 1 in England and above 2 in Belgium,

Portugal and Spain. In contrast with reductionist procedures that aim to reconstruct

variation from correlate markers left on individuals (such as antibody or reactive T

cells for susceptibility, or contact frequencies for exposure), we have embarked on a

holistic approach designed to infer the whole extent of individual variation from the

imprint it leaves on epidemic trajectories. Our estimates are therefore expected to be

higher and should ultimately be confronted with more direct measurements as these

become available. Adam at et[22] conducted a contact tracing study in Hong Kong and

estimated a coefficient of variation of 2.5 for the number of secondary infections

caused by individuals, attributing 80% of transmission to 20% of cases. This

statistical dispersion has been interpreted as reflecting a common pattern of contact

heterogeneity which has been corroborated by studies that specifically measure

mobility[23]. According to our inferences, 20% of individuals may be responsible for

47-94% infections depending on model and country. In parallel, there is accumulating

evidence of individual variation in the immune system′s ability to control SARS-

CoV-2 infection following exposure[24,25]. While our inferences serve their purpose of

improving accuracy in model predictions, diverse studies such as these are necessary

for developing interventions targeting individuals who may be at higher risk of being

infected and propagating infection in the community.

Country-level estimates of R_0 reported here (Figures 1, 2) are in the range 3-5 when

individual variation in susceptibility is factored and 4-9 when accounting for variation

in connectivity. The homogeneous version of our models would have estimated R_0

around 2.7 and 3.7, in line with other studies[26]. Estimates for England suggest lower

baseline R_0 and lower CV in comparison with Belgium. The net effect is a slightly

higher HIT in England which nevertheless we estimate around 25-27% depending on

characteristics of the individual variation. NPIs reveal less impact when heterogeneity

is contemplated (37-60%), appearing to inflate and agree with Flaxman et al[26] when

homogeneity assumptions are made (66-78%), although this does not affect the HIT

which relates to pre-pandemic societies.

More informative than reading these numbers, however, is to look at simulated

projections of how cases may unfold over future months (Figures 1, 2). In both

countries, when individual variation is considered, we foresee HIT being achieved

and the COVID-19 epidemic being mostly resolved by the end of 2020. Under the

homogeneous approximation, however, epidemic potential appears to justify

considerations of second lockdowns to flatten epidemic curves. Determining the level

of confidence on either scenario is highly relevant to policy decisions and hence of

central importance to public health. According to the Akaike information criterion

(Extended Data Table 5) data favours the models that account for individual variation

and a more natural resolution for the pandemic. Models that account for individual

variation in susceptibility score particularly well and amongst those that account for

variation in exposure those that allow this variation to reduce with social distancing

perform better.

Looking back, we conclude that NPIs had a crucial role in halting the growth of the

initial wave between February and April irrespective of individual variation. Although

the most extreme lockdown strategies may not be sustainable for longer than a month

or two, they proved effective at preventing overshoot, keeping cases within health

system capacities, and may have done so without impairing the development of herd

immunity.

1. Ferguson, N. M., et al. Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality and healthcare demand (Imperial College COVID-19 Response Team, 2020). 10.25561/77482.

2. Kissler, S. M., Tedijanto, C., Goldstein, E., Grad, Y. H. & Lipsitch, M. Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period. *Science* **368**, 860-868 (2020).

3.   Wu, J. T., Leung, K. & Leung, G. M. Nowcasting are forecasting the potential domestic

and international spread of the 2019-nCov outbreak originating in Wuhan, China: a

modelling study. *Lancet* **395**, 689-697 (2020).

4.   Kwok, K. O., Lai, F., Wei, W. I., Wong, S. Y. S. & Tang, J. Herd immunity – estimating

the level required to halt the COVID-19 epidemics in affected countries. *J. Infect.* **80**,

e32-e33 (2020).

5.   Diekmann, O., Heesterbeek, J. A. P. & Metz, J. A. J. On the definition and computation

of the basic reproduction ratio R_0 in models for infectious diseases in heterogeneous

populations. *J. Math. Biol.* **28**, 365-382 (1990).

6.   Pastor-Satorras, R. & Vespignani, A. Epidemic dynamics and endemic states in

complex networks. *Phys. Rev. E* **63**, 066117 (2001).

7.   Jarvis, C. I., et al. Quantifying the impact of physical distance measures on the

transmission of COVID-19 in the UK. *BMC Medicine* **18**, 206 (2020).

8.   Montalbán, A., Corder, R. M. & Gomes, M. G. M. Herd immunity under

individual variation and reinfection. *arXiv* 200800098v2. (2020).

9.   Gomes, M. G. M., et al. Individual variation in susceptibility or exposure to SARS-CoV-

2 lowers the herd immunity threshold. *medRvix* 10.1101/2020.04.27.20081893 (2020).

10. Gomes, M. G. M., White, L. J. & Medley, G. F. Infection, reinfection, and vaccination

under suboptimal immune protection: Epidemiological perspectives. *J. Theor. Biol.* **228**,

539-549 (2004).

11. Gomes, M. G. M., Gjini, E., Lopes, J. S., Souto-Maior, C. & Rebelo, C. A theoretical

framework to identify invariant thresholds in infectious disease epidemiology. *J. Theor.*

Biol. **395**, 97-102 (2016).

12. Pollán, M., et al. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide,

population-based seroepidemiological study. *Lancet* 10.1016/s0140-6736(20)31483-5

(2020).

13. Gonçalves, G. Herd immunity: recent uses in vaccine assessment. *Expert Rev. Vaccines*

7, 1493-1506 (2008).

14. Fine, P., Eames, K. & Heymann, D. L. "Herd immunity": a rough guide, *Clin. Infect.*

Dis. **52**, 911-916 (2011).

15. Ferrari, M. J., Bansal, S., Meyers, L. A. & Bjornstad, O. N. Network frailty and the

geometry of herd immunity. *Proc. R. Soc. B* **273**, 2743-2748 (2006).

16. Folegatti, P. M., et al. Safety and immunogenicity of the ChAdOx1 nCoV-19 vaccine

against SARS-CoV-2: a preliminary report of a phase 1/2, single-blind, randomised

controlled trial. *Lancet* 10.1016/S0140-6736(20)31604-4 (2020).

17. Zhu, F.-C., et al. Immunogenicity and safety of a recombinant adenovirus type-5-

vectored COVID-19 vaccine in healthy adults aged 18 years or older: a randomised,

double-blind, placebo-controlled, phase 2 trial. *Lancet* 10.1016/S0140-6736(20)31611-1

(2020).

18. Mossong, J., et al. Social contacts and mixing patterns relevant to the spread of

infectious diseases. *PLOS Med*. **5**, e74 (2008).

19. Prem, K., Cook, A. R. & Jit, M. Projecting social contact matrices in 152 countries using

contact surveys and demographic data. *PLOS Comput. Biol.* **13**, e1005697 (2017).

20. Zhang, J., et al. Changes in contact patterns shape the dynamics of the COVID-19

outbreak in China. *Science* **368**, 1481-1486 (2020).

21. Britton, T., Ball, F. & Trapman, P. A mathematical model reveals the influence of

population heterogeneity on herd immunity to SARS-CoV-2. *Science*

10.1126/science.abc6810 (2020).

22. Adam, D., et al. Clustering and superspreading potential of severe acute respiratory

syndrome coronavirus 2 (SARS-CoV-2) infections in Hong Kong. 10.21203/rs.3.rs-

29548/v1

23. Eubank, S., et al. Modelling disease outbreaks in realistic urban social networks. *Nature*

429, 180-184 (2004).

355 24. Grifoni, A., et al. Targets of T cell responses to SARS-CoV-2 coronavirus in humans

356   with COVID-19 disease and unexposed individuals. *Cell* **181**, 1489-1501.e15 (2020).

357 25. Le Bert, N., et al. SARS-CoV-2-specific T cell immunity in cases of COVID-19 and

358   SARS, and uninfected controls. *Nature* 10.1038/s41586-020-2550-z (2020).

359 26. Flaxman, S., et al. Estimating the effects of non-pharmaceutical interventions on

360   COVID-19 in Europe. *Nature* 10.1038/s41586-020-2405-7 (2020).

Fig. 1| SARS-CoV-2 transmission with individual variation. Variation in
susceptibility (top panels); variation in connectivity (CV reducing in proportion to
social distancing) (bottom panels). Susceptibility or connectivity factors implemented
as gamma distributions. Suppressed wave and subsequent dynamics in Belgium and
England (orange). Estimated epidemic in the absence of interventions revealing
overshoot (black). Blue bars are daily new cases. Controlled (R_c) and effective (R_{eff})
reproduction numbers are displayed on shallow panels underneath the main plots.
Blue shades represent social distancing (intensity reflected in R_c trends and shade
density). R_c values in the dotted portion of the orange lines do not interfere with the
fittings and are only used to illustrate how the epidemic may unfold beyond the data
analysed here. Consensus parameter values (Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per
day; and $\rho = 0.5$. Fraction of infected individuals identified as positive (reporting
fraction): 0.06 (Belgium); 0.024 (England). Basic reproduction number, coefficients
of variation and social distancing parameters estimated by Bayesian inference as
described in Methods (estimates in Extended Data Tables 1 and 3). Curves represent
mean model predictions from 10^4 posterior samples. Orange shades represent 95%
credible intervals. Vertical lines represent the expected time when herd immunity
threshold will be achieved.

Fig. 2| SARS-CoV-2 transmission in homogeneous populations. Suppressed wave
and subsequent dynamics in Belgium and England (orange). Estimated epidemic in
the absence of interventions revealing overshoot (black). Blue bars are daily new
cases. Controlled (R_c) and effective (R_{eff}) reproduction numbers are displayed on
shallow panels underneath the main plots. Blue shades represent social distancing
(intensity reflected in R_c trends and shade density). R_c values in the dotted portion of
the orange lines do not interfere with the fittings and are only used to illustrate how
the epidemic may unfold beyond the data analysed here. Consensus parameter values
(Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per day; and $\rho = 0.5$. Fraction of infected
individuals identified as positive (reporting fraction): 0.06 (Belgium); 0.024
(England); 0.09 (Portugal); 0.06 (Spain). Basic reproduction number, coefficients of
variation and social distancing parameters estimated by Bayesian inference as
described in Methods (estimates in Extended Data Table 4). Curves represent mean
model predictions from 10^4 posterior samples. Orange shades represent 95% credible
intervals.

402
403

Fig. 3| Herd immunity threshold with gamma-distributed susceptibility or exposure to infection. Curves generated with the SEIR model (Equation 1-4) assuming values of R_0 estimated for the study countries (Extended Data Tables 1 and 2) assuming gamma-distributed: susceptibility (top); connectivity (bottom) with constant CV. Herd immunity thresholds (solid curves) are calculated according to the formula $1 - (1/R_0)^{1/(1+CV^2)}$ for heterogeneous susceptibility and $1 - (1/R_0)^{1/(1+2CV^2)}$ for heterogeneous connectivity. Final sizes of the corresponding unmitigated epidemics are also shown (dashed).

METHODS

Model structure and underlying assumptions

The model presented here is a differential equation SEIR model, where susceptible individuals become exposed at a rate that depends on their susceptibility, the number of potentially infectious contacts they engage in, and the total number of infectious people in the population per time unit. Upon exposure, individuals enter an asymptomatic incubation phase, during which they slowly become infectious[27-30]. Thus, infectivity of exposed individuals is made to be $1/2$ of that of infectious ones ($\rho = 0.5$). After a few days, individuals develop symptoms – on average 4 days after the exposure to the virus ($\delta = 1/4$) – and thus become fully infectious[31-33]. They recover, i.e., they are no longer infectious 4 days after that ($\gamma = 1/4$), on average[34].

Efficacy of acquired immunity

We conducted the core of our analysis under the assumption that no reinfection occurs after recovery due to acquired immunity ($\sigma = 0$). To analyse the sensitivity of these results to leakage in immune response ($\sigma > 0$) we calculated herd immunity thresholds (HIT) as a function of coefficients of variation (CV) for different values of σ. The results displayed in Extended Data Figure 3 confirm the expectation that as the efficacy of acquired immunity decreases (σ increases) larger percentages of the population are infected before herd immunity is reached. Less intuitive is that there is an upper bound for how much it is reasonable to increase σ before the system enters a qualitatively different regime – the reinfection threshold[10,11] ($\sigma = 1/R_0$) – above which infection becomes stably endemic and the notion of herd immunity threshold

no longer applies. Respiratory viruses are typically associated with epidemics

dynamics below the reinfection threshold.

Effective reproduction number

The effective reproduction number (R_{eff}, also denoted by R_e or R_t by other authors)

is a time-dependent quantity which we calculate as the incidence of new infections

divided by the total number of active infections (affected by ρ for individuals in E)

multiplied by the average duration of infection (also affected by ρ for individuals in

E)

$$R_{eff}(t) = \frac{\int \lambda(x,t)x[S(x,t) + \sigma R(x,t)]\,dx}{\int \rho E(x,t) + I(x,t)\,dx}\left(\frac{\rho}{\delta} + \frac{1}{\gamma}\right). \tag{9}$$

Assortative mixing

In the main text we assumed random mixing among individuals, but human

connectivity patterns are assortative due societal structures and human behaviours. To

explore the sensitivity of our results to deviations from random mixing, we develop

an extended formalism that allows individuals to connect preferentially with those

with similar connectivity, formally $\lambda(x) =$

$(\beta/N)(\int y\,h(y-x)[\rho E(y) + I(y)]\,dy/\int yg(y)\,dy)$, where $h(y-x)$ is a normal

distribution on the difference between connectivity factors (Extended Data Figure 2).

Non-pharmaceutical interventions

We implemented non-pharmaceutical interventions (NPI) as a gradual decrease in

viral transmissibility in the population and thus a lowering of the controlled and

effective reproduction numbers (R_c and R_{eff}). Once containment measures are put in

place in each country, we postulate it takes 21 days until the maximum effectiveness

of social distancing measures is reached, approximately linearly. In the simulations

presented throughout we have held this condition (maximum "lockdown" efficacy)

for T_{max} days, where T_{max} is estimated for each study country in countrywide

analyses and fixed at 30 in regional analyses. Eventually, lockdowns were lifted and

social distancing measures progressively relaxed. This relaxation is implemented

linearly over a period of T_{lift} days, which is also estimated. We note however that the

estimation of epidemiological parameters is not affected by the shape assumed for the

relaxation of interventions beyond the date of the last data point (June 11[th] in Belgium

and July 1[st] in England). In particular, assuming that social distancing measures

remained constant from these dates onwards, for example, would make no difference

to the estimated parameters.

Bayesian Inference

The model laid out above is amenable to theoretical exploration as presented in the

main manuscript and provides a perfect framework for inference. Fundamentally, to

be able to reproduce the inception of any epidemic, we would need to estimate when

local transmission started to occur (t_0), and the pace at which individuals infected

each other in the very early stages of the epidemic (R_0). All countries, to different

extents and at different timepoints of the epidemic, enforced some combination of

social distancing measures. To fully understand the interplay between herd immunity

and the impact of NPIs, we then set out to estimate the time at which social distancing

measures started to have an impact on daily incidence (t_0^d), what their maximum

efficacy (d_{max}) was and how long it lasted (T_{max}), a parameter used to determine the

rate at which contacts resume as restrictions are lifted (T_{lift}, which more specifically

represents the time it would take to reach pre-pandemic contact intensity should the

lifting of restricting have linear effect), the basic reproduction number (R_0) and what

the underlying variance in heterogeneity is for susceptibility or exposure to infection.

In order to preserve identifiability, we made two simplifying assumptions: (*i*) the

fraction of infectious individuals reported as COVID-19 cases (reporting fraction) is

constant throughout the study period and is comparable between countries

proportionally to the number of tests performed per person; (*ii*) local transmission

starts (t_0) when countries/regions report 1 case per 5 million population in one day.

To calculate the reporting rates, we used the Spanish national serological survey[12] as a

reference and divided the total number of reported cases up to May 11[th] by the

estimated number of people that had been exposed to the virus. This gives us a

reporting rate for Spain around 6%. Unfortunately, there are no other national

serological surveys that could inform the proportion of the population infected in

other countries, so we had to extrapolate the reporting rate for those. Assuming the

reporting rate is highly dependent on the testing effort employed in each country,

reflected in the number of tests per individual, we estimate the reporting rate by

scaling the reporting rate recorded in Spain according to the ratio of PCR tests per

person in other countries relative to the Spanish reference of 0.9 tests per thousand

people (https://ourworldindata.org/coronavirus-testing). This produced estimated case

reporting rates (ratio of reported cases to infections) of 9% for Portugal, 6% for

Belgium (and Spain) and 2.4% for England.

Whist national case and mortality data is easily available for most countries, more

spatially resolute data is difficult to find in the public domain. Thus, we restricted our

analysis to countries for which disaggregated regional case data was easily available.

We collected the data at two time points. First, we compiled all available data from

the day the countries started reporting COVID-19 cases to the initial collection date

(May 20th) and later collated available data from May 21st to July 10th.

Parameter estimation was performed with the software MATLAB, using PESTO

(Parameter EStimation Toolbox)[35], and assuming the reported case data can be

accurately described by a Poisson process. We first fixed the beginning of local

transmission (parameter t_0) in each data series as the day in which reported cases

surpassed 1 in 5 million individuals. Next, we optimized the model for the set of

parameters $\theta = \{R_0, CV, t_0^d, d_{max}, T_{max}, T_{lift}\}$ by maximizing the logarithm of the

likelihood (LL) (Equation 10) of observing the daily reported number of cases in each

country $D = \{(k, \tilde{y}_k)\}_{k=0}^n$:

$$LL(\theta|D) = -\sum_{k=1}^n y(k, \theta) + \left(\sum_{k=1}^n \tilde{y}(k) \ln (y(k, \theta))\right) - ln\left(\prod_{k=1}^n \tilde{y}(k)!\right) \quad (10)$$

in which $y(k, \theta)$ is the simulated model output number of COVID-19 cases at day k

(with respect to t_0), and n is the total number of days included in the analysis for each

country.

To ensure that the estimated maximum is a global maximum, we performed 50 multi-

starts optimizations, and selected the combination of parameters resulting in the

maximal Loglikelihood as a starting point for 10^4 Markov Chain Monte-Carlo

iterations. From the resulting posterior distributions, we extract the median estimates

for each parameter and the respective 95% credible intervals for the set of parameters.

We used uniformly distributed priors with ranges {1-9, 0.0025-8,1-60, 0-0.7,1-90,60-

1000}.

When fitting the model to disaggregated data, we follow the procedure outlined above

but fix $T_{max} = 30$ days, $T_{lift} = 120$ days and estimate region-specific R_0 and CV,

with common t_0^d and d_{max}.

The countrywide fitting procedure was applied to 2 countries (Belgium and England)

and repeated for each of the 4 model variants considered here (homogeneous,

heterogeneous susceptibility, heterogeneous connectivity with constant CV, and

heterogeneous connectivity with CV reducing in proportion to social distancing). Our

study included two more countries (Portugal and Spain), where countrywide analyses

could not be performed due to the epidemic being geographically desynchronised. We

implemented stratified analyses on all 4 countries. In the fitting procedures using sub-

national data, we assumed regions had the same start date for interventions that

mitigate transmission (t_0^d), and that these measures produced the same maximum

impact on transmission (d_{max}) everywhere. Thus, the only region-specific parameters

to be estimated are $\{R_{0_i}, CV_i\}$. Parameter estimates obtained from each of the model

variants are displayed in Extended Data Table 1 (heterogeneity in susceptibility),

Extended Data Table 2 (heterogeneity in connectivity with constant CV), Extended

Data Table 3 (heterogeneity in connectivity with dynamic CV) and Extended Data

Table 4 (homogeneous model), are comparable to those obtained in other studies[7,26,37-

40]. Finally, we apply the Akaike information criterion (AIC) for each estimation

procedure to inform on the quality of each model's fit to the datasets of reported cases

(Extended Data Table 5). In all cases, heterogeneous models are preferred over the

homogeneous approximation. The three heterogeneous models are roughly equally

well supported by the data used in this study. Further research should complement

this with discriminatory data types and hybrid models to enable the integration of

different forms of individual variation.

Data and code availability

Datasets are publicly available at the respective national ministry of health websites

(41-45). Core models implemented in MATLAB available from:

https://github.com/mgmgomes1/covid

27. Wei, W. E., et al. Presymptomatic Transmission of SARS-CoV-2 — Singapore, January 23–March 16, 2020. MMWR Morb Mortal Wkly Rep [Internet]. 2020 Apr 10 [cited 2020 May 4];69(14):411–5. Available from: http://www.cdc.gov/mmwr/volumes/69/wr/mm6914e1.htm?s_cid=mm6914e1_w

28. To, K. K. W., et al. Temporal profiles of viral load in posterior oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-2: an observational cohort study. *Lancet Infect. Dis.* **20**, 565-74 (2020).

29. Arons, M. M., et al. Presymptomatic SARS-CoV-2 Infections and Transmission in a Skilled Nursing Facility. *N. Engl. J. Med.* **382**, 2081-2090 (2020).

30. He, X., et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat. Med.* **26**, 672-675 (2020).

31. Lauer, S. A., et al. The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Ann. Intern. Med.* **172**, 577-582 (2020).

32. Li, Q., et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N. Engl. J. Med.* **382**, 1199-1207 (2020).

33. Zhang, J., et al. Evolving epidemiology and transmission dynamics of coronavirus disease 2019 outside Hubei province, China: A descriptive and modelling study. *Lancet Infect. Dis.* **20**, 793-802 (2020).

34. Nishiura, H., Linton, N. M. & Akhmetzhanov, A. R. Serial interval of novel coronavirus (COVID-19) infections. *Int. J. Infect. Dis.* **93**, 284-6 (2020).

35. Stapor, P., et al. PESTO: Parameter EStimation TOolbox. *Bioinformatics* **34**, 705-707
(2018).

36. Prem, K., et al. The effect of control strategies to reduce social mixing on outcomes of
COVID-19 epidemic in Wuhan, China: a modelling study. *Lancet Public Health* **5**, e261-
e270.

37. Tian, H., et al. An investigation of transmission control measures during the first 50 days
of the COVID-19 epidemic in China. *Science* **368**, 638-642.

38. Kucharski, A. J., et al. Effectiveness of isolation, testing, contact tracing and physical
distancing on reducing transmission of SARS-CoV-2 in different settings: a
mathematical modelling study. *Lancet Infect. Dis.* 10.1016/s1473-3099(20)30457-6
(2020).

39. Salje, H., et al. Estimating the burden of SARS-CoV-2 in France. *Science* **369**, 208-211
(2020).

40. Di Domenico, L., Pullano, G., Sabbatini, C. E., Boelle, P.-Y. & Colizza, V. Expected
impact of lockdown in Île-de-France and possible exit strategies. *medRxiv*
10.1101/2020.04.13.20063933 (2020).

41. https://ourworldindata.org/coronavirus-testing#source-information-country-by-
country. Accessed on July 10[th] 2020.

42. https://cnecovid.isciii.es/covid19.

43. https://covid-19.sciensano.be/nl/covid-19-epidemiologische-situatie.

44. https://covid19.min-saude.pt/ponto-de-situacao-atual-em-portugal.

45. https://coronavirus.data.gov.uk.

Acknowledgements

We thank Jan Hasenauer and Antonio Montalbán for helpful discussions concerning
statistical inference and mathematics, respectively. R.M.C. and M.U.F. receive

Author contributions

606 M.G.M.G. conceived the study. R.A. and R.M.C. and M.G.M.G. performed the

607 analyses. All authors interpreted the data and wrote the paper.

Competing interests

610 The authors declare no competing interests.

Extended Data Fig. 1| SARS-CoV-2 transmission with individual variation in connectivity. Variation in connectivity (constant CV). Connectivity factors implemented as gamma distributions. Suppressed wave and subsequent dynamics in Belgium and England (orange). Estimated epidemic in the absence of interventions revealing overshoot (black). Blue bars are daily new cases. Controlled (R_c) and effective (R_{eff}) reproduction numbers are displayed on shallow panels underneath the main plots. Blue shades represent social distancing (intensity reflected in R_c trends and shade density). R_c values in the dotted portion of the orange lines do not interfere with the fittings and are only used to illustrate how the epidemic may unfold beyond the data analysed here. Consensus parameter values (Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per day; and $\rho = 0.5$. Fraction of infected individuals identified as positive (reporting fraction): 0.06 (Belgium); 0.024 (England). Basic reproduction number, coefficients of variation and social distancing parameters estimated by Bayesian inference as described in Methods (estimates in Extended Data Table 2). Curves represent mean model predictions from 10^4 posterior samples. Orange shades represent 95% credible intervals. Vertical lines represent the expected time when herd immunity threshold will be achieved.

Extended Data Fig. 2| Connectivity distributions with reducing coefficient of
variation in proportion to social distancing. Individual variation in connectivity is
originally implemented as a gamma distribution of mean 1 parameterised by the
coefficient of variation (CV) (black). Social distancing is initially implemented as a
reduction in connectivity by the same factor to every individual, from x to $(1 - d)x$
(top panels). A more general formulation where CV may reduce with social
distancing is implemented by modifying x to $(1 - d)[1 + (1 - d)(x - 1)]$ (bottom
panels).
642

Extended Data Fig. 3| Herd immunity threshold and epidemic final size with reinfection. Curves in the main panels generated with the SEIR model (Equation 1-4) assuming $R_0 = 3$ and gamma-distributed susceptibility (top) or connectivity (bottom) with constant CV. Efficacy of acquired immunity is captured by a reinfection parameter σ, potentially ranging between $\sigma = 0$ (100% efficacy) and $\sigma = 1$ (0 efficacy). This illustration depicts final sizes of unmitigated epidemics and associated HIT curves for 6 values of σ: $\sigma = 0$ (black); $\sigma = 0.1$ (green); $\sigma = 0.2$ (blue); $\sigma = 0.3$ (magenta); $\sigma = 1/3$ (red); and $\sigma = 0.4$ (orange);. Above $\sigma = 1/R_0$ (reinfection threshold (Gomes et al 2004; 2016)) the infection becomes stably endemic and there is no herd immunity threshold. Representative epidemics of the regime $\sigma \le 1/R_0$ are shown on the right while the regime $\sigma > 1/R_0$ is illustrated on top. All depicted dynamics are based on the rightmost CVs represented on the main panel.

Extended Data Fig. 4| Herd immunity threshold and epidemic final size with
gamma-distributed exposure to infection and assortative mixing. Curves in
central panel generated with the SEIR model (Equation 1-4) assuming $R_0 = 3$ and
gamma-distributed connectivity. Assortative mixing is implemented by imposing a
normal distribution for contact preferences such that individuals contact preferentially
with those with the similar contact degree (left). This illustration used normal
distributions with standard deviation $SD = 50$ (green); $SD = 10$ (blue); and $SD = 2$
(magenta). More assortative mixing leads to more skewed epidemics. Herd immunity
thresholds were calculated numerically as the percentage of the population no longer
susceptible when new outbreaks are effectively prevented (approximately when the
exposed fraction crosses the peak in the absence of mitigation). Final sizes of the
corresponding unmitigated epidemics are also shown. Representative epidemics are
depicted on the right based on the rightmost CVs represented on the main panel (with
vertical lines marking the point when herd immunity is achieved).
675

Extended Data Fig. 5| SARS-CoV-2 transmission at subnational levels in Belgium.
Suppressed wave and subsequent dynamics in Flanders and the rest of Belgium, with
individual variation in susceptibility (left) or exposure (right). Blue bars are daily new
cases. Shades represent social distancing (intensity reflected in shade density).
Susceptibility or exposure factors implemented as gamma distributions. Consensus
parameter values (Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per day; and $\rho = 0.5$.
Fraction of infected individuals identified as positive (reporting fraction): 0.06. Basic
reproduction number, coefficients of variation and social distancing parameters
estimated by Bayesian inference as described in Methods (estimates in Extended Data
Table 1 and 2). Curves represent mean model predictions from 10^4 posterior samples.
Orange shades represent 95% credible intervals. Red curves represent cumulative
infected percentages.
689

Extended Data Fig. 6| SARS-CoV-2 transmission at subnational levels in England.
Suppressed wave and subsequent dynamics in London, Northwest, Southeast and the
rest of England, with individual variation in susceptibility (left) or exposure (right).
Blue bars are daily new cases. Shades represent social distancing (intensity reflected
in shade density). Susceptibility or exposure factors implemented as gamma
distributions. Consensus parameter values (Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per
day; and $\rho = 0.5$. Fraction of infected individuals identified as positive (reporting
fraction): 0.024. Basic reproduction number, coefficients of variation and social
distancing parameters estimated by Bayesian inference as described in Methods
(estimates in Extended Data Table 1 and 2). Curves represent mean model predictions
from 10^4 posterior samples. Orange shades represent 95% credible intervals. Red
curves represent cumulative infected percentages.

Extended Data Fig. 7| SARS-CoV-2 transmission at subnational levels in Portugal.
Suppressed wave and subsequent dynamics in the North and Centre regions versus the
rest of Portugal, with individual variation in susceptibility (left) or exposure (right).
Blue bars are daily new cases. Shades represent social distancing (intensity reflected
in shade density). Susceptibility or exposure factors implemented as gamma
distributions. Consensus parameter values (Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per
day; and $\rho = 0.5$. Fraction of infected individuals identified as positive (reporting
fraction): 0.09. Basic reproduction number, coefficients of variation and social
distancing parameters estimated by Bayesian inference as described in Methods
(estimates in Extended Data Table 1 and 2). Curves represent mean model predictions
from 10^4 posterior samples. Orange shades represent 95% credible intervals. Red
curves represent cumulative infected percentages.

Extended Data Fig. 8| SARS-CoV-2 transmission at subnational levels in Spain.
Suppressed wave and subsequent dynamics in Madrid, Catalunya and the rest of
Spain, with individual variation in susceptibility (left) or exposure (right). Blue bars
are daily new cases. Shades represent social distancing (intensity reflected in shade
density). Susceptibility or exposure factors implemented as gamma distributions.
Consensus parameter values (Methods): $\delta = 1/4$ per day; $\gamma = 1/4$ per day; and $\rho =$
0.5. Fraction of infected individuals identified as positive (reporting fraction): 0.06.
Basic reproduction number, coefficients of variation and social distancing parameters
estimated by Bayesian inference as described in Methods (estimates in Extended Data
Table 1 and 2). Curves represent mean model predictions from 10^4 posterior samples.
Orange shades represent 95% credible intervals. Red curves represent cumulative
infected percentages and vertical red segments mark seroprevalences (95% CI)
according to a recent study[12].

Country/Region	R_0			CV			$1 - d_{max}$			t_0^d		
Belgium	5.01	**5.03**	5.05	3.84	**3.87**	3.89	0.40	**0.40**	0.40	1.00	**1.02**	1.08
Flanders	4.96	**5.00**	5.02	3.89	**3.91**	3.93	0.41	**0.41**	0.41	1.00	**1.02**	1.15
Rest	4.97	**5.01**	5.03	3.87	**3.89**	3.91						
Portugal												
North/Centre	4.27	**4.32**	4.36	3.69	**3.72**	3.74	0.34	**0.35**	0.35	7.51	**7.73**	8.00
Rest	3.54	**3.58**	3.61	3.72	**3.76**	3.79						
Spain												
Madrid	4.38	**4.39**	4.39	2.37	**2.37**	2.38	0.37	**0.37**	0.37	16.40	**16.41**	16.44
Catalunya	4.20	**4.21**	4.21	2.49	**2.50**	2.50						
Rest	3.96	**3.97**	3.97	3.80	**3.81**	3.82						
England	2.90	**2.91**	2.92	1.54	**1.54**	1.55	0.46	**0.46**	0.46	42.03	**42.19**	42.41
London	2.95	**2.96**	2.96	2.24	**2.24**	2.24	0.52	**0.53**	0.53	41.35	**41.51**	41.64
NorthWest	3.03	**3.04**	3.05	1.66	**1.67**	1.68						
SouthEast	2.80	**2.81**	2.82	2.07	**2.07**	2.07						
Rest	2.88	**2.88**	2.89	1.64	**1.64**	1.65						

741
742
743

Country/Region	R_0			CV			$1 - d_{max}$			t_0^d		
Belgium	7.13	**7.15**	7.16	2.87	**2.87**	2.87	0.27	**0.27**	0.28	1.00	**1.00**	1.02
Flanders	7.09	**7.11**	7.13	2.86	**2.87**	2.89	0.27	**0.27**	0.28	1.00	**1.01**	1.03
Rest	7.11	**7.13**	7.15	2.86	**2.87**	2.89						
Portugal												
North/Centre	5.06	**5.08**	5.09	3.24	**3.24**	3.24	0.25	**0.25**	0.25	7.21	**7.22**	7.24
Rest	5.68	**5.69**	5.69	2.79	**2.81**	2.83						
Spain												
Madrid	7.81	**7.83**	8.82	1.98	**1.99**	2.06	0.24	**0.26**	0.26	5.38	**7.02**	7.06
Catalunya	8.00	**8.02**	9.00	2.33	**2.34**	2.43						
Rest	7.97	**7.99**	8.96	3.58	**3.59**	3.72						
England	3.81	**3.82**	3.84	1.55	**1.55**	1.55	0.42	**0.42**	0.42	36.38	**37.49**	36.54
London	3.70	**3.70**	3.71	1.69	**1.69**	1.70	0.43	**0.43**	0.43	37.50	**37.52**	37.55
NorthWest	3.83	**3.83**	3.84	1.32	**1.32**	1.32						
SouthEast	3.58	**3.59**	3.59	1.66	**1.67**	1.68						
Rest	3.60	**3.60**	3.61	1.30	**1.31**	1.31						

752
753
754
755

Country/Region	R_0			CV			$1 - d_{max}$			t_0^d		
Belgium	8.88	**8.91**	8.94	3.89	**3.92**	3.95	0.55	**0.55**	0.55	1.00	**1.00**	1.00
Flanders	8.78	**8.82**	8.86	3.98	**4.00**	4.02	0.56	**0.56**	0.56	1.00	**1.01**	1.04
Rest	8.82	**8.86**	8.89	3.98	**4.00**	4.02						
Portugal												
North/Centre	6.65	**6.72**	6.80	3.75	**3.78**	3.81	0.56	**0.57**	0.57	5.84	**6.02**	6.19
Rest	5.98	**6.05**	6.13	4.09	**4.14**	4.19						
Spain												
Madrid	6.19	**6.20**	6.21	2.43	**2.43**	2.44	0.57	**0.57**	0.57	13.80	**13.81**	13.83
Catalunya	6.30	**6.32**	6.33	2.61	**2.62**	2.62						
Rest	6.34	**6.35**	6.36	3.80	**3.81**	3.82						
England	3.67	**3.67**	3.68	1.56	**1.57**	1.57	0.63	**0.63**	0.63	39.81	**39.82**	39.83
London	3.78	**3.79**	3.80	1.99	**2.00**	2.01	0.67	**0.67**	0.67	39.06	**39.20**	39.28
NorthWest	3.91	**3.92**	3.94	1.64	**1.65**	1.66						
SouthEast	3.67	**3.69**	3.70	1.89	**1.89**	1.90						
Rest	3.64	**3.65**	3.66	1.58	**1.58**	1.59						

764
765
766

Extended Data Table 4| Estimated parameters for the homogenous model.
Estimates generated from model fit to the national datasets are in the grey shaded
rows. The remaining rows provide the region-specific estimates. Best parameter
estimates are presented as a bold median bounded by the lower and upper ends for the
95% credible interval. Model runs are initiated on the day (t_0) when reported cases
surpassed 1 in 5 million individuals: Belgium (day 1); England (day 29); Portugal
(day 3); Spain (day 8).
774

Country/Region	R_0			$1 - d_{max}$			t_0^d		
Belgium	3.683	**3.698**	3.712	0.218	**0.219**	0.220	11.948	**12.054**	12.173
Flanders	3.235	**3.239**	3.308	0.208	**0.212**	0.213	16.324	**17.039**	17.064
Rest	3.235	**3.238**	3.307						
Portugal									
North/Centre	3.542	**3.578**	3.608	0.343	**0.345**	0.348	7.514	**7.725**	7.999
Rest	4.274	**4.321**	4.361						
Spain									
Madrid	4.113	**4.116**	4.120						
Catalunya	4.208	**4.214**	4.218	0.111	**0.111**	0.112	20.000	**20.000**	20.000
Rest	3.735	**3.751**	3.752						
England	2.690	**2.703**	2.708	0.339	**0.339**	0.340	48.091	**48.155**	48.244
London	2.307	**2.308**	2.310						
NorthWest	2.602	**2.603**	2.604	0.359	**0.360**	0.360	54.577	**54.578**	54.578
SouthEast	2.368	**2.370**	2.371						
Rest	2.502	**2.503**	2.504						

775
776
777

Extended Data Table 5| Model selection criteria. Displays the maximum
Loglikelihood obtained for each combination of model and data partitioning for each
country, as well as the Akaike information criterion. Models are labelled by a sort
name as follows: homog (homogenous); hetsus (heterogeneity in susceptibility);
hetcon (heterogeneity in connectivity with constant CV); hetdyn (heterogeneity in
connectivity with dynamic CV).
784

Country	Model	LL	AIC
Countrywide Data			
England	homog	1.6765E+06	-3.3529E+06
	hetsus	1.6793E+06	-3.3585E+06
	hetcon	1.6785E+06	-3.3571E+06
	hetdyn	1.6790E+06	-3.3579E+06
Belgium	homog	3.3204E+05	-6.6407E+05
	hetsus	3.3325E+05	-6.6649E+05
	hetcon	3.3269E+05	-6.6537E+05
	hetdyn	3.3278E+05	-6.6554E+05
Regional Data			
Portugal	homog	3.36E+05	-6.73E+05
	hetsus	3.55E+05	-7.11E+05
	hetcon	3.55E+05	-7.10E+05
	hetdyn	3.55E+05	-7.11E+05
Spain	homog	1.87E+06	-3.75E+06
	hetsus	1.96E+06	-3.91E+06
	hetcon	1.96E+06	-3.91E+06
	hetdyn	1.47E+06	-2.95E+06
England	homog	2.59E+06	-5.18E+06
	hetsus	2.63E+06	-5.25E+06
	hetcon	2.62E+06	-5.25E+06
	hetdyn	2.62E+06	-5.25E+06
Belgium	homog	3.74E+05	-7.48E+05
	hetsus	3.78E+05	-7.56E+05
	hetcon	3.78E+05	-7.55E+05
	hetdyn	3.78E+05	-7.55E+05