Smoking is associated with significantly increased green autofluorescence intensity

and asymmetry of the skin and the fingernails of natural populations, population at-

risk of developing stroke, and population of acute ischemic stroke

Mingchao Zhang[1,2,*], Yue Tao[1,2,*], Danhong Wu[3] , Weihai Ying[1,2,#]

[1]Med-X Research Institute and School of Biomedical Engineering, Shanghai Jiao

Tong University, Shanghai 200030, P.R. China; [2]Collaborative Innovation Center for

Genetics and Development, Shanghai 200043, P.R. China; [3]Department of Neurology,

Shanghai Fifth People's Hospital, Fudan University, Shanghai, P.R. China

* These two authors contributed equally to this work.

#: Corresponding author

Weihai Ying, Ph.D.

Professor, School of Biomedical Engineering and Med-X Research Institute

Shanghai Jiao Tong University

1954 Huashan Road

Shanghai, 200030, P.R. China

E-mail: weihaiy@sjtu.edu.cn

Abstract

Tobacco smoking is an important risk factor for multiple major diseases including cancer and stroke. It is of significance to search for the biomarkers of smoking for non-invasive, rapid and economic monitoring of the pathological changes of smokers' body. However, there has been no non-invasive method for monitoring the health state of smokers. Our current study has indicated that green autofluorescence (AF) of the fingernails and certain locations of the skin is a novel biomarker for smoking: Both the green AF intensity and the AF asymmetry of the fingernails and the skin of Dorsal Index Fingers of the smokers were significantly higher than those of the non-smokers in natural populations, the population at-risk of developing stroke and the acute ischemic stroke (AIS) patients. ROC analyses using the green AF intensity of either right or left Index Fingernails showed that the AUC values ranged from 0.8136 and 0.8644 for differentiating smokers and non-smokers in these populations. ROC analyses using the green AF intensity of the skin of the right Dorsal Index Fingers showed that the AUC values ranged between 0.7418 and 0.7478 for differentiating the smokers and the non-smokers in natural populations and the population at-risk of developing stroke. Collectively, our study has indicated that increased green AF intensity of the fingernails and certain locations of the skin is a novel biomarker for smoking. Based on this finding, pathological alterations of smokers' body may be monitored non-invasively, rapidly and economically, which could be highly valuable for the health management of the large population of tobacco smokers.

Keywords: Smoking; Biomarker; Autofluorescence; Fingernails; Skin

Introduction

Tobacco smoking is an important risk factor for multiple major diseases including cancer, AIS and cardiovascular diseases (6-8). In many Western countries smoking is responsible for one third of all cancer deaths {Sasco, 2004 #29692}. It is of significance to search for the biomarker of smoking for non-invasive and rapid monitoring of the pathological changes of smokers' body, which is highly valuable for the health monitoring of a large population of smokers. However, there has been no non-invasive method for monitoring the pathological changes of large population of smokers.

Human autofluorescence (AF) has been used for non-invasive diagnosis of diabetes and diabetes-related pathology [8]. The AF originates from the advanced glycation end-product (AGE)-modified collagen and elastin of dermis. NADH, FAD, keratins and melanin are major epidermal fluorophores [9,10]. It is established that such molecules as NADH and FAD can be significantly affected by oxidative stress.

Since tobacco smoking can induce oxidative stress in the body (11, 12), we hypothesized that tobacco smoking-induced changes of the AF of the skin may become a novel biomarker for tobacco smoking. In this study we tested this hypothesis, showing that smokers had significantly increased green AF intensity and asymmetry of the fingernails and certain locations of the skin is in natural populations, the population at-risk of developing stroke and the population of acute ischemic stroke (AIS).

Methods and Materials

Studies on Human Subjects

The study was conducted according to a protocol approved by the Ethics Committee of Shanghai Fifth People's Hospital affiliated to Fudan University. The human subjects in our study were divided into four groups: Group 1: The natural populations; Group 2: The healthy population; Group 3: The population at-risk of developing stroke; and Group 4: The population of the AIS patients who were hospitalized in the Department of Neurology, Shanghai Fifth People's Hospital affiliated to Fudan University. The age of Group 1, Group 2, Group 3 and Group 4 was 67.27±8.25, 60.94±12.98, 68.09±8.06, and 64.11±8.20 years of old, respectively.

Determinations of the AF of human subjects' skin and fingernails

A portable AF imaging equipment was used to determine the skin' AF of the human subjects. The excitation wavelength was 485 nm, and the emission wavelength was 500 - 550 nm.

Statistical analyses

All data are presented as mean $\pm$ SEM. Data were assessed by one-way ANOVA, followed by Student - Newman - Keuls *post hoc* test, except where noted. *P* values less than 0.05 were considered statistically significant.

Results

1) Both the green AF intensity and the AF asymmetry of the fingernails and the skin of Dorsal Index Fingers of the smokers in natural populations were significantly higher than those of the non-smokers

We determined the green AF intensity of the fingernails and twelve positions of the skin of natural populations. Compared with the green AF intensity of the fingernails and the skin of Dorsal Index Fingers of the non-smokers, the green AF intensity of both the fingernails and the skin of Dorsal Index Fingers of the smokers was significantly higher (Figs. 1A and 1B). In contrast, no significant difference was observed between the green AF intensity of the non-smokers and that of the smokers at other examined positions of the skin (Supplement Figs. 1A, 1B, 1C, 1D, and 1E).

Compared with the AF asymmetry of both the fingernails and the skin of Dorsal Index Fingers of the non-smokers, the AF asymmetry of both the fingernails and the skin of Dorsal Index Fingers of the smokers was also remarkably higher (Figs. 2A and 2B). In contrast, no significant difference was observed between the AF asymmetry of the non-smokers and that of the smokers at other examined positions of the skin (Supplement Figs. 2A, 2B, 2C, 2D and 2E).

2) Both the green AF intensity and the AF asymmetry of the fingernails and the skin of Dorsal Index Fingers of the smokers in the population at-risk of developing stroke were significantly higher than those of the non-smokers

We also determined the green AF intensity of the fingernails and twelve positions of the skin of the population at-risk of developing stroke. Compared with the green

AF intensity of the fingernails and the skin of Dorsal Index Fingers of the non-smokers, the green AF intensity of both the fingernails and the skin of Dorsal Index Fingers of the smokers was significantly higher (Figs. 3A and 3B). In contrast, no significant difference was observed between the green AF intensity of the non-smokers and that of the smokers at other examined positions of the skin (Supplement Figs. 3A, 3B, 3C, 3D, and 3E).

Compared with the AF asymmetry of the fingernails and the skin of Dorsal Index Fingers of the non-smokers, the AF asymmetry of both the fingernails and the skin of Dorsal Index Fingers of the smokers was also significantly higher (Figs. 4A and 4B). In contrast, no significant difference was observed between the AF asymmetry of the non-smokers and that of the smokers at other examined positions of the skin (Supplement Figs. 4A, 4B, 4C, 4D, and 4E).

3) Both the green AF intensity and the AF asymmetry of the fingernails and certain positions of the skin of the smokers in AIS population were significantly higher than those of the non-smokers

We determined the green AF intensity of the fingernails and twelve positions of the skin of the AIS patients. In nearly all of the positions examined, the AF intensity of both the non-smokers and the smokers in the AIS population was significantly higher than that of the healthy population (Figs. 5A, 5B, 5C and 5D; Supplement Figs. 5A, 5B and 5C). Compared with the green AF intensity of the non-smokers in their fingernails and the skin of Dorsal Index Fingers, Vetroforefingers and the left Centrametacarpus, the green AF intensity at these positions of the smokers was significantly higher (Figs. 5A, 5B, 5C and 5D). In contrast, no significant difference

was observed between the green AF intensity of the non-smokers and that of the smokers at other examined positions of the skin (Supplement Figs. 5A, 5B and 5C).

Compared with the AF asymmetry of the non-smokers in their fingernails and the skin of Centremetacarpus, the AF asymmetry of the smokers in these positions was also remarkably higher (Figs. 6A and 6B). In contrast, no significant difference was observed between the AF asymmetry of the non-smokers and that of the smokers at other examined positions of the skin (Supplement Figs. 6A, 6B, 6C, 6D, and 6E).

4) The green AF intensity of Index Fingernails and the skin of Dorsal Index Fingers holds great potential for differentiating smokers and non-smokers in natural populations, the population at-risk of developing stroke, and AIS patients

ROC analyses using the green AF intensity of either right or left Index Fingernails showed that the AUC was 0.8644 and 0.8266, respectively, for differentiating the smokers and the non-smokers in the natural populations (Fig. 7A). ROC analyses using the green AF intensity of the skin of either right or left Dorsal Index Fingers showed that the AUC was 0.7418 and 0.6495, respectively, for differentiating the smokers and the non-smokers in the natural populations (Fig. 7B).

ROC analyses using the green AF intensity of either right or left Index Fingernails showed that the AUC was 0.8608 and 0.8312, respectively, for differentiating the smokers and the non-smokers in the population at-risk of developing stroke (Fig. 7C). ROC analyses using the green AF intensity of the skin of either right or left Dorsal Index Fingers showed that the AUC was 0.7478 and

0.6437, respectively, for differentiating the smokers and the non-smokers in the population at-risk of developing stroke (Fig. 7D).

ROC analyses using the green AF intensity of either right or left Index Fingernails showed that the AUC was 0.8136 and 0.8631, respectively, for differentiating the smokers and the non-smokers in the AIS population (Fig. 7E). ROC analyses using the green AF intensity of the skin of either right or left Dorsal Index Fingers showed that the AUC was 0.6253 and 0.6273, respectively, for differentiating the smokers and the non-smokers in the AIS population (Fig. 7F).

Discussion

The major findings of our current study include: First, both the green AF intensity and the AF asymmetry of the fingernails and the skin of Dorsal Index Fingers of the smokers were significantly higher than those of the non-smokers in natural populations. Second, both the green AF intensity and the AF asymmetry of the fingernails and the skin of Dorsal Index Fingers of the smokers were significantly higher than those of the non-smokers in the population at-risk of developing stroke. Third, in the AIS population, the green AF intensity of the fingernails and the skin of Dorsal Index Fingers, Ventroforefingers and Centremetacarpus of the smokers was significantly higher than that of the non-smokers; and the asymmetry of the fingernails and the skin of Centremetacarpus of the smokers was significantly higher than that of the non-smokers. Fourth, ROC analyses using the green AF intensity of either right or left Index Fingernails showed that the AUC values ranged from 0.8136 and 0.8644 for differentiating the smokers and the non-smokers in natural populations, the population at-risk of developing stroke, and AIS patients. Fifth, ROC analyses

using the green AF intensity of the skin of the right Dorsal Index Fingers showed that the AUC values ranged between 0.7418 and 0.7478 for differentiating the smokers and the non-smokers in natural populations and the population at-risk of developing stroke. Collectively, our study has indicated that increased green AF intensity of the fingernails and certain locations of the skin is a novel biomarker for smoking.

Tobacco smoking is a major risk factor of multiple major diseases (6-8). It is of critical importance to search for the biomarkers of smoking for non-invasive and rapid monitoring of the pathological changes of smokers' body. The search is highly valuable for the health monitoring of a large population of smokers, which is a crucial part of health management of the general population. However, there has been no non-invasive method for monitoring the health state of smokers.

Our current study has provided evidence indicating that the green AF intensity of the Index Fingernails and the skin of certain locations such as Dorsal Index Fingers is a novel biomarker for smokers of natural populations, the population at-risk of developing stroke, and AIS patients: ROC analyses using the green AF intensity of these positions showed highly promising AUC values for differentiating the smokers and the non-smokers in these populations. Based on this finding, approaches for non-invasive, economic and efficient monitoring the heathy state of the smokers may be established, which may generate tremendous health benefits. With future studies on the AF of the skin and fingernails of smokers and non-smokers as well as applications of AI technology, it is expected that increasingly precise evaluations on the smokers' health status could be established.

Our previous studies have suggested that the 'Pattern of AF' is a novel biomarker that can be used for non-invasive diagnosis of several major diseases,

which forms a basis for establishing a new biomedical imaging technology (1-3). In order to improve this new technology, it is critical to further investigate the factors that can influence the AF in natural populations, populations at-risk of developing major diseases as well as patient populations. Our current study has indicated that smoking is an important factor that is associated with increased AF intensity and AF asymmetry of the fingernails and certain positions of the skin in these populations. These findings are highly valuable for understanding the mechanisms underlying the changes of the AF in these populations, which is required for establishing the AF-based models for disease diagnosis and evaluations of the health conditions of natural populations.

Our previous study has shown that both inflammation and oxidative stress can dose-dependently increase the epidermal green AF (1). It is established that smoking can cause increased inflammation and oxidative stress in human body (9-12). Therefore, we propose that the capacity of smoking to increase the AF in these populations may result from the capacity of smoking to increase inflammation and oxidative stress in the body of these populations.

It is noteworthy that in these populations, smoking was selectively associated with increased AF intensity of the fingernails and the skin of Dorsal Index Fingers. Since it takes a significant duration of time for the growth of nails, we propose that increased AF of the nails is a biomarker for the levels of chronic exposures of the nails to inflammation and oxidative stress in the body. Based on this proposal, it is reasonable to observe the increased AF in the fingernails of the people in these populations. Future studies are required to investigate the mechanisms underlying the AF of the nails.

Based on the observations of our previous studies, we have proposed that the positions with increased AF in the skin are associated with the 'channels' (or called Meridian) in traditional Chinese medicine. According to the Meridian theory, the skin of Dorsal Index Fingers is close to the Large Intestine Meridian. Therefore, our finding that smoking increases the AF of the skin of Dorsal Index Fingers has implicated that smoking may increase the inflammation and oxidative stress of the organs that are associated with the Large Intestine Meridian.

Our study has found that smoking can enhance not only the green AF intensity but also the AF asymmetry of the fingernails and certain positions of the skin in these populations. Our previous study has indicated that high levels of AF asymmetry are associated with major vascular diseases such as AIS (2). Therefore, we propose that smoking may have particularly strong capacity to increase vascular damage, which is consistent with previous reports (12).

Acknowledgment

The authors would like to acknowledge the financial support by two research grants from a Major Special Program Grant of Shanghai Municipality (Grant # 2017SHZDZX01) (to W.Y.).

References:

1.   Zhang, M., Li, Y., Maharjan, D. T., He, H., Tao, Y., Wu, D., Ying, W. (2019) Keratin-Based Epidermal Green Autofluorescence is a Common Biomarker of Organ Injury. *bioRxiv* 564112

2.   Wu, D., Zhang, M., Tao, Y., Li, Y., Zhang, S., Chen, X., Ying, W. (2018) Distinct pattern of autofluorescence of the skin and fingernails of acute ischemic stroke patients: A novel diagnostic biomarker for acute ischemic stroke. *bioRxiv* 310904

3.   Zhang, M., Tao, Y., Chang, Y., Li, Y., Chu, T., Ying, W. (2020) Selectively increased autofluorescence at certain locations of skin may become a novel diagnostic biomarker for lung cancer. *bioRxiv* 315440

4.   Wu, D., Tao, Y., Zhang, M., Li, Y., Shen, L., Li, Y., Ying, W. (2018) Selectively increased autofluorescence at fingernails and certain regions of skin: A potential novel diagnostic biomarker for Parkinson disease. *bioRxiv* 322222

5.   Qu, X., Li, Y., Tao, Y., Zhang, M., Wu, D., Guan S., Han, W., Ying, W. (2018) Distinct Patterns of the Autofluorescence of Body Surface: Potential Novel Diagnostic Biomarkers for Stable Coronary Artery Disease and Myocardial Infarction. *bioRxiv* 330985

6.   Larsson, S. C., Burgess, S., and Michaelsson, K. (2019) Smoking and stroke: A mendelian randomization study. *Ann Neurol* **86**, 468-471

7.   Markidan, J., Cole, J. W., Cronin, C. A., Merino, J. G., Phipps, M. S., Wozniak, M. A., and Kittner, S. J. (2018) Smoking and Risk of Ischemic Stroke in Young Men. *Stroke* **49**, 1276-1278

8.   Ambrose, J. A., and Barua, R. S. (2004) The pathophysiology of cigarette smoking and cardiovascular disease: an update. *J Am Coll Cardiol* **43**, 1731-1737

9.   McEvoy, J. W., Nasir, K., DeFilippis, A. P., Lima, J. A., Bluemke, D. A., Hundley, W. G., Barr, R. G., Budoff, M. J., Szklo, M., Navas-Acien, A., Polak, J. F., Blumenthal, R. S., Post, W. S., and Blaha, M. J. (2015) Relationship of cigarette smoking with inflammation and subclinical vascular disease: the Multi-Ethnic Study of Atherosclerosis. *Arterioscler Thromb Vasc Biol* **35**, 1002-1010

10.   Rom, O., Avezov, K., Aizenbud, D., and Reznick, A. Z. (2013) Cigarette smoking and inflammation revisited. *Respir Physiol Neurobiol* **187**, 5-10

11.   Dikalov, S., Itani, H., Richmond, B., Vergeade, A., Rahman, S. M. J., Boutaud, O., Blackwell, T., Massion, P. P., Harrison, D. G., and Dikalova, A. (2019) Tobacco smoking induces cardiovascular mitochondrial oxidative stress, promotes endothelial dysfunction, and enhances hypertension. *Am J Physiol Heart Circ Physiol* **316**, H639-H646

12.   Siasos, G., Tsigkou, V., Kokkou, E., Oikonomou, E., Vavuranakis, M., Vlachopoulos, C., Verveniotis, A., Limperi, M., Genimata, V., Papavassiliou, A. G., Stefanadis, C., and Tousoulis, D. (2014) Smoking and atherosclerosis: mechanisms of disease and new therapeutic approaches. *Curr Med Chem* **21**, 3936-3948

Figure Legends:

Fig. 1. The green AF intensity of the fingernails and the skin of Dorsal Index Fingers of the smokers in natural populations was significantly higher than that of the non-smokers. (A) In both right and left fingernails, the AF intensity of the smokers in natural populations was significantly higher than that of the non-smokers. (B) In the skin of both right and left Dorsal Index Fingers, the AF intensity of the smokers in natural populations was significantly higher than that of the non-smokers. The number of the non-smokers and the smokers was 280 and 47, respectively. ***, $p < 0.001$.

Fig. 2. The AF asymmetry of the fingernails and the skin of Dorsal Index Fingers of the smokers in natural populations was significantly higher than that of the non-smokers. (A) In both right and left fingernails, the AF asymmetry of the smokers in natural populations was significantly higher than that of the non-smokers. (B) In the skin of both right and left Dorsal Index Fingers, the AF asymmetry of the smokers in natural populations was significantly higher than that of the non-smokers. The number of the non-smokers and the smokers was 280 and 47, respectively. ***, $p < 0.001$.

Fig. 3. The green AF intensity of the fingernails and the skin of Dorsal Index Fingers of the smokers in the population at-risk of developing stroke was

significantly higher than that of the non-smokers. (A) In both right and left fingernails, the AF intensity of the smokers in the population at-risk of developing stroke was significantly higher than that of the non-smokers. (B) In the skin of both right and left Dorsal Index Fingers, the AF intensity of the smokers in the population at-risk of developing stroke was significantly higher than that of the non-smokers. The number of the healthy non-smokers, the non-smokers of the population at-risk of developing stroke, and the smokers of the population at-risk of developing stroke was 56,130, and 39, respectively. *, $p < 0.05$; ***, $p < 0.001$.

Fig. 4. The AF asymmetry of the fingernails and the skin of Dorsal Index Fingers of the smokers in the population at-risk of stroke was significantly higher than that of the non-smokers. (A) In both right and left fingernails, the AF asymmetry of the smokers in the population at-risk of developing stroke was significantly higher than that of the non-smokers. (B) In the skin of both right and left Dorsal Index Fingers, the AF asymmetry of the smokers in the population at-risk of developing stroke was significantly higher than that of the non-smokers. The number of the healthy non-smokers, the non-smokers in the population at-risk of developing stroke, and the smokers in the population at-risk of developing stroke was 56,130, and 39, respectively. ***, $p < 0.001$.

Fig. 5. The green AF intensity of the fingernails and the skin of Dorsal Index Fingers, Vetroforefingers and the left Centrametacarpus of the smokers in AIS population was significantly higher than that of the non-smokers. （A）The

green AF intensity of the fingernails of the smokers in AIS population was significantly higher than that of the non-smokers. （B）The green AF intensity of the skin of Dorsal Index Fingers of the smokers in AIS population was significantly higher than that of the non-smokers. (C) The green AF intensity of the skin of Vetroforefingers of the smokers in AIS population was significantly higher than that of the non-smokers. （D）The green AF intensity of the skin of left Centrametacarpus of the smokers in AIS population was significantly higher than that of the non-smokers. The number of the healthy non-smokers, the non-smokers in the AIS population, and the smokers in the AIS population was 56, 33 and 30, respectively. *, $p < 0.05$ **; $p < 0.01$ ***; $p < 0.001$.

Fig. 6. The AF asymmetry of the fingernails and the skin of Centrametacarpus of the smokers in AIS population was significantly higher than that of the non-smokers. （A）The AF asymmetry of the fingernails of the smokers in AIS population was significantly higher than that of the non-smokers. （B）The AF asymmetry of the skin of Dorsal Index Fingers of the smokers in AIS population was significantly higher than that of the non-smokers. The number of the healthy non-smokers, the non-smokers in the AIS population, and the smokers in the AIS population was 56, 33 and 30, respectively. *, $p < 0.05$ **; $p < 0.01$ ***; $p < 0.001$.

Fig. 7. The green AF intensity of Index Fingernails and the skin of Dorsal Index Fingers holds great potential for differentiating the smokers and the non-

smokers in natural populations, the population at-risk of developing stroke, and AIS patients.

(A) ROC analyses using the green AF intensity of either right or left Index Fingernails showed that the AUC was 0.8644 and 0.8266, respectively, for differentiating the smokers and the non-smokers in the natural populations. (B) ROC analyses using the green AF intensity of the skin of the either right or left Dorsal Index Fingers showed that the AUC was 0.7418 and 0.6495, respectively, for differentiating the smokers and the non-smokers in the natural populations. (C) ROC analyses using the green AF intensity of either right or left Index Fingernails showed that the AUC was 0.8608 and 0.8312, respectively, for differentiating the smokers and non-smokers in the population at-risk of developing stroke. (D) ROC analyses using the green AF intensity of the skin of the either right or left Dorsal Index Fingers showed that the AUC was 0.7478 and 0.6437, respectively, for differentiating the smokers and the non-smokers in the population at-risk of developing stroke. (E) ROC analyses using the green AF intensity of either right or left Index Fingernails showed that the AUC was 0.8136 and 0.8631, respectively, for differentiating the smokers and the non-smokers in the AIS population. (F) ROC analyses using the green AF intensity of the skin of the either right or left Dorsal Index Fingers showed that the AUC was 0.6253 and 0.6273, respectively, for differentiating the smokers and the non-smokers in the AIS population. The number of the healthy non-smokers, the non-smokers in the AIS population, and the smokers in the AIS population was 56, 33 and 30, respectively.

Legends of Supplemental Figures:

Supplemental Fig. 1. The green AF intensity of the skin at other examined positions of the smokers in natural populations was not significantly different from that of the non-smokers. The number of the non-smokers and the smokers was 280 and 47, respectively.

Supplemental Fig. 2. The AF asymmetry of the skin at other examined positions of the smokers in natural populations not significantly different from that of the non-smokers. The number of the non-smokers and the smokers was 280 and 47, respectively.

Supplemental Fig. 3. The green AF intensity of the skin of at other examined positions of the smokers in the population at-risk of stroke not significantly different from that of the non-smokers. The number of the healthy non-smokers, the non-smokers in the population at-risk of developing stroke, and the smokers in the population at-risk of developing stroke was 56,130, and 39, respectively.

Supplemental Fig. 4. The AF asymmetry of the skin at other examined positions of the smokers in the population at-risk of stroke not significantly different from that of the non-smokers. The number of the healthy non-smokers, the non-smokers of the population at-risk of developing stroke, and the smokers of the population at-risk of developing stroke was 56,130, and 39, respectively.

Supplemental Fig. 5. The green AF intensity of the skin at other examined positions of the smokers in AIS population not significantly different from that of the non-smokers. The number of the healthy non-smokers, the non-smokers in the AIS population, and the smokers in the AIS population was 56, 33 and 30, respectively.

Supplemental Fig. 6. The AF asymmetry of the skin at other examined positions of the smokers in AIS population not significantly different from that of the non-smokers. The number of the healthy non-smokers, the non-smokers in the AIS population, and the smokers in the AIS population was 56, 33 and 30, respectively.

■ Non-smokers
■ Smokers

*** *** ***

AF Intensity
(% of Non-smokers)

Right   Left

Index Fingernail