

17 cumulative mortality rate was 1.73%, the age range of diagnosed cases was
18 5days-85years old, and the male to female ratio was 1.09:1.The confirmed cases of
19 COVID-19 in Henan province were mainly imported cases from Hubei, accounting for
20 87.74%, of which the highest number was 70.50% in Zhumadian. The contact cases and
21 local cases increased in a fluctuating manner over time.**Significance:** In this paper,
22 epidemiological characteristics of COVID-19 in Henan province from the outbreak to the
23 effective control within 60 days were analyzed, and effective and distinctive prevention
24 and control measures in various cities were summarized, so as to provide a favorable
25 reference for the further formulation and implementation of epidemic prevention and
26 control and a valuable theoretical basis for effectively avoiding the second outbreak.

27 **Importance**

28 Epidemic prevention and control in China has entered the normalized this new stage, this
29 article analyze the epidemiological characteristics of COVID - 19 in henan province, and
30 summarizes the prefecture level and effective disease prevention and control means and
31 measures, the normalized private provide the theoretical reference to the prevention and
32 control work, formulate and carry out, at the same time in other countries and regions of
33 epidemic prevention and control work can also be used for reference.

34 **Key words:** Henan Province;COVID - 19;Epidemic;The prevention and control

35 **1. Introduction**

36 In December 2019, Wuhan, Hubei province, China reported a number of cases of

37 viral pneumonia of unknown cause[1], whose rapid spread has aroused great concern
38 from the whole society.The virus was confirmed to be a novel coronavirus with a unique
39 genome by sequencing the lower respiratory tract samples collected by the expert group
40 and named as "2019 Novel Corona virus" (2019-nCoV-2),designated by the International
41 Committee for the Classification of Viruses as SARS-CoV-2.The virus novel
42 SARS-CoV-2 spread throughout the world within three months, causing a pandemic and
43 posing a serious threat to human health[2].On February 11, 2020, WHO named the novel
44 coronavirus disease as COVID-19 and declared the infectious disease a public health
45 emergency[3].The COVID-19 is reported to have spread through human-to-human
46 transmission[4].

47 So far, there is a lack of immunity against the COVID-19 in humans[5].At present,
48 the known source of COVID-19 infection is novel Coronavirus infection (including
49 asymptomatic infection), which is transmitted by respiratory droplets, close contact,
50 aerosol and fecal mouth. Respiratory droplets and close contact are the main transmission
51 routes, and people are generally susceptible.Based on the known epidemiological
52 investigation, the incubation period of this disease is 1-14d, generally 3-7d.Fatigue, dry
53 cough and fever were the main symptoms of common infected persons. A small number
54 of patients were accompanied by nasal congestion, runny nose, sore throat, diarrhea and
55 other symptoms. CT examination showed chest imaging characteristics.Most of the
56 patients with severe illness developed dyspnea or/and hypoxemia 7 days after
57 onset.Critically ill patients may rapidly progress to acute respiratory distress syndrome,

58 septic shock, multiple organ failure and other symptoms. It should be noted that most of
59 the patients with severe and critical diseases had moderate to low fever or no obvious
60 fever symptoms[6-8].

61 Henan province is adjacent to Hubei province, with Xinyang, Nanyang and
62 Zhumadian as the junction. Zhengzhou, as the capital city of Henan Province, has a large
63 flow of people, so the four cities are COVID-19 high incidence areas in Henan
64 Province. On January 21, 2020 health committee issued reports from Henan province
65 Zhengzhou in Henan province found the first imported COVID - 19 patients, reported on
66 June 26, 24 when Henan cumulative COVID - 19 confirmed cases, 1276 cases, cure the
67 hospital 1254, accumulative total 22 cases died, in addition to March 11, the new case of
68 overseas (Italy) input COVID - 19 confirmed cases, confirmed cases in the province all
69 reset! In this paper, according to the vertical distribution from south to north of Henan
70 province south (Nanyang, Xinyang, Zhumadian), selected (Zhengzhou City), in the north
71 (Anyang, Puyang 6 city COVID - 19 outbreak of epidemiological investigation and
72 analysis of the prevention and control, in order to further to carry out the command of the
73 epidemic prevention and control work, prevent secondary kickback provide theoretical
74 basis for disease[9].

75 **2. Materials and methods**

76 **2.1 Data classification**

77 The data investigated in this paper include age, gender, incubation period,
78 cumulative confirmed cases, cumulative cured and discharged cases, daily new

79 cases/cured cases, number of permanent residents in Henan province and cities, gender
80 and age structure of Henan province and cities, source of confirmed cases and types of
81 confirmed cases. In this paper, the source types of cases were divided into three categories:
82 imported cases (the patient had a history of overseas travel before diagnosis); Contact
83 cases (contact with confirmed or suspected patients before diagnosis); Local case (the
84 patient has not left the city before diagnosis and has no contact history). There are four
85 types of cases: common (mild) cases (mild clinical symptoms, visible imaging/no
86 manifestations of pneumonia); Severe cases (on the basis of the common type, oxygen
87 saturation $\leq 93\%$ under resting state, partial arterial oxygen pressure PaO₂/ oxygen
88 absorption concentration FiO₂ ≤ 300 mmHg, pulmonary imaging showed significant
89 lesion progression within 24-48 hours & GT;50%, etc.); Critical cases (including
90 respiratory failure requiring mechanical ventilation, shock, and other organ failure
91 requiring ICU care); Deaths. The calculated data include: cumulative incidence of
92 COVID-19 (1/100,000) = cumulative confirmed cases of COVID-19 in a city (county
93 included)/permanent resident population in a city by the end of 2018; Cumulative cure
94 rate (%) = cumulative number of cured and discharged COVID-19 cases in a
95 city/cumulative number of confirmed COVID-19 cases in a city; Cumulative mortality
96 rate (%) = The number of COVID-19 deaths in a city/the number of COVID-19
97 confirmed cases in a city.

98 **2.2 Data Sources**

99 According to Chapter third of Henan Statistical Yearbook 2019, the number of

100 permanent residents in Henan province and the 6 cities (including counties) required in
101 this paper is determined. According to the official website of Henan Provincial Health
102 Commission, "The Latest Situation of COVID-19 in Henan Province" column, the overall
103 data of COVID-19 epidemic in Henan province were collected, namely, the number of
104 new confirmed cases/cured and discharged cases, the cumulative number of confirmed
105 cases/cured and discharged cases/death cases, and the number of severe and critical
106 cases. Age, sex, incubation period, source type of COVID-19 cases, number of new
107 confirmed cases, number of cured and discharged cases, and number of deaths in the city
108 were collected through the details of confirmed COVID-19 cases published on the official
109 websites of the municipal Health Commission.

110 **2.3 Analytical Methods**

111 Use Excel2016 for data collation and analysis. The maximum value, minimum value
112 and mean value were used to describe the measurement data, and the selection rate and
113 proportion were used to describe the counting data. Statistical charts and tables were used
114 to describe the overall trend of COVID-19 epidemic in Henan province and other cities,
115 as well as the sex and age distribution ratio, case source type and case type composition
116 ratio.

117 **3. Results**

118 **3.1 Basic epidemiological characteristics of COVID-19 in Henan Province**

119 **3.1.1 The basic situation of the epidemic in Henan Province.**

120 Since the first confirmed imported case was reported in Zhengzhou on January 21,

121 2020, the daily number of newly confirmed cases in the province has fluctuated with
122 time. On February 4, 2020, the province's new confirmed cases of peak number of 109
123 people, then the entire province new daily total number of confirmed cases of fluctuation
124 decline over time, the Henan province on February 24 for the first time, no new
125 confirmed cases, cumulative COVID - 19 confirmed cases, 1272 cases of, March 11,
126 Zhengzhou in case of input outside (Italy). A total of 1,276 confirmed COVID-19 cases
127 have been reported in Henan province as of June 17, 2020, and the distribution of
128 COVID-19 cases in each city is shown in Fig.1. The cumulative incidence was 1.33 per
129 100,000.

130 On January 29, 2020, a COVID-19 cured and discharged case was reported for the
131 first time in Henan province. From February 2 to March 8, the total number of
132 COVID-19 cured and discharged cases in Henan province rose with time fluctuation,
133 reaching a peak of 103 cases on February 20. From March 9 to 21, 2020, 1 new
134 COVID-19 patient was cured and discharged from hospital. By March 19, all confirmed
135 COVID-19 hospitalized cases in the province had been cleared to zero, with a total of
136 1,250 COVID-19 patients cured and discharged from hospital, with a cure rate of
137 98.19%.

138 On January 26, 2020, there was one death case of COVID-19 for the first time in
139 Nanyang, Henan province. By March 21, 2020, there had been 22 COVID-19 in Henan
140 province, with a fatality rate of 1.73%. There were 1148 common cases in The province,
141 accounting for 90.25%. There were 64 severe cases, accounting for 5.03%. The total

142 number of critical cases was 38, accounting for 2.99%(Fig.2).

143 **3.1.2 Basic characteristics of confirmed cases in Henan Province**

144 Due to limited information on confirmed COVID-19 cases published on the official
145 websites of health commissions of various cities, a total of 600 cases in 4 cities
146 (Zhengzhou, Xinyang, Nanyang and Puyang City) with comprehensive information were
147 selected for statistical analysis. Among the 600 confirmed cases, 315 were male and 285
148 were female, with a sex ratio of 1.11 to 1. The age of confirmed cases ranged from 5 days
149 to 85 years, with the oldest from Zhumadian and the youngest from Xinyang. The
150 average age of confirmed cases was 45 years old, the median age was 46 years old and
151 the majority age was 56 years old. Therefore, it was analyzed that COVID-19 population
152 is generally susceptible to COVID-19, but the middle-aged and elderly population is
153 more likely to be 40-60 years old. Among the confirmed cases of COVID-19, 325 were
154 imported, of which 277 were imported from Hubei, accounting for 85.23%. 204 contact
155 cases, accounting for 34%; Local type 73 cases accounted for 12.17%(Fig.3).

156 **3.2 COVID-19 epidemic situation at the municipal level**

157 **3.2.1 Variation trend of COVID-19 in each city**

158 As of 24:00 on June 17, 2020, A total of 157 confirmed COVID-19 cases have been
159 reported in Zhengzhou City, with a cumulative incidence of 142/100,000. Xinyang City
160 has reported 273 confirmed COVID-19 cases, with a cumulative incidence of
161 3.61/100,000. In Nanyang, 154 confirmed COVID-19 cases have been reported, with a
162 cumulative incidence rate of 1.35 per 100,000. In Zhumadian, 139 confirmed COVID-19

163 cases have been reported, with a cumulative incidence of 175/100,000. Anyang City has
164 reported 52 confirmed cases of COVID-19, with a cumulative incidence of 0.8 per
165 100,000. Puyang City has reported a total of 17 confirmed COVID-19 cases, with a
166 cumulative incidence of 0.5 per 100,000. Six cities accounted for 62.26% of the province's
167 total number of confirmed COVID-19 cases. Xinyang had the highest incidence of
168 COVID-19, followed by Zhumadian, Zhengzhou, Nanyang and Anyang City. Puyang
169 City had the lowest incidence of COVID-19. According to the data, the COVID-19
170 epidemic situation in Henan province as a whole showed a downward trend from north to
171 south based on the analysis of the geographical location and the flow of people in each
172 city. As the capital city of Henan province, Zhengzhou has a complex population and a
173 large number of people moving around. The overall COVID-19 situation is higher than
174 other central regions. On February 23, 2020 first without new Xinyang COVID - 19
175 confirmed cases, February 24, the province has no new cases confirmed, as of March 10,
176 2020, the cumulative reported COVID - 19 confirmed cases, 1272 cases of, March 11
177 people outside input with confirmed cases of Zhengzhou to find one, then as of June 17,
178 2020, the province had no new confirmed cases, cumulative COVID - 19, 1276 cases of
179 confirmed cases.

180 **3.2.2 Age distribution and sex ratio of confirmed cases**

181 By gender, the male-to-female ratio of confirmed COVID-19 cases was 1.38:1 in
182 Xinyang, 0.78:1 in Nanyang, 1.58:1 in Zhumadian, 1.14:1 in Zhengzhou and 0.55:1 in
183 Puyang (Anyang City only disclosed the number of confirmed cases, and did not disclose

184 the detailed information of confirmed cases, so the following contents will not be
185 analyzed in detail).According to the analysis of the sex ratio among cities, the cumulative
186 incidence of COVID-19 among males in Zhengzhou, Xinyang and Zhumadian is higher
187 than that among females. Among them, the cumulative incidence of COVID-19 among
188 males in Xinyang is the highest (4.08 per 100,000).Puyang had the lowest incidence of
189 COVID-19 among males (0.34/100,000).The cumulative incidence of COVID-19 among
190 women in Nanyang and Puyang is higher than that of men, and the highest incidence of
191 COVID-19 among women in Xinyang is 3.11/100,000, while the lowest incidence of
192 COVID-19 among women in Puyang is 1.33/100,000.

193 According to the age analysis, in order to make a correct and objective analysis of
194 the age composition structure, the age is divided according to the age structure in Henan
195 Statistical Yearbook 2019.The age of confirmed COVID-19 cases ranged from 5 days
196 (Xinyang) to 88 years (Zhumadian), and the age of confirmed COVID-19 cases was
197 concentrated between 15 and 64 years old, accounting for 87.42%. The cumulative
198 incidence of COVID-19 in this age group was the highest (511/100,000) in Xinyang.The
199 cumulative incidence of COVID-19 at age ≤ 14 years was the lowest, with Nanyang
200 having the lowest incidence (0.10/100,000).The results are shown in Table 1.1.The
201 average age of confirmed COVID-19 cases was 42.8 years old in Zhengzhou City, 47.3
202 years old in Xinyang, 44.4 years old in Nanyang, 43.2 years old in Zhumadian, and 39.5
203 years old in Puyang.

204 **3.2.3 Source of confirmed cases**

205 As of 24:00 on June 17, 2020, the source type ratio of confirmed COVID-19 cases in
206 each city (input: contact: local) was 4.62:3.79:1 in Xinyang, 5.73:3.60:1 in Nanyang,
207 8.17:2.42:1 in Zhumadian, 3.96:1.33:1 in Zhengzhou City and 1.5:1.5:1 in
208 Puyang. According to the data, the main sources of confirmed COVID-19 cases are
209 imported cases, among which the highest proportion of imported cases is 70.50% in
210 Zhumadian, followed by Zhengzhou. In Xinyang, imported cases and contact cases
211 accounted for 89.05%, and the ratio of imported cases and contact cases was 1.22:1. The
212 results are shown in Table 1.2. According to the public information of the epidemic
213 notification in Henan Province, confirmed imported cases were selected from Xinyang,
214 Zhengzhou, Zhumadian and Nanyang City for detailed investigation, and the imported
215 cases were subdivided into confirmed imported cases from Hubei and confirmed
216 non-imported cases. The results are shown in Table 2. Hubei accounted for 87.74% of the
217 total imported cases, while Hubei accounted for the majority of COVID-19 imported
218 cases.

219 **3.3 Epidemic Prevention and control**

220 **3.3.1 National epidemic prevention and control**

221 In order to do a good job in the prevention and control of the COVID-19 epidemic in
222 China and safeguard the health and life safety of the people, the National Health
223 Commission has drawn up a novel Coronavirus Pneumonia Prevention and control plan
224 to prevent the spread of the epidemic in light of the epidemic situation and the latest
225 research progress. The health administration departments at all levels are required to

226 coordinate with the CDC, allocate funds and materials for epidemic prevention and
227 control in a timely manner, carry out joint prevention and control work, timely find,
228 timely report, timely feedback and timely analysis. Medical institutions at all levels and of
229 various types are responsible for the detection and reporting of cases, isolation, diagnosis,
230 treatment and clinical management. While preventing and treating cases, they care about
231 people's mental health problems and do a good job in health education and risk
232 communication with the public.

233 **3.3.2 Main prevention and control measures of Henan Provincial Government**

234 In the face of COVID-19, the Henan Provincial government responded positively,
235 quickly implemented the guidance given in the document of the CPC Central Committee,
236 and added a number of important prevention and control measures with Henan
237 characteristics. First, increase publicity. The government has made the public aware of the
238 seriousness of COVID-19 through TV and radio 24/7, and sent text messages on
239 epidemic prevention and control to the public through mobile phone platforms. Second,
240 strengthen supervision. Since the outbreak of the epidemic, Henan Province has given full
241 play to big data computing. The "Health code" of Henan Province has been recognized by
242 31 provinces, autonomous regions and municipalities, and 13.98 million people have
243 applied for it. Of citizens to move data through the cloud and the ticket real-name records
244 for screening, grassroots community unit (streets, bring, village, etc.) strict screening area
245 of Hubei province, Xinyang, Nanyang, Zhumadian, Zhengzhou COVID - 19 passengers
246 serious area, take the necessary means to monitor its own home quarantine is greater than

247 14 d, stop all provincial traffic operations, public supervision and prevention and control
248 of epidemic reporting platform to encourage the populace to supervise to report.Third,
249 fight the epidemic comprehensively.All units at all levels of the health care system will
250 cancel the Spring Festival holiday in 2020, and all staff will arrive at their posts on time
251 to maintain 24-hour communications and respond to the epidemic at any time.130
252 designated hospitals for MEDICAL treatment of COVID-19 have been identified for
253 timely treatment of COVID-19.

254 **3.3.3 Specific epidemic prevention and control measures**

255 All cities in Henan province promptly launched the first-level response to major
256 health emergencies, and adopted epidemic prevention and control measures with the
257 characteristics of each city.COVID - 19 high incidence area in Henan province
258 (Zhengzhou, Xinyang, Nanyang and Zhumadian City) as an example, the Zhengzhou to
259 open special line 12320 provide consultation service for 24 h free COVID - 19, stop all
260 outside of Zhengzhou passenger service at the same time reduce the city public traffic
261 divisions, each district area residents swept a yard code registration management system,
262 such as have not obey the management focuses on the compulsory isolation.Set up 10
263 card point in Xinyang, check the traffic of vehicles at the same time for a full sanitizers,
264 temperature detection was carried out on the staff to ensure that no one car, leakage of
265 Hubei direction into the vehicles and personnel are advised to return within the territory
266 of Xinyang, found that patients with fever timely report to the epidemic prevention and
267 control command and arrange to specify the quantities further isolation.All entertainment

268 venues are closed for business. No food in the catering industry, no matter big or small, is
269 allowed to enter public places. Only when there is no abnormality can you enter public
270 places with a mask and a temperature measurement. Nanyang postponed the work of
271 government organs, the resumption of work of various enterprises and the opening of
272 schools, delayed the opening time of administrative examination and approval services
273 under the line, implemented centralized isolation and centralized observation system, and
274 adopted rewarding reporting measures to encourage citizens to supervise surrounding
275 personnel, so as to control the epidemic in an all-round and multi-angle way. Zhumadian
276 and its subsidiaries closed four fire stations, suspended all traffic operations, closed
277 farmers' markets and entertainment venues. In addition to the above four cities, all other
278 cities have imposed compulsory control on the traffic operation within the city, including
279 subordinate counties, and suspended all kinds of public service places at all levels and all
280 kinds of external purchase and distribution activities, so as to effectively ensure the
281 internal non-proliferation and external import prevention.

282 **4. Discussion**

283 COVID-19 is a new and highly infectious disease that will break out in 2020. It has
284 had a great impact on the whole society. It has not only seriously affected people's life
285 and mental health, but also seriously affected people's production and life. As a province
286 with a large population and strong personnel mobility, Henan province is close to Hubei
287 Province. As of June 17, 2020, Henan province had a total of 1,273 confirmed cases, with
288 a cumulative incidence rate of 1.32/100,000, higher than that of other provinces and cities

289 outside Hubei, which is closely related to the geographical location of Henan
290 province. Since the first confirmed case was reported in Henan province, the number of
291 new confirmed cases per day in general showed an inverted v-shaped trend, reaching a
292 peak of 109 cases on February 20, and the epidemic reached a stable control after
293 February 24. There was no new case except one imported case from abroad on March
294 11. Study shows that the population migration and COVID - 19 cumulative disease rate is
295 positively correlated, Zhengzhou as the capital city as the main transport hub city of
296 Henan province, the population of complex personnel fluidity big range, Zhumadian City
297 has 4 place with Wuhan high-speed and next to, Nanyang, Xinyang and Nanyang City in
298 Henan province and Hubei province at the junction, turnover main direction is in Hubei
299 province, due to the convenient transportation only 30 min drive, so both the city
300 personnel will be in Hubei province as the first choice for migrant workers, entertainment,
301 so the above four city of Henan province COVID - 19 outbreaks. Anyang and Puyang
302 City are in the northernmost part of Henan Province, far away from Hubei Province, so
303 the epidemic has been least affected[10].

304 In terms of the source of confirmed COVID-19 cases, Hubei imported cases
305 dominated the initial stage of the outbreak. Contact cases and local cases fluctuated over
306 time, indicating that the outbreak shifted from external import to internal spread. However,
307 the total proportion of contact cases and local cases was 42.80%, indicating that there was
308 no large-scale spread of the epidemic in Henan province. It is worth noting that one
309 imported case occurred in Zhengzhou on March 11, and the confirmed imported cases

310 from central Hubei accounted for 20.38% of the total cases in Zhengzhou City. Therefore,
311 Zhengzhou City should strengthen the import of other non-Hubei and foreign regions
312 while preventing imported cases from Hubei.

313 Studies by the COVID-19 Epidemiology Group of the Chinese Center for Disease
314 Control and Prevention showed that the sex ratios of confirmed COVID-19 cases in
315 China, Hubei province and Wuhan were 1.06:1, 1.04:1 and 0.99:1,
316 respectively[11].Zhong Nanshan group of patients in gender studies showed that 41.9%
317 of female patients accounted[12], Xi 'an, the centers for disease control and prevention
318 reported male patients accounted for 54.3%[13]^[13], and Lu Jiahai[14],Kang[15]^[15]to
319 Guangzhou confirmed cases found women more than men, the prompt COVID - 19 men
320 and women are prone to infection in the gender, but confirmed cases of gender
321 distribution is different in different areas.For example, male patients in Zhengzhou City,
322 Xinyang and Zhumadian City were more than female patients, while female patients in
323 Puyangand Nanyang City were more than male patients.

324 Since January 21, the province has reported 1,273 confirmed COVID-19 cases
325 (including one imported from abroad) and 22 deaths. On March 15, as the last COVID-19
326 patient in Xinyang was discharged from hospital on March 14, all local hospitalized
327 patients in the province have been cured and discharged, realizing zero cases!At present,
328 the COVID-19 epidemic in Our province has been effectively controlled, but do not relax
329 your vigilance!With the return of overseas students and the return of overseas staff, while
330 preventing the spread of asymptomatic infected persons in Our province, we should

331 strictly control the import of external cases, strictly control the risk of imported cases,
332 strengthen the epidemic prevention and control work in our province, and earnestly
333 achieve the goal of "internal anti-diffusion, external anti-input".

334 **Funding:** This study was funded by the National Natural Science Foundation Project of
335 China (No. U1704182)

336 **Conflict of Interest:** The author declares no conflict of interest.

337 **Acknowledgment:** We thank Dr. Honglei Zhu for his guidance on figure.1 in the
338 manuscript.

339

340 **References:**

341 [1] Commission WMH. Report on current pneumonia epidemic situation in the city (in Chinese).

342 2020.<http://wjw.wuhan.gov.cn/front/web/showDetail/2019123108989> (accessed March 2, 2020) .

343 [2] Rothan HA, Byrareddy SN. The epidemiology and pathogenesis of coronavirus disease (COVID-19)

344 outbreak. *J Autoimmun* 2020; 109:102433.

345 [3] WHO. Novel Coronavirus (2019-ncov) - Situation Report - 22, 2020.

346 https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200211-sitrep-22-ncov.pdf?sfvrsn%3d49b1_2.(accessed March 2, 2020).

347

348 [4] Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in Wuhan, China,

349 of Novel Coronavirus-Infected Pneumonia. *N Engl J Med* 2020; 382:1199-207.

350 [5] AH A, F AO. Novel SARS-CoV-2 outbreak and COVID19 disease; a systemic review on the global

351 pandemic.*Genes & Diseases* 2020.

- 352 [6] Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019
353 novel coronavirus in Wuhan, China. *Lancet (London, England)* 2020; 395.
- 354 [7] Riou J, Althaus CL. Pattern of early human-to-human transmission of Wuhan 2019 novel coronavirus
355 (2019-nCoV), December 2019 to January 2020. *Euro Surveill* 2020; 25.
- 356 [8] China NHCO. Notification on the issuance of the COVID-19 Protocol (trial seventh edition). 2020.
- 357 [9] Province HCOH. Update on COVID-19 in Henan province as of 24:00, March 21. 2020.
- 358 [10] Baidu migration: Destinations moved out of Hubei Province. 2020.
- 359 [11] CDC. CERM. Epidemiological characteristics of COVID - 19. *Chinese Journal of Epidemiology*
360 2020.
- 361 [12] Guan W, Ni Z, Hu Y. Clinical characteristics of 2019 novel coronavirus infection in China. 2020.
- 362 [13] Yao B, Kun L, Zhijun C, Baozhong C, Zhongjun S. Early transmission dynamics of COVID-19 in
363 Shanxi Province. *Chinese Journal of Nosocomiology* 2020; 30:834-8.
- 364 [14] Xuanzhuo W, Conghui L, Zhihui L, Huan H, Xiaomin C, Qianlin L, et al. Preliminary analysis of the
365 early epidemic and temporal and spatial distribution of COVID-19 in Guangdong Province. *Journal of*
366 *Tropical Medicine* 2020; 20:427-30.
- 367 [15] M K, J W, W M. Coronavirus - COVID-19; Evidence and characteristics of human-to-human
368 transmission of SARS-CoV-2 (Updated February 17, 2020). *Medical Letter on the CDC & FDA* 2020.
- 369

370 **Figure captions**

371 **Figure.1** Distribution of COVID-19 in Henan Province, China

372 The figure above indicates the cumulative confirmed COVID-19 cases in Henan
373 Province, China till June 17, 2020. The different colours represent the overall incidence of
374 confirmed COVID-19 cases within the general population within each city of Henan
375 Province, China. The five-pointed star stands for Zhengzhou, the capital of Henan
376 province.

377 **Figure.2** Daily statistics of the epidemic in Henan Province

378 **Figure.3** Age ratio of confirmed COVID-19 cases in Henan Province

379 Here is a breakdown of age structure (<60 years old, 20 years old for a period;> 60 years
380 old, 10 years old for a period).

Table 1.1 Confirmed cases and cumulative incidence of COVID-19 in six longitudinal cities in Henan province as of June 17, 2020

	Xinyang City		Nanyang City		Zhumadian City	
	confirmed case[n(%)]	cumulative incidence(1/100000)	confirmed case[n(%)]	cumulative incidence(1/100000)	confirmed case[n(%)]	cumulative incidence(1/100000)
total	274(100.0)	3.61	156(100.0)	1.35	139(100.0)	1.75
male patients	159(58.03)	4.08	67(42.95)	1.17	84(60.43)	2.13
female patients	115(41.97)	3.11	86(55.13)	1.50	53(38.13)	1.46
missing data	0	—	3(1.92)	—	2(1.44)	—
≤14years old	5(1.82)	0.30	3(1.92)	0.10	5(3.60)	0.28
15-64years old	246(89.78)	5.11	132(84.62)	1.84	120(86.33)	2.40
≥65years old	23(8.39)	2.02	18(11.54)	1.27	14(10.07)	1.28
missing data	0	—	3(1.92)	—	0	—
	Zhengzhou City		Puyang City		Anyang City	
	confirmed case[n(%)]	cumulative incidence(1/100000)	confirmed case[n(%)]	cumulative incidence(1/100000)	confirmed case[n(%)]	cumulative incidence(1/100000)
total	157(100.0)	1.42	17(100.0)	0.5	53(100.0)	0.8
male patients	83(52.86)	1.5	6(35.29)	0.34	—	—
female patients	73(46.49)	1.34	11(64.71)	1.33	—	—
missing data	1(0.65)	—	0	—	—	—
≤14years old	6(3.82)	0.28	1(5.88)	0.11	—	—
15-64years old	134(85.35)	1.77	14(82.35)	0.6	—	—
≥65years old	16(10.19)	1.3	2(11.76)	0.49	—	—

missing data	1(0.64)	—	0	—	—	—
--------------	---------	---	---	---	---	---

381 Note: This table only collects statistics from the official website, and the age structure is
 382 based on the age of Statistical yearbook of Henan Province.
 383 —:Data missing or not statistically significant
 384

Table 1.2 Source types of confirmed COVID-19 cases

	Xinyang City		Nanyang City		Zhumadian City	
	confirmed case	percentage [n(%)]	confirmed case	percentage [n(%)]	confirmed case	Percentage [n(%)]
imported case	134	49.08	85	55.19	98	70.50
contact case	110	40.29	53	34.42	29	20.86
local type of case	29	10.62	15	9.74	12	8.63
missing data	—	—	1	0.65	—	—
	Zhengzhou City		Puyang City		Anyang City	
	confirmed case	percentage [n(%)]	confirmed case	percentage [n(%)]	confirmed case	percentage [n(%)]
imported case	99	63.06	6	35.29	—	—
contact case	33	21.02	7	41.18	—	—
local type of case	25	15.92	4	23.53	—	—
missing data	—	—	—	—	—	—

385 Note: This table only collects statistics from the official website, and the age structure is
 386 based on the age of Statistical yearbook of Henan Province.
 387 —:Data missing or not statistically significant


388

Table 2 Analysis of imported cases in areas with high epidemic incidence in Henan

	Province			
	Xinyang City	Zhengzhou City	Zhumadian City	Nanyang City
	confirmed case[n(%)]	confirmed case[n(%)]	confirmed case[n(%)]	confirmed case[n(%)]
Imported case from Hubei	128(46.72)	69(43.95)	92(66.19)	76(49.03)
Non - Hubei imported case	9(3.28)	32(20.38)	6(4.32)	4(2.58)

389

390



391

Figure.1 Distribution of COVID-19 in Henan Province, China




Figure.2 Daily statistics of the epidemic in Henan Province

392
393
394
395




Figure.3 Age ratio of confirmed COVID-19 cases in Henan Province

396