The Impact of COVID-19 on Medical Practice: A Nationwide Survey of Dermatologists and Healthcare Providers

Mohammed Shanshal[1, *], **Hayder Saad Ahmed**[2, ^], **Hayder Asfoor**[3, *], **Raad Ibrahim Salih**[2, ^], **Shehab Ahmed Ali**[4, ^], **Yusif k. Aldabouni** [1, ^]

[1] Baghdad Teaching Hospital, Department of Dermatology and Venereology, [2] University of Tikrit, College of Medicine, Department of Dermatology and Venereology, [3] University of Kerbala, College of Medicine, Department of Medicine, [4] Al- iskandria Hospital, Department of Dermatology and Venereology

* Specialist dermatologist, MBChB, FABMS – Dermatology

^ Specialist dermatologist, MBChB, FIBMS – Dermatology

Corresponding author:

Mohammed Shanshal, MBChB, FABMS – Dermatology
Specialist Dermatologist at Baghdad Teaching Hospital, Department of Dermatology and Venereology
Email: doctor.dermatology30@yahoo.com
ORCID Id: https://orcid.org/0000-0003-2877-5919

Word count: 2227
Abstract word count: 199
Highlights: 51
Number of figures: 5
Number of tables: 5
Number of references: 24
Conflicts of interest: None
Funding sources: None
Patient privacy and ethical use of images:
The patients in this manuscript have given written informed consent for publication of their case details, including the use of images.

key words:
PPE-related dermatoses, COVID-19 pandemic, COVID-19 and dermatology, COVID-19 cutaneous manifestations, Telemedicine

Abstract

Background: The COVID-19 pandemic has dramatically changed medical practice worldwide. It posed a significant impact on different health services, including dermatology.

Methods: A Cross-sectional observational study of 200 healthcare providers and 100 dermatologists (survey 1 and 2, respectively) were conducted.

Objectives: To determine the prevalence of occupational skin diseases among healthcare providers working amid the pandemic, and to demonstrate the outbreak's impact on dermatology practice.

Results: Most healthcare providers (83%) reported hygiene-related hand dermatitis. The rates of PPE- related dermatoses were estimated to be 73%, including pressure injuries (51.9%), acne (33.1%), non-gloves contact dermatitis (29.9%), nonspecific rash (17.5%), urticaria (9.1%) and skin infections (3.2%). The emerging COVID-19-related cutaneous manifestations were recognized by 20% of surveyed dermatologists, including maculopapular rash (41.67%), urticaria (37.50%), chilblain (25%) and vasculitis (16.67). Telemedicine was provided by 73% of the dermatologists, and 89% reported minimal use of immunosuppressive drugs amid the pandemic.

Conclusions: This article highlights the emergence of hygiene-related hand dermatitis and PPE-related dermatoses among healthcare providers working in the COVID-19 era. It also provides an appreciation of the major impact of COVID-19 on different aspects of dermatology practice in Iraq, and how the dermatologists adapt to these unfamiliar circumstances to meet the challenges.

Highlights:

● COVID-19 is associated with an ongoing emergence of occupational skin disease among healthcare providers

● COVID-19 posed a significant impact on medical practice, including the epidemiology of diseases, the use of telemedicine, and modification of management plans

● Dermatologists play a crucial role in recognizing the cutaneous manifestations associated with COVID-19 infection.

Introduction

An ongoing outbreak of coronavirus disease 2019 (COVID-19) caused by SARS-CoV-2, started in December 2019. It was first identified in the Chinese city of Wuhan. The virus then propagated and began to hit other regions around the world until it was declared a global pandemic by the World Health Organization (WHO) on March 11, 2020 [1].

The emerging COVID-19 virus is phylogenetically linked to viruses that cause severe acute respiratory syndrome (SARS) and Middle East respiratory syndrome (MERS). COVID-19 causes variable degrees of illness, ranging from fever, cough, dyspnea, fatigue, and diarrhea to critical cases of severe acute respiratory distress syndrome [2].

The COVID-19 pandemic has brought the world to a standstill and placed considerable challenges on healthcare workers, including dermatologists. Telemedicine has gained particular importance, where many of the dermatological consultations were transitioned into teledermatology services. The current outbreak has led to the ongoing emergence of personal protective equipment (PPE)- related dermatoses and hygiene-related hand dermatitis, particularly among healthcare providers. In addition, due to a shortage of vaccination programs

imposed by the COVID-19 lockdown, there is a significant risk of other infectious disease outbreaks, such as measles.

Objectives and methods

A cross-sectional observational study involving two online surveys were conducted for data collection. A 5-item electronic survey was created and sent to 276 randomly selected healthcare providers working in the emergency departments, inpatient wards, and outpatient clinics of the major hospitals in Iraq. The survey assessed the frequency of occupational skin diseases due to hygienic work practices and PPE use among healthcare providers working during the current pandemic. The participants were asked about their gender, age, as well as the frequency of hygiene-related hand dermatitis and skin complications related to PPE use.

For survey 2, a 12-item electronic questionnaire was sent to 156 randomly selected dermatologists practicing amid the COVID-19 era. The dermatologists were asked about the impact of the current outbreak on various aspects of dermatology practice, including the provided medical services, infection control methods, disease epidemiology, and the currently applied medical and surgical management plans.

Results

In the first survey, a total of 200 responses were collected from healthcare providers working amid the COVID-19 pandemic with a response rate of (72.46%). The mean age of respondents was 31.3 years (4.95 standard deviation), of whom 64% were female and 36% were male (**Table 1**). Of the participants, 166 (83%) reported hygiene-related hand dermatitis due to overuse of detergents and disinfectants, while 147 (73.5%) reported cutaneous side effects due to the use of PPE. Types and frequency of PPE- related dermatoses are summarized in **Table 2.**

Table 1:

Occupational skin diseases among health care providers working amid COVID-19

	Male n (%)	Female n (%)	Total n (%)
Response rate	72 (36%)	128 (64%)	200 (72.46%)
Hygiene- related hand dermatitis	43 (25.9%)	123 (74%)	166 (83%)
PPE-related dermatoses	36 (24.9%)	111 (75.5%)	147 (73.5%)

Table 2:

Types and frequency of PPE-related dermatoses

	Male n (%)	Female n (%)	Total n (%)
Pressure injuries	14 (17.5)	66 (82.5%)	80 (51.9 %)
Acne	8 (15.68)	43 (84.3%)	51 (33.1 %)
Non-gloves CD	11 (24%)	35 (76%)	46 (29.9 %)
Non-specific rash	5 (18.5%)	22 (81.5%)	27 (17.5%)
Urticaria	3 (21.42%)	11 (78.57)	14 (9.1%)
Skin infections	3 (60%)	2 (40%)	5 (3.2%)

In the second survey, the questionnaire was completed by 100 practicing dermatologists, with a 64% response rate. The survey questions and results are illustrated in **Table 3**. Most dermatologists (68%) reported using immunosuppressive drugs at a lower rate and 81% endeavored to shift their patients towards more conservative options, such as topical agents and phototherapy. To decrease the risk of infection spread, 72% of responding dermatologists reported providing some forms of teledermatology services, whereas 71% continued to perform procedures/biopsies at a lower rate for selected urgent cases. Of the participants, 20% of dermatologists reported diagnosing one or more of COVID-19 related skin manifestations, including

maculopapular rash, urticaria, chilblain and vasculitis (**Table 4**). Aggravation of certain preexisting dermatologic diseases like psoriasis, atopic dermatitis and rosacea have been reported by 51% of the surveyed dermatologists (**Table 5**).

Table 3

Impact of COVID-19 on dermatology practice: Survey questions and results

	Yes n= (%)	No n= (%)
Do you continue to perform dermatological procedures and biopsies?	71 (yes, but at limited capacity) 16 (yes, as before the outbreak)	13
Are you still using immunosuppressive to treat skin conditions?	68 (yes, but only for limited cases when highly indicated) 11 (yes, as before the outbreak)	21
Do you diagnose one of the COVID-19 skin manifestations?	20	80
Due to COVID-19 related psychological upset, do you notice an increase in relapse rate of some chronic skin diseases?	51	49
Do you try to switch immunosuppressive to more conservative options like topical agents/ phototherapy?	81	19
Do you get COVID-19 infection during your work? (PCR confirmed case only)	25	75
Do you provide online/ phone consultations (teledermatology)?	72	28

Did you work at the respiratory triage or any other COVID-19 related unit?	19	81
Did you diagnose a case of measles during the current pandemic?	14	86

Table 4

Cutaneous manifestations of COVID-19 diagnosed by dermatologists

	n	%
Erythematous maculopapular rash	10	41.67%
Urticaria	9	37.50%
Chilblain	6	25%
Vasculitis	4	16.67%

Table 5

Dermatoses aggravated by COVID-19 related psychological stress

	n	%
Psoriasis	33	64.70%
Atopic dermatitis	30	58.82%
Rosacea	9	17.64%
Vitiligo	5	9.80%
alopecia areata	3	5.88%

Discussion

✚ **Impact of COVID-19 on medical services**

In this article, we aim to review the impact of COVID-19 pandemic on dermatology practice in Iraq. Although the population of Iraq has exceeded 40 million in 2020 based on the latest United Nation data [3], available statistics shows that the number of physicians is 7.08 per 10,000 populations which is extremely low compared to developed countries [4].

On 24 February 2020, Iraq announced the first confirmed case of SARS-CoV-2 infection. Since then, the spread of COVID-19 has increased exponentially. By early September, about 260,370 confirmed cases and 7,512 deaths were reported in Iraq [5].

The COVID-19 pandemic has targeted every single aspect of medical practice, including dermatology. As Baghdad became the epicenter of the COVID-19 pandemic in mid-March, most dermatology clinics were closed down or continued to operate at a minimal capacity. The objective was to provide essential medical care for emergency cases such as Stevens-Johnson syndrome/toxic epidermal necrolysis, erythroderma, acute angioedema, and lesions worrisome for skin cancer.

With the acute shortage of PPE, medical practitioners clenched their teeth with each patient, wondering if they had been exposed to the virus already. COVID-19 patients may present with a wide range of dermatological manifestations with or without typical respiratory signs [6] [7].

Of the 100 dermatologists involved in the survey, 20% reported diagnosing one or more COVID-19 related skin manifestations. Erythematous maculopapular rash (41.67%), urticaria (37.50%), chilblain (25%), and vasculitis (16.67%) were the most commonly diagnosed manifestations in decreasing frequency (**Figure 1**).

Figure 1: **COVID-19 related cutaneous manifestations**
(A) The approximate frequency of diagnosed COVID-19 related skin manifestations reported by dermatologists. **(B)** 25 years old male with erythematous maculopapular eruption involving the extremities **(C)** 12 years old girl with generalized urticarial eruption.

🞣 Impact of COVID-19 on infection control

The dermatology team tried to reduce face-to-face consultations in public hospitals and private practice clinics and limit them to emergency cases. Most non-emergent presentations such as melasma, acne, vitiligo, and stable psoriasis have been diverted to telemedicine platforms to avoid the unwarranted risk of spreading infection. About 25 dermatologists reported contracting the virus during their medical practice.

The survey revealed that during the outbreak, 72 % of practicing dermatologists provided teledermatology consultations through telephone and/or online social

network applications. Since dermatology diagnosis depends primarily on visual characteristics, dermatologists have always been leaders in providing telemedicine services even before the crisis, leading to greater success with this experience.

In order to reduce the risk of infection spread while attending dermatology clinics, several measures have been implemented including stressing commitment to social distancing and wearing a mask, the pre-examination measurement of the patient's temperature, and exclusion of the respiratory symptoms at the hospital entrance respiratory triage unit, in addition to the meticulous disinfection of equipment [8] (**Figure 2**).

Figure 2: **COVID-19 pre-examination triage steps**
Flowchart represents the pre-examination triage to decrease the risk of infection spread while attending the dermatology clinic and utilizing teledermatology for initial +/- follow-up assessment.

Impact of COVID-19 on skin disease epidemiology

> **Hand dermatitis**

Due to frequent handwashing, prolonged use of gloves, and overuse of disinfectants, the incidence of hand dermatitis among healthcare staff and the general population has dramatically increased (**Figure 3**).

In this survey, hand dermatitis was reported in 83% of healthcare providers working during the current pandemic. Another study reported a comparable result of gloves-related hand dermatitis (88.5%) among healthcare providers working in Hubei Province hospitals amid the outbreak [9]. Smith DR et al. conducted a pre-outbreak epidemiological study in mainland China where the reported frequency of hand dermatitis ranged from 14.3% to 23.8% across hospital departments, far from those recorded during the COVID-19 outbreak [10].

Several measures can be used to decrease the incidence of hand dermatitis, including the use of mild soaps (syndets) that are equally effective in preventing the viral spread, and frequent use of moisturizers and physical barriers, especially after handwashing to protect the skin and decrease irritation. Depending on the severity of the lesions, established hand dermatitis should be treated with emollients and/or topical steroids [11].

Figure 3: **Hygiene-related hand dermatitis**

(A) (B) Eczematous contact dermatitis **(C)** Papular contact dermatitis **(D)** Post-inflammatory hyperpigmentation following severe disinfectants-related dermatitis

Personal protective equipment (PPE)-related dermatoses

Prolonged contact with PPE such as masks, protective gowns, face shields, and goggles has led to an unprecedented emergence of PPE-related dermatoses among healthcare providers working amidst the current pandemic. PPE-induced skin injuries are attributed to mechanical friction, long-term occlusion, maceration, and allergic contact reactions [12].

Among the surveyed healthcare providers, 147 (73.5%) reported PPE-related dermatoses. Of these, 111 (75.5%) occurred in women and 36 (24.9%) in men. The median age was 31 years (range 24-47 years; standard deviation 4.94 years).

The frequency of PPE-related dermatosis reported in this survey was as follows: pressure injuries (51.9%), acne (33.1%), non-gloves contact dermatitis (29.9%), nonspecific rash and itching (17.5%), urticaria (9.1%), and skin infections (3.2%) (**Figure 4**). Folliculitis can occur at the site of skin friction, and the inflamed follicles may become infected with bacteria, especially Staphylococcus. Friction caused by surgical masks can also trigger reactivation of the herpes simplex virus.

Kaihui Hu et al. explored the frequency of PPE-related cutaneous complications among healthcare workers in the Hubei Province during the COVID-19 era. Adverse

skin reactions associated with the prolonged wearing of PPE, including N95 mask, latex glove and protective clothing, were 95.1%, 88.5% and 60.7%, respectively [8]. During the SARS outbreak, Foo et al. investigated the prevalence of PPE-related dermatosis among Singapore health care workers. The study revealed that 35.5% of the staff who used masks regularly reported acne (59.6%), facial itch (51.4%), and rash (35.8%) while 21.4% who used gloves regularly reported adverse skin reactions, including dry skin (73.4%), itching (56.3%), and rash (37.5%) [13]. Bhoyrul B et al. analyzed incident case reports from UK dermatologists of non-glove PPE-related dermatoses between 1993 and 2013; 15.5% of the affected patients were health and social workers [14].

General measures to reduce skin damage associated with the PPE include wearing suitable PPE, avoiding prolonged use wherever possible and working in a cooler environment. In the areas of maximum tension, dressing can be applied to the skin to redistribute pressure and reduce friction. Frequent use of moisturizers is advisable to protect and repair the skin barrier, and adequate cleansing is necessary to prevent secondary infections [15].

Mild pressure injuries with intact epidermis usually heal spontaneously without intervention. Small bullae can be kept intact, while a larger one can be aspirated to ease the associated pressure. Eczematous lesions are treated with moisturizers and/or topical steroids, depending on the severity of the lesions. Antihistamines may be used for associated pruritus. For resistant cases, a patch test may be necessary to identify the allergen involved. Mask-related frictional acne can be treated with topical benzoyl peroxide, retinoid, and antimicrobials, similar to acne vulgaris [16].

Figure 4: **Spectrum of PPE-related dermatosis among healthcare provides working in COVID-19 era**

(A) Pressure marks and frictional acne **(B)** Mask –induced contact dermatitis (**C**) and **(D)** Folliculitis and herpes simplex reactivation triggered by surgical mask friction **(E)** and **(F)** gown-related contact dermatitis and urticaria **(G)** Pressure injuries and nonspecific rash **(H)** mask-related frictional acne **(I)** Goggles-related nasal bridge dermatitis and hyperpigmentation.

➢ **Chronic skin disease relapse**

It is currently enigmatic how COVID-19 can affect those with chronic skin conditions such as psoriasis and how it can affect these individuals' treatment. The psychological impact of the crisis, patients' reluctance to attend their clinic appointment, limited use of immunosuppressive drugs, and shortage of drugs lead to a perceptible increase in the rate of relapse and exacerbation of several chronic skin diseases [17]. Relapse rates of psoriasis, atopic dermatitis, rosacea, vitiligo, and alopecia areata were noticeably increased as observed by 64.70%, 58.82%, 17.64% 9.80%, and 5.88% of dermatologists, respectively.

Patients should be strongly advised against stopping their medications without consulting their physicians [18]. The importance of supporting patients with chronic dermatologic conditions during the current crisis cannot be overemphasized.

➢ **Preventable infectious diseases**

The WHO expressed great concern about the spread of previously controllable infectious diseases like measles. Derangement of immunization services caused by the ongoing COVID-19 pandemic is likely to further to decrease measles vaccination coverage by an additional 20%, leaving the most susceptible children at risk for highly infectious disease outbreak [19]. About 14% of the surveyed dermatologists

recorded new cases of measles amid the current pandemic; hence, health authorities are called to emphasize the importance of identifying and reaching children with missed vaccine doses to avoid an inevitable outbreak.

▪ Impact of COVID-19 on dermatologic treatments

Several expert opinions have recommended restricting the use of immunosuppressive agents during the current pandemic [20] [21]. Most dermatologists in the survey reported reduced use of non-biologic immunosuppressants and limited their prescription to selected cases, particularly when alternative options are exhausted, and only after COVID-19 symptoms have been excluded. In this condition, the dose should be kept to the lowest acceptable level that still adequately controls the disease.

Because of the uncertainty of whether biologics may place patients at a higher risk of COVID-19 or more dramatic disease course, the initiation of biologic therapy in the current crisis is unadvisable. Alternative therapeutic approaches, particularly in vulnerable patients, may be considered [22]. Patients already receiving scheduled biological therapy, who do not exhibit respiratory symptoms or test positive for COVID-19 may continue their treatment while underscoring the importance of COVID-19 preventive measures, like social distancing and self-quarantining [23].

✚ Impact of COVID-19 on surgical dermatology

Most of the non-essential dermatological procedures and biopsies were postponed to minimize the risk of spreading COVID-19 infection. Skin surgery was limited to a few undeferrable biopsies and high-risk skin cancer removal when the risk of delay exceeded the risk of exposure to the virus **(Figure 5)**. Therefore, some hospitals began to apply specific internal protocols to protect their patients and staff, including testing patients for COVID-19 before their scheduled surgeries and procedures, providing full PPE to all operating staff, and solicitous intraoperative environment sterilization and waste disposal [24].

Figure 5: COVID-19 and surgical dermatology

A dramatic decrease in the number of skin biopsies at Baghdad Teaching Hospital in response to the increasing numbers of COVID-19 patients.

Working on frontlines

Baghdad has been hit particularly hard by COVID-19, where almost half of the confirmed cases were recorded. The dermatology staff felt the call of duty and were actively involved in the respiratory triage at the emergency department, where they continued to help patients and support their fellow physicians. Nineteen percent of the dermatologists surveyed reported working in the respiratory triage or other COVID-19 related units.

Conclusion

The COVID-19 related preventive measures are associated with an upsurge in occupational skin diseases related to excessive handwashing, overuse of disinfectants, and prolonged utilization of personal protective equipment. The pandemic's psychological impact, the fear of attending hospitals, and the shortage of medications have led to an increase in the relapse rate of common chronic skin diseases like psoriasis and atopic dermatitis. The dermatological management plans were overshadowed by the current public health crisis, where dermatologists continued to collaborate with their physician fellows in the battle against COVID-19 infection.

Figure legends:

Figure 1: **COVID-19 related cutaneous manifestations**

(A) The approximate frequency of diagnosed COVID-19 related skin manifestations reported by dermatologists. **(B)** 25 years old male with erythematous maculopapular eruption involving the extremities **(C)** 12 years old girl with generalized urticarial eruption.

Figure 2: **COVID-19 pre-examination triage steps**

Flowchart represents the pre-examination triage to decrease the risk of infection spread while attending the dermatology clinic and utilizing teledermatology for initial +/- follow-up assessment.

Figure 3: **Hygiene-related hand dermatitis**

(A) (B) Eczematous contact dermatitis **(C)** Papular contact dermatitis **(D)** Post-inflammatory hyperpigmentation following severe disinfectants-related dermatitis

Figure 4: **Spectrum of PPE-related dermatosis among healthcare provides working in COVID-19 era**

(A) Pressure marks and frictional acne **(B)** Mask –induced contact dermatitis (**C)** and **(D)** Folliculitis and herpes simplex reactivation triggered by surgical mask friction **(E)** and **(F)** gown-related contact dermatitis and urticaria **(G)** Pressure injuries and nonspecific rash **(H)** mask-related frictional acne **(I)** Goggles-related nasal bridge dermatitis and hyperpigmentation.

Figure 5: COVID-19 and surgical dermatology

A dramatic decrease in the number of skin biopsies at Baghdad Teaching Hospital in response to the increasing numbers of COVID-19 patients.

Acknowledgement

Special thanks to doctor (Jhan Adwer Darzi) for his great effort in the proofreading and editing of manuscript, we also thank doctor (Mohammad Fawzi Al Sultan) for his contribution in the data collection.

References

1- Bi Q, Wu Y, Mei S, Ye C, Zou X, Zhang Z, Liu X, Wei L, Truelove SA, Zhang T, Gao W. Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study. The Lancet Infectious Diseases. 2020 Apr 27.

2- Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, Liu L, Shan H, Lei CL, Hui DS, Du B. Clinical characteristics of coronavirus disease 2019 in China. New England journal of medicine. 2020 Apr 30;382(18):1708-20.

3- Madigan LM, Micheletti RG, Shinkai K. How Dermatologists Can Learn and Contribute at the Leading Edge of the COVID-19 Global Pandemic. JAMA dermatology. 2020 Apr 30.

4- Worldometer, Iraq Population (LIVE); accessed on September 6 2020. Available from https://www.worldometers.info/world-population/iraq-population/

5- World Health Organization (WHO): THE GLOBAL HEALTH OBSERVATORY, Medical doctors (per 10 000 population); last updated February 12 2020. Accessed on September 7 2020. Available from:

https://www.who.int/data/gho/data/indicators/indicator-details/GHO/medical-doctors-(per-10-000-population)

6- Ministry of Health of Iraq, The Daily COVID-19 Update; accessed on September 6 2020. Available from "The daily epidemiological situation of registered infections of the emerging coronavirus in Iraq".

7- Bai Y, Yao L, Wei T, Tian F, Jin DY, Chen L, Wang M. Presumed asymptomatic carrier transmission of COVID-19. Jama. 2020 Apr 14;323(14):1406-7.

8- Zhang H, Long H, Ma L, Wang G, Mu QR, Ran YP, Liu QZ, Xiao SX, Zhang XJ, Zhang JZ, Zhang FR. Consensus on pre-examination and triage in clinic of dermatology during outbreak of COVID-19 from Chinese experts. International Journal of Dermatology and Venereology. 2020 May 12.

9- Hu K, Fan J, Li X, Gou X, Li X, Zhou X. The adverse skin reactions of health care workers using personal protective equipment for COVID-19. Medicine. 2020 Jun 12;99(24):e20603.

10- Smith DR, Wei N, Zhao L, Wang RS. Hand dermatitis among nurses in a newly developing region of Mainland China. International journal of nursing studies. 2005 Jan 1;42(1):13-9.

11- Kampf G, Loeffler H. Prevention of irritant contact dermatitis among health care workers by using evidence-based hand hygiene practices: a review. Industrial health. 2007;45(5):645-52.

12- Gefen A, Ousey K. Update to device-related pressure ulcers: SECURE prevention. COVID-19, face masks and skin damage. Journal of Wound Care. 2020 May 2;29(5):245-59.

13- Foo CC, Goon AT, Leow YH, Goh CL. Adverse skin reactions to personal protective equipment against severe acute respiratory syndrome–a descriptive study in Singapore. Contact dermatitis. 2006 Nov;55(5):291-4.

14- Bhoyrul B, Lecamwasam K, Wilkinson M, Latheef F, Stocks SJ, Agius R, Carder M. A review of non-glove personal protective equipment-related occupational dermatoses reported to EPIDERM between 1993 and 2013. Contact dermatitis. 2019 Apr;80(4):217-21.

15- Zhou NY, Yang L, Dong LY, Li Y, An XJ, Yang J, Yang L, Huang CZ, Tao J. Prevention and Treatment of Skin Damage Caused by Personal Protective Equipment: Experience of the First-Line Clinicians Treating 2019-nCoV Infection. International Journal of Dermatology and Venereology. 2020 Apr 13.

16- Gheisari M, Araghi F, Moravvej H, Tabary M, Dadkhahfar S. Skin Reactions to Non-glove Personal Protective Equipment: An Emerging Issue in the COVID-19 Pandemic. Journal of the European Academy of Dermatology and Venereology. 2020 Apr 17.

17- Darlenski R, Tsankov N. Covid-19 pandemic and the skin-What should dermatologists know?. Clinics in Dermatology. 2020 Mar 24.

18- Rosenbaum L. The untold toll—the pandemic's effects on patients without Covid-19. N Engl J Med. 2020 Jun 11; 382:2368-2371

19- UNICEF Iraq, UNICEF and WHO sound the alarm on the health dangers of children in Iraq missing routine immunization during the Covid-19 pandemic; 26 April 2020. Available from:

https://www.unicef.org/iraq/press-releases/unicef-and-who%C2%A0sound-alarm-%C2%A0health-dangers-children-iraq-missing-routine

20- Conforti C, Giuffrida R, Dianzani C, Di Meo N, Zalaudek I. COVID-19 and psoriasis: is it time to limit treatment with immunosuppressants? A call for action. Dermatologic Therapy. 2020 Mar 11.

21- Shah P, Zampella JG. Use of systemic immunomodulatory therapies during the coronavirus disease 2019 (COVID-19) pandemic. Journal of the American Academy of Dermatology. 2020 Jun 1;82(6):e203-4.

22- American Academy of Dermatology (AAD). Guidance on the use of biologic agents during COVID-19 outbreak, 18 March, 2020. Available from:

https://assets.ctfassets.net/1ny4yoiyrqia/PicgNuD0IpYd9MSOwab47/023ce3cf6e

b82cb304b4ad4a8ef50d56/Biologics_and_COVID-19.pdf

23- Brownstone ND, Thibodeaux QG, Reddy VD, Myers BA, Chan SY, Bhutani T, Liao

W. Novel coronavirus disease (COVID-19) and biologic therapy in psoriasis:

infection risk and patient counseling in uncertain times. Dermatology and therapy.

2020 Apr 16:1-1.

24- Der Sarkissian SA, Kim L, Veness M, Yiasemides E, Sebaratnam DF.

Recommendations on dermatologic surgery during the COVID-19 pandemic.

Journal of the American Academy of Dermatology. 2020 Jul 1;83(1):e29-30.