Body mass index multiple regression formula testable by

all nine Bradford Hill causality criteria: Artificial

intelligence analytics applied to global burden of disease

data relating to the obesity epidemic

David K Cundiff, MD (0000-0002-3206-9665), independent researcher,[1,3] Chunyi Wu, PhD (0000-0002-

2186-3433), Research Epidemiologist/Statistician[2,3]

[1] Long Beach, California, USA

[2] Area Specialist Lead in Epidemiology and Statistics, Michigan Medicine, Ann Arbor, Michigan, USA

[3] Volunteer collaborators with the Institute of Health Metrics and Evaluation, Seattle, Washington, USA

Correspondence to: David K Cundiff davidkcundiff@gmail.com Phone: 1-562-438-880

Word count: abstract=350, text=2983

Key Points

Question: Can worldwide global burden of disease (GBD) data analysis derive a body mass index (BMI) multiple regression formula testable by Bradford Hill causality criteria?

Findings: A multiple regression derived formula including population attributable fractions (PAFs) of 20 dietary risk factors, physical activity, childhood severe underweight, and sex satisfied all nine Bradford Hill causality criteria. The BMI formula also plausibly predicted the long-term BMI outcomes related to various dietary and physical activity scenarios.

Meaning: GBD data analysis of BMI and associated risk factors may infer causality with overweight/obesity and possibly for health outcomes in nutritional epidemiology generally.

Abstract

Importance Artificial intelligence analytics may be applied to global burden of disease (GBD) data on body mass index (BMI) and associated risk factors.

Objective Rigorously quantify the interactions of dietary and other risk factors that result in adult BMI

Design We formatted global burden of disease data relevant to body mass index and associated risk factors. We empirically explored the univariate and multiple regression correlations of BMI risk factors with worldwide BMI to derive a BMI multiple regression formula (BMI formula).

Setting Worldwide

Participants Institute of Health Metrics and Evaluation staff and volunteer collaborators analyzed over 12,000 GBD risk factor surveys of people from 195 countries and synthesized the data into representative cohort BMI and risk factor values.

Main outcome measures the performances of the BMI formula when tested with all nine Bradford Hill causality criteria each scored on a 0-5 scale: 0=negative to 5=very strong support

Results In the BMI formula derived, all foods were expressed in kilocalories/day (kcal/day). We adjusted BMI formula risk factor coefficients to equate with their population attributable fractions (PAFs in percent's) relating to worldwide mean BMI. BMI increasing foods had "+" signs and BMI decreasing foods had "-" signs. Total PAF of BMI formula=80.96%. BMI formula=(0.37%*processed meat + 4.23%*red meat + 0.02%*fish + 2.24%*milk + 5.67%*poultry + 1.77%*eggs + 0.34%*alcohol + 0.99%*sugary beverages + 0.04%*corn + 0.72%*potatoes + 8.48%*saturated fatty acids + 3.89%*polyunsaturated fatty acids + 0.27%*trans fatty acids - 2.99%*fruit - 4.07%*vegetables - 0.37%*nuts and seeds - 0.45%*whole grains - 1.49%*legumes - 8.62%*rice - 0.10%*sweet potatoes - 7.45% physical activity (METs/week) - 20.38%*child underweight + 6.02%*sex (male=1, female=2))*0.05012 + 21.77. (BMI formula versus BMI: r=0.907 (95% CI: 0.903 to 0.911) $p<0.0001$, Bradford Hill causality criteria test scores (0-5): (1) strength=5, (2) experimentation=5, (3) consistency=5,

48 (4) dose-response=5, (5) temporality=5, (6) analogy=4, (7), plausibility=5, (8) specificity=5, and (9)

49 coherence=5. Total score=44/45.

50 **Conclusions and relevance** Nine Bradford Hill causality criteria strongly supported a causal relationship

51 between the BMI formula derived and mean BMIs of worldwide cohorts. The artificial intelligence

52 methodology introduced could inform individual, clinical, and public health strategies regarding

53 overweight/obesity prevention/treatment and other health outcomes.

Introduction

Comparable risk assessment (CRA) methodology by the World Health Organization[1] and the Institute of Health Metrics and Evaluation (IHME)[2] based on systematic literature reviews provides our current analytic framework for relating risk factors to health outcomes in population health. However, medical journal reports include a relatively small fraction of the world's nearly 8 billion people. In an editorial in the BMJ titled, "Implausible results in human nutrition research", influential Stanford University meta researcher, Dr. John Ioannidis detailed why, in his words, "definitive solutions won't come from another million observational papers or small randomized trials."[3] Dr. Ioannidis called for radical reform of nutritional epidemiology methodologies used to influence food/agricultural policies and to produce dietary guidelines for clinicians and the public.[4]

According to IBM Cloud Education, "At its simplest form, artificial intelligence (AI) is a field, which combines computer science and robust datasets, to enable problem-solving."[5] This paper's analysis dataset formatted from 1.4 gigabytes of global burden of disease (GBD) worldwide data and BMI multiple regression derived risk factor formula (BMI formula) satisfy this definition of AI. The intent is to answer Dr. Ioannidis' call for innovation in epidemiology, particularly nutritional epidemiology.

Satisfying Bradford Hill causality criteria are considered validating in epidemiological research.[6] We used Bradford Hill causality criteria to test the BMI formula. We hypothesized that AI analytics with GBD big data would contribute to quantifying the determinants of BMI and thereby to better understanding the dietary and other risk factors driving the obesity epidemic.

Methods

As volunteer collaborators with the IHME we received raw GBD ecological data (≈1.4 Gigabytes) on mean BMIs of male and female cohorts 15-49 years old and 50-69 years old from each year 1990-2017 from 195 countries and 365 subnational locations (n=1120 cohorts). We also utilized GBD data on exposures to 32 risk factors and covariates potentially related to BMI.

Food risk factors came from surveys of individuals as g/day. IHME dietary covariate data originally came from Food and Agriculture Organization data on animal and plant food commodities available percapita in countries worldwide—as opposed to consumption data from participant interviews.[7] Supplementary Table 1 lists the relevant GBD risk factors, covariates, and other available variables with definitions of those risk factor exposures.[8]

GBD worldwide citations of over 12,000 surveys constituting ecological data inputs for this analysis are available from IHME.[7] The main characteristics of IHME GBD data sources for BMI, the protocol for the GBD study, and all risk factor values have been published by IHME GBD data researchers and discussed elsewhere.[9-12] These included detailed descriptions of categories of input data, potentially important biases, and methodologies of analysis. We did not clean or pre-process any of the GBD data. GBD cohort risk factor and BMI data from the IHME had no missing records. The updated 2019 raw data with variables we used for this analysis may be obtained by volunteer researchers collaborating with IHME.[13]

To maximally utilize the available data, we averaged the values for ages 15-49 years old together with 50-69 years old for BMI and for each risk factor exposure for each male and female cohort for each year. Finally, for each male and female cohort, data from all 28 years (1990-2017) on mean BMI and on each of the risk factor exposures were averaged using the computer software program R.

To weigh the country and subnational data according to population, internet searches (mostly Wikipedia)

yielded the most recent population estimates for countries and subnational states, provinces, and regions.

World population data from the World Bank and the Organisation for Economic Co-operation and

Development could not be used because they did not include all 195 countries or any subnational data.

Using the above-described formatted dataset of risk factors, covariates, and BMIs, a software program in

R generated a population-weighted analysis dataset. Each male or female cohort in the population-

weighted analysis dataset represented approximately 1 million people (range: < 100,000 to 1.5 million).

The analysis dataset had n=7846 cohorts (rows of data), half male and half female, representative of over

seven billion people.

Supplementary Table 2 details how we converted omega-3 fatty acid g/day to fish g/day using data on the

omega-3 fatty acid content of frequently eaten fish from the National Institutes of Health Office of

Dietary Supplements (USA).[14] As shown in Supplementary Table 3, we converted all of the animal and

plant food data, including alcohol and sugary beverage consumption, from g/day to kcal/day. For the g to

kcal conversions, we used the Nutritionix track app,[15] which tracks types and quantities of foods

consumed. Saturated fatty acids (SFA) risk factor (0-1 portion of the entire diet) was not available with

GBD data from 2017, so GBD SFA risk factor data from 2016 was used. Polyunsaturated fatty acid

(PUFA) and trans fatty acid (TFA) GBD risk factor data from 2017 (0-1 portion of the entire diet) were

also utilized. These fatty acid data were converted to kcal/day by multiplying the fatty acid 0-1 portion of

the entire diet by the kcal/day available for each cohort.

Statistical methods

To determine the strengths of the risk factor correlations with mean BMIs of population weighted

worldwide cohorts (7846 cohorts) or subgroups of cohorts (e.g., continents, socio-demographic quartiles,

etc.), we utilized Pearson correlation coefficients: r, 95% confidence intervals (95% CIs), and p values.

In seeking to derive the optimal BMI multiple regression formula from worldwide data, we determined

our methodology as we proceeded by experimenting with strategies to optimize the functioning of

candidate BMI formulas derived by

1.   including as many as possible of the available dietary variables,

2.   combining dietary variables if appropriate, and

3.   including physical activity and other plausibly informative variables.

Appendix 1 explains the use of Bradford Hill causality criteria to assess whether the risk factors in the

BMI formula derived accurately modeled the worldwide mean BMI. Briefly, we tested the BMI formula

output with the Bradford Hill causality criteria (1) strength, (2) experimentation, (3) consistency, (4) dose

response, (5) temporality, (6 analogy, (7) plausibility, (8) specificity, and (9) coherence. For each

criterion, we used a 0-5 scale to assess the magnitude of support of the BMI formula output being

causally related to the BMIs worldwide (0=no support of causality to 5=very strong support of causality,

total possible points=45.

In determining the variables to include and exclude in worldwide BMI formula, we set the statistical

threshold for a variable to enter and to remain in the formula at $p<0.25$. We used SAS and SAS Studio

statistical software 9.4 (SAS Institute, Cary, NC) for the data analysis.

Results

Table 1 shows the basic statistics and univariate correlations of mean cohort BMI worldwide with mean

dietary and other risk factor values. Whole grains, legumes, rice, and sweet potatoes negatively correlated

with BMI. We designated them, "BMI decreasing foods." The six animal foods (processed meat, red

meat, fish, milk, poultry, and eggs), alcohol, sugary beverages, corn availability, potato availability, added

SFA, added PUFA, and added TFA all positively correlated with BMI. We designated them, "BMI

increasing foods."

In Supplementary Table 4, which shows BMI risk factor to risk factor correlations, corn availability

(kcal/day percapita, a covariate) correlated moderately strongly with sugary beverages (r=0.419, 95% CI

0.400 to 0.437, p<0.0001), suggesting that high fructose corn syrup may have accounted for the

unexpected positive correlation of corn with BMI.

According to the International Potato Center, potato availability (a covariate), which positively correlated

with BMI, included ≥50% highly processed potato products worldwide,[16] likely accounting for the

positive correlation of potatoes with BMI.

The strong positive correlations of fruits, vegetables, and nuts and seeds with BMI (Table 1) were

unexpected. These findings suggested likely multicollinearity of fruits, vegetables, and nuts and seeds

with BMI increasing foods (when an independent variable is highly correlated with one or more known or

unknown other independent variables). Supplementary Table 4 shows the univariate correlations of each

of the BMI formula risk factors with each other risk factor. Fruits, vegetables and nuts and seeds strongly

positively correlated with 10 of the 13 dietary variables that positively correlated with BMI (six animal-

based foods, alcohol, SFA, PUFA, and TFA). Fruits, vegetables and nuts and seeds negatively correlated

with the four other plant-based foods (whole grains, legumes, rice, and sweet potatoes), except fruits was

not significantly correlated with whole grains. Together, these findings strongly suggested that

multicollinearities accounted for the positive correlations of fruits, vegetables, and nuts and seeds with

BMI.

Supplementary Table 5 shows the Excel spreadsheet calculations, which were coordinated with SAS

multiple regressions involved in the derivation of the worldwide BMI formula. Appendix 2 detailed the

steps in the BMI (dependent variable) versus BMI risk factors (independent variables) multiple regression

formula derivation. As seen in Supplementary Table 5, described in Appendix 2, and contrasted with the

current method of deriving PAFs,[1, 2, 17] the BMI formula versus BMI variance (R^2=0.8096) * 100

determined the total PAF (80.96%) accounted for by the BMI formula. The 23 risk factor coefficients

could then be equated to the PAFs of those risk factors. The resulting BMI formula is shown below (all

foods in kcal/day):

190   BMI formula = (0.37% * processed meat + 4.23% * red meat + 0.020% * fish + 2.24% *

191   milk + 5.67% * poultry + 1.77% * eggs + 0.338% * alcohol + 0.99% * sugary beverages

192   + 0.04% * corn + 0.72% * potatoes + 8.48% * saturated fatty acids + 3.89% *

193   polyunsaturated fatty acids + 0.27% * trans fatty acids - 2.99% * fruit - 4.07% *

194   vegetables - 0.37% * nuts and seeds - 0.45 * whole grains - 1.49% * legumes - 8.62% *

195   rice - 0.10% * sweet potatoes - 7.45% * physical activity (METs/week) - 20.38% * child

196   underweight (≥2 SD below mean BMI) + 6.02% * sex (male=1, female=2)) * 0.05012 +

197   21.77.

199   BMI formula output analyzed by Bradford Hill causality criteria

200   As mentioned in the methods and detailed in Appendix 1, the nine Bradford Hill causality criteria tested

201   the functionality of the BMI formula:

1. Strength score=5 for r>0.500, $p<0.0001$—The BMI formula's output regressed with mean BMI of worldwide cohorts gave r= 0.907 (95% CI: 0.903 to 0.911) $p<0.0001$), R^2=0.8096, total percent weight=80.96%.

2. **Experiment score=5 for all 20 bootstrap trial BMI formulas with r>0.500, $p<0.0001$)**— Table 2 shows bootstrapping the BMI formula related to mean worldwide BMI with 20 trials (repeated sampling from the worldwide analysis dataset with replacements[18]). Each trial had 100 randomly selected cohorts to generate a unique BMI formula. As Table 2 shows, the mean values for BMI and the risk factor PAFs were all quite close to the mean values for BMI and BMI risk factors in the worldwide BMI formula (last column).

3. **Consistency score=5 for the mean of absolute differences of BMI formula outputs and mean BMI < 0.300 kg/M^2**—In Table 3, we used 37 subsets of worldwide BMI and risk factor data to test the consistency of BMI formula outputs, utilizing the 20 bootstrap trials (#2 Experiment) to generate 80% confidence intervals for each subset. Table 3 shows the average absolute difference between the 37 subgroups mean BMIs and BMI formula outputs was 0.252 kg/M^2.

4. **Dose-response (Biological gradient) score=5 for the mean of absolute differences of BMI formula outputs and mean BMI for the dose-response quartiles< 0.300 kg/M^2**—Table 3 also shows that the mean of the BMI absolute differences between the BMI formula estimates and mean BMIs in the four dose-response quartiles was 0.220 kg/M^2.

5. **Temporality score=5 for the BMI trend formula versus BMI trend r>0.500, $p<0.0001$**—As is shown in Supplementary Table 6 and detailed in Appendix 1, a multiple regression analysis with the worldwide BMI trend 1990-2007 (dependent

variable) versus 22 of the 23 risk factor trends (independent variables excluding sex)

generated a BMI trend formula (All foods are in Kcal/day.):

BMI trend formula= -0.00 * processed meat + 2.52 * red meat - 0.15 * fish +

0.22 * milk + 5.46 * poultry + 1.89 * eggs + 0.07 * alcohol + 0.12 * sugary

beverages - 0.02 * corn + 0.49 * potatoes - 0.01 * saturated fatty acids + 1.01 *

polyunsaturated fatty acids - 0.01 * trans fatty acids + 3.89 * fruit + 5.92 *

vegetables + 0.11 * nuts and seeds + 0.05 * whole grains - 0.88 * legumes + 3.28

* rice + 3.09 * sweet potatoes + 4.91 * physical activity (METs/week) - 0.93 *

child underweight ($\geq$ 2 SD below mean BMI).

As with the BMI formula, the coefficients of the BMI risk factor trends formula were

equated to their PAFs. BMI trend formula r=0.592, 95% CI: 0.577 to 0.606,

$p<0.0001$, and total PAF=35.02%.

6. Analogy score=4—The three metabolic risk factors correlated with BMI to test

analogy were systolic blood pressure (SBP), low density lipoprotein cholesterol

(LDL-C), and fasting plasma glucose (FPG).

• BMI correlated with SBP: r=0.102, 95% CI: 0.080 to 0.124, $p<0.0001$.

• BMI correlated with FPG: r=0.558, 95% CI: 0.542 to 0.573, $p<0.0001$.

• BMI correlated with LDL-C: r=0.756, 95% CI: 0.746 to 0.765, $p<0.0001$.

Since BMI versus SBP r<0.500, Bradford Hill causality score=4.

7. Plausibility: Score=5—Based on systematic medical literature reviews, physical

activity inversely correlated with BMI,[19] and BMI directly correlated with intakes

of sugary beverages,[20] alcohol,[21] and animal foods.[22] The relationship of adult

BMI with early childhood severe underweight has not been reported worldwide.

Since people in poor countries with high infant/childhood malnutrition have fewer

resources to obtain animal foods, fatty acids, and alcohol and more need for

physical exercise than in developed countries, it is highly plausible that childhood

severe underweight negatively correlated with lower BMI in adulthood.

8. Specificity: Score=5 for the BMI formula being unique.—The BMI formula was
   specific to worldwide BMI and would have been different from risk factor
   formulas modeling SBP, FPG, LDL-C, or any health outcome.

9. Coherence: Score=5—As evidenced by the near perfect score 39/40 on the first eight
   criteria, the BMI formula accurately modeled worldwide mean BMI—total causality
   criteria score: 44/45**.**

260   Table 5 shows BMI formula estimates for various patterns of diet and/or other BMI formula risk factors.

261   For instance, increasing physical activity by adding a run for 1 hour/day at six miles/hour on average

262   along with decreasing BMI increasing foods by isocalorically shifting 25% of BMI increasing foods

263   (Kcal/day) to BMI decreasing foods was projected to reduce the mean cohort BMI from 26.66 kg/M^2 to

264   mean BMI=22.26 kg/M^2.

Discussion

In this analysis of long-term worldwide BMI and GBD risk factor data, the nine Bradford Hill causality criteria all supported that the BMI formula derived was causally related to mean worldwide BMI of cohorts of 15-69-year-old people. Now we should consider including the quantification of BMI risk factors demonstrated here into US Department of Health and Human Services (USDHHS)/ US Department of Agriculture (USDA) health policy discussions and clinical nutrition guideline deliberations.

The Supplemental Nutrition Assistance Program (SNAP—formerly Food Stamps) spent an estimated 22.6% of its $73 billion/year budget[23] on payments to low-income Americans for "sweetened beverages, prepared desserts, salty snacks, candy, and sugar."[24] Additionally, the USDA has subsidized crops that go primarily for animal feed or that are processed into sugars while not subsidizing fruits and vegetables.[25] While the USDA recognizes the relatively low intake of fruits and vegetables in the USA and sponsors a publicity campaign to increase fruits and vegetables consumption,[26] USDA expenditures should promote reduced prices of BMI decreasing foods and increased prices of BMI increasing foods.

A review of the literature on food costs relative to nutrient quality found that the median costs of fruits and vegetables (€0.82/100 kcal) was quite high relative meat/eggs/fish (€0.64/100 kcal), fresh dairy (€0.32/100 kcal), nuts (€0.25/100 kcal), and starches (€0.14/100 kcal).[27] This suggests that people who can afford fruits and vegetables are people who also can afford high quantities of the BMI increasing foods, particularly government subsidized animal products. The economics of food, including the high cost of fruits and vegetables may contribute to overweight and obesity in the USA and in other affluent countries.

291 Following a low-carbohydrate, high-fat diet has been demonstrated to cause modest short term weight

292 loss in obese people,[28] However, the BMI formula projects that the long-term effect of a low-carb, high

293 fat diet would be a mean cohort BMI of 29.05 kg/M^2 (Table 5).

Limitations

The GBD data on animal foods, plant foods, alcohol, and fatty acids were not comprehensive and

comprised only 1191.4 Kcal/day on average worldwide. Subnational data were available on only four

countries. Because the data formatting and statistical methodology were new, this was necessarily a post

hoc analysis and no pre-analysis protocol or Bradford Hill causality criteria scoring system was possible.

As detailed in the Foresight Report on obesity,[29] obesity is affected by a complex system of interacting

factors besides diet, physical activity, and childhood feeding patterns, and sex. So genes,[30] gut

microbiome,[31] ultra-processing of food,[32, 33] and other influences on BMI were outside of the purview of

this analysis.

Generalizability

Given the strength and consistency of the relationship of the mean cohort BMI with the BMI formula

estimates (Tables 1- 4), the findings should be generalizable to people all over the world. However, over

70% of the cohorts in analysis dataset had low to moderate BMI levels (BMI < 23 kg/M^2) the results

could be further refined for relatively high SDI countries by deriving a BMI formula from the four

countries that have subnational data (UK, USA, Mexico, and Japan n=730 cohorts) or the cohorts with

mean BMI $\geq$ 23 (n=2256 cohorts).

Conclusions

313 Nine Bradford Hill causality criteria strongly supported that the BMI formula derived with the 23 risk

314 factors in proportion to their PAFs accurately modeled worldwide cohort mean BMI. While this study

315 dealt only with dietary and other available risk factors for BMI (overweight/obesity), the AI methodology

316 introduced could easily apply to estimating PAFs of multiple dietary and other risk factors that pertain to

317 dozens of non-communicable diseases, for which the IHME have GBD data.

Table 1. BMI risk factor basic statistics□ (n=7846)

BMI versus BMI risk factors worldwide n=7846 □	Mean	SD	Min	Max	r	95% CI low	95% CI high	p	R²
BMI kg/M²	21.77	2.290	17.95	29.39					
Processed meat Kcal/day	5.334	9.720	0.20	68.77	0.594	0.579	0.608	<.0001	0.353
Red meat Kcal/day	50.27	45.13	3.21	235.95	0.656	0.643	0.668	<.0001	0.430
Fish Kcal/day	9.985	36.52	0.40	370.36	0.102	0.080	0.124	<.0001	0.010
Milk Kcal/day	25.04	27.05	1.06	146.8	0.676	0.664	0.688	<.0001	0.457
Poultry Kcal/day available	44.32	50.08	1.06	411.9	0.809	0.801	0.816	<.0001	0.654
Eggs Kcal/day available	19.36	14.71	0.79	69.64	0.684	0.672	0.695	<.0001	0.467
Alcohol Kcal/day	81.03	57.33	4.25	429.8	0.146	0.124	0.168	<.0001	0.021
Sugary beverages Kcal/day	298.4	152.4	72.91	1472	0.131	0.109	0.152	<.0001	0.017
Corn Kcal/day available	34.72	48.28	0.16	305.2	0.076	0.054	0.098	<.0001	0.006
Potatoes Kcal/day available	84.04	74.60	3.07	533.9	0.210	0.188	0.231	<.0001	0.044
Saturated fatty acids Kcal/day	190.6	63.94	70.79	481.1	0.703	0.692	0.714	<.0001	0.495
PUFAs Kcal/day	81.10	73.44	2.93	381.3	0.730	0.720	0.740	<.0001	0.533
Trans fatty acids Kcal/day	13.23	13.65	1.99	77.76	0.475	0.458	0.492	<.0001	0.226
Fruits Kcal/day	40.21	22.50	3.58	161.4	0.616	0.602	0.630	<.0001	0.380
Vegetables Kcal/day	79.76	43.12	9.48	304.2	0.511	0.495	0.527	<.0001	0.261
Nuts and seeds Kcal/day	8.414	8.357	0.05	103.0	0.474	0.457	0.491	<.0001	0.225
Whole grains Kcal/day	55.65	30.93	1.14	235.1	-0.203	-0.224	-0.182	<.0001	0.041
Legumes Kcal/day	51.74	32.23	0.51	194.7	-0.384	-0.402	-0.364	<.0001	0.147
Rice Kcal/day available	141.9	116.3	1.42	461.8	-0.557	-0.572	-0.542	<.0001	0.310
Sweet potatoes Kcal/day available	22.76	35.95	0.02	364.7	-0.148	-0.169	-0.126	<.0001	0.022
Physical activity METs	4714	1368	1609	7669	-0.443	-0.461	-0.425	<.0001	0.196
Child underweight by >2SD	0.1862	0.1707	0.0039	0.5300	-0.794	-0.802	-0.786	<.0001	0.631
Sex male 1 and female 2	1.50	0.500	1.00	2.00	0.124	0.102	0.145	<.0001	0.015
Total Kcal/day available	2574	418	1579	3898	0.840	0.834	0.847	<.0001	0.706
Stop breast feeding <6 months	0.1193	0.0555	0.0159	0.2400	0.799	0.791	0.807	<.0001	0.638
Sodium g/d	4.45	2.34	1.33	9.21	-0.017	-0.039	0.005	0.1268	0.000
Calcium g/d	0.301	0.179	0.081	1.04	0.765	0.756	0.774	<.0001	0.585
Dietary fiber g/d	9.21	3.15	2.72	22.68	0.309	0.289	0.329	<.0001	0.096
Fasting plasma glucose mmol/L	4.30	0.350	3.32	5.58	0.557	0.542	0.572	<.0001	0.311
LDL cholesterol mmol/L	2.35	0.400	1.27	3.25	0.756	0.746	0.765	<.0001	0.571
Systolic BP mm Hg	133.9	4.320	123.4	147.9	0.102	0.080	0.124	<.0001	0.010
Socio-demographic index (0-1)	0.543	0.174	0.112	0.900	0.733	0.722	0.743	<.0001	0.537

□ See Supplementary Table 1 for definitions

Table 2. Bootstrap validation experiment: 20 BMI formulas each with 100 random cohorts

Worldwide BMI and BMI risk factors	RCT1	RCT2	RCT3	RCT4	RCT5	RCT6	RCT7	RCT8	RCT9	RCT 10	RCT 11	RCT 12	RCT 13
BMI kg/M^2	21.71	22.04	21.68	21.92	21.57	21.42	21.79	22.18	21.75	21.85	21.49	21.88	21.71
Processed meat	0.33	0.32	0.34	0.34	0.28	0.31	0.32	0.39	0.57	0.36	0.21	0.26	0.28
Red meat	3.95	3.73	4.92	3.47	3.80	3.70	3.45	4.89	5.61	3.69	3.56	3.20	3.16
Fish	0.03	0.02	0.03	0.01	0.64	0.03	0.03	0.03	0.04	0.02	0.02	0.03	0.49
Milk	2.13	2.29	2.40	2.19	2.04	1.95	1.89	2.39	2.66	2.14	1.68	1.83	1.43
Poultry	6.62	6.24	6.60	4.82	5.36	4.33	5.92	6.72	5.41	4.84	4.48	6.55	4.15
Eggs	1.81	1.68	1.84	1.45	2.17	1.70	1.57	1.85	2.10	1.65	1.59	1.46	1.39
Alcohol	0.09	0.74	0.03	0.28	0.05	0.04	0.77	0.59	1.65	0.38	0.04	0.00	0.01
Sugary beverages	1.90	2.25	6.36	0.78	4.79	2.44	0.43	3.96	0.30	1.55	2.04	3.43	2.90
Corn	0.04	0.00	0.09	0.01	0.26	0.02	0.14	0.27	0.00	0.01	0.32	0.28	0.32
Potatoes	0.23	0.10	0.60	0.67	0.69	0.15	0.11	1.77	1.26	0.68	1.39	0.02	1.34
SFA	8.56	7.14	8.55	6.96	9.12	9.81	7.44	8.52	8.80	7.75	6.66	7.10	5.97
PUFA	4.30	3.97	4.31	3.16	4.09	3.59	4.09	4.12	3.96	3.66	2.84	3.30	2.84
TFA	0.33	0.29	0.23	0.23	0.26	0.14	0.29	0.30	0.38	0.26	0.19	0.35	0.14
Fruits	3.18	2.55	3.99	3.25	3.88	2.39	1.95	3.38	2.44	2.00	2.31	2.82	2.65
Vegetables	3.22	3.39	4.70	5.83	3.67	2.95	4.38	3.79	5.06	4.04	5.51	4.46	1.87
Nuts and seeds	0.22	0.32	0.51	0.30	0.55	0.24	0.36	0.59	0.60	0.25	0.07	0.13	0.13
Whole grains	0.14	0.40	0.24	0.85	0.26	0.68	0.18	0.01	0.12	0.29	1.32	0.53	0.17
Legumes	2.86	1.41	0.71	1.38	1.44	1.82	1.44	1.06	1.65	1.48	1.12	1.71	1.33
Rice	7.24	6.91	7.76	9.80	10.26	8.25	6.35	9.19	5.84	4.31	12.17	8.20	7.82
Sweet potatoes	0.05	0.29	0.12	0.20	0.04	0.07	0.22	0.12	0.17	0.17	0.02	0.30	0.05
Physical activity	6.65	10.11	3.69	9.29	0.66	5.71	14.65	2.63	3.81	10.17	7.80	7.37	6.38
Child underweight	8.86	8.41	5.17	5.47	9.68	21.13	23.33	15.63	20.69	27.27	22.28	15.50	23.20
Sex M=1, F=2	21.20	19.02	13.78	18.92	15.67	9.49	1.65	7.23	5.06	4.36	4.29	10.32	9.07
Total % weights	83.94	81.57	76.96	79.67	79.66	80.95	80.97	79.45	78.17	81.35	81.91	79.14	77.11

Worldwide BMI and BMI risk factors	RCT 14	RCT 15	RCT 16	RCT 17	RCT 18	RCT 19	RCT 20	Mean RCTs 1-20	SD	80% CI low	80% CI high	World mean n=7846
BMI kg/M2	21.79	21.65	21.81	21.81	22.01	21.76	21.79	21.78	0.13	21.49	22.04	21.77
Processed meat	0.30	0.53	0.37	0.41	0.28	0.31	0.55	0.36	0.09	0.26	0.55	0.37
Red meat	2.65	4.77	3.57	4.55	2.47	3.49	5.71	3.95	0.84	2.65	5.61	4.23
Fish	0.02	0.02	0.01	0.03	0.02	0.56	0.70	0.12	0.23	0.01	0.64	0.02
Milk	1.84	2.29	1.79	2.51	1.58	1.80	2.50	2.10	0.33	1.58	2.51	2.24
Poultry	4.55	4.91	4.87	5.66	4.66	4.52	6.87	5.43	0.87	4.33	6.72	5.67
Eggs	1.47	1.51	1.33	2.01	1.32	1.61	2.08	1.69	0.25	1.33	2.10	1.77
Alcohol	0.01	2.10	0.66	0.67	0.01	0.19	1.32	0.46	0.56	0.01	1.65	0.34
Sugary beverages	2.58	1.31	0.79	1.11	0.99	0.61	0.29	1.90	1.55	0.30	4.79	0.99
Corn	0.37	0.00	0.54	0.11	0.22	0.35	0.00	0.15	0.16	0.00	0.37	0.04
Potatoes	0.49	0.06	1.49	0.26	0.16	1.71	0.21	0.68	0.56	0.06	1.71	0.72
SFA	6.28	7.85	6.26	9.46	5.28	5.56	10.62	7.77	1.41	5.56	9.81	8.48
PUFA	3.09	3.93	3.53	4.87	2.99	2.81	5.02	3.73	0.62	2.84	4.87	3.89
TFA	0.36	0.29	0.24	0.31	0.34	0.28	0.29	0.28	0.07	0.14	0.38	0.27
Fruits	2.93	3.55	3.77	2.45	2.58	2.01	2.41	2.88	0.61	2.00	3.88	2.99
Vegetables	2.89	3.97	2.99	2.68	3.26	4.29	1.73	3.88	1.15	1.87	5.51	4.08
Nuts and seeds	0.32	0.39	0.38	0.32	0.26	0.27	0.33	0.33	0.14	0.13	0.59	0.37
Whole grains	0.35	0.74	0.06	0.01	0.11	0.36	0.27	0.39	0.33	0.01	0.85	0.45
Legumes	0.68	1.18	0.55	2.30	0.95	0.78	1.61	1.39	0.52	0.68	2.30	1.49
Rice	10.38	6.31	11.98	3.53	5.90	10.30	9.13	8.26	2.25	4.31	11.98	8.62
Sweet potatoes	0.12	0.11	0.03	0.18	0.12	0.10	0.18	0.13	0.08	0.03	0.29	0.10
Physical activity	8.54	8.15	7.04	8.47	12.49	7.95	5.06	7.48	3.10	2.63	12.49	7.45
Child underweight	19.16	19.70	22.65	22.31	28.86	24.16	12.46	18.18	6.77	5.47	27.27	20.38
Sex M=1, F=2	10.56	6.31	5.80	7.13	6.59	6.62	10.47	9.21	5.20	4.29	19.02	6.02
Total % weights	79.92	80.01	74.68	81.34	81.42	80.63	79.83	80.74	2.17	76.96	81.91	80.96

Table 3. Consistency measured with 37 subgroups of worldwide GBD data

Subsets of worldwide data on risk factors related to BMI	Final BMI formula n	Mean BMI	BMI formula output	Absolute differ-ences	RCT 1-20 Mean	80% CI low	80% CI high	BMI output out of 80% CI
SDI quartile 1	1926	24.486	24.496	0.010	24.455	24.138	24.738	
SDI quartile 2	2096	21.827	22.158	0.331	22.185	21.900	22.523	1
SDI quartile 3	2220	20.401	19.931	0.471	19.976	19.584	20.338	1
SDI quartile 4	1604	20.306	20.546	0.240	20.554	20.329	20.903	1
Africa	1682	21.666	21.429	0.237	21.447	21.121	21.772	
Asia	4188	20.478	20.492	0.014	20.515	20.180	20.780	
Europe	880	24.313	24.604	0.292	24.557	24.356	24.730	1
North America	558	25.912	25.494	0.418	25.477	25.016	25.810	1
Oceana	54	24.433	25.205	0.772	25.033	24.721	25.430	1
South America	468	23.463	24.128	0.665	24.222	23.788	24.594	1
Four countries with subnational data	730	24.848	24.795	0.052	24.775	24.296	25.110	
Quartile 1 BMI increasing foods	1967	24.089	24.365	0.275	24.362	24.120	24.584	1
Q2	1834	21.962	21.961	0.000	21.943	21.639	22.290	
Q3	2050	20.607	20.167	0.440	20.188	19.885	20.462	1
Q4	1995	20.494	20.682	0.188	20.738	20.446	21.066	
Quartile 1 BMI decreasing foods	1475	20.612	20.273	0.339	20.312	19.955	20.650	
Q2	2275	21.125	21.049	0.076	21.056	20.731	21.345	
Q3	2171	22.405	22.377	0.028	22.410	22.117	22.772	
Q4	1925	22.695	23.084	0.389	23.067	22.846	23.411	
Quartile 1 physical activity	2371	20.480	20.311	0.169	20.314	20.062	20.555	
Q2	1573	21.663	21.881	0.217	21.912	21.658	22.230	
Q3	1998	21.325	21.324	0.001	21.343	21.026	21.727	
Q4	1904	23.923	23.963	0.039	23.974	23.553	24.369	
Male	3923	21.485	21.420	0.065	21.388	21.138	21.566	
Female	3923	22.051	22.120	0.068	22.180	21.858	22.501	
Bootstrap trial 1	100	21.715	21.741	0.025	21.755	21.503	22.013	
Bootstrap trial 2	100	22.040	21.986	0.055	22.011	21.724	22.254	
Bootstrap trial 3	100	21.676	21.744	0.068	21.769	21.479	22.019	
Bootstrap trial 4	100	21.926	21.944	0.018	21.954	21.659	22.206	
UK	66	24.994	25.245	0.250	25.155	24.739	25.497	
USA	376	26.017	25.809	0.208	25.684	25.177	26.352	
Japan	158	21.892	22.888	0.996	22.898	22.254	23.327	1
Mexico	130	24.985	23.953	1.032	24.222	23.679	24.977	1
Quartile 1 Dose-response: BMI formula versus BMI	1928	24.763	24.949	0.186	24.936	24.617	25.230	
Q2	1469	22.568	22.480	0.088	22.525	22.187	22.805	
Q3	2647	20.872	21.069	0.197	21.069	20.884	21.325	1
Q4	1802	19.227	18.818	0.409	18.859	18.443	19.220	1
Mean of BMI - BMI formula absolute val				0.252				
Mean of BMI-BMI formula dose response				0.220				
Total out of 80% CI range								13

 Table 4. Trends of BMI and the BMI risk factors (1990-2017)

BMI trend versus risk factor trends, n=7846 cohorts	Mean	SD	Min	Max	R	95% CI	95% CI	P	R²
BMI trend	0.076	0.031	-0.018	0.216					
Processed meat Kcal/day trend	-0.051	0.156	-0.812	1.680	-0.009	-0.031	0.013	0.44	0.000
Red meat Kcal/day trend	0.556	1.084	-3.291	4.731	0.410	0.392	0.428	<.0001	0.168
Fish Kcal/day trend	-0.224	0.953	-0.130	9.790	-0.158	-0.180	-0.137	<.0001	0.025
Milk Kcal/day trend	0.190	0.299	-2.018	2.503	0.206	0.185	0.227	<.0001	0.043
Poultry Kcal/day trend	1.146	1.042	-0.871	7.655	0.420	0.402	0.439	<.0001	0.177
Eggs Kcal/day trend	0.300	0.492	-0.877	1.275	0.484	0.467	0.501	<.0001	0.234
Alcohol Kcal/day trend	0.031	0.164	-0.914	1.993	0.289	0.268	0.309	<.0001	0.083
Sugary beverages Kcal/day trend	0.289	1.139	-5.401	14.001	0.125	0.103	0.147	<.0001	0.016
Corn Kcal/day available trend	-0.028	0.606	-4.355	2.809	0.141	0.119	0.162	<.0001	0.020
Potatoes Kcal/day available trend	0.246	1.837	-12.59	11.031	0.272	0.251	0.292	<.0001	0.074
SFA Kcal/day trend	0.210	0.925	-3.660	2.095	-0.050	-0.072	-0.028	<.0001	0.002
PUFA Kcal/day trend	0.996	0.959	-3.026	5.473	0.286	0.266	0.306	<.0001	0.082
TFA kcal/day trend	-0.210	0.446	-2.681	0.013	-0.066	-0.088	-0.044	<.0001	0.004
Fruits Kcal/day trend	0.623	0.604	-2.078	4.104	0.481	0.464	0.498	<.0001	0.232
Vegetables Kcal/day trend	1.399	1.614	-7.961	8.465	0.396	0.377	0.415	<.0001	0.157
Nuts and seeds Kcal/day trend	0.276	0.265	-0.338	3.124	0.123	0.101	0.144	<.0001	0.015
Whole grains Kcal/day trend	0.173	0.586	-2.604	8.463	0.104	0.082	0.126	<.0001	0.011
Legumes Kcal/day trend	0.370	0.979	-3.240	5.401	-0.298	-0.318	-0.277	<.0001	0.089
Rice Kcal/day available trend	-0.787	1.292	-5.051	3.496	-0.393	-0.412	-0.374	<.0001	0.155
Sweet potatoes Kcal/day available trend	-0.460	1.339	-4.818	3.905	-0.499	-0.516	-0.482	<.0001	0.249
Physical activity METs trend	4.724	6.826	-33.86	28.629	0.379	0.360	0.398	<.0001	0.144
Child underweight >2SD trend	-0.004	0.004	-0.014	0.001	0.045	0.023	0.067	<.0001	0.002

329 **Table 5. BMI formula predictions with different diet or other risk factor scenarios** β

Risk factor scenario	BMI kg/M^2	BMI kg/M^2 predicted	80 % CI low	80 % CI high
World (n=7846 cohorts mean 1990-2017)	21.77	21.77	21.33	22.13
World with all children severely underweight		16.90	15.79	18.05
World with no children severely underweight		22.88	22.35	23.38
USA	26.67	26.08	25.30	26.43
USA 25% kcal shifted to BMI decreasing foods†		23.41	22.31	23.93
USA physical activity mean plus 1 hour/day run		24.94	24.05	25.67
USA physical activity mean + 1 hour run and 25% shift to BMI decreasing foods†		22.26	21.63	23.00
USA no red or processed meat		25.36	24.67	25.72
USA no sugary beverages		26.03	25.17	26.34
USA vegetarian		24.13	23.44	24.47
USA vegan		23.47	22.67	23.97
EAT-Lancet diet		22.86	21.67	23.53
Low Carb Mediterranean Diet		29.05	27.92	29.54

330 β BMI formula estimates based on 28 years of following dietary and risk factor patterns

331 † Kcal/day 13 BMI increasing foods isocalorically shifted to the 7 BMI decreasing foods in the BMI

332 formula, distributed proportionally.

335 **Supplementary Table 1. Definitions of IHME GBD risk factors and covariates related to BMI**

Variables	Definition
Alcohol	Any alcohol consumption (g/day)
Body-mass index	Body mass index (BMI) (kg/m^2)—the dependent variable of interest
Child underweight	Proportion of children – 3 SD to – 2 SD of the WHO 2006 standard weight-for-age curve (0-1)
Corn	Corn availability percapita (g/day), a covariate
Discontinued breast feeding	Proportion of children aged 6-23 months who do not receive any breast milk
Eggs	Eggs availability percapita (g/day) a covariate
Fasting plasma glucose	Fasting plasma glucose (mmol/L)
Fish	This variable expressed in g/day was derived by determining the weight of fish in g corresponding to 1 g of omega-3 fatty acids (eicosapentaenoic acid and docosahexaenoic acid) by averaging the fish g per 1 g of omega-3 fatty acids 20 species of fish= 117.04 g/day fish/1 g/day omega-3 fatty acids (Supplementary Table 2)
Fruits	Consumption of fruits (includes fresh, frozen, cooked, canned, or dried fruit but excludes fruit juices and salted or pickled fruits) (g/day)
Kilocalories available /day	The mean number of kilocalories percapita available per day to people in each location (kcal/day available), a covariate
LDL cholesterol	Serum low-density lipoprotein cholesterol (mmol/L)
Legumes	Consumption of beans, lentils, pulses (g/day)
Milk	Consumption of milk including non-fat, low-fat, and full-fat milk but excluding soy milk and other plant derivatives (g/day)
Nuts and seeds	Consumption of nuts and seeds (g/day)
Physical activity	Average weekly physical activity at work, home, transport-related and recreational measured by MET min per week. Less than 3000 METs per week constitutes low physical activity.
Poultry	Poultry availability percapita (g/day), a covariate
Potatoes	Potatoes availability percapita (g/day), a covariate
Processed meat	Consumption of any processed meat (includes meat preserved by smoking, curing, salting, or addition of chemical preservatives, including bacon, salami, sausages, or deli or luncheon meats like ham, turkey, and pastrami (g/day)
Red meat	Consumption of red meat (includes beef, pork, lamb, and goat but excludes poultry, fish, eggs, and all processed meats) (g/day)
Rice	Rice availability percapita (g/day), a covariate
Seafood omega-3 fatty acids	Seafood omega-3 fatty acids (eicosapentaenoic acid and docosahexaenoic acid) in tablet or fish form (g/day)
Socio-demographic index	SDI is a composite indicator of development status that was originally constructed for GBD 2015, and is derived from components that correlate strongly with health outcomes. It is the geometric mean for indices of the total fertility rate among women younger than 25 years, mean education for those aged 15 years or older, and lag-distributed income per capita. The resulting metric ranges from 0 to 1, with higher values corresponding to higher levels of development.
Sugar-sweetened beverages	Consumption of any beverage with ≥50 calories of sugar per one-cup serving, including carbonated beverages, sodas, energy drinks, fruit drinks but excluding 100% fruit and vegetable juices (g/day)
Sweet potatoes	Sweet potato availability percapita (g/day), a covariate
Systolic blood pressure	Systolic blood pressure (mm Hg)
Total sugar	Total sugar availability percapita (g/day), a covariate

Vegetables	Consumption of frozen, cooked, canned, or dried vegetables (including legumes but excluding salted or pickled, juices, nuts and seeds, and starchy vegetables such as potatoes or corn) (g/day)
Whole grains	Consumption of whole grains (bran, germ, and endosperm in their natural proportions) from breakfast cereals, bread, rice, pasta, biscuits, muffins, tortillas, pancakes, and others (g/day)

Supplementary Table 2. Omega-3 Fatty Acid g to fish g calculation¶

Fish	DHA g/3 ounce fish	EPA g/3 ounce fish	Omega-3 Fatty Acids (DHA _ EPA) g/3 ounce fish mean	Fish 3 ounces = 85.02 g	Fish (g) per omega-3 Fatty Acids (g)=columns E/F
Salmon Atlantic farmed	1.24	0.59			
Salmon Atlantic wild	1.22	0.35			
Herring Atlantic	0.94	0.77			
Sardines canned in tomato sauce drained	0.74	0.45			
Mackerel Atlantic	0.59	0.43			
Salmon pink canned drained	0.63	0.28			
Trout rainbow wild	0.44	0.40			
Oysters eastern wild	0.23	0.30			
Sea bass	0.47	0.18			
Shrimp	0.12	0.12			
Lobster	0.07	0.10			
Tuna light canned in water drained	0.17	0.02			
Tilapia	0.11				
Scallops	0.09	0.06			
Cod Pacific	0.1	0.04			
Tuna yellowfin	0.09	0.01			
Mean DHA and EPA Omega-3 Fatty Acids g/3 ounce fish	0.4531	0.2733			
Calculations total Omega-3 FA g to fish g			0.7264	85.02	117.043

¶ Data on omega-3 fatty acid content of varieties of fish came from the National Institutes of Health Office of Dietary Supplements (USA)

344 **Supplementary Table 3. Calculations of kcal/day from g/day of animal and plant foods¶**

Foods	Food sub-categories	kcal/serving	g/serving	kcal/g
Milk (2% fat)		122	244	0.5
Fish		218	170	1.28
Eggs		72	50	1.44
Poultry		187	85	2.91
Red meat		247	85	2.91
Processed meat				
	Salami	222	59	3.76
	Pastrami	104	71	1.46
	Ring baloney	86	28	3.07
	Pepperoni	94	100	0.94
Average processed meat		126.5	64.5	1.96
Fruits		97	162	0.60
Vegetables		59	91	0.65
Legumes		249	179	1.39
Nuts		172	28	6.14
Seeds				
	Flax seeds	55	10	5.5
	Chia seeds	58	12	4.83
	Fennel seeds	34.5	10	3.45
	Hemp seeds	55.3	10	5.53
Average of seeds		50.7	10.5	4.83
Average of nuts and seeds		111.4	19.25	5.78
Corn		99	103	0.96
Potatoes		161	173	0.93
Sweet potatoes		115	151	0.76
Rice		205	158	1.3
Whole grains		120	52	2.31

345 ¶ Source: NutritionIX app

Supplementary Table 4. Pearson correlations between BMI formula risk factors from Table 2

Correlations between BMI risk factors (r)	Red meat	Fish	Milk	Poul-try	Eggs	Alco-hol	Sug-ary bever-ages	Corn	Pota-toes	SFA	PUFA
Processed meat KC/d	0.647	0.286	0.756	0.662	0.569	0.576	-0.108	-0.050	0.159	0.624	0.699
Red meat KC/d	1.000	0.118	0.676	0.664	0.784	0.466	0.046	-0.143	0.123	0.704	0.694
Fish KC/d	0.118	1.000	0.109	0.163	0.393	0.294	-0.163	-0.018	-0.058	0.038	0.163
Milk KC/d	0.676	0.109	1.000	0.673	0.525	0.385	0.072	-0.107	0.202	0.766	0.680
Poultry KC/d	0.664	0.163	0.673	1.000	0.628	0.351	0.142	0.011	0.102	0.701	0.837
Eggs KC/d	0.784	0.393	0.525	0.628	1.000	0.341	0.021	-0.099	0.030	0.595	0.631
Alcohol KC/d	0.466	0.294	0.385	0.351	0.341	1.000	-0.090	-0.107	-0.011	0.287	0.379
Sugary beverages KC/d	0.046	-0.163	0.072	0.142	0.021	-0.090	1.000	0.419	-0.025	0.045	0.113
Corn KC/d	-0.143	-0.018	-0.107	0.011	-0.099	-0.107	0.419	1.000	0.158	-0.133	-0.034
Potatoes KC/d	0.123	-0.058	0.202	0.102	0.030	-0.011	-0.025	0.158	1.000	0.211	0.086
SFA KC/d	0.704	0.038	0.766	0.701	0.595	0.287	0.045	-0.133	0.211	1.000	0.635
PUFAs KC/d	0.694	0.163	0.680	0.837	0.631	0.379	0.113	-0.034	0.086	0.635	1.000
TFA KC/d	0.187	-0.008	0.428	0.446	0.203	0.152	0.198	0.269	-0.091	0.233	0.433
Fruits KC/d	0.371	0.098	0.490	0.606	0.366	0.058	0.198	0.102	0.243	0.470	0.589
Vegetables KC/d	0.382	0.272	0.349	0.345	0.513	0.191	-0.167	-0.203	-0.047	0.224	0.385
Nuts and seeds KC/d	0.386	0.072	0.579	0.504	0.333	0.232	-0.146	-0.112	0.187	0.471	0.527
Whole grains KC/d	-0.112	0.091	-0.254	-0.023	-0.077	0.066	0.204	0.428	-0.058	-0.165	-0.063
Legumes KC/d	-0.404	0.061	-0.194	-0.220	-0.416	0.140	0.197	0.269	0.019	-0.389	-0.145
Rice KC/d	-0.351	-0.070	-0.565	-0.389	-0.271	-0.129	-0.148	-0.250	-0.377	-0.378	-0.368
Sweet potatoes KC/d	0.005	-0.074	-0.395	-0.240	0.049	-0.063	-0.218	-0.028	0.239	-0.130	-0.234
Physical activity METs	-0.068	-0.163	-0.430	-0.396	-0.114	0.105	-0.103	0.028	0.022	-0.309	-0.431
Child underweight	-0.736	-0.189	-0.403	-0.618	-0.803	-0.131	-0.065	-0.083	-0.215	-0.584	-0.622

Correlations between BMI risk factors (r)	TFA	Fruits	Veget-ables	Nuts-seeds	Whole grains	Le-gumes	Rice	Sweet pota-toes	Physi-cal activity	Child under weight	sex m/f
Processed meat KC/d	0.463	0.379	0.290	0.504	-0.023	-0.197	-0.391	-0.194	-0.212	-0.411	-0.086
Red meat KC/d	0.187	0.371	0.382	0.386	-0.112	-0.404	-0.351	0.005	-0.068	-0.736	-0.230
Fish KC/d	-0.008	0.098	0.272	0.072	0.091	0.061	-0.070	-0.074	-0.163	-0.189	-0.023
Milk KC/d	0.428	0.490	0.349	0.579	-0.254	-0.194	-0.565	-0.395	-0.430	-0.403	-0.018
Poultry KC/d	0.446	0.606	0.345	0.504	-0.023	-0.220	-0.389	-0.240	-0.396	-0.618	0.000
Eggs KC/d	0.203	0.366	0.513	0.333	-0.077	-0.416	-0.271	0.049	-0.114	-0.803	0.000
Alcohol KC/d	0.152	0.058	0.191	0.232	0.066	0.140	-0.129	-0.063	0.105	-0.131	-0.432
Sugary beverages KC/d	0.198	0.198	-0.167	-0.146	0.204	0.197	-0.148	-0.218	-0.103	-0.065	-0.311
Corn KC/d	0.269	0.102	-0.203	-0.112	0.428	0.269	-0.250	-0.028	0.028	-0.083	0.000
Potatoes KC/d	-0.091	0.243	-0.047	0.187	-0.058	0.019	-0.377	0.239	0.022	-0.215	0.000
SFAKC/d	0.233	0.470	0.224	0.471	-0.165	-0.389	-0.378	-0.130	-0.309	-0.584	0.007
PUFAs KC/d	0.433	0.589	0.385	0.527	-0.063	-0.145	-0.368	-0.234	-0.431	-0.622	-0.013
TFA KC/d	1.000	0.387	0.284	0.312	0.087	0.044	-0.261	-0.322	-0.412	-0.121	0.096
Fruits KC/d	0.387	1.000	0.386	0.579	0.012	-0.028	-0.319	-0.096	-0.391	-0.493	0.116
Vegetables KC/d	0.284	0.386	1.000	0.547	-0.349	-0.157	-0.230	-0.095	-0.208	-0.453	-0.066
Nuts and seeds KC/d	0.312	0.579	0.547	1.000	-0.211	-0.023	-0.352	-0.074	-0.302	-0.293	-0.030
Whole grains KC/d	0.087	0.012	-0.349	-0.211	1.000	0.100	0.548	-0.039	0.268	0.089	-0.064
Legumes KC/d	0.044	-0.028	-0.157	-0.023	0.100	1.000	-0.050	-0.054	-0.068	0.506	-0.142
Rice KC/d	-0.261	-0.319	-0.230	-0.352	0.548	-0.050	1.000	0.070	0.386	0.374	0.000
Sweet potatoes KC/d	-0.322	-0.096	-0.095	-0.074	-0.039	-0.054	0.070	1.000	0.327	-0.167	0.000
Physical activity	-0.412	-0.391	-0.208	-0.302	0.268	-0.068	0.386	0.327	1.000	0.135	-0.257
Child underweight	-0.121	-0.493	-0.453	-0.293	0.089	0.506	0.374	-0.167	0.135	1.000	-0.017

Red: Fruits, vegetables, and nuts and seeds correlated positively with BMI increasing foods

Blue: Fruits, vegetables, and nuts and seeds correlated positively with themselves

Green: Fruits, vegetables, and nuts and seeds correlated negatively with BMI decreasing foods

Supplementary Table 5. Derivation of the BMI formula

Row		Column A		Column B		Column C		Column D		Column E		Column F
		BMI risk factors		Mean		BMI v RF R^2		Adjust for fatty acids		Multiple regression parameter estimates		Column B Column C Column D Column F
1	+	Processed meat	*	5.334	*	0.3527	*	1	*	0.0034	=	0.0064
2	+	Red meat	*	50.27	*	0.4304	*	1	*	0.0034	=	0.0736
3	+	Fish	*	9.985	*	0.0104	*	1	*	0.0034	=	0.0004
4	+	Milk	*	25.04	*	0.4573	*	1	*	0.0034	=	0.0389
5	+	Poultry available	*	44.32	*	0.6539	*	1	*	0.0034	=	0.0985
6	+	Eggs available	*	19.36	*	0.4672	*	1	*	0.0034	=	0.0308
7	+	Alcohol	*	81.03	*	0.0213	*	1	*	0.0034	=	0.0059
8	+	Sugary beverages	*	298.36	*	0.0170	*	1	*	0.0034	=	0.0173
9	+	Corn available	*	34.72	*	0.0058	*	1	*	0.0034	=	0.0007
10	+	Potatoes available	*	84.04	*	0.0439	*	1	*	0.0034	=	0.0125
11	+	SFA	*	190.58	*	0.4946	*	0.46	*	0.0034	=	0.1474
12	+	PUFAs	*	81.096	*	0.5330	*	0.46	*	0.0034	=	0.0676
13	+	TFA	*	13.23	*	0.2256	*	0.46	*	0.0034	=	0.0047
14	+	Fruits	*	40.21	*	0.3798	*	1	*	0.0034	=	0.0519
15	+	Vegetables	*	79.76	*	0.2612	*	1	*	0.0034	=	0.0708
16	+	Nuts and seeds	*	8.414	*	0.2248	*	1	*	0.0034	=	0.0064
17	-	Whole grains	*	55.65	*	0.0412	*	1	*	0.0034	=	0.0078
18	-	Legumes	*	51.74	*	0.1471	*	1	*	0.0034	=	0.0259
19	-	Rice available	*	141.86	*	0.3104	*	1	*	0.0034	=	0.1497
20	-	Sweet potatoes available	*	22.76	*	0.0218	*	1	*	0.0034	=	0.0017
21	-	Physical activity METs								*	0.12949	=	0.1295
23	-	Child underweight by >2SD								*	0.35420		0.3542
22	+	Sex male 1 and female 2								*	0.10455	=	0.1046
24		Sum of parameter estimates											1.4071
25		R^2 of BMI formula											0.8585
26		R^2 of BMI formula / Sum of parameter estimates											0.6101
27		F1-F23 * F27 * 100 to convert to percent weights											61.01

359   **Supplementary Table 5. Derivation of the BMI formula continued**

Row		Column G BMI risk factors		Column H Un-adjusted Percent weights		Column I BMI risk factors with fruits, vegetables, and nuts and seeds with negative signs		Column J Percent weights (J1- J23) * 0.94299 (J27)
1	+	Processed meat	*	0.3903	+	Processed meat	*	0.37
2	+	Red meat	*	4.4882	+	Red meat	*	4.23
3	+	Fish	*	0.0216	+	Fish	*	0.02
4	+	Milk	*	2.3753	+	Milk	*	2.24
5	+	Poultry available	*	6.0115	+	Poultry available	*	5.67
6	+	Eggs available	*	1.8764	+	Eggs available	*	1.77
7	+	Alcohol	*	0.3585	+	Alcohol	*	0.34
8	+	Sugary beverages	*	1.0532	+	Sugary beverages	*	0.99
9	+	Corn available	*	0.0416	+	Corn available	*	0.04
10	+	Potatoes available	*	0.7654	+	Potatoes available	*	0.72
11	+	SFA	*	8.9947	+	SFA	*	8.48
12	+	PUFAs	*	4.1245	+	PUFAs	*	3.89
13	+	TFA	*	0.2848	+	TFA	*	0.27
14	+	Fruits	*	3.1683	-	Fruits	*	2.99
15	+	Vegetables	*	4.3224	-	Vegetables	*	4.08
16	+	Nuts and seeds	*	0.3923	-	Nuts and seeds	*	0.37
17	-	Whole grains	*	0.4751	-	Whole grains	*	0.45
18	-	Legumes	*	1.5789	-	Legumes	*	1.49
19	-	Rice available	*	9.1354	-	Rice available	*	8.61
20	-	Sweet potatoes available	*	0.1029	-	Sweet potatoes available	*	0.10
21	-	Physical activity METs	*	7.9004	-	Physical activity METs	*	7.45
22	-	Child underweight by >2SD	*	21.6105	-	Child underweight by >2SD	*	20.38
23	+	Sex male 1 and female 2	*	6.3788	+	Sex male 1 and female 2	*	6.02
24		Sum of parameter estimates		85.8512		Sum of parameter estimates		80.96
25		r of unadjusted BMI formula		0.9266		r of adjusted BMI formula		0.8998
26		R^2 of unadjusted BMI formula		0.8585		R^2 of adjusted BMI formula		0.8096
27						Sum percent weights (J24) / unadjusted sum percent weights (H24)		0.9430
28		H26 * 100 to convert to total percent weights=H24		85.8512		J26 * 100 to convert to percent weights=J24		80.96

Supplementary Table 6. BMI trend versus risk factor trends formula derivation (all variables standardized)

Row		Column A BMI trend and risk factor trends (change/year)		Column B Mean		Column C BMI trend v RF trends R^2		Column D Adjustment for fatty acids		Column E Multiple regression parameter estimates		Column F Column B * Column C * Column D * Column E
		BMI kg/M^2 trend		0.07603								
1	-	**Processed meat KC/d trend**	*	0.0513	*	0.0001	*	1.00	*	1.1427	=	0.0000
2	+	**Red meat KC/d trend**	*	0.5556	*	0.1682	*	1.00	*	1.1427	=	0.1068
3	-	**Fish KC/d trend**	*	0.2244	*	0.0251	*	1.00	*	1.1427	=	0.0064
4	+	**Milk KC/d trend**	*	0.1901	*	0.0425	*	1.00	*	1.1427	=	0.0092
5	+	**Poultry KC/d trend**	*	1.1457	*	0.1768	*	1.00	*	1.1427	=	0.2315
6	+	**Eggs KC/d trend**	*	0.2996	*	0.2340	*	1.00	*	1.1427	=	0.0801
7	+	**Alcohol KC/d trend**	*	0.0307	*	0.0833	*	1.00	*	1.1427	=	0.0029
8	+	**Sugary beverages KC/d trend**	*	0.2890	*	0.0156	*	1.00	*	1.1427	=	0.0051
9	-	**Corn KC/d trend**	*	0.0284	*	0.0198	*	1.00	*	1.1427	=	0.0006
10	+	**Potatoes KC/d trend**	*	0.2461	*	0.0740	*	1.00	*	1.1427	=	0.0208
11	-	**SFA KC/d trend**	*	0.2101	*	0.0025	*	0.46	*	1.1427	=	0.0003
12	+	**PUFA KC/d trend**	*	0.9963	*	0.0820	*	0.46	*	1.1427	=	0.0429
13	+	**TFA trend**	*	0.2096	*	0.0044	*	0.46	*	1.1427	=	0.0005
14	+	**Fruits KC/d trend**	*	0.6234	*	0.2317	*	1.00	*	1.1427	=	0.1651
15	+	**Vegetables KC/d trend**	*	1.3992	*	0.1570	*	1.00	*	1.1427	=	0.2511
16	+	**Nuts and seeds KC/d trend**	*	0.2762	*	0.0150	*	1.00	*	1.1427	=	0.0047
17	+	**Whole grains KC/d trend**	*	0.1729	*	0.0108	*	1.00	*	1.1427	=	0.0021
18	-	**Legumes KC/d trend**	*	0.3696	*	0.0886	*	1.00	*	1.1427	=	0.0374
19	+	**Rice KC/d trend**	*	0.7869	*	0.0886	*	1.00	*	1.1427	=	0.1389
20	+	**Sweet potatoes KC/d trend**	*	0.4603	*	0.0886	*	1.00	*	1.1427	=	0.1311
21	+	**Physical activity METs trend**	*							*	0.20825	=	0.20825
22	-	**Child underweight >2SD trend**	*							*	0.03926	=	0.03926
23												1.48519

Supplementary Table 6. BMI trend versus risk factor trends formula derivation continued

Row		Column G BMI trend and risk factor trends		Column H Preliminary risk factor coefficients (Column F repeated)		Column I BMI trend and risk factor trends		Column J Column H: H27 * (H1-H22) = BMI trend formula percent weights
	+	BMI kg/M^2 trend		0.07603	+	BMI kg/M^2 trend		0.0760
1	-	Processed meat KC/d trend	*	0.00000	-	Processed meat KC/d trend	*	0.00
2	+	Red meat KC/d trend	*	0.10681	+	Red meat KC/d trend	*	2.52
3	-	Fish KC/d trend	*	0.00644	-	Fish KC/d trend	*	0.15
4	+	Milk KC/d trend	*	0.00923	+	Milk KC/d trend	*	0.22
5	+	Poultry KC/d trend	*	0.23146	+	Poultry KC/d trend	*	5.46
6	+	Eggs KC/d trend	*	0.08011	+	Eggs KC/d trend	*	1.89
7	+	Alcohol KC/d trend	*	0.00292	+	Alcohol KC/d trend	*	0.07
8	+	Sugary beverages KC/d trend	*	0.00514	+	Sugary beverages KC/d trend	*	0.12
9	-	Corn KC/d trend	*	0.00064	-	Corn KC/d trend	*	0.02
10	+	Potatoes KC/d trend	*	0.02082	+	Potatoes KC/d trend	*	0.49
11	-	SFA KC/d trend	*	0.00027	-	SFA KC/d trend	*	0.01
12	+	PUFA KC/d trend	*	0.04291	+	PUFA KC/d trend	*	1.01
13	+	TFA trend	*	0.00048	+	TFA trend	*	0.01
14	+	Fruits KC/d trend	*	0.16508	+	Fruits KC/d trend	*	3.89
15	+	Vegetables KC/d trend	*	0.25107	+	Vegetables KC/d trend	*	5.92
16	+	Nuts and seeds KC/d trend	*	0.00474	+	Nuts and seeds KC/d trend	*	0.11
17	+	Whole grains KC/d trend	*	0.00213	+	Whole grains KC/d trend	*	0.05
18	-	Legumes KC/d trend	*	0.03741	-	Legumes KC/d trend	*	0.88
19	+	Rice KC/d trend	*	0.13891	+	Rice KC/d trend	*	3.28
20	+	Sweet potatoes KC/d trend	*	0.13109	+	Sweet potatoes KC/d trend	*	3.09
21	+	Physical activity METs trend	*	0.20825	+	Physical activity METs trend	*	4.91
22	-	Child underweight >2SD trend	*	0.03926	-	Child underweight >2SD trend	*	0.93
23		Sum		1.48519		Total percent weights		35.02
24		BMI trend formula r		0.59178				
25		BMI trend formula R^2		0.35020				
26		BMI trend formula R^2 (H25)/sum (H23)		0.23579				
27		H26 * 100 for percent weights		23.57947				

Appendix 1. Bradford Hill causality criteria based assessment methodology for BMI formula detailed

The Bradford Hill causality criteria, enumerated by the English occupational physician and epidemiologist Sir Austin Bradford Hill, are the gold standard assessment tools to test causality of risk factors related to health outcomes.[3] The relevant causality criteria included #1 strength, #2 experimentation, #3 consistency, #4 dose-response (biological gradient), #5 temporality, #6 analogy, #7 plausibility, (8) specificity, and #9 coherence.

A literature search revealed no published methodological precedents for statistically modeling and validating the relationship between mean cohort BMIs of worldwide countries and subnational regions/provinces/states and their corresponding dietary and other risk factors. In considering many candidate methodologies involving univariate and multiple regression analysis, we sought a methodology with good performance with as many Bradford Hill causality criteria as possible. Of the candidate statistical modeling strategies, the methodology for deriving the BMI multiple regression formula that will be detailed in Appendix 2 functioned best. So, after deriving the BMI formula, we formulated the following Bradford Hill causality criteria scoring methodology for this unprecedented purpose.

Bradford Hill causal criteria testing methodology applied to current study

The nine original Bradford Hill criteria were each scored as:

- "5" very strongly supporting causality,
- "4" strongly supporting causality,

• "3" moderately strongly supporting causality,

• "2" supporting causality,

• "1" weakly supporting causality, and

• "0" not supporting causality

The scoring for each Bradford Hill causal criterion was as follows:

1. Strength: The correlation coefficient, r, of the worldwide multiple regression derived

BMI formula with BMI (dependent variable) and BMI risk factors (independent

variables) assessed strength.

Scoring of strength:

5=BMI formula regressed with BMI r $\geq$0.50 and p<0.0001

4=BMI formula regressed with BMI 0.50>r$\geq$0.40 and p<0.0001

3=BMI formula regressed with BMI 0.40>r$\geq$0.30 and p<0.0001

2=BMI formula regressed with BMI 0.30>r$\geq$0.20 and p<0.0001

1=BMI formula regressed with BMI 0.20>r$\geq$0.10 and p<0.0001

0=BMI formula regressed with BMI r< 0.10 or $p$$\geq$0.0001

2. Experiment: Dr. Hill thought that evidence drawn from experimentation, including in

epidemiologic studies, may lead to the strongest support for causal inference.[3] We used

the bootstrap method to test Hill's "experiment" criterion. Random number generation of

20 subgroups each with 100 cohorts (with replacements) derived 20 standardized BMI

formulas to compare with the standardized worldwide BMI formula.

Scoring of experiment:

5=In all 20 bootstrap trials each with 100 random cohorts, RCT1-RCT20 of BMI formulas

regressed with RCT1-RCT20 of BMIs, respectively: r $\geq$0.50 and *p*<0.0001

4= In at least 15 out of 20 bootstrap trials each with 100 random cohorts, RCT1-RCT20 of BMI

formulas regressed with RCT1-RCT20 of BMIs, respectively: r $\geq$0.50 and *p*<0.0001.

3= In at least 15 out of 20 bootstrap trials each with 100 random cohorts, RCT1-RCT20 of BMI

formulas regressed with RCT1-RCT20 of BMIs, respectively: r $\geq$0.40 and *p*<0.0001.

2= In at least 10 out of 20 bootstrap trials each with 100 random cohorts, RCT1-RCT20 of BMI

formulas regressed with RCT1-RCT20 of BMIs, respectively: r $\geq$0.30 and *p*<0.0001.

1= In at least 5 out of 20 bootstrap trials each with 100 random cohorts, RCT1-RCT20 of BMI

formulas regressed with RCT1-RCT20 of BMIs, respectively: r $\geq$0.30 and *p*<0.0001.

0=None of the above.

3. Consistency: For the purposes of this study, consistency between BMI and BMI formula

output was determined by comparing the mean BMI and the mean BMI formula output in

each of the following 37 subgroups:

A. We divided the world's population by quartiles of socio-demographic index (SDI)—

see Supplementary Table 1 for definition of SDI.

B. A variable, "continents," allowed for analyses of the cohorts from countries from

each of the six inhabited continents.

C. The four countries (UK, USA, Mexico, and Japan) with subnational data on BMI and

the risk factors were grouped together to compare the BMI formula output with the

overall mean BMI in those four countries.

D. Based on the total kcal/day of the 13 foods that increased BMI (six animal foods,

sugary beverages, alcohol, corn, potatoes, SFA, PUFA, and TFA), a combination

variable was constructed and the world's population divided into quartiles from the

highest to lowest total kcal/day of BMI increasing foods.

E. Similarly to D above, based on the total kcal/day of all seven foods that decreased

BMI (fruits, vegetables, nuts and seeds, whole grains, legumes, rice, and sweet

potatoes), we divided the world's population into quartiles from the highest to lowest

in BMI decreasing foods (kcal/day).

F. Based on physical activity (METs/week), we divided the world's population into

quartiles from the highest to the lowest.

G. We evaluated dose response by dividing the BMI formula output into quartiles from

the highest to lowest.

H. The four countries with subnational data were individually evaluated.

I. The first four of the 20 random number generated database subgroups were included

in the consistency analysis.

J. Male and female cohorts were individually assessed.

K. We assessed dose-response with the BMI formula outputs after the BMI formula was

harmonized with the mean and SD of worldwide mean BMI. BMI formula outputs

were divided into quartiles.

For each of the 37 subgroups, the absolute differences between the means of BMI and the

BMI formula output were totaled (e.g., continent Africa BMI formula output – mean BMI

for Africa, etc.).

Scoring of consistency for mean cohort BMI compared with BMI formula output for each

of the 37 subgroups:

5=The mean of the absolute differences between mean BMI and BMI formula output was ≤ 0.30

kg/M^2.

4=The mean of the absolute differences between mean BMI and BMI formula output was ≤ 0.40

kg/M^2.

3=The mean of the absolute differences between mean BMI and BMI formula output was ≤ 0.50

kg/M^2.

2=The mean of the absolute differences between mean BMI and BMI formula output was ≤ 0.60

kg/M^2.

1=The mean of the absolute differences between mean BMI and BMI formula output was ≤ 1.0

kg/M^2.

0=The mean of the absolute differences between mean BMI and BMI formula output was > 1.0

kg/M^2.

4. Dose-response (biological gradient): Dr. Hill thought that a clear dose-response effect on

the incidence of disease with exposure to a single risk factor was the clearest evidence of

a causal relationship. In this analysis levels of a multivariable regression derived BMI

formula outputs in quartiles were related to mean BMIs in those quartiles (Table 3.

subgroups #34-#37).

As with consistency, scoring of dose-response was based on this mean absolute

difference of BMI and BMI formula output when the BMI formula output was divided

into quartiles:

5= The BMI formula output versus mean BMI absolute differences from each of the four

quartiles averages ≤ 0.30 kg/M^2.

4= The BMI formula output versus mean BMI absolute differences from each of the four

quartiles averages ≤ 0.40 kg/M^2.

3= The BMI formula output versus mean BMI absolute differences from each of the four

quartiles averages ≤ 0.50 kg/M^2.

2= The BMI formula output versus mean BMI absolute differences from each of the four

quartiles averages ≤ 0.60 kg/M^2.

1= The BMI formula output versus mean BMI absolute differences from each of the four

quartiles averages ≤ 1.0 BMI units.

0= The BMI formula output versus mean BMI absolute differences from each of the four

quartiles averages > 1.0 kg/M^2.

5. Temporality: Dr. Hill said, "Temporality refers to the necessity that the cause precedes

the effect in time."[3] Dr. Hill was an occupational physician before the current availability

in nutritional epidemiology of data on trends over 28 years of 20 components of

worldwide diets along with the global BMI trend. Consequently, now it is fair to test

temporality by deriving a standardized multiple regression formula with BMI trend,

measured by the slope of the least squared regression line (LSRL) over 1990-2017, as the

dependent variable. The independent variables consisted of the LSRL trends over 1990-

2017 of the same dietary components, physical activity, and childhood underweight as in

the original BMI formula—but not sex. The signs of the risk factors in the BMI trend

formula were determined as follows:

a. If the sign of the risk factor in Table 4 (BMI basic trend statistics) was "+" and

506   the r was "+", then the sign in the BMI trend formula was +.

b. If the sign of the risk factor in Table 4 was "-" and the r was "-," then the sign in

the BMI trend formula was "+". If the risk factor mean value was trending down

and the correlation coefficient of the risk factor with the BMI trend was negative,

this risk factor tread would be in alignment with an up-trending BMI.

c. If the sign of the risk factor in Table 4 was "-" and the r was "+," then the sign in

the BMI trend formula was "-".

d. If the sign of the risk factor in Table 4 was "+" and the r was "-," then the sign in

the BMI trend formula was "-".

Supplementary Table 6 shows the derivation of the BMI trend formula with standardized

BMI trend (dependent variable) versus standardized risk factor trends (independent

variables). As with the derivation of the BMI formula (Supplementary Table 5), the

coefficients of the risk factors in the formula were adjusted to equate to the trend percent

weights.

Scoring of temporality: The multiple regression derived BMI trend formula output versus

the BMI trend r (correlation coefficient) determined the score:

$5 = r \geq 0.50$ and $p < 0.0001$.

$4 = 0.50 > r \geq 0.40$ and $p < 0.0001$.

$3 = 0.40 > r \geq 0.30$ and $p < 0.0001$.

$2 = 0.30 > r \geq 0.20$ and $p < 0.0001$.

$1 = 0.20 > r \geq 0.10$ and $p < 0.0001$.

$0 = r < 0.10$ or $p \geq 0.0001$

6. Analogy: High BMI was among the four metabolic risk factors that were strongly associated

with cardiovascular diseases, cancers, and other non-communicable diseases. The other

major metabolic risk factors for non-communicable diseases were high fasting plasma

glucose (FPG). low density lipoprotein cholesterol (LDL-C), and high systolic blood pressure

(SBP).

We tested analogy by the strength of the correlations of BMI with FPG, BMI with LDL-C,

and BMI with SBP in univariate analysis.

Scoring of analogy:

5=BMI correlated with all three other metabolic risk factors showed r$\geq$0.500, p<0.0001

539   for all three.

540   4= BMI correlated with the three other metabolic risk factors showed r$\geq$0.500, p<0.0001

541   for two of the three.

542   3= BMI correlated with the three other metabolic risk factors showed r$\geq$0.400, p<0.0001

543   for two of the three.

544   2= BMI correlated with the three other metabolic risk factors showed r$\geq$0.300, p<0.0001

545   for two of the three.

546   1=BMI correlated with the three other metabolic risk factors showed r$\geq$0.200, p<0.0001

547   for two of the three.

0= None of the above.

7. Plausibility: To test plausibility that the BMI formula could accurately predict mean BMI

of cohorts, we looked to find if any of our findings were at variance with the

preponderance of studies published. We searched the medical literature particularly for

systematic reviews of the relationships of foods and other variables with BMI.

Scoring of plausibility:

5=None of the BMI formula risk factor percent weight coefficients was at variance with

the preponderance of the medical literature about risk factors for high BMI.

4= One of the BMI formula risk factor percent weight coefficients was at variance with

the preponderance of the medical literature about risk factors for high BMI.

3= Two of the BMI formula risk factor percent weight coefficients were at variance with

the preponderance of the medical literature about risk factors for high BMI.

2=Three of the BMI formula risk factor percent weight coefficients were at variance with

the preponderance of the medical literature about risk factors for high BMI.

1=Four of the BMI formula risk factor percent weight coefficients were at variance with

the preponderance of the medical literature about risk factors for high BMI.

0= Five or more of the BMI formula risk factor percent weight coefficients were at

variance with the preponderance of the medical literature about risk factors for high BMI.

8. Specificity: Either the BMI formula derived is unique or not. Scoring of specificity:

5=The BMI formula derived is unique and can model no other metabolic risk factor, other

risk factor, or health outcome.

0= The BMI formula derived is not unique and can model at least one other metabolic risk

factor, other risk factor, or health outcome.

9. Coherence: According to Dr. Hill, "…cause and effect interpretation of our data should not

seriously conflict with the generally known facts of the natural history and biology of the

disease."[3] In this analysis of BMI associated with BMI formula estimates, coherence was the

numerical total score of the above eight relevant causality criteria each on a 0-5 scale. The

maximum score was 40.

Scoring of coherence:

5=Score on the first eight Bradford Hill causation criteria=35-40.

4=Score on the first eight Bradford Hill causation criteria=30-34.

3=Score on the first eight Bradford Hill causation criteria=25-29.

2=Score on the first eight Bradford Hill causation criteria=20-24.

1=Score on the first eight Bradford Hill causation criteria=15-24.

0=Score on the first eight Bradford Hill causation criteria<15.

583
584

Appendix 2. The derivation of the BMI formula

Population attributable fraction (PAF) derivation systematic literature review methodology by WHO and IHME

Because of wide variations in definitions of risk factors and health outcomes in reports of risk factor-health outcome correlations, the World Health Organization in 2002 published a new framework, termed comparative risk assessment (CRA), for quantifying deaths and burden of disease caused by risk factors.[15] Goals were to increase the rigor of population health estimates and to improve the compatibility of the estimates of different groups of investigators studying different populations. CRA can be divided into six key steps:[16]

1. Inclusion of risk–outcome pairs in the analysis;
2. estimation from literature reviews of relative risk as a function of exposure;
3. estimation from literature reviews of exposure levels and distributions;
4. determination from literature reviews of the counterfactual level of exposure, the level of exposure with minimum risk called the theoretical minimum risk exposure level (TMREL);
5. computation of population attributable fractions (PAFs) and attributable burden; and
6. estimation from literature reviews of mediation of different risk factors through other risk factors

Originally, most PAFs were calculated for single risk factors related to single health outcomes (e.g., cigarette smoking and lung cancer). This methodology has now evolved to encompass 550 risk factor-health outcome pairs in the latest risk factor article published by the IHME.[16] Additionally, multiple risk

factors with PAFs for the same health outcome have been reported. Data inputs for CRAs have depended

on systematic literature reviews rather than GBD data on risk factors and health outcomes.

For the purposes of risk factor-health outcome pair analysis, high BMI has been considered a risk factor

for cardiovascular outcomes, cancer outcomes, etc., but not itself a health outcome for modeling with

CRA.

Population attributable fraction (PAF) derivation AI analytics methodology for this analysis

The methodology in this paper is offered as an alternative to CRA to model PFAs for the risk factors for

BMI. The case will be made that PAFs can be derived directly from IHME GBD data. For validation, the

BMI formula derived was tested with the nine Bradford Hill causality criteria (Appendix 1).

Supplementary Table 5 shows the BMI multiple regression formula derivation with mean cohort BMI

(dependent variable) versus

1. a 20 dietary risk factor combination variable,

2. physical activity,

3. child underweight, and

4. sex (independent variables).

To maximize the functionality of the BMI formula when tested with the nine Bradford Hill causality

criteria, all 20 dietary variables underwent the following four empirically derived adjustments:

1. multiplied times their mean kcal/day values (Column B),

2. multiplied times their R^2 values in univariate correlation with BMI (Column C),

3. SFA, PUFA, and TFA multiplied times 0.46, an adjustment for fatty acids extracted (i.e.,

oils and solid fats) versus coming from within dietary foods (Column D) In determining

the portion of SFA, PUFA, and TFA added in addition to these fatty acids in the animal

and plant foods, we used an adjustment factor (fatty acids * 0.46), adapted from the

website "Our World in Data."[17] This adjustment factor differentiated the fatty acids in

individual foods (54% of the total) from the added fatty acids (46% of the total), and

4. the parameter estimates from the BMI versus risk factors multiple regression analysis

(Column F).

To derive the PAFs of each of the risk factors in the BMI formula, we used the following steps:

1. Assign the signs of the risk factors in Column A according to the signs of the risk factor

correlations with BMI, including leaving positive signs for fruits, vegetables, and nuts and seeds.

2. Calculate the adjusted dietary risk factor coefficients (Column F (F1-F23) = Column B (B1-B23)

* Column C (C1-C23 * Column D (D1=D23) * Column E (E1-E23)).

3. Copy Column F coefficients (F1-F23) to Column H (H1-H23) for a further adjustment.

4. Multiply the copied coefficients in Column H (H1-H23) times the R^2 value of the entire BMI

formula correlated with BMI (F26=H26=0.8585) and divide the result by the sum of the 23

adjusted coefficients from step 3=1.4071 (F24) to get the BMI formula adjustment factor

(F26=0.6101).

5. Multiply (F26=0.6101) * 100 to give a multiplier (F27=61.01) to derive PAFs for the coefficients

in Column H (H1-H23).

6. Multiply (F27=61.01) times the coefficients in Column H to get the risk factor PAFs (H1-H23)

and the not multicollinearity adjusted BMI formula total percent weight (H24=85.85%).

Note that the above BMI formula contains the multicollinearities of fruits, vegetables, and nuts and

seeds (i.e., fruits, vegetables, and nuts and seeds correlate positively with 10 BMI increasing foods

and correlate negatively with the four other plant foods). Consequently, we need to adjust the BMI

formula to account for these multicollinearities by the following:

1. Add Column I and Column J that initially have the same BMI risk factors and coefficients as

Column G and Column H.

2. Switch the signs on fruits, vegetables, and nuts and seeds from "+" in Column H to "-" in

Column J to adjust for the multicollinearities.

3. Run the multicollinearities adjusted BMI formula in step 2 on SAS software.

4. Check the r of the resulting multicollinearity adjusted BMI formula correlated with BMI

(r=0.8998 (J25)).

5. Calculate the multicollinearity adjusted BMI formula R^2 by squaring the r (R^2=0.8096 (J26)).

6. Multiply J26=0.8096 times 100 to derive an adjusted total PAF for the final BMI formula

(J28=80.96%).

7. Divide the final BMI formula total percent weight (J28=80.96%) by the sum of the non-

collinearity adjusted BMI formula Column H coefficients (H24=85.85) to derive a BMI

formula coefficient multiplier (J27=0.9432) to account for the multicollinearities.

8. Multiply the Column J coefficients (J1-J23) by the multicollinearity adjusting BMI formula

coefficient multiplier (J27=0.9432) to derive the final BMI formula coefficients (J1-J23) as

PAFs, totaling the final BMI formula total percent weight (J24=80.96%).

9. Test the validity of the BMI formula with the nine Bradford Hill causality criteria (Appendix

1).

679
680

Article Information

Corresponding Author: David K. Cundiff, Independent researcher, davidkcundiff@gmail.com

- **Authors' contributions:** DKC acts as guarantor; conceived and designed the study, acquired and analysed the data, interpreted the study findings, drafted the manuscript, critically reviewed and edited the manuscript and tables, and approved the final version for publication.

  CW designed software programs in R to format and population weight the data, aided with the SAS statistical analysis, critically reviewed the manuscript, and approved the final version for publication.

- **Conflict of Interest Disclosures:** None reported. Both authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

- **Funding/Support:** This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. The Bill and Melinda Gates Foundation funded the acquisition of the data for this analysis by the IHME. The data were provided to the authors as volunteer collaborators with IHME.

-

  Role of the Funder/Sponsor: While IHME GBD faculty and staff by virtue of the Bill and Melinda Gates Foundation grants provided the raw data for this analysis, they did not vet the analysis or sponsor the manuscript.

706
Additional Contributors: Martin Sebera, from the Department of Kinesiology, Faculty of Sports Studies

Masaryk University, Czech Republic, critiqued statistical aspects of the manuscript and provided useful

input. Pavel Grasgruber, from Masaryk University, Czech Republic, provided suggestions after reviewing

the manuscript. We thank Scott Glenn and Brent Bell from IHME who supplied us with the GBD risk

factor exposure data for the risk factors and for BMI data.

Data sharing statement: The raw, unformatted data used in this analysis is now out of date. The 2019

GBD data on all the variables in this analysis may be obtained from the IHME by volunteer collaborating

researchers. The formatted database, SAS codes and Excel spreadsheets on which this analysis is based

are posted on the Mendeley data repository: https://data.mendeley.com/v1/datasets/publish-

confirmation/g6b39zxck4/6

STROBE checklist: This report follows the STrengthening the Reporting of OBservational studies in

Epidemiology (STROBE) guidelines for reporting global health estimates.[34]

722
723
724

References

1.   Murray C, Lopez A, Rodgers A, et al. The World Health Report 2002: Reducing Risks, Promoting Healthy Life. World Health Organization. https://apps.who.int/iris/bitstream/handle/10665/42510/WHR_2002.pdf

2.   Murray CJL, Aravkin AY, Zheng P, et al. Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet*. 2020;396(10258):1223-1249. doi:10.1016/S0140-6736(20)30752-2

3.   Ioannidis JPA. Implausible results in human nutrition research. *BMJ : British Medical Journal*. 2013;347doi:10.1136/bmj.f6698

4.   Ioannidis JA. The challenge of reforming nutritional epidemiologic research. *JAMA*. 2018;320(10):969-970. doi:10.1001/jama.2018.11025

5.   Artificial Intelligence (AI). IBM Cloud Education. Accessed August 7, 2021, https://www.ibm.com/cloud/learn/what-is-artificial-intelligence

6.   Hill AB. THE ENVIRONMENT AND DISEASE: ASSOCIATION OR CAUSATION? *Proc R Soc Med*. 1965;58(5):295-300.

7.   Global Burden of Disease Study 2017 (GBD 2017) Data Input Sources Tool. Institute of Health Metrics and Evaluation. Updated 2019. Accessed August 10, 2019, http://ghdx.healthdata.org/gbd-2017/data-input-sources

8.   GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017: Table of risk factor definitions. *The Lancet*. 2018;392(10159):1923-94.

9.   PROTOCOL FOR THE GLOBAL BURDEN OF DISEASES, INJURIES, AND RISK FACTORS STUDY (GBD) Version 3.0; Issue 26. Institute for Health Metrics and Evaluation. Updated February 2018. Accessed August 2, 2019, http://www.healthdata.org/sites/default/files/files/Projects/GBD/GBD_Protocol.pdf

10.   Naghavi M, Abajobir AA, Abbafati C, et al. Global, regional, and national age-sex specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet*. 2017;390 (10100):1151-1210.

11.   Flaxman AD, Lee YY, Vos T, et al. *An Integrative Metaregression Framework for Descriptive Epidemiology*. University of Washington Press; 2015. http://www.guilles.website/wp-content/uploads/2017/11/Metaregression-AFlaxman.pdf

12.   GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. . *The Lancet*. 2018;392(10159):1923-94.

13.   Call for Collaborators. . Institute for Health Metrics and Evaluation at the University of Washington. Updated 2019. Accessed April 25, 2019, http://www.healthdata.org/gbd/call-for-collaborators

14.   Omega 3 Fatty Acids: Fact Sheet for Health Professionals (The Office of Dietary Supplements) (2018).

15.   Nutritionix Track App. Syndigo LLC Updated 2019. Accessed April 25, 2019, https://www.nutritionix.com/

16.   POTATO PROCESSING AND USES. International Potato Center. Updated 2020.
Accessed May 28, 2020, https://cipotato.org/potato/potato-processing-uses/
17.   Ritchie H, Roser M. Diet Compositions. *Our World in Data*. 2017;
18.   Bloom JOJ. Bootstrap confidence intervals (Class 24, 18.05). MIT. Accessed July 25,
2021, https://ocw.mit.edu/courses/mathematics/18-05-introduction-to-probability-and-statistics-
spring-2014/readings/MIT18_05S14_Reading24.pdf
19.   Ruotsalainen H, Kyngäs H, Tammelin T, Kääriäinen M. Systematic review of physical
activity and exercise interventions on body mass indices, subsequent physical activity and
psychological symptoms in overweight and obese adolescents. *Journal of Advanced Nursing*.
2015;71(11):2461-2477. doi:10.1111/jan.12696
20.   Te Morenga L, Mallard S, Mann J. Dietary sugars and body weight: systematic review
and meta-analyses of randomised controlled trials and cohort studies. *BMJ* 2013;346:e7492.
doi:10.1136/bmj.e7492
21.   Traversy G, Chaput J-P. Alcohol Consumption and Obesity: An Update. *Curr Obes Rep*.
2015;4(1):122-130. doi:10.1007/s13679-014-0129-4
22.   Turner-McGrievy G, Mandes T, Crimarco A. A plant-based diet for overweight and
obesity prevention and treatment. *J Geriatr Cardiol*. 2017;14(5):369-374.
doi:10.11909/j.issn.1671-5411.2017.05.002
23.   United States Department of Agriculture FY 2019 BUDGET SUMMARY (US Office of
Budget and Program Analysis) (2019).
24.   Why is Government Subsidizing Junk Food?--22.6 percent of SNAP dollars are spent on
unhealthy food. Foundation for Economic Education. Updated February 25, 2017. Accessed
August 11, 2019, https://fee.org/articles/why-is-government-subsidizing-junk-food/
25.   Wakamo B. We Subsidize the Wrong Kind of Agriculture. Inequality.org. Updated JUNE
21, 2018. https://inequality.org/great-divide/subsidize-wrong-kind-agriculture/
26.   Fruits and Veggies Now Chock Full of Marketing Power (2017).
27.   Darmon N, Drewnowski A. Contribution of food prices and diet cost to socioeconomic
disparities in diet quality and health: a systematic review and analysis. *Nutrition reviews*.
2015;73(10):643-660. doi:10.1093/nutrit/nuv027
28.   Ebbeling CB, Feldman HA, Klein GL, et al. Effects of a low carbohydrate diet on energy
expenditure during weight loss maintenance: randomized trial. *BMJ*. Nov 14 2018;363:k4583.
29.   FORESIGHT--Tackling Obesities: Future Choices –Follow Up Actions. Government
Office for Science (UK). Updated 2019. Accessed August 5, 2019,
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file
/295885/07-1526-obesity-action-plan.pdf
30.   Mărginean CO, Mărginean C, Meliț LE. New Insights Regarding Genetic Aspects of
Childhood Obesity: A Minireview. *Front Pediatr*. 2018;6:271-271.
doi:10.3389/fped.2018.00271
31.   Castaner O, Goday A, Park Y-M, et al. The Gut Microbiome Profile in Obesity: A
Systematic Review. *Int J Endocrinol*. 2018;2018:4095789-4095789. doi:10.1155/2018/4095789
32.   Monteiro CA, Moubarac J-C, Levy RB, Canella DS, Louzada MLdC, Cannon G.
Household availability of ultra-processed foods and obesity in nineteen European countries.
Public Health Nutrition. 2017;21(1):18-26. doi:10.1017/S1368980017001379
33.   Monteiro CA, Cannon G, Levy RB, et al. Ultra-processed foods: what they are and how
to identify them. *Public Health Nutrition*. 2019;22(5):936-941.
doi:10.1017/S1368980018003762

34.   Stevens GA, Alkema L, Black PRE, et al. Guidelines for Accurate and Transparent
Health Estimates Reporting: the GATHER statement. *The Lancet*. 2016;388(10062):e19-e23.
816