Body mass index multiple regression formula testable by eight Bradford Hill causality criteria: Worldwide ecological cohort data analysis to inform dietary guidance

David K Cundiff (0000-0002-3206-9665), independent researcher,[1,3] Chunyi Wu (0000-0002-2186-3433), Research Epidemiologist/Statistician[2,3]

[1] Long Beach, California, USA

[2] Area Specialist Lead in Epidemiology and Statistics, Michigan Medicine, Ann Arbor, Michigan, USA

[3] Volunteer collaborators with the Institute of Health Metrics and Evaluation, Seattle, Washington, USA

Correspondence to: David K Cundiff davidkcundiff@gmail.com Phone: 01-562-438-8805

Word count: abstract=399, text=3603

Abstract

Objectives (1) Derive a worldwide body mass index (BMI) multiple regression formula (BMI

formula), (2) Quantify the percent weights attributable to dietary and other risk factors from this

BMI formula, and (3) Test relevant dietary guidelines and diets using the BMI formula generated

BMI estimates

Design BMI and risk factor cohort data from 1990-2017 from the Institute of Health Metrics and

Evaluation (IHME) were formatted and population-weighted. We empirically explored the

univariate and multiple regression correlations of risk factors with BMI to maximize BMI

formula functionality.

Setting Worldwide

Ecological cohort data Over 12,000 Global Burden of Disease (GBD) risk factor surveys of

people 15-69 years old from 195 countries analysed and synthesised into representative cohort

risk factor values by staff and volunteer collaborators with the IHME

Main outcome measures Performance of a worldwide BMI formula when tested with eight

Bradford Hill causality criteria, each scored with 0-5 scale: 0=negative to 5=very strong support

(40 possible score)

Results In the BMI formula derived, all foods were expressed in kilocalories/day (kcal/day), and

all risk factor coefficients were adjusted to equate with their percent weight impacts on BMI.

BMI formula =0.29*processed meat +11.87*red meat +0.02*fish +3.74*milk +11.09*poultry

+5.46*eggs +1.95*alcohol +6.29*sugary beverages +0.29*corn +3.53*potatoes +2.71 *

saturated fatty acids+ 0.64*polyunsaturated fatty acids+0.06*trans fatty acids-0.37*fruit -

0.52*vegetables -0.03*nuts and seeds -0.12 *whole grains -0.47*legumes-8.26*rice -18*sweet

potatoes -9.93*physical activity (METs/week) +5.78*total kcals/day available-4.13*child

underweight+0.92*discontinued breast feeding. BMI increasing foods have + signs in the BMI

formula and BMI decreasing foods have - signs. Bradford Hill causality criteria test scores (0-5):

(1) strength=5, (2) consistency=5, (3) dose-response=5, (4) temporality=5, (5) analogy=4, (6)

plausibility=5, (7) experimentation=5 and (8) coherence=5. Total=39/40. Based on the USA

BMI trend data, the mean adult BMI in 1990 was 25.45, is projected to be 28.13 in 2020, and

31.81 in 2050. BMI formula based mean BMI estimates included the following: Dietary

Guidelines for Americans=26.34, USA with 50% reduction of BMI increasing food intake

=23.34, USA with 25% reduction of BMI increasing food intake and increasing physical activity

by running at 6 mph for 1 hour/day on average=23.67, vegetarian diet=22.54, low carbohydrate,

high fat diet=31.76.

Conclusions Eight relevant Bradford Hill causality criteria strongly supported that the BMI

formula derived was causally related to mean BMIs of worldwide cohorts, subgroups, and

individual risk factor patterns. The statistical analysis methodology introduced could inform

individual, clinical, and public health strategies regarding overweight/obesity

prevention/treatment and other health outcomes.

Introduction

The scientific validity of the Dietary Guidelines for Americans for 2015-2020 was challenged by Journalist Nina Teicholz in the *BMJ*.[1] The Center for Science in the Public Interest called for the *BMJ* to retract the article. The peer reviewers selected to adjudicate the far reaching dispute concluded that, "Teicholz's criticisms of the methods used by Dietary Guidelines for Americans Committee are within the realm of scientific debate."[2] Regarding scientific study of treatments of overweight/obesity, *JAMA* published a representative randomised trial comparing a low fat diet with a low carbohydrate diet. It reported no difference in the very modest weight loss achieved at one year.[3] This suggests that current short term dietary intervention trials (< 5 years) are meaningless in determining the long-term relationship of diet (> 20 years) with overweight/obesity.

Influential Stanford University meta researcher, Dr. John Ioannidis, called for radical reform of all nutritional epidemiology methodologies used to influence food/agricultural policies and to produce dietary guidelines for clinicians and the public.[4] Currently, no methodology for relating body mass index (BMI) long-term or BMI change/year to food intake and physical activity has been generally accepted as rigorous, replicable, and scientifically valid. This study analysing worldwide data for dietary and other risk factors for BMI attempts to answer Dr. Ioannidis' call for a more rigorous and reliable nutritional epidemiology methodology to base public health policies and individual dietary guidance.

The first objective of this analysis was to derive a multiple regression risk factor formula using worldwide ecological data on BMIs of male and female cohorts (dependent variable) and dietary

and other risk factors (independent variables). Secondly, we used Bradford Hill causality criteria

to test the BMI formula containing worldwide diet and other risk factors. Satisfying Bradford

Hill causality criteria is considered validating in epidemiological research.[5] Third we wanted to

apply the BMI formula to assorted diet and other risk factor patterns to derive mean BMI

estimates associated with long-term (>20 years) following of those diet and other risk factor

patterns. We only hypothesised that the objectives were achievable.

Methods

As volunteer collaborators with the Institute of Health Metrics and Evaluation (IHME), we

received raw Global Burden of Disease (GBD) ecological data (≈1.4 Gigabytes) on mean BMIs

of male and female cohorts 15-49 years old and 50-69 years old from each year 1990-2017 from

195 countries and 365 subnational locations (n=1120 cohorts). We also utilised GBD data on

exposures to 32 risk factors and covariates potentially related to BMI. IHME dietary covariate

data originally came from Food and Agriculture Organization surveys of animal and plant food

commodities available percapita—as opposed to consumed per capita—in countries worldwide.[6]

Food risk factors came as g/day consumed percapita. Other variables were utilised as part of

deriving the BMI formula and testing the BMI formula according to the Bradford Hill causality

criteria: physical activity (METs/day), kilocalories per day available percapita (kcal/day, a

covariate), severe child underweight (2 SD below the mean weight for age), and discontinued

breast feeding before 6 months. Other available variables tested included socio-demographic

index, LDL-cholesterol, fasting plasma glucose and, systolic blood pressure. Supplementary

Table 1 lists the relevant GBD risk factors, covariates, and other available variables with

definitions of those risk factor exposures.[7]

GBD worldwide citations of over 12,000 surveys constituting ecological data inputs for this

analysis are in Appendix 1.[6,8] The main characteristics of IHME GBD data sources for BMI and

all risk factor values have been published by IHME GBD data researchers and discussed

elsewhere.[9-11] These included detailed descriptions of categories of input data, potentially

important biases, and methodologies of analysis. We did not clean or pre-process any of the

GBD data. GBD cohort risk factor and BMI data from the IHME had no missing records.

Because this was a post hoc analysis and the GBD data for the study came from IHME, no ethics

committee approval or institutional review board review was needed for this statistical analysis.

The raw data for this analysis may be obtained by volunteer researchers collaborating with

IHME.[12]

IHME made available GBD BMI, risk factor, and covariate exposure data for each year from

1990-2017 for male and female cohorts from ages 15-49 years old and from 50-69 years old. To

maximally utilise the available data, we averaged the values for ages 15-49 years old together

with 50-69 years old for BMI and for each risk factor exposure for each male and female cohort

for each year. Finally, for each male and female cohort, data from all 28 years (1990-2017) on

mean BMI and on each of the risk factor exposures were averaged using the computer software

program R.

To weigh the country and subnational data according to population, internet searches (mostly

Wikipedia) yielded the most recent population estimates for countries and subnational states,

provinces, and regions. World population data from the World Bank and the Organisation for

Economic Co-operation and Development could not be used because they did not include all 195

countries or any subnational data.

Using the above described formatted dataset of risk factors, covariates, and BMIs, a software

program in R generated a population-weighted analysis dataset. Each male or female cohort in

the population-weighted analysis dataset represented approximately 1 million people (range: <

100,000 to 1.5 million). The analysis dataset had n=7886 cohorts (rows of data), half male and

half female, representative of over seven billion people. For example, India with about 1.234

billion people had 617 rows of the same data for males and 617 rows for females. Maldives, with

about 445,000 people, had a single row of data for males and another for females. Without

population-weighting the data, cohorts in India and Maldives each would have had one row in

the analysis dataset, invalidating the analysis results.

This report follows the STrengthening the Reporting of OBservational studies in Epidemiology

(STROBE) guidelines for reporting global health estimates.[13]

Supplementary Table 2 details how omega-3 fatty acid g/day was converted to fish g/day using

data on the omega-3 fatty acid content of frequently eaten fish from the National Institutes of

Health Office of Dietary Supplements (USA).[14] As shown in Supplementary Table 3, we

converted all of the animal and plant food data, including alcohol and sugary beverage

consumption, from g/day to kcal/day. For the g to kcal conversions, we used the Nutritionix track

app,[15] which tracks types and quantities of foods consumed. Saturated fatty acids risk factor (0-1

portion of the entire diet) was not available with GBD data from 2017, so GBD saturated fatty

acids risk factor data from 2016 was used. Polyunsaturated fatty acid and trans fatty acid GBD

risk factor data from 2017 (0-1 portion of the entire diet) was also utilised. These fatty acid data

were converted to kcal/day by multiplying by the kcal/day available for each cohort.

Statistical methods

To determine the strengths of the risk factor correlations with mean BMIs of population

weighted worldwide cohorts (7886 cohorts) or subgroups of cohorts (e.g., continents, socio-

demographic quartiles, etc.), we utilised Pearson correlation coefficients: r, 95% confidence

intervals (95% CIs), and *P* values.

In this first of its kind, post hoc, exploratory analysis; the methodology was determined as we

proceeded by experimenting with strategies to optimise the functioning of the BMI formula

derived. We sought to derive a BMI multiple regression formula from worldwide data with the

following characteristics:

1. including as many as possible of the available animal and plant food variables,

2. including physical activity and other plausibly informative available ecological variables,

and

3. combining dietary variables if appropriate to utilise more variables and improve BMI

formula strength (i.e., R^2 with BMI) and consistency at predicting subgroup mean BMIs.

Appendix 2 further details the statistical methodology and explains the use of Bradford Hill

causality criteria to assess whether the worldwide BMI was causally related to the risk factors in

the BMI formula. Briefly, we tested the BMI formula output with the Bradford Hill causality

criteria (1) strength, (2) consistency, (3) dose response, (4) temporality, (5) analogy, (6)

plausibility, (7) experimentation, and (8) coherence. For each criterion, we used a 0-5 scale to

assess the magnitude of support of the BMI formula output being causally related to the BMIs

worldwide (0=no support of causality to 5=very strong support of causality).

In determining the variables to include and exclude in worldwide BMI formula, we set the

statistical threshold for a variable to enter and to remain in the formula at $P < 0.25$. We used

SAS and SAS Studio statistical software 9.4 (SAS Institute, Cary, NC) for the data analysis.

Results

Table 1 shows the basic statistics and univariate correlations of mean BMI with dietary and other risk factors and with the BMI formula derived from worldwide cohorts. Table 2 shows the univariate correlations of each of the 24 BMI formula risk factors with each other. All six animal foods (processed meat, red meat, fish, milk, poultry, and eggs), alcohol, sugary beverages, corn availability, potato availability, saturated fatty acids, polyunsaturated fatty acids, and trans fatty acids all positively correlated with BMI. Table 2 shows that corn availability (kcal/day percapita, a covariate) correlated moderately strongly with sugary beverages (r=0.419, 95% CI 0.400 to 0.437, $P<0.0001$), suggesting that high fructose corn syrup may account for the positive correlation with BMI. Potato availability (kcal/day percapita, a covariate), which positively correlated with BMI, included ≥50% highly processed potato products worldwide according to the International Potato Center.[16] Four starchy plant foods (whole grains, legumes, rice, and sweet potatoes) all negatively correlated with BMI.

The strong positive correlations of fruits, vegetables, and nuts and seeds with BMI were unexpected since nutritional literature broadly supports these foods as non-obesogenic. It suggested possible multicollinearity (when an independent variable is highly correlated with one or more known or unknown other independent variables). Supplementary Table 4 demonstrates the multicollinearity of fruits, vegetables, and nuts and seeds with BMI increasing foods:

- BMI increasing foods (animal foods + alcohol + sugary beverages + corn + potatoes) as a group were very strongly correlated with BMI (r= 0.597, 95% CI 0.583 to 0.611, $P< 0.0001$).

204   •   Fruits, vegetables, nuts and seeds grouped together were strongly positively correlated

205   with BMI (r= 0.655, 95% CI 0.642 to 0.677, *P*< 0.0001) but also with BMI increasing

206   foods (r= 0.323, 95% CI 0.304 to 0.343, *P*< 0.0001).

Supplementary Table 4 also shows that the top sociodemographic index (SDI) cohort quartile's

mean BMI was about 3.6 BMI units higher than the three lowest SDI quartiles (24.54 versus

20.88), meaning that the more developed countries had much higher mean BMIs. BMI increasing

foods in the top SDI quartile exceeded intake in the three lowest quartiles by 58% (905/kcal/day /

571 kcal/day=1.58). Fruits, vegetables, and nuts and seeds in the top SDI quartile exceeded

intake in the three lower quartiles by 1.9 to 1 (199/kcal/day / 106 kcal/day=1.88). Because of the

multicollinearities of fruits, vegetables and nuts and seeds with the 10 BMI increasing foods, we

grouped fruits, vegetables and nuts and seeds with BMI decreasing foods (including whole

grains, legumes, rice, and sweet potatoes) in a combination variable used to derive the BMI

formula, gives the coefficients of risk factors equated to their percent weights (shown in

Supplementary Table 5):

219   BMI formula =0.29*processed meat +11.87*red meat +0.02*fish +3.74*milk

220   +11.09*poultry +5.46*eggs +1.95*alcohol +6.29*sugary beverages +0.29*corn

221   +3.53*potatoes +2.71 * saturated fatty acids+ 0.64*polyunsaturated fatty

222   acids+0.06*trans fatty acids-0.37*fruit -0.52*vegetables -0.03*nuts and seeds -

223   0.12 *whole grains -0.47*legumes-8.26*rice-0.18*sweet potatoes -9.93*physical

224   activity (METs/week) +5.78*total kcals/day available-4.13*child

225   underweight+0.92*discontinued breast feeding. Percent weights of risk factors

226   were equated to risk factor coefficients. Total percent weights=78.64. (BMI

formula R^2=0.7864, r=0.887, 95% CI 0.882 to 0.891, *P*<0.0001, + signs mean

BMI increasing and - signs mean BMI decreasing).

As detailed in the methods and Appendix 2, the standardised BMI formula coefficients of the

dietary and other risk factors were all adjusted to equate to the percent weights of risk factors in

the BMI formula.

As shown in Supplementary Table 5, the 13 BMI increasing foods and seven BMI decreasing

foods underwent three adjustments: (1) multiplied times their mean kcal/day values (Column C),

(2) multiplied times their R^2 values in univariate correlation with BMI (Column E), and (3)

multiplied times the ratio of the mean kcal/day of the second, third, and fourth sociodemographic

index (SDI) quartiles divided by the mean kcal/day of the first SDI quartile (Column G). Table 3

demonstrates that these adjustments improved the consistency of the BMI formula as measured

by the closeness of fit of the multiple regression derived BMI formula outputs with the mean

BMIs of 37 subgroups of the worldwide data.

BMI formula output analysed by Bradford Hill causality criteria

Eight Bradford Hill causality criteria tested the functionality of the BMI formula. With the

methodology detailed in Appendix 2, we assessed the relationship of the worldwide BMI

formula output versus the mean BMI as follows:

1. Strength=5—The correlation coefficient of the BMI formula regressed with BMI: r=

0.887 (95% CI: 0.882 to 0.891), *P*< 0.0001.

2. Consistency score=5— Table 3 shows the average absolute differences between subgroup

mean BMIs and BMI formula estimates of the BMI formula compared with two other

variations of the BMI formula. The BMI formula with fruits, vegetables, and nuts and

seeds grouped with BMI decreasing foods as negatively correlated with BMI had a

slightly closer fit than a comparison BMI formula with fruits, vegetables, and nuts and

seeds grouped with BMI increasing foods positively correlated with BMI (average BMI -

BMI formula estimate (absolute difference): 0.249 versus 0.255 BMI units). Without the

three adjustments to each of the food variables, the average BMI - BMI formula estimate

absolute difference was 0.368 BMI units.

3. Dose-response (Biological gradient) score=5—Table 3 shows that the mean of the BMI

absolute differences between the BMI formula estimates and mean BMIs and in the four

dose-response quartiles was 0.302 BMI units.

4. Temporality score=5—a multiple regression with the worldwide BMI trend 1990-2007

(dependent variable) versus the risk factor trends (independent variables) generated a

BMI trend formula versus risk factor BMI trends: r= 0.615, (95% CI: 0.601 to 0.628),

P<0.0001). Fifteen out of 24 risk factor trends contributed to the BMI trend formula. The

derivation of standardised BMI trend versus risk factors trends formula with sign

concordant coefficients adjusted to equal the percent weights of the significant risk

factors shown in Supplementary Table 7 was as follows:

269
BMI trend formula=4.84 * Red meat + 0.63 * Milk + 4.61 * Poultry + 2.62 * Eggs + 3.91

* Alcohol + 2.69 * Sugary beverages + 0.40 * corn + 3.60 * Potatoes + 3.90 * PUFA -

0.65 * Legumes - 3.08 * Rice - 0.80 * Sweet potatoes + 6.06 * kcal available (total

percent weight 37.79, R^2=0.3779).

5. Analogy score=4—Systolic blood pressure had 15/24 risk factors concordant in

sign with BMI. SBP correlated weakly with the BMI formula (r= 0.024, 95% CI

0.002 to 0.046, *P* =0.0361). The low density lipoprotein cholesterol (LDL-C) and

the fasting plasma glucose (FPG) formulas had 23/24 and 21/24 risk factors

concordant in sign with the BMI formula, respectively. The LDL-C and the FPG

were both strongly correlated with BMI (r=0.757, 95% CI 0.747 to 0.766, *P* <

0.0001 and (r=0.565, 95% CI 0.550 to 0.580, *P* < 0.0001, respectively). They

were also strongly correlated with the BMI formula (r=0.757, 95% CI 0.747 to

0.766, *P* < 0.0001 and r=0.581, 95% CI 0.566 to 0.595), *P* < 0.0001, respectively.

6. Plausibility: Score=5—Based on systematic medical literature reviews, physical activity

inversely correlated with BMI[17] and BMI directly correlated with intakes of sugar,[18]

alcohol,[19] and animal foods.[20] The relationship of adult BMI with early childhood severe

underweight has not been reported worldwide. Since people in poor countries have less

animal foods and alcohol and more physical exercise than in developed countries, it is

plausible that childhood severe underweight positively correlates with lower BMI in

adulthood.

7. Experiment: Score=5—Table 4 demonstrates cross validation of the BMI formula with a

very high degree of reproducibility in the percent weight distributions from 20 randomly

selected subgroups of the dataset (n=100 for each, which are shown in Supplementary

Table 7).

8. Coherence: Score=5—As evidenced by the near perfect score 34/35 on the first seven

criteria, the Bradford Hill causality criteria overall were strongly supportive that the

worldwide BMI formula accurately modeled the 24 risk factors that led to increased and

decreased BMI—total causality criteria score=39/40.

Relative to other countries, the USA had one of the world's highest levels of kcal/day of BMI

increasing foods (exclusive of fatty acids), 1015 kcal/day and below average consumption of

BMI decreasing foods, 326 kcal/day, corresponding to an average adult BMI=26.66. Based on

the USA BMI trend data, the mean adult BMI in 1990 was 25.45 and is projected to be 28.13 in

2020, and 30.81 in 2050. Table 5 shows BMI formula estimates for various relevant patterns of

diet and/or other BMI formula risk factors. Following the Dietary Guidelines for Americans

2015 -2020 would result in a mean adult BMI of 26.34, which is almost the mean USA adult

BMI from 1990 - 2017 (26.66). A major increase in exercise on a regular basis alone would

decrease the BMI formula estimated mean long term BMI. In addition to the current mean level

of physical activity (3853 METs/week), running two hours/day on average at 6 miles per hour

(8400 METs/week)[21] would reduce the estimated mean BMI by 2.68 BMI units to 23.98.

Discussion

The eight Bradford Hill criteria used in this analysis of long term worldwide data all support that the BMI formula is causally related to adult BMI.

Short of extreme and unrealist for most people increases in exercise to control weight, dietary change from the USA current dietary patterns or even the USA dietary guidelines recommended diet would be required to reach the normal range for BMI (BMI $\geq$ 18.5 and BMI < 25). For example, shifting 50% of kcal/day from BMI increasing foods to BMI decreasing foods equates to a BMI formula estimate=23.26 BMI units over the long term. While following a low carbohydrate, high fat diet has been demonstrated to cause modest short term weight loss in obese people,[22] the BMI formula projects that the long-term effect (> 20 years) would be obesity (projected mean BMI=31.76).

The perhaps counterintuitive findings that fruits, vegetables, and nuts and seeds were positively correlated with BMI (r=0.655, 95% CI 0.642 to 0.677, $P < 0.0001$) may be largely explained by the multicollinearity between fruits, vegetables, and nuts and seeds and BMI increasing foods (r=0.323, 95% CI 0.304 to 0.343, $P < 0.0001$, Table 2).The high cost of fruits, vegetables, and nuts and seeds may account for much of this multicollinearity both worldwide and within wealthy countries that have high economic inequality levels like the USA. A systematic review of the literature on food cost relative to nutrient quality found that the median costs of starches (€0.14/100 kcal) was quite low relative to fruits and vegetables (€0.82/100 kcal), meat/eggs/fish (€0.64/100 kcal), fresh dairy (€0.32/100 kcal), and nuts (€0.25/100 kcal).[23]

The findings in this study, particularly those in Table 5, should influence government food policy

decisions. The Supplemental Nutrition Assistance Program (SNAP—formerly Food Stamps)

spends an estimated 22.6% of its $73 billion/year budget[26] on payments to low-income

Americans for "sweetened beverages, prepared desserts, salty snacks, candy, and sugar."[27]

Additionally, the US Department of Agriculture (USDA) subsidises crops that go primarily for

animal feed or that are processed into sugars while not subsidizing fruits and vegetables.[24] While

the USDA recognises the relatively low intake of fruits and vegetables in the USA and sponsors

a publicity campaign to increase fruits and vegetables,[25] USDA expenditures should promote

reduced prices of BMI decreasing foods and increased prices of BMI increasing foods.

The strong correlation of discontinuation of breastfeeding with BMI (r=0.802, 95% CI 0.794 to

0.810, P<0.0001) was in accord with a meta-analysis of breast feeding related to subsequent

childhood and adult BMI by Jeanne Stolzer.[26] However, the estimated benefit in reducing adult

BMI of breastfeeding in this study was very modest (0.19 BMI unit, see Table 5). Breastfeeding

for at least six months is recommended by the American Academy of Pediatrics[27,28] and the

National Institute of Clinical Excellence.[29]

Limitations

The GBD data on animal and plant foods were not comprehensive and comprised only 1199

kcal/day on average. Subnational data were available on only four countries. Because the data

formatting and statistical methodology were new, this was necessarily a post hoc analysis and no

pre-analysis protocol was possible. As detailed in the Foresight Report on obesity,[30] obesity is

affected by a complex system of interacting factors besides diet, physical activity, and breast

feeding. So genes,[31] gut microbiome,[32] ultra processing of food,[33,34] and other influences on BMI

were outside of the purview of this analysis.

Generalisability

360   Given the strength and consistency of the relationship of the BMI formula in modeling BMI

361   (SDI quartiles, continents, and subnationals, etc., see Tables 3 and 5), the findings should be

362   generalisable to people all over the world.

Conclusion

Eight Bradford Hill causality criteria strongly supported that the worldwide obesity epidemic is causally related to the 24 risk factors in proportion to their coefficients in the BMI formula. The findings in this study should be considered by health policymakers drafting dietary guidelines for healthy weight management. While this study deals only with dietary and other risk factors for BMI (overweight/obesity), the methodology introduced could easily apply to estimating percent weights of multiple dietary and other risk factors that pertain to dozens of non communicable diseases, for which the IHME have GBD data.

Table 1.BMI risk factor basic statistics and Bradford Hill criterion strength

BMI and risk factors: n=7886	Mean	Std Dev	Mini-mum	Maxi-mum	r	95% CI	*P*	R²
BMI kg/M2	21.79	2.31	17.95	29.39				
BMI formula	21.79	2.31	18.15	30.32	0.887	0.882 to 0.891	<.0001	0.791
Processed kcal/day	5.49	9.94	0.20	68.8	0.604	0.590 to 0.618	<.0001	0.365
Red meat kcal/day	50.70	45.45	3.21	235.9	0.662	0.649 to 0.674	<.0001	0.438
Fish kcal/day	10.01	36.43	0.40	370.4	0.103	0.081 to 0.124	<.0001	0.011
Milk kcal/day	25.37	27.39	1.06	146.8	0.684	0.672 to 0.696	<.0001	0.468
Poultry kcal/day avail	45.06	51.03	1.06	411.9	0.813	0.805 to 0.820	<.0001	0.661
Eggs kcal/day avail	19.47	14.76	0.79	69.6	0.688	0.676 to 0.699	<.0001	0.473
Alcohol kcal/day	81.71	58.06	4.25	430	0.166	0.145 to 0.188	<.0001	0.028
Sugary beverages kcal/day	297.65	152.33	73	1472	0.119	0.097 to 0.141	<.0001	0.014
Corn kcal/day avail	34.67	48.16	0.16	305.2	0.073	0.051 to 0.095	<.0001	0.005
Potatoes kcal/day avail	84.16	74.43	3.07	533.9	0.211	0.189 to 0.232	<.0001	0.044
Saturated FA % kcal available	191.27	64.49	70.79	481.1	0.710	0.699 to 0.721	<.0001	0.504
Polyunsaturated fatty acids kcal%	82.24	74.98	2.93	381.31	0.737	0.727 to 0.747	<.0001	0.543
Trans fatty acids % kcal	13.4	13.82	1.99	77.76	0.488	0.471 to 0.504	<.0001	0.238
Fruits kcal/day	40.39	22.59	3.58	161.4	0.622	0.609 to 0.636	<.0001	0.387
Vegetables kcal/day	80.14	43.36	9.48	304.2	0.520	0.503 to 0.536	<.0001	0.270
Nuts and seeds kcal/day	8.51	8.44	0.05	103.0	0.486	0.469 to 0.503	<.0001	0.237
Whole grains kcal/day	55.65	30.85	1.14	235.1	-0.201	-0.222 to -0.179	<.0001	0.040
Legumes kcal/day	51.66	32.16	0.51	194.7	-0.384	-0.403 to -0.365	<.0001	0.148
Rice kcal/day avail	141.23	116.38	1.42	461.8	-0.561	-0.576 to -0.545	<.0001	0.314
Sweet potatoes kcal/day avail	22.67	35.88	0.02	364.7	-0.152	-0.173 to -0.130	<.0001	0.023
Physical activity METs	4709	1367	1609	7669	-0.445	-0.462 to -0.427	<.0001	0.198
Kcal available/day	2579	421	1579	3898	0.843	0.837 to 0.850	<.0001	0.711
Child/infant underweight	0.19	0.17	0.00	0.53	-0.794	-0.802 to -0.786	<.0001	0.631
Discontinued breast feeding< 6 months	0.12	0.06	0.02	0.24	0.802	0.794 to 0.810	<.0001	0.644
Sodium g/day	4.45	2.33	1.33	9.21	-0.019	-0.041 to -0.003	0.099	0.000
Calcium g/day	0.3	0.18	0.08	1.04	0.771	0.762 to 0.779	<.0001	0.594
Dietary fiber g/day	9.22	3.15	2.72	22.68	0.310	0.290 to 0.330	<.0001	0.096
Sugar available kcal/day	447.1	283.8	48.3	1249	0.423	0.405 to 0.441	<.0001	0.179
Fasting plasma glucose mmol/dl	4.31	0.35	3.32	5.58	0.565	0.550 to 0.580	<.0001	0.32
Systolic BP mm/Hg	133.9	4.33	123.4	147.9	0.087	0.065 to 0.109	<.0001	0.008
LDL cholesterol mmol/dl	2.36	0.4	1.27	3.25	0.757	0.747 to 0.766	<.0001	0.573
Socio-demographic index	0.54	0.17	0.11	0.9	0.738	0.728 to 0.748	<.0001	0.544

378 **Table 2. Correlations between the risk factors that are in Table 1**

Variables (foods in kcal/day)	Process ed meat	Red meat	Fish	Milk	Poultry	Eggs	Alcohol
Processed meat	1.000						
Red meat	0.656	1.000					
Fish	0.281	0.118	1.000				
Milk	0.763	0.682	0.109	1.000			
Poultry	0.675	0.670	0.162	0.684	1.000		
Eggs	0.574	0.786	0.392	0.532	0.633	1.000	
Alcohol	0.592	0.478	0.292	0.402	0.372	0.352	1.000
Sugary beverages	-0.118	0.037	-0.163	0.059	0.126	0.014	-0.098
Corn	-0.052	-0.144	-0.018	-0.108	0.008	-0.100	-0.108
Potatoes	0.160	0.124	-0.058	0.202	0.104	0.032	-0.007
Saturated fatty acids	0.633	0.709	0.039	0.772	0.709	0.601	0.304
Poly unsaturated fatty acids	0.711	0.699	0.162	0.691	0.844	0.635	0.400
Trans fatty acids	0.479	0.203	-0.006	0.445	0.465	0.217	0.174
Fruits	0.391	0.380	0.099	0.499	0.613	0.374	0.075
Vegetables	0.308	0.392	0.271	0.362	0.360	0.519	0.209
Nuts and seeds	0.520	0.399	0.073	0.590	0.519	0.344	0.253
Whole grains	-0.022	-0.111	0.091	-0.250	-0.023	-0.076	0.065
Legumes	-0.199	-0.404	0.061	-0.197	-0.222	-0.418	0.133
Rice	-0.396	-0.357	-0.071	-0.568	-0.396	-0.277	-0.139
Sweet potatoes	-0.197	0.000	-0.075	-0.395	-0.242	0.045	-0.068
Physical activity METs	-0.215	-0.072	-0.163	-0.431	-0.397	-0.119	0.097
Kcal /day available	0.602	0.649	0.098	0.775	0.754	0.644	0.231
Child/infant 2SD underweight	-0.416	-0.737	-0.189	-0.409	-0.618	-0.804	-0.141
Discontinued breast feeding <6 mo	0.583	0.812	0.219	0.639	0.727	0.847	0.307

379
380
381
382 **Table 2. Correlations between the risk factors that are in Table 1 (continued)**

Variables (foods in kcal/day)	Sugary bever- ages	Corn	Pot- atoes	Fruits	Vege- tables	Nuts and seeds	Whole grains
Corn	0.419	1.000					
Potatoes	-0.026	0.158	1.000				
Fruits	0.189	0.100	0.243	1.000			
Vegetables	-0.173	-0.203	-0.044	0.394	1.000		
Nuts and seeds	-0.154	-0.114	0.188	0.586	0.556	1.000	
Whole grains	0.204	0.428	-0.058	0.012	-0.346	-0.208	1.000
Legumes	0.199	0.269	0.018	-0.032	-0.160	-0.028	0.100
Rice	-0.142	-0.248	-0.378	-0.325	-0.237	-0.359	0.547
Sweet potatoes	-0.215	-0.027	0.238	-0.100	-0.099	-0.079	-0.039
Physical activity METs	-0.099	0.028	0.021	-0.393	-0.212	-0.306	0.268
Kcal /day available	0.114	-0.044	0.182	0.640	0.615	0.601	-0.187
Child/infant 2SD underweight	-0.060	-0.081	-0.216	-0.497	-0.457	-0.300	0.089
Discontinued breast feeding <6 mo	0.133	-0.064	0.132	0.564	0.480	0.373	-0.125

383
384

Table 2. Correlations between the risk factors that are in Table 1 (continued)

Variables (foods in kcal/day)	Legumes	Rice	Sweet potatoes	Physical activity	Kcal	Child/ infant under- weight
Rice	-0.047	1.000				
Sweet potatoes	-0.053	0.072	1.000			
Physical activity METs	-0.066	0.388	0.328	1.000		
Kcal /day available	-0.250	-0.429	-0.257	-0.500	1.000	
Child/infant 2SD underweight	0.507	0.377	-0.164	0.138	-0.621	1.000
Discontinued breast feeding <6 mo	-0.406	-0.419	-0.062	-0.227	0.727	-0.836

Table 3. Bradford Hill criteria consistency and dose response: Mean BMI and BMI formula output differences in 37 subgroups

Subsets of worldwide data on risk factors related to BMI	n	Mean BMI	BMI formula output with mean and SD equated with mean BMI	BMI and BMI formula‡ output difference	BMI and BMI formula ¶ output difference	BMI and BMI formula† output difference
Socio-demographic index quartile 1	1966	24.54	24.75	0.210	0.243	0.289
SDI quartile 2	2096	21.82	21.69	0.130	0.210	0.402
SDI quartile 3	2220	20.40	20.30	0.099	0.104	0.107
SDI quartile 4	1604	20.31	20.31	0.007	0.079	0.012
Africa	1682	21.67	21.25	0.411	0.009	0.519
Asia	4188	20.47	20.46	0.013	0.065	0.058
Europe	880	24.31	24.66	0.343	0.388	0.234
North America	592	25.97	26.00	0.032	0.079	0.312
Oceana	54	24.43	24.66	0.229	0.277	0.206
South America	474	23.48	24.24	0.761	0.667	0.463
Four countries with subnational data	730	25.18	25.45	0.267	0.300	0.964
Quartile 1 BMI increasing foods	2007	24.14	24.70	0.565	0.572	0.654
Q2	1834	21.96	21.86	0.096	0.103	0.274
Q3	2050	20.60	20.31	0.293	0.281	0.205
Q4	1995	20.49	20.28	0.212	0.226	0.224
Quartile 1 BMI decreasing foods	1475	20.61	20.16	0.454	0.570	0.197
Q2 BMI decreasing foods	2275	21.12	21.03	0.092	0.114	0.114
Q3 BMI decreasing foods	2211	22.48	22.74	0.258	0.294	0.239
Q4 BMI decreasing foods	1925	22.69	22.82	0.125	0.199	0.112
Quartile 1 physical activity	2371	20.48	20.56	0.081	0.089	0.019
Q2 physical activity	1573	21.66	21.13	0.533	0.598	0.554
Q3 physical activity	2018	21.37	21.55	0.175	0.198	0.308
Q4 physical activity	1924	23.95	24.07	0.118	0.136	0.125
Male	3943	21.51	21.91	0.406	0.401	0.368
Female	3943	22.07	21.65	0.423	0.419	0.382
Random controlled trial #1	100	21.92	21.96	0.038	0.054	0.061
RCT #2	100	22.00	21.76	0.236	0.226	0.198
RCT #3	100	21.67	21.78	0.109	0.111	0.149
RCT #4	100	21.89	21.81	0.074	0.069	0.064
UK	66	24.99	25.13	0.132	0.132	0.393
USA	376	26.66	26.74	0.082	0.132	0.380
Japan	158	21.89	22.87	0.978	0.964	2.888
Mexico	130	24.99	24.99	0.006	0.062	0.606
Quartile 1 Dose-response: BMI formula versus BMI	1862	24.82	25.33	0.509	0.538	0.663
Q2 Dose-response	1802	22.44	22.06	0.379	0.350	0.539
Q3 Dose-response	1773	21.10	20.84	0.266	0.034	0.201
Q4 Dose-response	2449	19.51	19.56	0.057	0.158	0.128
Mean of BMI - BMI formula absolute values				0.249	0.255	0.368

Mean of BMI-BMI formula dose response absolute values				0.302	0.284	0.493

‡In combination variables grouped by BMI increasing foods and BMI decreasing foods, each food was multiplied times (1) its kcal/day (2) its coefficient of determination (R^2) with BMI, and (3) its ratio of kcal/day in SDI quartiles 2-4 /SDI quartile 1. Fruits, vegetables, and nuts/seeds grouped with plants that negatively correlate with BMI.

ɣ As with ‡ but with fruits, vegetables, and nuts/seeds grouped with plants that positively correlate with BMI

† As with § but with food risk factor unadjusted but with food risk factors unadjusted

Table 4. Cross validation: summary percent weight data for 20 RCT derived BMI formulas↓

Risk factors	Mean	Std Dev	Minimum	Maximum
Processed meat kcal/day	0.26	0.09	0.11	0.43
Red meat kcal/day	9.82	3.05	5.41	14.38
Fish kcal/day	0.18	0.51	0.01	1.97
Milk kcal/day	3.13	1.22	1.38	4.88
Poultry kcal/day available	9.10	2.88	5.35	15.53
Eggs kcal/day available	4.66	1.09	2.50	7.73
Alcohol kcal/day	1.78	2.08	0.01	7.92
Sugary beverages kcal/day	7.24	1.51	0.01	14.62
Corn kcal/day available	0.80	0.70	0.00	2.00
potatoes kcal/day available	2.79	1.94	0.01	7.81
Saturated fatty acids kcal/day	3.09	2.09	0.90	8.18
Poly unsaturated fatty acids kcal/day	0.71	0.38	0.18	1.50
Trans fatty acids kcal/day	0.09	0.05	0.03	0.18
Fruits kcal/day	0.46	0.21	0.12	0.88
Vegetables kcal/day	0.63	0.30	0.23	1.31
Nuts and seeds kcal/day	0.03	0.02	0.01	0.07
Whole grains kcal/day	0.13	0.02	0.00	0.39
Legumes kcal/day	0.56	3.51	0.09	1.54
Rice kcal/day available	8.89	2.65	3.40	13.19
Sweet potatoes kcal/day available	0.32	0.32	0.02	1.13
Physical activity METs	9.76	4.09	4.24	13.29
Kcal available/day	7.89	3.23	3.68	13.56
Child/infant 2SD underweight	5.41	4.51	2.49	8.29
Discontinued breast feeding <6 mo	1.19	0.84	0.20	2.96
Total percent weights	78.90	2.88	74.65	83.04

↓ See Supplementary Table 7 for the data on the 20 RCTs each with n=100

Table 5. BMI formula estimates with common dietary and other risk factor scenariosβ

Testing common dietary and other risk factor scenarios with standardised BMI formula estimates	BMI kg/M²	BMI formula estimate
World (n=7886 cohorts)	21.79	21.71
United Kingdom (n=66 cohorts)	24.99	25.09
USA (n=376 cohorts)	26.66	26.67
Following USA Dietary Guidelines 2015-2020 recommendations		26.34
USA with 25% reduction of BMI increasing food intake†		25.04
USA with 50% reduction of BMI increasing food intake†		23.34
USA mean physical activity plus 2 hours/day running at 6 mph		23.98
USA mean physical activity plus 1 hour/day running at 6 mph and 25% reduction of BMI increasing food intake†		23.67
USA with no red or processed meat†		24.81
USA with no sugary beverage intake†		26.31
USA vegetarian (no meat, poultry, fish)†		22.54
USA vegan (no meat, poultry, fish, dairy or eggs)†		21.11
USA breast feeding continued at least 6 months in all		26.68
USA breast feeding discontinued before 6 months in all		26.87
USA no occurrence of child severe underweight		26.73
USA all children with severe underweight (theoretically)		25.54
EAT-Lancet diet		22.47
Low Carbohydrate Mediterranean Diet		31.76

β BMI formula estimates based on 28 years of following dietary and/or risk factor patterns

† Kcal/day isocalorically shifted to the 7 BMI decreasing foods in the BMI formula, distributed equally

Acknowledgments

• **Authors' contributions:** DKC acts as guarantor; conceived and designed the study, acquired

and analysed the data, interpreted the study findings, drafted the manuscript, critically reviewed

and edited the manuscript and tables, and approved the final version for publication.

CW designed software programs in R to format and population weight the data,

aided with the SAS statistical analysis, critically reviewed the manuscript, and

approved the final version for publication.

The corresponding author attests that all listed authors meet authorship criteria and that no others

meeting the criteria have been omitted.

Copyright: The retain the copyright to the paper

The Patient and Public Involvement

● When and how were patients/public first involved in the research?

IHME acquired, catalogued and extracted information from over 12,000 surveys from

government and non government agencies in order to compile the GBD database.

● How were the research question(s) developed and informed by their priorities,

experience, and preferences?

The diverse surveys contributing to the GBD database had many different priorities and

intentions unrelated to this post hoc analysis. The current analysis research questions were

developed to give researchers, policymakers, and the public the methodological tool to quantify

the BMI impacts of dietary and other risk factor patterns.

● How were patients/public involved in

(a) the design and conduct of the study?

The way patients/public were involved in the collection of the surveys has not been

systematically studied and reported on. Patients/public are the intended beneficiaries of

this analysis of the GBD data.

(b) choice of outcome measures?

The outcome measures chosen all related to human health and to adult BMI.

(c) recruitment to the study?

The recruitment methods varied by the survey.

● How were (or will) patients/ public be involved in choosing the methods and agreeing

plans for dissemination of the study results to participants and linked communities?

IHME and all the health surveyors that contributed will make these decisions. This analysis of

the data will be available to all by open access.

451•   **Competing interests statement:** Both authors have completed the ICMJE uniform disclosure

form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the

submitted work; no financial relationships with any organisations that might have an interest in

the submitted work in the previous three years; no other relationships or activities that could

appear to have influenced the submitted work.

Contributors: Martin Sebera, from the Department of Kinesiology, Faculty of Sports Studies

Masaryk University, Czech Republic, critiqued statistical aspects of the manuscript and provided

useful input. Pavel Grasgruber, from Masaryk University, Czech Republic, provided suggestions

after reviewing the manuscript.

Transparency declaration: The manuscript is an honest, accurate, and transparent account of

the study being reported; that no important aspects of the study have been omitted; and that any

discrepancies from the study as planned have been explained.

Ethics: Studies based solely on data from IHME GBD database do not need approval from any

bioethics committee.

469•   **Funding:** This research received no specific grant from any funding agency in the public,

470   commercial or not-for-profit sectors. The Bill and Melinda Gates Foundation funded the

471   acquisition of the data for this analysis by the IHME. The data were provided to the authors as

472   volunteer collaborators with IHME.

473•

474
Details of the role of the study sponsors: While IHME GBD faculty and staff by virtue of

Gates Foundation grants provided the raw data for this analysis, they did not vet the analysis or

sponsor the manuscript.

Statement of independence of researchers from funders: The researchers have received no

funding. Gates Foundation funded IHME to collect and analyse the GBD data.

Dissemination declaration: Dissemination of this manuscript to the participants of the more

than 12,000 surveys is not possible individually, but the information will become in the public

domain.

Disclosures: We thank Scott Glenn and Brent Bell from IHME who supplied us with the GBD

risk factor exposure data for the risk factors and for BMI data.

487•

Ethics: Studies were based solely on data from the IHME GBD database and do not need

approval from any bioethics committee.

Participant informed consents: Not applicable.

Author access to data: As volunteer collaborators with the Institute of Health Metrics and

Evaluation, we received about 1.4 gigabytes of raw data on BMI and 32 relevant risk factors for

BMI and other health outcomes.

Data sharing statement: The raw, unformatted data used in this analysis is now out of date. The

2019 GBD data on all the variables in this analysis may be obtained from the IHME by volunteer

collaborating researchers. Our data formatting software code in R and SAS and our formatted

database are available on request to researchers.

Protocol, submitted as a supplementary file: Not applicable.

STROBE checklist: Submitted.

Patient consent*:* Not applicable.

Clinical trial registration: Not applicable.

References

References

1.   Teicholz N. **The scientific report guiding the US dietary guidelines: is it scientific?**

BMJ. 2015; **351**. http://www.bmj.com/bmj/351/bmj.h4962.full.pdf

2.   **Outcome of Post-publication review of article by Nina Teicholz**. *BMJ.* 2015; **351:**

h4962. https://www.bmj.com/content/351/bmj.h4962/rr-48

3.   Gardner CD, Trepanowski JF, Del Gobbo LC, et al. **Effect of Low-Fat vs Low-**

Carbohydrate Diet on 12-Month Weight Loss in Overweight Adults and the

Association With Genotype Pattern or Insulin Secretion: The DIETFITS

Randomized Clinical Trial. *JAMA.* 2018; **319** (7)**:** 667-679.

https://jamanetwork.com/journals/jama/fullarticle/2673150

4.   Ioannidis JA. **The challenge of reforming nutritional epidemiologic research** *JAMA*

2018; **320** (10)**:** 969-70. http://dx.doi.org/10.1001/jama.2018.11025

5.   Hill AB. **THE ENVIRONMENT AND DISEASE: ASSOCIATION OR**

CAUSATION? *Proceedings of the Royal Society of Medicine.* 1965; **58** (5)**:** 295-300.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1898525/

6.   Global Burden of Disease Study 2017 (GBD 2017) Data Input Sources Tool. Seattle,

Washington: Institute of Health Metrics and Evaluation Accessed August 10, 2019.

http://ghdx.healthdata.org/gbd-2017/data-input-sources

7.   **GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative**

risk assessment of 84 behavioural, environmental and occupational, and metabolic

risks or clusters of risks for 195 countries and territories, 1990-2017: a systematic

analysis for the Global Burden of Disease Study 2017: Table of risk factor

definitions. *The Lancet.* 2018; **392** (10159)**:** 1923-94.

https://www.thelancet.com/action/showFullTableHTML?isHtml=true&tableId=tbl1&pii=

S0140-6736%2818%2932225-6

8.   PROTOCOL FOR THE GLOBAL BURDEN OF DISEASES, INJURIES, AND RISK

FACTORS STUDY (GBD) Version 3.0; Issue 26. Seattle, Washington: Institute for

Health Metrics and Evaluation Accessed August 2, 2019.

http://www.healthdata.org/sites/default/files/files/Projects/GBD/GBD_Protocol.pdf

9.   Naghavi M, Abajobir AA, Abbafati C, et al. **Global, regional, and national age-sex**

specific mortality for 264 causes of death, 1980-2016: a systematic analysis for the

Global Burden of Disease Study 2016. *The Lancet.* 2017; **390** (10100)**:** 1151-1210.

https://doi.org/10.1016/S0140-6736(17)32152-9

10.   Flaxman AD, Lee YY, Vos T, et al. An Integrative Metaregression Framework for

Descriptive Epidemiology. Seattle, WA: University of Washington Press, 2015.

11.   GBD 2017 Risk Factor Collaborators. **Global, regional, and national comparative risk**

assessment of 84 behavioural, environmental and occupational, and metabolic risks

or clusters of risks for 195 countries and territories, 1990-2017: a systematic

analysis for the Global Burden of Disease Study 2017. . *The Lancet.* 2018; **392**

(10159)**:** 1923-94. https://doi.org/10.1016/S0140-6736(18)32225-6

12.   Call for Collaborators. Seattle, Washington: Institute for Health Metrics and Evaluation

at the University of Washington 2019; Accessed April 25, 2019.

http://www.healthdata.org/gbd/call-for-collaborators

13.   Stevens GA, Alkema L, Black PRE, et al. **Guidelines for Accurate and Transparent**

Health Estimates Reporting: the GATHER statement. *The Lancet.* 2016; **388**

(10062)**:** e19-e23. https://doi.org/10.1016/S0140-6736(16)30388-9

14.   Omega 3 Fatty Acids: Fact Sheet for Health Professionals. The Office of Dietary

Supplements 2018; Accessed September 1, 2018.

https://ods.od.nih.gov/factsheets/Omega3FattyAcids-HealthProfessional/

15.   Nutritionix Track App. Syndigo LLC Accessed April 25, 2019.

https://www.nutritionix.com/

16.   POTATO PROCESSING AND USES. Lima, Peru: International Potato Center

Accessed May 28, 2020. https://cipotato.org/potato/potato-processing-uses/

17.   Ruotsalainen H, Kyngäs H, Tammelin T, Kääriäinen M. **Systematic review of physical**

activity and exercise interventions on body mass indices, subsequent physical

activity and psychological symptoms in overweight and obese adolescents. *Journal of*

Advanced Nursing. 2015; **71** (11)**:** 2461-2477.

https://onlinelibrary.wiley.com/doi/abs/10.1111/jan.12696

18.   Te Morenga L, Mallard S, Mann J. **Dietary sugars and body weight: systematic review**

and meta-analyses of randomised controlled trials and cohort studies. *BMJ* 2013;

346: e7492. https://www.bmj.com/content/bmj/346/bmj.e7492.full.pdf

19.   Traversy G, Chaput J-P. **Alcohol Consumption and Obesity: An Update**. *Current*

obesity reports. 2015; **4** (1)**:** 122-130.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4338356/

20.   Turner-McGrievy G, Mandes T, Crimarco A. **A plant-based diet for overweight and**

obesity prevention and treatment. *Journal of geriatric cardiology : JGC.* 2017; **14** (5)**:**

369-374. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5466943/

21.   AINSWORTH BE, HASKELL WL, HERRMANN SD, et al. **Compendium of Physical**

Activities: A Second Update of Codes and MET Values. *Medicine & Science in Sports*

& Exercise. 2011 **43** (8)**:** 1575-1581. https://journals.lww.com/acsm-

msse/Fulltext/2011/08000/2011_Compendium_of_Physical_Activities__A_Second.25.as

px

22.   Ebbeling CB, Feldman HA, Klein GL, et al. **Effects of a low carbohydrate diet on**

energy expenditure during weight loss maintenance: randomized trial. *BMJ.* 2018;

363: k4583.

23.   Darmon N, Drewnowski A. **Contribution of food prices and diet cost to**

socioeconomic disparities in diet quality and health: a systematic review and

analysis. *Nutrition reviews.* 2015; **73** (10)**:** 643-660.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4586446/

24.   Wakamo B. We Subsidize the Wrong Kind of Agriculture. Inequality.org Accessed.

https://inequality.org/great-divide/subsidize-wrong-kind-agriculture/

25.   Tagtow A. Fruits and Veggies Now Chock Full of Marketing Power. Washington, DC

2017; Accessed June 24, 2020. https://www.usda.gov/media/blog/2015/05/07/fruits-and-

veggies-now-chock-full-marketing-power

26.   Stolzer JM. **Breastfeeding and obesity: a meta-analysis**. *Open Journal of Preventive*

Medicine. 2011; **1** (3)**:** 88-93.

http://www.scirp.org/journal/paperinformation.aspx?paperid=8509

27.   AAP Reaffirms Breastfeeding Guidelines. The American Academy of Pediatrics 2012;

Accessed August 10, 2019. https://www.aap.org/en-us/about-the-aap/aap-press-

room/Pages/AAP-Reaffirms-Breastfeeding-Guidelines.aspx

28.   **Breastfeeding and the Use of Human Milk**. *Pediatrics.* 2012; **129** (3)**:** e827-e841.

doi:10.1542/peds.2011-3552

29.   Quality statement 5: Breastfeeding. National Institute of Clinical Excellence 2013;

Accessed August 10, 2019. https://www.nice.org.uk/guidance/qs37/chapter/Quality-

statement-5-Breastfeeding

30.   FORESIGHT--Tackling Obesities: Future Choices -Follow Up Actions. London,

England: Government Office for Science (UK) October 2007; Accessed August 5, 2019.

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_

data/file/295885/07-1526-obesity-action-plan.pdf

31.   Mărginean CO, Mărginean C, Meliţ LE. **New Insights Regarding Genetic Aspects of**

Childhood Obesity: A Minireview. *Frontiers in pediatrics.* 2018; **6:** 271-271.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6180186/

32.   Castaner O, Goday A, Park Y-M, et al. **The Gut Microbiome Profile in Obesity: A**

Systematic Review. *International journal of endocrinology.* 2018; **2018:** 4095789-

4095789. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5933040/

33.   Monteiro CA, Moubarac J-C, Levy RB, Canella DS, Louzada MLdC, Cannon G.

Household availability of ultra-processed foods and obesity in nineteen European

countries. *Public Health Nutrition.* 2017; **21** (1)**:** 18-26.

https://www.cambridge.org/core/article/household-availability-of-ultraprocessed-foods-

and-obesity-in-nineteen-european-countries/D63EF7095E8EFE72BD825AFC2F331149

34.   Monteiro CA, Cannon G, Levy RB, et al. **Ultra-processed foods: what they are and**

how to identify them. *Public Health Nutrition.* 2019; **22** (5)**:** 936-941.

https://www.cambridge.org/core/article/ultraprocessed-foods-what-they-are-and-how-to-

identify-them/E6D744D714B1FF09D5BCA3E74D53A185

 Supplementary Table 1. Definitions of IHME GBD risk factors and covariates related to BMI

Variables	Definition
Alcohol	Any alcohol consumption (g/day)
Body-mass index	Body mass index (BMI) (kg/m^2)—the dependent variable of interest
Child underweight	Proportion of children - 3 SD to - 2 SD of the WHO 2006 standard weight-for-age curve (0-1)
Corn	Corn availability percapita (g/day), a covariate
Discontinued breast feeding	Proportion of children aged 6-23 months who do not receive any breast milk
Eggs	Eggs availability percapita (g/day) a covariate
Fasting plasma glucose	Fasting plasma glucose (mmol/L)
Fish	This variable expressed in g/day was derived by determining the weight of fish in g corresponding to 1 g of omega-3 fatty acids (eicosapentaenoic acid and docosahexaenoic acid) by averaging the fish g per 1 g of omega-3 fatty acids 20 species of fish= 117.04 g/day fish/1 g/day omega-3 fatty acids (Supplementary Table 2)
Fruits	Consumption of fruits (includes fresh, frozen, cooked, canned, or dried fruit but excludes fruit juices and salted or pickled fruits) (g/day)
Kilocalories available /day	The mean number of kilocalories percapita available per day to people in each location (kcal/day available), a covariate
LDL cholesterol	Serum low-density lipoprotein cholesterol (mmol/L)
Legumes	Consumption of beans, lentils, pulses (g/day)
Milk	Consumption of milk including non-fat, low-fat, and full-fat milk but excluding soy milk and other plant derivatives (g/day)
Nuts and seeds	Consumption of nuts and seeds (g/day)
Physical activity	Average weekly physical activity at work, home, transport-related and recreational measured by MET min per week. Less than 3000 METs per week constitutes low physical activity.
Poultry	Poultry availability percapita (g/day), a covariate
Potatoes	Potatoes availability percapita (g/day), a covariate
Processed meat	Consumption of any processed meat (includes meat preserved by smoking, curing, salting, or addition of chemical preservatives, including bacon, salami, sausages, or deli or luncheon meats like ham, turkey, and pastrami (g/day)
Red meat	Consumption of red meat (includes beef, pork, lamb, and goat but excludes poultry, fish, eggs, and all processed meats) (g/day)
Rice	Rice availability percapita (g/day), a covariate
Seafood omega-3 fatty acids	Seafood omega-3 fatty acids (eicosapentaenoic acid and docosahexaenoic acid) in tablet or fish form (g/day)
Socio-demographic index	SDI is a composite indicator of development status that was originally constructed for GBD 2015, and is derived from components that correlate strongly with health outcomes. It is the geometric mean for indices of the total fertility rate among women younger than 25 years, mean education for those aged 15 years or older, and lag-distributed income per capita. The resulting metric ranges from 0 to 1, with higher values corresponding to higher levels of development.
Sugar-sweetened beverages	Consumption of any beverage with ≥50 calories of sugar per one-cup serving, including carbonated beverages, sodas, energy drinks, fruit drinks but excluding 100% fruit and vegetable juices (g/day)
Sweet potatoes	Sweet potato availability percapita (g/day), a covariate
Systolic blood pressure	Systolic blood pressure (mm Hg)
Total sugar	Total sugar availability percapita (g/day), a covariate

Vegetables	Consumption of frozen, cooked, canned, or dried vegetables (including legumes but excluding salted or pickled, juices, nuts and seeds, and starchy vegetables such as potatoes or corn) (g/day)
Whole grains	Consumption of whole grains (bran, germ, and endosperm in their natural proportions) from breakfast cereals, bread, rice, pasta, biscuits, muffins, tortillas, pancakes, and others (g/day)

627 **Supplementary Table 2. Omega-3 Fatty Acid g to fish g calculation¶**

Fish	DHA g/3 ounce fish	EPA g/3 ounce fish	Omega-3 Fatty Acids (DHA _ EPA) g/3 ounce fish mean	Fish 3 ounces = 85.02 g	Fish (g) per omega-3 Fatty Acids (g)=columns E/F
Salmon Atlantic farmed	1.24	0.59			
Salmon Atlantic wild	1.22	0.35			
Herring Atlantic	0.94	0.77			
Sardines canned in tomato sauce drained	0.74	0.45			
Mackerel Atlantic	0.59	0.43			
Salmon pink canned drained	0.63	0.28			
Trout rainbow wild	0.44	0.40			
Oysters eastern wild	0.23	0.30			
Sea bass	0.47	0.18			
Shrimp	0.12	0.12			
Lobster	0.07	0.10			
Tuna light canned in water drained	0.17	0.02			
Tilapia	0.11				
Scallops	0.09	0.06			
Cod Pacific	0.1	0.04			
Tuna yellowfin	0.09	0.01			
Mean DHA and EPA Omega-3 Fatty Acids g/3 ounce fish	0.4531	0.2733			
Calculations total Omega-3 FA g to fish g			0.7264	85.02	117.043

628 ¶ Data on omega-3 fatty acid content of varieties of fish came from the National Institutes of
629 Health Office of Dietary Supplements (USA)
630

Supplementary Table 3. Calculations of kcal/day from g/day of animal and plant foods¶

Foods	Food sub-categories	kcal/serving	g/serving	kcal/g
Milk (2% fat)		122	244	0.5
Fish		218	170	1.28
Eggs		72	50	1.44
Poultry		187	85	2.91
Red meat		247	85	2.91
Processed meat				
	Salami	222	59	3.76
	Pastrami	104	71	1.46
	Ring baloney	86	28	3.07
	Pepperoni	94	100	0.94
Average processed meat		126.5	64.5	1.96
Fruits		97	162	0.60
Vegetables		59	91	0.65
Legumes		249	179	1.39
Nuts		172	28	6.14
Seeds				
	Flax seeds	55	10	5.5
	Chia seeds	58	12	4.83
	Fennel seeds	34.5	10	3.45
	Hemp seeds	55.3	10	5.53
Average of seeds		50.7	10.5	4.83
Average of nuts and seeds		111.4	19.25	5.78
Corn		99	103	0.96
Potatoes		161	173	0.93
Sweet potatoes		115	151	0.76
Rice		205	158	1.3
Whole grains		120	52	2.31

¶ Source: NutritionIX app

Supplementary Table 4. Multicollinearity of fruits, vegetables, and nuts and seeds

High BMI risk factors (n=7886 cohorts)	Mean Kcal/day	BMI kg/M² r	95% CI	P	BMI increasing r †	95% CI	P
BMI kg/m²	21.79				0.724	0.713 to 0.734	<0.0001
BMI increasing r	654	0.597	0.583 to 0.611	<0.0001			
Fruits + vegetables + nuts and seeds	129	0.655	0.642 to 0.677	<0.0001	0.323	0.304 to 0.343	<0.0001
Fruits	40.4	0.622	0.609 to 0.636	<0.0001	0.498	0.482 to 0.515	<0.0001
Vegetables	80.1	0.520	0.503 to 0.536	<0.0001	0.150	0.128 to 0.171	<0.0001
Nuts and seeds	8.51	0.486	0.469 to 0.503	<0.0001	0.279	0.258 to 0.299	<0.0001
Top socio-demographic index (SDI) quartile n=1996	**Kcal/day**	**BMI kg/M² r**	**95% CI**	**P**	**BMI increasing r †**	**95% CI**	**P**
BMI kg/m²	24.54				0.364	0.325 to 0.402	<0.0001
BMI increasing	905	0.364	0.325 to 0.402	<0.0001			
Fruits + vegetables + nuts and seeds	199	0.104	0.060 to 0.148	<0.0001	-0.306	0.346 to 0.266	<0.0001
Fruits	61.1	0.296	0.255 to 0.336	<0.0001	-0.022	-0.007 to -0.095	0.32
Vegetables	120	-0.064	-0.108 to -0.020	0.0043	-0.388	-0.425 to -0.350	<0.0001
Nuts and seeds	17.5	0.314	0.273 to 0.353	0.0018	-0.118	-0.161 to -0.074	<0.0001
Bottom three SDI quartiles (n-=5920 cohorts)	**Kcal/day**	**BMI kg/M² r**	**95% CI**	**P**	**BMI increasing r †**	**95% CI**	**P**
BMI kg/m²	20.88	`			0.364	0.342 to 0.386	<0.0001
BMI increasing	571	0.364	0.342 to 0.386	<0.0001			
Fruits + vegetables + nuts and seeds	106	0.536	0.517 to 0.554	<0.0001	0.108	0.083 to 0.133	<0.0001
Fruits	33.5	0.466	0.446 to 0.486	<0.0001	0.429	0.408 to 0.450	<0.0001
Vegetables	66.8	0.395	0.374 to 0.417	0.0043	-0.103	-0.128 to -0.077	0.0186
Nuts and seeds	5.52	-0.040	-0.066 to -0.015	0.0018	-0.072	-0.098 to -0.047	<0.0001

† BMI increasing foods include all six animal products, alcohol, sugary beverages, corn (high fructose corn syrup included), and potatoes (highly processed potato products included). Saturated fatty acids, PUFAs, and TFAs also correlate positively with BMI. However, for the purposes of this table, they were not included with BMI increasing foods, because animal foods contain most of the fatty acids.

SupplementaryTable 5. Derivation of the BMI formula (variables all standardized)

A	B	C	D	E	F	G	H	I	J	K
Variables in BMI formula		Mean		R2 with BMI		SDI Q 2-4 / SDI Q 1		Foods and physical activity multiple regression derived parameter estimates		Foods and physical activity synthesis parameter estimates= Columns C*E*G*I absolute values
Processed meat kcal/day	*	5.486	*	0.365	*	0.099	*	0.02972	=	0.00586
Red meat kcal/day	*	50.70	*	0.438	*	0.364	*	0.02972	=	0.24003
Fish kcal/day	*	10.02	*	0.011	*	0.106	*	0.02972	=	0.00033
Milk kcal/day	*	25.37	*	0.468	*	0.214	*	0.02972	=	0.07553
Poultry kcal/day availability	*	45.06	*	0.661	*	0.253	*	0.02972	=	0.22416
Eggs kcal/day availability	*	19.47	*	0.473	*	0.402	*	0.02972	=	0.11011
Alcohol kcal/day	*	81.71	*	0.028	*	0.587	*	0.02972	=	0.03948
Sugary beverages kcal/day	*	297.7	*	0.014	*	1.013	*	0.02972	=	0.12722
Corn kcal/day availability	*	34.67	*	0.005	*	1.081	*	0.02972	=	0.00590
Potatoes kcal/day availability	*	84.16	*	0.044	*	0.644	*	0.02972	=	0.07137
Saturated fatty acids	*	191.3	*	0.504	*	0.653	*	0.0008705	=	0.05481
Polyunsaturated fatty acids	*	82.24	*	0.543	*	0.334	*	0.0008705	=	0.01297
Trans fatty acids	*	13.39	*	0.238	*	0.471	*	0.0008705	=	0.00131
Fruits kcal/day	*	40.39	*	0.387	*	0.549	*	0.0008705	=	0.00747
Vegetables kcal/day	*	80.14	*	0.270	*	0.555	*	0.0008705	=	0.01046
Nuts and seeds kcal/day	*	8.511	*	0.237	*	0.315	*	0.0008705	=	0.00055
Whole grains kcal/day	*	55.65	*	0.040	*	1.248	*	0.0008705	=	0.00243
Legumes kcal/day	*	51.65	*	0.148	*	1.432	*	0.0008705	=	0.00951
Rice kcal/day availability	*	141.2	*	0.314	*	4.320	*	0.0008705	=	0.16697
Sweet potatoes kcal/day availability	*	22.67	*	0.023	*	7.977	*	0.0008705	=	0.00362
Physical activity METs								0.20087	=	0.20087
Total Kcal/day availability										
Child/infant underweight										
Discontinued breast feed <6 months										
Sums										1.37098
Sums										1.37098
BMI formula R2										0.73530
BMI formula R2 / sum of parameter estimates										0.53633

A	L	M	N	O	P	Q	R	S
Variables in BMI formula	**Adjust column K μ**	**Add other risk factors**	**Adjust column M by multi-plying by M28 = 0.1211**	**Com-bine columns L and N**		**Check result with statistics software SAS Studio**		**Final BMI formula percent weights: Column O * 0.9208 (O30) *100**
Processed meat kcal/day	0.00315			0.00315	+	Processed meat	*	0.29
Red meat kcal/day	0.12874			0.12874	+	Red meat	*	11.87
Fish kcal/day	0.00018			0.00018	+	Fish	*	0.02
Milk kcal/day	0.04051			0.04051	+	Milk	*	3.74
Poultry kcal/day	0.12022			0.12022	+	Poultry	*	11.09
Eggs kcal/day	0.05906			0.05906	+	Eggs	*	5.45
Alcohol kcal/day	0.02117			0.02117	+	Alcohol	*	1.95
Sugary beverages kcal/day	0.06823			0.06823	+	Sugary beverages	*	6.29
Corn kcal/day	0.00317			0.00317	+	Corn	*	0.29
Potatoes kcal/day	0.03828			0.03828	+	Potatoes	*	3.53
Saturated fatty acids	0.02940			0.02940	+	SFA	*	2.71
Polyunsaturated fatty acids	0.00696			0.00696	+	PUFA	*	0.64
Trans fatty acids	0.00070			0.00070	+	TFA	*	0.06
Fruits kcal/day	0.00401			0.00401	-	Fruits	*	0.37
Vegetables kcal/day	0.00561			0.00561	-	Vegetables	*	0.52
Nuts and seeds kcal/day	0.00030			0.00030	-	Nuts and seeds	*	0.03
Whole grains kcal/day	0.00130			0.00130	-	Whole grains	*	0.12
Legumes kcal/day	0.00510			0.00510	-	Legumes	*	0.47
Rice kcal/day	0.08955			0.08955	-	Rice	*	8.26
Sweet potatoes kcal/day	0.00194			0.00194	-	Sweet Potatoes	*	0.18
Physical activity	0.10773			0.10773	-	Physical activity	*	9.93
Total Kcal/day		0.5408	0.063	0.06270	+	Total kcal	*	5.78
Child/infant underweight		0.3865	0.045	0.04481	-	Child underweight	*	4.13
Discontinued breast feed <6 months		0.0861	0.010	0.00999	+	Discontinued breast feeding	*	0.92
Sums	0.7353	1.0133	0.118	0.85280				78.64
Sums		1.0133		0.85280				
BMI formula R2		0.1175		0.78643				0.79
BMI formula R2 / sum of parameter estimates		0.1160		0.92218				100.00

648 μ Adjust column K by K28 =0.5314 (foods and physical activity BMI formula R^2 (K27) /parameter estimate sums (K26)
649

651 Supplementary Table 6. Spreadsheet derivation of BMI trend formula

A	B	C	D	E	F	G	H	I
BMI trend risk factor trends		Mean		Risk factor with BMI17fts R2		Parameter estimates from multiple regression		Column C * Column E * Column G= preliminary parameter estimates
Processed meat kcal/day trend	*	5.486	*	0.0001	*		=	0.000
Red meat kcal/day trend	*	50.695	*	0.1650	*	0.01645	=	0.138
Fish kcal/day trend	*	10.015	*	0.0251	*		=	0.000
Milk kcal/day trend	*	25.372	*	0.0425	*	0.01645	=	0.018
Poultry kcal/day trend	*	45.059	*	0.1768	*	0.01645	=	0.131
Eggs kcal/day trend	*	19.466	*	0.2322	*	0.01645	=	0.074
Alcohol kcal/day trend	*	81.713	*	0.0826	*	0.01645	=	0.111
Sugary beverages kcal/day trend	*	297.648	*	0.0156	*	0.01645	=	0.077
Corn kcal/day available trend	*	34.669	*	0.0198	*	0.01645	=	0.011
Potatoes available kcal/day available trend	*	84.160	*	0.0739	*	0.01645	=	0.102
Saturated fatty acids trend	*	191.27	*					
Polyunsaturated fatty acids trend	*	82.24	*	0.0820		0.01645		0.111
Trans fatty acids trend	*	13.4	*					
Fruits kcal/day trend	*	40.388	*	0.2301	*			0.000
Vegetables kcal/day trend	*	80.141	*	0.1559	*			0.000
Nuts and seeds kcal/day trend	*	8.510	*	0.0152	*			0.000
Whole grains kcal/day trend	*	55.647	*	0.0107	*		=	0.000
Legumes kcal/day trend	*	51.657	*	0.0887	*	0.00404	=	0.019
Rice kcal/day available trend	*	141.230	*	0.1536	*	0.00404	=	0.088
Sweet potatoes available kcal/day trend	*	22.667	*	0.2487	*	0.00404	=	0.023
Physical activity mets trend								
Kcal available/day trend					*	0.17225	=	0.17225
Child/infant 2SD underweight trend								
Discontinued breast feeding <6 mo trend								
Sums								1.07414
Sum								1.07414
R^2								0.37790
Trend R^2 / trend BMI formula parameter estimates sum								0.35182

652
653
654

Spreadsheet derivation of BMI trend formula continued

A	I	J	K	L	M	N
BMI trend risk factor trends	**Column C * Column E * Column G= preliminary parameter estimates**	**Column I adjusted to equate to parameter estimates sum to R2 of the BMI trend formula**	**K**	**Final BMI trend formula to check percent weights with SAS Studio**	**M**	**Column J * 100 to derive percent weights of risk factors**
Processed meat kcal/day trend	0.000			pmeat17KCts		
Red meat kcal/day trend	0.138	0.048	+	rmeat17KCts	*	4.841
Fish kcal/day trend	0.000	0.000		fish17KCts		
Milk kcal/day trend	0.018	0.006	+	milk17KCts	*	0.624
Poultry kcal/day trend	0.131	0.046	+	poultry16KCts	*	4.610
Eggs kcal/day trend	0.074	0.026	+	eggs16KCts	*	2.616
Alcohol kcal/day trend	0.111	0.039	+	Alcohol17KCts	*	3.908
Sugary beverages kcal/day trend	0.077	0.027	+	Sugarb17KCts	*	2.692
Corn kcal/day available trend	0.011	0.004	+	corn16KCts	*	0.397
Potatoes available kcal/day available trend	0.102	0.036	+	potatoes16KCts	*	3.600
Saturated fatty acids trend		0.000				
Polyunsaturated fatty acids trend	0.111	0.039	+	PUFA17KCts	*	3.904
Trans fatty acids trend		0.000				
Fruits kcal/day trend	0.000	0.000		fruits17KCts		
Vegetables kcal/day trend	0.000	0.000		Vegetables17KCts		
Nuts and seeds kcal/day trend	0.000	0.000		nutsseeds17KCts		
Whole grains kcal/day trend	0.000	0.000		wgrains17KCts		
Legumes kcal/day trend	0.019	0.007	-	legumes17KCts	*	0.651
Rice kcal/day available trend	0.088	0.031	-	rice16KCts	*	3.084
Sweet potatoes available kcal/day trend	0.023	0.008	-	swtpot16KCts	*	0.801
Physical activity mets trend		0.000		PAMETs17ts		
Kcal available/day trend	0.172	0.061	+	KCal2016ts	*	6.060
Child/infant 2SD underweight trend				Childunwt17ts		
Discontinued breast feeding <6 mo trend				discbreastF17ts		
Sums	1.074	0.378				37.79
Sum	1.074					0.37790
R^2	0.378					100.00
Trend R^2 / trend BMI formula parameter estimates sum	0.352					

Supplementary Table 7. Cross validation experiment with 20 randomised controlled trials of n=100

Risk factors	RCT1 Percent Weights	RCT2 Percent Weights	RCT3 Percent Weights	RCT4 Percent Weights	RCT5 Percent Weights
Processed meat kcal/day	0.39	0.22	0.14	0.22	0.11
Red meat kcal/day	13.27	8.57	6.41	7.71	5.41
Fish kcal/day	0.00	0.01	0.01	0.02	0.01
Milk kcal/day	5.96	3.58	1.38	2.72	1.42
Poultry kcal/day available	13.66	8.22	5.35	8.90	5.53
Eggs kcal/day available	5.43	3.72	2.50	4.35	3.72
Alcohol kcal/day	0.56	1.58	0.30	1.76	0.93
Sugary beverages kcal/day	0.01	8.49	13.37	9.67	11.67
Corn kcal/day available	0.01	0.16	0.19	1.96	1.04
potatoes kcal/day available	2.17	0.83	4.55	2.36	2.60
Saturated fatty acids kcal/day	1.79	7.70	3.12	1.44	3.71
Polyunsaturated fatty acids kcal/day	0.51	1.58	0.88	0.48	0.86
Trans fatty acids kcal/day	0.05	0.20	0.08	0.07	0.07
Fruits kcal/day	0.29	0.73	0.56	0.28	0.79
Vegetables kcal/day	0.45	1.05	0.59	0.63	0.89
Nuts and seeds kcal/day	0.03	0.06	0.03	0.02	0.05
Whole grains kcal/day	0.18	0.26	0.04	0.06	0.01
Legumes kcal/day	0.46	1.10	0.65	0.24	0.47
Rice kcal/day available	3.40	12.76	9.29	7.81	8.14
Sweet potatoes kcal/day available	0.05	0.74	0.18	0.53	0.06
Physical activity METs	13.08	13.10	7.90	9.28	7.11
Kcal available/day	6.52	3.72	10.22	8.31	14.90
Child/infant 2SD underweight	4.86	2.49	7.62	6.03	10.51
Discontinued breast feeding <6 mo	1.52	0.52	0.42	0.45	2.34
Total percent weights in BMI formula	74.65	81.39	75.77	75.28	82.35

Supplementary Table 7. continued

Risk factors	RCT6 Percent weights	RCT7 Percent weights	RCT8 Percent weights	RCT9 Percent weights	RCT10 Percent weights
Processed meat kcal/day	0.27	0.34	0.26	0.39	0.33
Red meat kcal/day	8.14	11.88	8.69	13.49	14.38
Fish kcal/day	0.01	0.04	0.02	0.04	0.03
Milk kcal/day	2.89	3.98	1.89	3.47	4.88
Poultry kcal/day available	6.69	11.11	5.89	7.89	15.53
Eggs kcal/day available	4.41	5.84	4.14	6.21	7.18
Alcohol kcal/day	0.11	4.39	0.62	7.92	1.26
Sugary beverages kcal/day	10.44	4.11	14.62	0.34	1.79
Corn kcal/day available	0.01	0.00	0.82	1.49	0.01
potatoes kcal/day available	1.82	1.97	1.80	2.15	5.32
Saturated fatty acids kcal/day	8.18	1.87	1.97	3.01	2.42

Polyunsaturated fatty acids kcal/day	1.22	0.41	0.54	0.56	0.45
Trans fatty acids kcal/day	0.13	0.07	0.07	0.06	0.04
Fruits kcal/day	0.71	0.33	0.36	0.43	0.35
Vegetables kcal/day	1.31	0.37	0.41	0.77	0.41
Nuts and seeds kcal/day	0.07	0.02	0.03	0.06	0.03
Whole grains kcal/day	0.30	0.16	0.00	0.22	0.08
Legumes kcal/day	1.54	0.39	0.47	0.57	0.46
Rice kcal/day available	11.87	11.51	8.24	8.93	3.53
Sweet potatoes kcal/day available	0.61	0.03	0.79	0.32	0.10
Physical activity METs	4.24	11.48	11.91	9.73	11.59
Kcal available/day	7.47	4.25	6.31	7.51	3.68
Child/infant 2SD underweight	5.93	4.09	4.88	4.10	4.02
Discontinued breast feeding <6 mo	1.92	1.47	2.96	1.65	0.62
Total percent weights in BMI formula	74.65	81.39	75.77	75.28	82.35

664 **Supplementary Table 7. continued**

Risk factors	RCT11 Percent weights	RCT12 Percent weights	RCT13 Percent weights	RCT14 Percent weights	RCT15 Percent weights
Processed meat kcal/day	0.13	0.18	0.28	0.35	0.16
Red meat kcal/day	6.31	6.46	12.56	11.78	6.90
Fish kcal/day	0.01	0.02	0.02	0.01	0.01
Milk kcal/day	1.70	2.12	3.83	2.99	2.35
Poultry kcal/day available	7.81	6.50	11.94	11.25	7.17
Eggs kcal/day available	2.82	3.48	5.55	5.44	2.83
Alcohol kcal/day	0.21	0.06	0.60	0.13	1.35
Sugary beverages kcal/day	10.33	8.84	4.75	4.17	10.87
Corn kcal/day available	1.32	0.33	1.21	0.81	1.29
potatoes kcal/day available	1.73	2.12	7.81	2.91	0.01
Saturated fatty acids kcal/day	2.90	2.49	2.40	2.90	3.65
Polyunsaturated fatty acids kcal/day	0.58	0.70	0.56	0.44	0.88
Trans fatty acids kcal/day	0.05	0.15	0.03	0.08	0.17
Fruits kcal/day	0.28	0.43	0.34	0.50	0.75
Vegetables kcal/day	0.64	0.68	0.40	0.61	0.84
Nuts and seeds kcal/day	0.01	0.02	0.02	0.03	0.04
Whole grains kcal/day	0.16	0.39	0.22	0.21	0.06
Legumes kcal/day	0.38	0.89	0.77	0.29	0.65
Rice kcal/day available	9.30	10.10	8.34	9.08	7.09
Sweet potatoes kcal/day available	0.12	0.18	0.04	0.25	0.67
Physical activity METs	9.01	10.97	6.78	9.80	10.34
Kcal available/day	13.56	11.05	4.34	7.80	12.42
Child/infant 2SD underweight	8.29	8.08	4.01	4.05	7.08
Discontinued breast feeding <6 mo	0.32	0.20	0.58	0.47	1.52
Total percent weights in BMI formula	77.97	76.44	77.38	76.37	79.09

667 **Supplementary Table 7. continued**

Risk factors	RCT16 Percent weights	RCT17 Percent weights	RCT18 Percent weights	RCT19 Percent weights	RCT20 Percent weights
Processed meat kcal/day	0.33	0.21	0.29	0.43	0.26
Red meat kcal/day	13.14	7.99	7.48	14.07	11.69
Fish kcal/day	0.02	0.02	0.01	1.97	1.28
Milk kcal/day	3.65	2.02	4.49	4.10	3.12
Poultry kcal/day available	11.06	5.98	9.24	11.22	11.02
Eggs kcal/day available	4.90	3.25	4.17	7.73	5.61
Alcohol kcal/day	4.34	1.52	0.01	4.10	3.80
Sugary beverages kcal/day	1.58	12.85	7.01	3.37	6.49
Corn kcal/day available	2.00	0.42	1.95	0.17	0.77
potatoes kcal/day available	2.98	0.18	3.88	6.20	2.33
Saturated fatty acids kcal/day	1.20	5.43	1.81	0.90	2.87
Polyunsaturated fatty acids kcal/day	0.29	1.50	0.49	0.18	1.07
Trans fatty acids kcal/day	0.05	0.18	0.06	0.03	0.07
Fruits kcal/day	0.23	0.88	0.35	0.12	0.47
Vegetables kcal/day	0.32	1.17	0.54	0.23	0.38
Nuts and seeds kcal/day	0.02	0.05	0.03	0.01	0.04
Whole grains kcal/day	0.03	0.00	0.04	0.12	0.02
Legumes kcal/day	0.17	0.68	0.40	0.09	0.56
Rice kcal/day available	7.18	12.12	9.14	6.79	13.19
Sweet potatoes kcal/day available	0.02	1.13	0.04	0.22	0.21
Physical activity METs	13.29	4.32	12.97	9.16	9.16
Kcal available/day	8.12	9.24	7.45	6.10	4.78
Child/infant 2SD underweight	4.37	5.50	6.14	3.27	2.89
Discontinued breast feeding <6 mo	1.29	2.87	0.49	1.22	0.95
Total percent weights in BMI formula	80.57	79.47	78.50	81.80	83.04

669   Appendix 1. Worldwide surveys contributing to the IHME GBD risk
670   factor data.

671   Online only.

Appendix 2. Bradford Hill causality criteria based assessment methodology detailed

A literature search revealed no published methodological precedents for statistically modeling the correlation between mean BMIs of worldwide countries and subnational regions/provinces/states and their corresponding dietary and other risk factors. Since the Bradford Hill causality criteria[1] (enumerated by the English occupational physician and epidemiologist Sir Austin Bradford Hill) are the gold standard assessment tools to test causality of risk factors related to health outcomes, we explored the data looking for mean BMI to risk factor correlations that could be tested with the Bradford Hill criteria.

The relevant causality criteria included #1 strength, #2 consistency, #3 dose-response (biological gradient), #4 temporality, #5 analogy, #6 plausibility, #7experimentation, and #8 coherence. The non applicable criterion is specificity. Since diet, physical activity, and other risk factors may affect many non-communicable disease health outcomes, specificity is not relevant. As Dr. Bradford Hill said, "In short, if specificity exists we may be able to draw conclusions without hesitation; if it is not apparent, we are not thereby necessarily left sitting irresolutely on the fence."[1]

In considering many candidate methodologies involving univariate and multiple regression analysis, we sought a technique that produced the worldwide BMI versus risk factors multiple regression formula which most accurately predicted the mean BMI of subgroups (e.g., continents, socio-economic index quartiles, etc.). Such a BMI formula with quantifiable high accuracy in predicting the mean BMI of subgroups would necessarily score high with the

Bradford Hill criteria strength, consistency, and dose-response (biological gradient). Of the

candidate statistical modeling strategies, the methodology that created the worldwide BMI

formula that most accurately estimated subgroup mean BMIs had the following steps:

1. Transform all the food group risk factors (i.e., alcohol, total sugar, plant foods,

animal foods, and fatty acids) from g/day to kilocalories/day (kcal/day).

2. Multiply each food group risk factor by its mean kcal/day consumption.

3. Multiply each risk factor by the R^2 (coefficient of determination) of its

univariate correlation with BMI.

4. Adjust for multicollinearities (any animal or plant risk factors in the

combination variables that have significant multicollinearity (when an

independent variable is highly correlated with one or more of the other

independent variables in a multiple regression equation).

5. After performing #2, #3, and #4 above, form combination dietary variables

with individual foods each expressed in kcal/day to be added together as the

data suggests are appropriate:

712   a. For example: processed meat * processed meat kcal/day * processed

713   meat R^2 * processed meat kcal/day in bottom three SDI quartiles/top

714   quartile + red meat * red meat R^2 * red meat kcal/day * red meat

715   kcal/day in bottom three SDI quartiles/ kcal/day in the top quartile +

716   …+ eggs * eggs kcal/day, etc.

717     b. Explore methods to adjust for multicollinearities based in part on maximizing

718       the closeness of fit of the multiple regression derived BMI formula outputs

719       with the mean BMIs of 37 subgroups of the worldwide data.

720   6. Add physical activity before performing multiple regression.

721   7. Perform multiple regression analysis with the food combination variables in #5 and

722     physical activity to generate a BMI formula. (e.g., BMI formula= combination

723     variable$_a$ * parameter estimate$_a$ + combination variables$_b$ * parameter estimate$_b$ …+

724     physical activity * parameter estimate of physical activity.

725   8. Explore integrating the variables kcal available, child underweight, and discontinued

726     breast feeding into the BMI formula.

727   9. Discard any of these variables that's sign in regression with BMI switches from

728     univariate to the multivariate analysis (e.g., + in univariate and - in multivariate

729     formula).

730   10. To simplify the calculations, use the Excel spreadsheet and transfer back and forth to

731     SAS Studio.

732   11. To test the BMI formula with subgroups, match the BMI formula mean and standard

733     deviation (SD) with that of mean worldwide BMI:

734     a. Adjust the SD of the BMI formula output to equal the SD of the worldwide

735       BMI by multiplying the BMI formula output times the BMI SD / BMI

736       formula SD.

737     b. Adjust the BMI formula constant to equal the worldwide population's mean

738       BMI by adding the difference between the mean BMI and the BMI formula

739       constant.

12. Explore the BMI formula functionality with different Bradford Hill criteria with

standardising the BMI and BMI risk factors or with all variables non standardised.

We estimated percent weights attributable to each risk factor with the standardised BMI formula

from the above methodology with the following additions:

1. Equate the sum of the individual food and other risk factor parameter estimates to the

total R^2 of the BMI formula:

a. Total the absolute values of the parameter estimates for each food and any non

food risk factors.

b. Divide the worldwide BMI formula R^2 by the sum of the risk factor parameter

estimates.

c. Multiply the result of b above times each of the individual risk factor parameter

estimates.

2. Multiply the results in #1 above times 100 to generate percent weights for each risk

factor. The sum of the risk factor percent weights will equal the BMI formula R^2 times

100.

To cross validate the generated worldwide BMI formula experimentally, create 20 random

subgroups of 100 cohorts with the macro function in SAS. Determine each of the 20 subgroup

BMI formulas, and, from those 20 BMI formulas, create a table with: mean, SD, minimum, and

maximum for each of the risk factors.

Bradford Hill causal criteria testing methodology applied to current study

The relevant eight of the nine original Bradford Hill criteria were each scored as:

"5" very strongly supporting causality,

"4" strongly supporting causality,

"3" moderately strongly supporting causality,

"2" supporting causality,

"1" weakly supporting causality, and

"0" not supporting causality

The scoring for each Bradford Hill causal criterion was as follows:

1. Strength: The correlation coefficient, r, of the worldwide multiple regression derived

   BMI formula with BMI (dependent variable) and BMI risk factors (independent

   variables) assessed strength.

   Scoring of strength:

   5=BMI formula regressed with BMI r $\geq$0.50 and P<0.0001

   4=BMI formula regressed with BMI 0.50>r$\geq$0.40 and P<0.0001

   3=BMI formula regressed with BMI 0.40>r$\geq$0.30 and P<0.0001

   2=BMI formula regressed with BMI 0.30>r$\geq$0.20 and P<0.0001

   1=BMI formula regressed with BMI 0.20>r$\geq$0.10 and P<0.0001

   0=BMI formula regressed with BMI r< 0.10 or $P$$\geq$0.0001

2. Consistency: For the purposes of this study, consistency between BMI and BMI formula

output was determined by comparing the mean BMI and the mean BMI formula output in

each of the following 37 subgroups:

1) A variable socio-demographic index (SDI)—see Supplementary Table 1 for

definition of SDI—divided the world's population by quartiles of SDI.

2) A variable "continents" allowed for analyses of countries from each of the six

inhabited continents.

3) The four countries (UK, USA, Mexico, and Japan) with subnational data on BMI and

the risk factors were grouped and assessed to compare the BMI formula output with

the overall mean BMI in those countries.

4) Based on the total kcal/day of all foods that increased BMI, a combination variable

was constructed and the world's population divided into quartiles from the highest to

lowest total kcal/day.

5) Similarly, based on the total kcal/day of all foods that decreased BMI, we divided the

world's population into quartiles from the highest to lowest by total kcal/day.

6) Based on physical activity (METs/week), the world's population was divided into

quartiles.

7) We evaluated dose response by dividing the BMI formula output into quartiles from

the highest to lowest.

8) The four countries with subnational data were individually evaluated.

9) The first four of the 20 random number generated database subgroups were included

in the consistency analysis.

10) Male and female were separately assessed.

11) Assessing dose-response was done in quartiles by the BMI formula outputs after the

BMI formula was harmonized with the mean and SD of worldwide mean BMI. See

the dose-response criteria (#4).

For each of the 37 subgroups, the absolute differences between the means of BMI and the

BMI formula output in were totaled (e.g., continent Africa BMI formula output - mean

BMI for Africa, etc.).

Scoring of consistency for BMI compared with BMI formula output for each of the 37

subgroups:

5=The mean of the absolute differences between mean BMI and BMI formula output was

≤ 0.40 units.

4=The mean of the absolute differences between mean BMI and BMI formula output was

≤ 0.50 units.

3=The mean of the absolute differences between mean BMI and BMI formula output was

≤ 0.60 units.

2=The mean of the absolute differences between mean BMI and BMI formula output was

≤ 0.70 units.

1=The mean of the absolute differences between mean BMI and BMI formula output was

≤ 1.0 units.

0=The mean of the absolute differences between mean BMI and BMI formula output was

> 1.0 units.

3. Dose-response (biological gradient): Dr. Bradford Hill thought that a clear dose-response

effect on the incidence of disease with exposure to a single risk factor was the clearest

evidence of a causal relationship. More recently, it has been realized that cause and effect

relationships are often more complex. In this analysis of dose-response, instead of using

single risk factor levels related to BMI, levels of a multivariable regression derived BMI

formula outputs in quartiles were related to mean BMIs in those quartiles. Dose-response

of BMI formula estimates versus mean BMI were included in the testing of consistency

above.

Scoring of dose-response (biologic gradient) was based on this mean absolute difference

when the BMI formula output was divided into quartiles:

5= The BMI formula output versus mean BMI absolute differences from each of the four

quartiles averages ≤ 0.40 BMI units.

4= The BMI formula output versus mean BMI absolute differences from each of the four

quartiles averages ≤ 0.50 units.

3= The BMI formula output versus mean BMI absolute differences from each of the four

quartiles averages ≤ 0.60 BMI units.

2= The BMI formula output versus mean BMI absolute differences from each of the four

quartiles averages ≤ 0.70 BMI units.

1= The BMI formula output versus mean BMI absolute differences from each of the four

quartiles averages ≤ 1.0 BMI units.

0= The BMI formula output versus mean BMI absolute differences from each of the four

quartiles averages > 1.0 BMI units

4. Temporality: The dictionary defines the noun temporality as: "The state of existing

within or having some relationship with time." Bradford Hill said, "Temporality refers to

the necessity that the cause to precede the effect in time. This criterion is unarguable,

insofar as any claimed observation of causation must involve the putative cause C

preceding the putative effect D. It does not, however, follow that a reverse time order is

evidence against the hypothesis that C can cause D. Rather, observations in which C

followed D merely shows that C could not have caused D in these instances; they provide

no evidence for or against the hypothesis that C can cause D in those instances in which it

precedes D."

Dr. Bradford Hill was an occupational physician before the current availability in

nutritional epidemiology of data on trends over 28 years of 18 components of worldwide

diets along with the global BMI trend. Consequently, now it is fair to test temporality by

deriving a standardised multiple regression formula with BMI trend, measured by the

slope of the least squared regression line (LSRL) over 1990-2017, as (dependent

variable). The independent variables would consist of the LSRL trends over 1990-2017

of the same dietary components, physical activity, and other variables as in the original

BMI formula. We considered the strength (r) of the BMI trend formula versus the BMI

trend an appropriate measure of temporality. Any risk factor was excluded from the trend

BMI formula if its sign (+ or -) did not match the sign in the original BMI formula.

With a standardised multiple regression methodology similar to that of deriving risk

factor percent weights above, a standardised BMI trend (dependent variable) versus

875  standardised risk factor trends (independent variables) formula was derived with the risk

876  factor parameter estimates adjusted to equate to trend percent weights.

877  Scoring of temporality: The Pearson correlation coefficient, r, of the resulting multiple

878  regression derived BMI trend versus risk factors trends formula:

879  $5 = r \geq 0.50$ and $P < 0.0001$.

880  $4 = 0.50 > r \geq 0.40$ and $P < 0.0001$.

881  $3 = 0.40 > r \geq 0.30$ and $P < 0.0001$.

882  $2 = 0.30 > r \geq 0.20$ and $P < 0.0001$.

883  $1 = 0.20 > r \geq 0.10$ and $P < 0.0001$.

884  $0 = r < 0.10$ or $P \geq 0.0001$

886 5. Analogy: High BMI is among the four metabolic risk factors that are strongly associated with

887  cardiovascular diseases, cancers, and other non-communicable diseases. The other major

888  metabolic risk factors for non-communicable diseases are high systolic blood pressure (SBP),

889  high low density lipoprotein cholesterol (LDL-C), and high fasting plasma glucose (FPG).

891  We tested analogy by the number of the diet, physical activity, and other variables in the

892  percent weight of risk factor BMI formula that were also in the percent weight of risk factor

893  formulas of the other metabolic factors and including only risk factors concordant in

894  direction (+ or -) with the coefficients of percent weights of risk factors in the BMI formula.

895  Scoring of analogy:

5=At least three-quarters of BMI formula risk factors are also in the risk factor formulas

of all three of the other metabolic factors (SBP, LDL-C, and FPG) and have concordant

signs (+ or -) for each risk factor.

4= At least three-quarters of BMI formula risk factors are also in the risk factor formulas

of two of the three other metabolic factors (SBP, LDL-C, and FPG) and have concordant

signs (+ or -) for each risk factor.

3= At least three-quarters of BMI formula risk factors are also in the risk factor formulas

of one of the three other metabolic factors (SBP, LDL-C, and FPG) and have concordant

signs (+ or -) for each risk factor.

2= At least two-thirds of BMI formula risk factors are also in the risk factor formulas of

one of the three other metabolic factors (SBP, LDL-C, and FPG) and have concordant

signs (+ or -) for each risk factor.

1=At least half of BMI formula risk factors are also in the risk factor formulas of one of

the three other metabolic factors (SBP, LDL-C, and FPG) and has concordant signs (+ or

-) for each risk factor.

0= None of the above.

6. Plausibility: To test plausibility, we looked to find if any of our findings were at variance

with the preponderance of studies published. We searched the medical literature

particularly for systematic reviews of the relationships of foods and other variables with

BMI.

Scoring of plausibility:

5=None of the current findings were at variance with the preponderance of the medical

literature.

4= One of the current findings was at variance with the preponderance of the medical

literature.

3= Two of the current findings were at variance with the preponderance of the medical

literature.

2=Three of the current findings were at variance with the preponderance of the medical

literature.

1=Four of the current findings were at variance with the preponderance of the medical

literature.

0= Five or more of the current findings were at variance with the preponderance of the

medical literature.

7. Experiment: Dr. Bradford Hill thought that evidence drawn from experimentation,

including in epidemiologic studies, may lead to the strongest support for causal

inference.[1]

We used a cross validation method to assess Bradford Hill's "experiment" criterion.

Random number generation of 20 subgroups each with 100 cohorts derived 20

standardised BMI formulas to compare with the standardised worldwide BMI formula.

Scoring of experiment:

5=At least 15 subgroups BMI formulas included all same risk factors with the same signs

of coefficients as the worldwide BMI formula.

4=At least 10 subgroups of BMI formulas included all same risk factors with the same

signs of coefficients as the worldwide BMI formula.

3=All 20 subgroups of BMI formulas included at least three-quarters of same risk factors

with the same signs of coefficients as the worldwide BMI formula.

2=At least 10 subgroups of BMI formulas included at least three-quarters of same risk

with the same signs of coefficients factors as the worldwide BMI formula.

1= At least 10 subgroups of BMI formulas included at least half of same risk factors with

the same signs of coefficients as the worldwide BMI formula.

0=None of the above.

8. Coherence: According to Dr. Bradford Hill, "…cause and effect interpretation of our data

should not seriously conflict with the generally known facts of the natural history and

biology of the disease."[1] In this analysis of BMI associated with BMI formula estimates,

coherence was the numerical total score of the above seven relevant causality criteria

each on a 0-5 scale. The maximum score was 40.

Scoring of coherence:

5=Score on the first seven Bradford Hill causation criteria=35-40.

4=Score on the first seven Bradford Hill causation criteria=30-34.

3=Score on the first seven Bradford Hill causation criteria=25-29.

2=Score on the first seven Bradford Hill causation criteria=20-24.

1=Score on the first seven Bradford Hill causation criteria=15-24.

0=Score on the first seven Bradford Hill causation criteria<15.

Reference

1. Hill AB. **THE ENVIRONMENT AND DISEASE: ASSOCIATION OR CAUSATION?**

Proceedings of the Royal Society of Medicine. 1965; **58**(5):295-300.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1898525/

969
970