A multi-phenotype genome-wide association study of clades causing tuberculosis in a Ghanaian- and South African cohort

Stephanie J. Müller [a, b, †,1], Haiko Schurz[a, b], Gerard Tromp[a, b], Gian D. van der Spuy [a, b], Eileen G. Hoal [a], Paul D. van Helden [a], Ellis Owusu-Dabo [c], Christian G. Meyer [d, e], Thorsten Thye [f], Stefan Niemann [g], Robin M. Warren [a], Elizabeth Streicher [a], Marlo Möller [a], Craig Kinnear [a]

[a] DSI-NRF Centre of Excellence for Biomedical Tuberculosis Research, South African Medical Research Council Centre for Tuberculosis Research, Division of Molecular Biology and Human Genetics, Faculty of Medicine and Health Sciences, Stellenbosch University, Cape Town, South Africa

[b] South African Tuberculosis Bioinformatics Initiative (SATBBI), Faculty of Medicine and Health Sciences, Stellenbosch University, Cape Town, South Africa

[c] School of Public Health, College of Health Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana

[d] Institute of Tropical Medicine, Eberhard-Karls University Tübingen, Germany

[e] Faculty of Medicine, Duy Tan University, Da Nang, Vietnam

[f] Bernhard Nocht Institute for Tropical Medicine, Hamburg, Germany

[g] German Centre for Infection Research (DZIF), Partner site Hamburg-Lübeck-Borstel, Borstel, Germany

* Correspondence: Stephanie Julia Müller

E-mail address: steph15pitts@gmail.com, Telephone number: +27 83 541 8269

[1] Present address:

IBM Research Africa, 45 Juta street, Braamfontein, Johannesburg, South Africa

Abstract

Despite decades of research and advancements in diagnostics and treatment, tuberculosis remains a major public health concern, particularly in low- and middle-income countries. New bioinformatics and computational methods are needed to interrogate the intersection of host- and bacterial genomes and identify novel targets for anti-tuberculosis drugs. Host genotype datum and paired infecting bacterial isolate information were analysed for associations using a multinomial logistic regression framework implemented in SNPTest. Two geographically distinct cohorts were evaluated: a cohort of 947 participants self-identifying as belonging to a five-way admixed South African population and a Ghanaian cohort consisting of 3 311 participants. We report potential associations between host genetic variants and multiple members of the *Mycobacterium tuberculosis* complex (MTBC). Although none of the variants analyzed in the South African cohort passed the GWAS cut-off for significance, 32 single nucleotide polymorphisms were identified in the Ghanaian cohort as being statistically significantly associated with risk for infection with strains of different members of the MTBC. Further analysis revealed that two of these SNPs were directly genotyped, and the rest were imputed using the 1000 Genomes Phase 3 reference panel. The availability of paired host-pathogen data is imperative for investigating strain-specific interactions between MTBC and its host. As demonstrated by this study, the implementation of a multinomial logistic regression using paired host-pathogen data may prove valuable for further research investigating the complex relationships driving infectious disease.

Keywords

Multi-phenotype GWAS; *Mycobacterium tuberculosis* complex; Imputation; South Africa; Ghana

1. Introduction

Tuberculosis (TB), a disease primarily affecting the lungs, is caused by pathogenic members of the *Mycobacterium tuberculosis* complex (MTBC) such as *M. africanum* (*M. africanum*) and *M. tuberculosis* (*M. tb*). Infection alone, however, is not sufficient to cause disease (Bellamy, 1998; Kinnear et al., 2017; Möller and Hoal, 2010). Each branch of the MTBC comprises several clades of specific strains with variable virulence and disease-causing mechanisms (Brites and Gagneux, 2015; Gagneux, 2012; Hoal et al., 2017). While *M. africanum* is the main cause of TB in West-African countries including Ghana, *M. tb* is responsible for TB cases in most other parts of the world (Gagneux and Small, 2007). In addition to socio-economic- and environmental factors (Pratiwi, 2016; Seddon et al., 2013), predisposing diseases (Cheng et al., 2016; Lim et al., 2017), and the genetic make-up of the human host (Chimusa et al., 2014; Kinnear et al., 2017; Oki et al., 2011; Png et al., 2012; Salie et al., 2014; Thye et al., 2012, 2010) have been shown to play pivotal roles in determining susceptibility to the disease.

Several associations between genomic loci and susceptibility to infectious diseases such as TB (Bellamy, 1998; Herb et al., 2007; Thye et al., 2010), malaria (Rockett et al., 2014) and HIV (Pastinen et al., 1998) have been reported using candidate-gene association studies, linkage studies, and genome-wide association studies (GWAS). Using these approaches, a number of genes encoding proteins of the host immune system have been associated with susceptibility to TB, including human leukocyte antigens (HLAs), *NRAMP1*, mannose binding lectin (*MBL*), IFN-gamma (*IFN*-ɣ), and Vitamin D Receptor (*VDR*) (Bellamy, 1998; Salie et al., 2014; Søborg et al., 2003; Yim and Selvaraj, 2010). While several studies have investigated the genetic association with TB (Chimusa et al., 2014; Hoal et al., 2017; Hong et al., 2017; Kinnear et al., 2017; Schurz et al., 2019a), few have investigated the association between gene variants and susceptibility to particular strains of the MTBC (McHenry et al., 2019; Omae et al., 2017).

Candidate gene studies compare allelic and genotyping frequencies of a specific genetic marker between a group of unrelated cases and controls. In a large cohort of 1 916 sputum-positive Ghanaian TB patients genotyped for the *ALOX5* g.760G>A variant, individuals who were heterozygous for the polymorphism were found to be at increased risk for developing TB (Herb et al., 2007). Furthermore, patients harboring the exonic variant (g.760A) had a greater association (OR= 1.70; [95% CI: 1.2–2.6]) with infection caused by *M. africanum* West Africa-2 (Herb et al., 2007). Modelling a recessive mode of inheritance, a protective

association (OR= 0.60; [95% CI: 0.4–0.9]) was identified between the occurrence of TB and

the *MBL2* G57E variant in another cohort of Ghanaian patients (Thye et al. 2011). TB patients

belonging to the Ewe population were significantly more likely to be infected with *M.*

africanum (OR= 3.02; [95% CI: 1.67–5.47]) and further stratification by lineage revealed that

the association was strongly driven by infection with members of *M. africanum* West Africa-1

(Asante-Poku et al., 2015). *HLA* types are also known to be important in the immune response

to pathogens.

In a South African candidate-gene association study of *HLA* alleles and the *M. tb* strain

responsible for active TB, the *HLA-B27* allele was found to decrease risk for an additional

disease episode due to a Beijing strain, following multiple episodes of disease caused by a

Beijing strain (Salie et al., 2014). In addition, specific *HLA* types were found to be associated

with disease caused by the different strains investigated (Salie et al., 2014). Finally, Caws and

colleagues were investigated the susceptibility of the human host to different *M. tb* strains.

Using a candidate-gene approach, a cohort of 237 adult Vietnamese TB patients were analysed.

The authors concluded that for this cohort, individuals carrying the C allele of the toll-like

receptor-2 (*TLR2*) T597C polymorphism were significantly more likely to develop TB caused

by mycobacteria belonging to the East-Asian/Beijing strain family (OR = 1.57 [95% CI 1.15-

2.15]) (Caws et al., 2008).

A limitation of the candidate-gene study design however, is that it requires an *a priori*

hypothesis regarding which genes to target in the association analysis. To address this

limitation, GWAS have become a popular alternative for identifying genetic associations with

disease. Through genotyping of many common genetic variants, GWA studies enable a global

interrogation of a host's genome for associations to disease, without the limitation of

predefined candidate genes (Hirschhorn and Daly, 2005). While genome-wide associations

between the human host and TB have been extensively studied in several populations, the

susceptibility of the host to different members of the MTBC has only in recent years gained

some attention.

The first GWAS to investigate genetic susceptibility to strains of different MTBC lineages

aimed to identify genome-wide associations with TB onset, stratifying a Thai cohort by

infecting MTBC lineage, and the age at onset (Omae et al., 2017). The study initially attempted

to identify age-related associations between five MTBC lineages and two age-stratified groups

of TB participants, namely 219 young cases (under the age of 45), and 467 old cases (over the age of 45). To reduce the complexity of the association tests, the MTBC lineages were tested as one lineage versus a collective of all other lineages. After applying Bonferroni corrections, none of the genotyped single nucleotide polymorphisms (SNPs) reached genome-wide significance for association with either of the age-groups for any of the five MTBC lineages. However, when reducing the five lineages to two groups consisting only of 'Beijing' and 'non-Beijing' cases, and testing for age-related association to TB, the authors identified a single SNP on chromosome 1p13, rs1418425, reported to have a significant association to non-Beijing infected cases classified in the "old" age category ($P = 1.58 \times 10^{-7}$; OR = 1.62 [95% C.I.: 1.35-1.93]). The authors were able to replicate the SNP in two independent cohorts, further demonstrating the importance of performing GWAS with a specific focus on pathogen lineage (Omae et al., 2017).

Another study investigated the coevolution of *M. tb* and it's human host. The article is based on the hypothesis that longstanding coexistence between the human genome and *M. tb* lineage may reduce the risk of progressing to active TB or minimize the severity of disease. The authors investigated TB severity (as measured by the TBScore) in two cohorts from Uganda with paired *M. tb*-human DNA available to determine if interactions between *M. tb* lineage and human genetic variants exist. Although no association was found between lineage and disease severity, an interaction between a single nucleotide polymorphism (SNP) in SLC11A1 and the L4-Ugandan lineage were identified in both cohorts. In addition several IL12B polymorphisms were associated with disease severity (McHenry et al., 2019).

In order to improve our understanding of the genetic susceptibility to the MTBC clades, this study leveraged genome-wide genotyping data from the host and pathogen data to perform a genome-wide screen for clade-specific genetic associations in cohorts originating from two distinct populations.

2. Results

2.1. Participant recruitment, sample collection, and SNP genotyping

All participants recruited for the South African and Ghanaian cohorts provided blood and sputum samples for SNP genotyping, and MTBC strain identification, respectively. The South African cohort comprised 947 participants successfully genotyped for 397 337 variants, while the Ghanaian cohort consisted of 3 311 participants successfully genotyped for 783 338

165   variants (Table 1). Of the Ghanaian samples included in this study, 1 359 were TB cases and

166   69% of the participants were male (Table 1). Principal Component Analysis of the Ghanaian

167   cohort showed contributing ethnicities from the Akan, Ga-Adangbe, Exe, and several other

168   ethnic groups from northern Ghana (Thye et al. 2012).

170   2.2. Defining MTBC clades and superclades

171   2.2.1.   South Africa

The MTBC strains obtained in the South African cohort contained strains of eight of the 12

lineages, namely Beijing, CAS (represented as CAS1 in the infection database), Haarlem,

Haarlem-like, Low-copy Clade (LCC), T, and Quebec (Fig. 2A). During the grouping strategy,

Beijing and CAS were clustered to form the "BeijingCAS1" superclade (Fig. 2B). Similarly,

Haarlem, Haarlem-like, and LCC cases were clustered to form the "HaarlemsLCC" superclade

(Fig. 2B). A clade denoted as "Other" was also present in the South African cohort but does

not appear on the phylogenetic tree (Fig. 1) and thus was kept as a distinct member during the

grouping strategy. The T superclade was excluded from subsequent analysis due to low

frequency in the cohort after clustering into superclades, leaving five superclades represented

by this cohort (Fig. 2B). MTBC clade distributions were dominated by the LAM clade and

closely followed by Beijing. Superclade distributions showed similar frequencies for LAM,

BeijingCAS1, and HaarlemsLCC, while the Quebec and "Other" clades were the least

abundant (Fig. 2B).

186   2.2.2.   Ghana

The strains obtained in the Ghanaian cohort contained 12 clade annotations obtained from

spoligotyping. The afri-181 and afri-438 clades were represented by *M. africanum* on the

phylogenetic tree and were subsequently grouped with EAI at the point of divergence, and

named as the EAI_afri superclade (Fig. 1). Beijing and CAS were merged, as were Haarlem

and X, into the "BeijingCAS', and "HaarlemX" superclades, respectively. T and U clades were

also clustered (Fig. 1). The "Ghana-2" clade was kept as a distinct superclade, while LAM and

CAM were grouped based on the similarity in their spoligotyping patterns as illustrated in

Stucki *et al.* 2016. This cohort thus contained 12 clades and six superclades after clustering

(Fig. 2C, and Fig. 2D, respectively).

After grouping the clades into superclades, the strains of the EAI_afri and LAM_CAM

superclades occurred most frequently in the cohort with both superclades having a frequency

greater than 400 in the dataset (Fig. 2D). The HaarlemX, and T_U superclades had a frequency
of approximately 160, and 250 in the cohort, respectively, while the BeijingCAS and Ghana-2
superclades were in least abundance (Fig. 2D).

2.3. Genotype data quality control, haplotype phasing, and genotype imputation
For the South African cohort, analysis using Genotype Harmonizer yielded a dataset of 947
participants with 356 165 variants. After quality control filters were applied, 919 participants
and 239 612 variants remained. For the Ghanaian cohort, similar analysis yielded a dataset of
3 311 participants with 713 223 variants, and genotype QC filters resulted in a further reduction
to 617 409 variants. No additional QC steps were used for the IH protocol, while the results of
the additional data preparation steps for both cohorts imputed using the MIS tool are detailed
in Table 2.

2.4. Selection of high-quality imputed genotype data
Imputation results are presented for Chromosomes 1, and X for the South African cohort, while
for the Ghanaian cohort, Chromosome X data was not available, and thus the imputation of
two autosomes are reported Table 3.

SNP densities were calculated for the proportion of SNPs with a quality score metric greater
than 0.45. For the South African cohort, the SIS workflow using the AGR resource imputed
the highest proportion of SNPs, whereas for the Ghanaian cohort, the MIS workflow using the
CAAPA resource imputed the greatest proportion of SNPs with a quality metric greater than
0.45 (Table 3). For both chromosomes 1 and X, imputation using either the 1000G or the
CAAPA resource with the MIS performed the worst for the South African cohort (Schurz et
al., 2019b) with the maximum median quality score only reaching 0.82 at a MAF of 50%. In
comparison, the SIS-AGR workflow outperformed all other workflows, and the result
correlated with the AGR imputing the highest SNP density for chromosome 1 and chromosome
X (Schurz et al., 2019b).

For the South African cohort, genotype datum imputed with the AGR reference panel was
selected as the dataset with the highest imputation quality across all reference panels assessed.
After removal of monomorphic sites and filtering on an INFO score of 0.45, 28 566 283 SNPs
for 919 samples remained. After removing the 136 related individuals identified prior to
imputation, and filtering for SNP- and sample missingness, and MAF, a dataset of 7 145 406

variants for 783 participants remained. Of the 525 clade-matched samples, 445 were extracted from the dataset of samples passing QC and used in the association analysis.

For the Ghanaian cohort, despite the CAAPA resource imputing the greatest SNP density (Fig. 3D and Fig. 4D), the IH workflow using the 1000G reference panel imputed the highest quality of SNPs per MAF bin but was very closely followed by the other workflows and reference panels from the 20-30% MAF bin upwards (Fig. 5). Thus, the IH dataset, imputed with the 1000G reference panel was selected as the best dataset. After filtering monomorphic SNPs, INDELS, and variants not reaching the INFO score threshold, 25 968 622 SNPs remained for 3 239 samples. After filtering for MAF, SNP- and sample missingness, and removal of 93 related individuals identified pre-imputation, the dataset comprised of 5 275 890 variants for 1 273 clade-matched samples.

2.5. Covariable data

For the South African cohort, the covariables age, sex, and ancestry proportions were available for all samples. Ancestry proportions were for the European, African, San, South-Asian, and East-Asian ancestries. Of the 445 clade-matched samples that passed the post-imputation QC, ancestry proportions were available for 357 samples as generated previously using the Affymetrix genotype datum for this cohort and ADMIXTURE software (Daya et al., 2013). For the remaining 88 samples, ancestry proportions were calculated from genotype datum generated by the MEGA array using ADMIXTURE. The East-Asian ancestry, being the smallest contributing ancestry proportion, was not included as a covariable in the analysis. Variances were calculated for each of the four remaining ancestry proportions and determined to be 0.027 (San), 0.035 (African), 0.014 (European), and 0.009 (South-Asian). As the variances were greater than the minimum cut-off of 0.001, they were included as covariables in the analysis.

For the Ghanaian cohort, age, sex, and ethnicity were included as covariables. Further examination of admixture for the Ghanaian dataset revealed that the cohort was not highly admixed, and thus ethnicity in the form of principal components was used. One sample passing QC filters did not have one of the covariables and was excluded from the dataset leaving 1 272 samples for the association analysis. The variance in the PCs provided for the cohort was calculated to be 0.0002 (PC1), 0.0003 (PC2), and 0.0004 (PC3) and therefore determined to be insufficient for inclusion in the association analysis as covariables.

2.6. Genome-wide Association Analyses

2.6.1.   South Africa

An MLR was conducted for the 445 superclade-matched South African samples using

SNPTEST under an additive model. All results were reported using the LAM superclade as the

baseline. Although none of the SNPs passed the GWAS cut-off for significance, 4 631 SNPs

had an LRT p-value less than 0.0005 and eleven SNPs had an LRT p-value less than 1×10^{-6}

(Table 4). Odds ratios (OR) are reported (Table 4) and standard errors of the odds ratios are

shown in Fig. 6A. A single SNP, rs9389610, located on chromosome 6, had a p-value of

1.60×10^{-7}. Individuals with the A allele of this SNP were twice as likely to be have TB due to

infection with a member of the BeijingCAS1 superclade (OR: 2.19) than due to either the

HaarlemsLCC (OR: 1.07) or LAM superclades. Individuals with the A allele were only slightly

more at risk of being infected with a member of the 'Other' superclade (OR: 2.78), when

compared to the BeijingCAS1 superclade (OR: 2.19), and were very unlikely to be infected

with a member of the Quebec superclade (OR: 0.25).

For the four SNPs located on chromosome 5, individuals with the risk allele doubled the

chances of being infected with the Quebec superclade (OR: ~2) while halving the risk of being

infected with a member of the "Other" superclade (OR: ~0.5) (Table 4). Lastly, for the six

SNPs located on chromosome 17, the risk allele was shown to double the risk of being infected

with a member of the LAM superclade than with the HaarlemsLCC (OR: ~0.50) superclade.

Individuals with the risk allele of these six SNPs were also equally at risk of being infected

with a member of the BeijingCAS1 or LAM superclade and were twice as likely to be infected

with the member of the "Other" superclade (OR: ~ 2) when compared to the BeijingCAS1

superclade (OR: ~ 1) (Table 4).

The VEP tool was used to retrieve gene annotations for SNPs of interest. While these SNPs are

unlikely to have a direct effect on the gene expression itself, the SNP may be in linkage

disequilibrium with other nearby SNPs which do have a direct effect on the gene. For the South

African cohort, the most significantly associated SNP was rs9389610 (g.139039029G>A),

located on chromosome 6. This SNP is an imputed SNP and its two closest directly genotyped

SNPs were rs4896385 (g.139011266G>T), and rs7742202 (g.139074280A>G). The rs4896385

SNP is located in *NHSL-1*, while the rs7742202 SNP is located in *GVQW2*. Neither of these

genes have been previously shown to be involved in the pathogenesis of TB. The four SNPs

located on chromosome 5 (Table 4) were annotated to the StAR Related Lipid Transfer Domain

Containing 4 gene *(STARD4)* using the VEP tool, while the six SNPs located on chromosome

17 were annotated to the *TANC2* gene.

2.6.2. Ghana

Using SNPTEST under an additive model, an MLR was also conducted for 1 272

superclade-matched Ghanaian samples. For this cohort, all association results were reported

using the LAM_CAM superclade as the baseline. In summary, a total of 32 SNPs had an LRT

p-value less than 1×10^{-6} (Table 5) and were significantly associated with the MTBC

superclades.

Several imputed SNPs were shown to dramatically increase the risk of being infected by a

particular superclade. For example, the risk allele of the SNP rs577800201 (g.20476046C>T)

was shown to increase the risk of being infected with the EAI_afri superclade by 93 times,

compared to the baseline LAM_CAM superclade, and was annotated by the VEP to map to

ACSM2A. The risk allele of the SNP rs374315920 (g.38496435C>T) located on chromosome

17, was also found to increase an individual's risk of being infected with the EAI_afri

superclade by more than 500 times, as compared to the LAMCAM reference superclade, and

the MLR specific for this SNP was highly significant with an LRT p-value of 1.68×10^{-255}.

Using the VEP tool, this SNP was annotated to lie within the retinoic acid receptor alpha

(RARA) gene. Since these SNPs were imputed, their associations should be considered

tentative, because they may have been incorrectly imputed into this study dataset.

Further analysis revealed that of the 32 SNPs identified here, two, namely the intergenic variant

rs529920 on chromosome 6, and the intron variant rs41472447 on chromosome 12, were

directly genotyped. The risk allele of the SNP rs529920 (g.153835125A>G) was found to half

the risk of being infected with a member of the BeijingCAS and EAI_afri superclades when

compared to the LAM_CAM baseline superclade. Individuals with the risk allele for this SNP

were also equally at risk of being infected with a member of the Ghana2, HaarlemX, or T_U

superclades (OR: 1.04-1.24) (Table 5). Additionally, this SNP has an MAF of 0.4721 in the

study cohort, which is similar to the MAF observed in African populations in the 1000G

(MAF: 0.571) and in gnomAD genome (MAF:0.582). However, this SNP has no known gene

consequence to date.

Similarly, the risk allele of the intron variant rs41472447 (g. 41708761A>G) was found to

nearly triple the risk of infection with a member of the BeijingCAS superclade (OR: 2.56, C.I.:

(1.48–4.41)) or the Ghana2 superclade (OR:2.94, C.I.: (1.71–5.08)). In contrast, the risk allele

halved the risk of infection with the T_U superclade (OR:0.59, C.I.: 0.44-0.79) and had no

effect on the risk for infection with members of the EAI_afri or HaarlemX superclades

(Table 5). This variant had an MAF of 0.2461 in the study dataset which is similar to the MAF

observed in African populations in the 1000G (MAF: 0.200) and in the gnomAD genome

(MAF:0.165). Furthermore, rs41472447 maps to the *PDZRN4* gene in dbSNP, but has to date

not been linked to susceptibility to tuberculosis.

3.   Discussion

TB is a highly infectious disease affecting millions of people each year. The genetic

susceptibility of the host to developing the disease has been extensively studied using linkage

analysis, candidate gene studies, and GWAS. Furthermore, a number of selected genes have

been investigated for their contribution to genetic susceptibility of the host to strains of

different lineages of the MTBC. However, to date, no genome-wide analysis of genetic markers

affecting susceptibility to strains of different MTBC lineages has been performed.

3.1. Imputation

While several reference panels exist to facilitate genotype imputation, most of these panels

focus on representing populations of European ancestry, and little representation has been

made for African populations. Therefore, the present study focused on evaluating the quality

of imputation attainable for the five-way admixed South African population and the Ghanaian

cohort using the 1000G, AGR, and CAAPA reference panels.The more admixed a population

is, the greater the heterogeneity in its haplotype block structure. This genetic complexity

requires large reference panels with suitable ancestry to facilitate accurate genotype imputation

(Lin et al., n.d.). The five-way admixed South African population contains genetic

contributions from Bantu-speaking Africans, Europeans, KhoeSan, and South- and East-

Asians (Daya et al. 2013; De Wit et al. 2010). Imputation was previously performed for this

population; however, it was done using the 1000G Phase 1 (The 1000 Genomes Project

Consortium 2012) and the HapMap3 release 2 (The International HapMap 3 Consortium 2010)

reference panels. The 1000G panel has been expanded substantially since whereas the

HapMap3 reference panel, representing individuals mostly of European ancestry (Chimusa et

al. 2014), has since been deprecated.

When evaluating the Ghanaian cohort, although imputation of the 1000G reference panel with
the IH method performed the best, there was very little difference in the median quality scores
for the different workflows seen for SNPs with an MAF of 10-50% (Fig. 5). For rare variants
(MAF 0-5%) however, the IH method outperformed all others with a median quality score
above 0.75, whereas both analyses with the MIS produced a median score below the cut-off of
0.45. Thus, for the Ghanaian cohort, all reference panels and methods tested could be
considered viable options for imputing common variants with an MAF of 10-50% but should
be considered carefully for variants with an MAF below 10%.

In contrast to the AGR which contains no individuals recruited from West-African countries,
the CAAPA resource contains 88 individuals recruited from the West-African country of
Nigeria (Mathias et al., 2016). Thus, it was unsurprising that the CAAPA resource performed
well when imputing SNPs with a MAF above 10% (Fig. 5). From an MAF of 20-50%, the
CAAPA resource performed similarly to the other three reference panels and may thus be
considered suitable for imputing cohorts of West-African ancestry, such as the Ghanaian cohort
used in this study.

3.2. Multi-phenotype GWAS
In this study, the MLR functionality within SNPTEST enabled the genome-wide investigation
of genetic markers for association to a number of MTBC superclades. Although none of the
SNPs passed the GWAS p-value cut-off, the most significant associations for the South African
cohort imputed with the AGR reference panel were reported. This was likely due to the small
sample size resulting in a subsequent reduction in statistical power. In contrast, 32 SNPs passed
the GWAS cut-off for the Ghanaian cohort and may be considered as potential targets for
further investigation of host-directed therapies suitable for individuals of West-African
descent. None of the SNPs with an LRT p-value less than 0.0005 in either cohort were found
in the other, demonstrating the population-specific association of SNPs with the strains of
different MTBC superclades, which has been previously shown in the investigation of TLRs
and their association with cases of TB in populations of different ethnicities (Schurz et al.
2015).

3.3. Potential drug targets

For the Ghanaian cohort, 32 SNPs with significant LRT p-values were identified as being associated with the MTBC superclades investigated (Table 5). Nine of the SNPs located on chromosome 12 mapped to the *PDZRN4* gene. For these nine SNPs, the risk allele increased the chances of individuals being infected with the BeijingCAS superclades 2.5 times, and in the region of three times for the Ghana2 superclade, while the risk allele halved the chances of being infected with the T_U superclade. Due to the low frequencies of the BeijingCAS and Ghana2 superclades observed for this cohort (Fig. 2D), it is possible that these odds ratios were inflated because of small sample sizes. Notably, the two SNPs which were directly genotyped, and not imputed, were found to be significantly associated with infection with particular MTBC superclades. This provided crucial evidence that despite the vast amount of imputed genotype data included in the MLR, the association analysis was able to detect two directly genotyped SNPs with potentially significant associations with the MTBC superclades.

Few studies have described the direct influence of *M. tb* infection on the expression of the *STARD4* and *RARA* genes, and no studies have investigated the outcomes of infection with strains belonging to different MTBC clades on these genes. The *STARD4* encodes the StAR-related lipid transfer protein which plays a crucial role in the transmembrane trafficking of lipids (such as cholesterol) - an important source of energy for *M. tb* (Soccio et al. 2002). Infection of macrophages with pathogens such as *M. tb* stimulates the process of lipid droplet formation (Daniel et al. 2011). It has been hypothesised that *M. tb* initiates this process in order to secure a reliable source of carbon to fuel bacterial growth (Brzostek et al. 2009). Additionally, the accumulation of cholesterol in the bacterial cell wall drastically reduces the permeability of the cell wall, subsequently reducing the penetrating capability of the anti-TB drug Rifampicin (Brzostek et al. 2009). However, a recent study has contradicted the notion that lipid droplet formation is a bacteria-driven process. Instead, it was proposed that the formation of lipids is an immune system-activated process, and does not occur as a result of direct stimulation by *M. tb*, but rather via the IFN- ɣ, H1F-alpha-dependent pathway of the host immune system (Knight et al. 2018).

All-trans retinoic acid (*RARA)*, the active form of Vitamin A, plays an essential role in the normal functioning of the adaptive and innate immune systems. The oral administration of retinoic acid to rats resulted in inhibition of the *M. tb* growth, following *in vitro* infection (Yamada et al. 2007), thus making this gene a potential target for anti-TB therapies. The results

437 of this study have highlighted several SNPs which possibly significantly increased the risk of
438 individuals with Ghanaian ethnicity to being infected with the endemic TB strain of
439 *M. africanum*. Given the burden of disease, and the dominance of *M. africanum* strains in
440 Ghana, it may be a worthwhile exploring the functional effect of these SNPs on the biological
441 processes described.
442

3.4. Recommended improvements for future multi-phenotype GWAS
Several limitations may have affected this study. Obtaining a suitable sample size is a problem
inherent in GWA studies making use of logistic regression modelling. Furthermore, the many
phenotypes being analysed in this study, demanded a sufficient number of cases for each class.
The frequency of each MTBC clade however, is dependent not only on the host, but is also
affected by the virulence of the bacterium. Thus, with all these considered, the sample sizes
included in the MLR should be sufficient for inclusion in the analysis but will likely also reflect
the distribution in the population. Another limitation of the study is that at the time of analysis,
the AGR reference panel was not publicly available for download to a local machine. Thus, its
use in this study could only be facilitated via the SIS, a freely accessible online imputation
server. Through this, we were able to obtain high-quality imputed data for the South African
dataset, but it was necessary to be mindful when drawing comparisons as the other workflows
made use of different imputation software.
456

With the current trajectory of the TB epidemic, novel methods are needed to augment current
therapies for TB and combat the disease. This study provides the groundwork for future GWAS
wishing to investigate the relationship between the host and the many members of the MTBC
causing disease. Furthermore, the SNPs identified in this study may be evaluated in functional
studies to assess their viability as targets for host-directed therapies. This study would not have
been possible were it not for the collection of paired samples of blood and sputum from study
participants. Thus, future studies of this kind will require that both samples be collected from
participants in order to perform this association analysis. Although it was necessary to exclude
low frequency MTBC cases at the superclade level to prevent the reduction in statistical power
of the association test, this may have brought in a weakness in interpreting the odds ratios
derived from the model. Thus, the odds ratios derived may only be interpreted at the superclade
level and does not provide further granularity to association with specific clades. Therefore,
future studies employing this method should consider excluding low-frequency clades before
clustering as well as Bayesian analyses that allow inclusion of prior probability distribution for

strain prevalence. Additionally, incorporation of more diverse reference panels, such as the
AGR, and new algorithms for imputation could improve association results.
473
4. Materials and methods
4.1. Study design
To perform genome-wide association analyses between human host genotypes and the
infecting member of the MTBC, host genotypes with paired MTBC isolate information were
sourced for two geographically distinct cohorts. Paired host genotype and isolate information
was available for a South African cohort, and a Ghanaian cohort. As these two cohorts are
geographically distinct, and possess vastly different admixture profiles, we did not aim to
replicate our findings within these two cohorts.
482
The South African cohort consisted of study participants recruited in the Western Cape
Province of South Africa during the period of January 1993 through December 2004.
Participants were recruited from suburbs where the TB incidence was high (28.9 % in 2005)
and the prevalence of HIV was a low 2% (Kritzinger et al., 2009; Shisana et al., 2012). All
study participants in this cohort self-identified as belonging to a five-way admixed South
African population, were HIV-negative, and provided written informed consent. Blood
samples were collected for SNP genotyping of the host and sputum samples were collected for
bacterial culture on Loewenstein-Jensen (L-J) media. For the Ghanaian cohort, participants
were enrolled between September 2001 and July 2004 at Korle Bu Teaching Hospital in Accra,
Komfo Anokye Teaching Hospital in Kumasi, and at 15 additional hospitals and polyclinics in
Accra and Kumasi, as well as regional district hospitals. All cases were HIV-negative and
confirmed to have pulmonary TB by sputum microscopy, performing solid mycobacterial
cultures using L-J media and also by two independent radiologists (Thye et al., 2012, 2010).
All subsequent research was conducted in accordance with the principles expressed in the
Declaration of Helsinki (WHO, 2001).
498
4.2. Host SNP genotyping
4.2.1. South Africa
SNP genotyping was performed using DNA extracted from blood samples. All participants in
the South African cohort were genotyped using the GeneChip Human Mapping 500K SNP
array which contains 500 000 SNP markers (Affymetrix, California, United States), while a

subset of this cohort was also genotyped using the Infinium Multi-Ethnic Genotyping Array

(MEGA), which is comprised of 1.7 million SNP markers (Illumina, California, United States).

Genotype-calling was performed using the Affymetrix Power Tools pipeline (V1.10.0) as

previously described (De Wit et al., 2010; Schurz et al., 2019a). Genotype datum was made

available in PLINK format (Purcell and Chang, n.d.; Chang et al. 2015). Following standard

genotyping quality control (QC), ancestry proportions for the South African cohort on both the

Affymetrix and MEGA arrays were estimated (Daya et al. 2013, Schurz et al. 2018) using the

unsupervised algorithm implemented in ADMIXTURE (Alexander et al. 2009).

4.2.2.   Ghana

DNA extracted from blood samples was genotyped using the Affymetrix SNP 6.0 array at the

Affymetrix Services Laboratory in California, and at ATLAS Biolabs GmbH in Berlin.

Genotype datum was made available in PLINK format, genotypes were called using the

Birdseed version 2 algorithm and ancestry proportions in the form of principal components

were derived using the Eigenstrat software (Thye et al., 2012).

4.3. Bacterial SNP genotyping

For the South African cohort, MTBC isolates were genotyped using spoligotyping and IS*6110*

Restriction Fragment Length Polymorphism (RFLP) methods as previously described

(van der Spuy et al. 2009). For the Ghanaian cohort, MTBC isolates extracted from sputum

samples were cultured on L-J media at the Kumasi Centre for Collaborative Research (Owusu-

Dabo et al., 2006) and strains were identified using IS*6110* RFLP and spoligotyping (Supply

et al., 2006). All MTBC isolate information were captured on manually-curated infection

databases for archiving and made available for this study.

4.4. Defining MTBC clades and superclades

The term "superclades" was used to describe the grouping of clades using a SNP-based

phylogenetic tree of the MTBC. Where a common progenitor was shared, MTBC clades were

grouped into superclades near a point of divergence (Fig. 1) (Dippenaar, 2014). This was

performed to reduce the number of clades of low frequency, as low frequency groups are

known to induce an unfavourable collinearity effect on logistic regression models (Bergtold et

al., 2011). Additionally, clustering of clades into superclades mitigated the class imbalance

which would negatively affect the statistical models. Clades not amalgamated with other clades

were also referred to as "superclades" after clustering and those not represented on the
phylogenetic tree, were kept as distinct superclades and not clustered with any of the members
on the existing tree, except when a suitable reference phylogeny was found. After clustering,
superclades with a frequency less than ten in the study dataset were excluded from subsequent
analyses.
542

4.5. Genotype data quality control, haplotype phasing, and genotype imputation
Quality control, haplotype phasing and genotype imputation were performed using the methods
described in Schurz *et al.* (2019). Briefly, genotype data were iteratively filtered for 2% SNP
genotype missingness, 10% sample missingness and 5% SNP minor allele frequency (MAF),
until no additional samples or variants were removed. Additionally, variants lacking
chromosome or base pair information were updated using the 1000 Genomes Phase 3 (1000G)
reference panel and the dbSNP (Sherry et al., 2001) database. Imputation processes are
strengthened by sample size. Thus, all available samples, regardless of relatedness or whether
the sample had matching MTBC information, were included in the imputation. Related
individuals were noted, but not removed in order to maximise the number of haplotypes
available for the imputation process. The genotype QC procedure concluded with a sex
concordance check as well as strand-alignment to the 1000G reference panel [human genome
build 37 (Sudmant et al., 2015)] using the Genotype Harmonizer (version 1.4.15) tool (Deelen
et al., 2014).
557

Following the initial QC, haplotype phasing was performed using the ShapeITv2 (Delaneau et
al., 2008) software set at default parameters. Although some studies have reported that pre-
phasing reduces imputation accuracy (Roshyara et al. 2016), it is known to significantly speed
up the computationally intensive process of genotype imputation (Kanterakis et al. 2015; B.
Howie et al. 2012). To maximise the number of variants tested in the association analysis, the
cleaned genotype data were imputed using five protocols - and three different reference panels
- to determine which panel best served the given dataset in imputing missing variants, as
detailed in Schurz *et al.* (2019).
566

567   The In-House (IH) protocol made use of the 1000G reference panel with the IMPUTE2
568   imputation software (Howie et al., 2009). The Sanger Imputation Server (SIS) (McCarthy et
569   al., 2016) makes use of the Positional Burrows-Wheeler Transformation (PBWT) algorithm

(Durbin, 2014), with two options for the reference panel: the African Genome Resource (AGR) and the 1000G. Lastly, the Michigan Imputation Server (MIS) (Das et al., 2016) makes use of the Minimac3 algorithm (Howie et al., 2009) and provides access to imputation using the 1000G and the Consortium on Asthma among African-ancestry Populations in the Americas (CAAPA) (Mathias et al., 2016) reference panels. Of note, both the CAAPA and AGR reference panels were not publicly accessible for download at the time of this study and thus could only be accessed using these two online imputation server platforms. The key differences between the three protocols using the 1000G reference panel was the imputation software used, as well as additional strict QC filters imposed on the study dataset by the MIS and SIS methods.

4.6. Selection of high-quality imputed genotype data

The quality control procedure for the imputed data was implemented as described in Schurz *et al.* (2019). Briefly, following five imputation protocols, imputed data were filtered using a genotype calling threshold of 0.7, and the internal quality metric produced by the imputation process (Schurz et al., 2019b). SNPs with an INFO or R-squared (Rsq) value greater than 0.45 were prioritised for the association analysis and filtered iteratively for a maximum of 2% SNP genotype missingness, 10% sample missingness, and 5% SNP MAF using PLINK. Related individuals identified prior to imputation were removed followed by a second round of iterative filters for SNP- and sample missingness and MAF (Schurz et al., 2019b). Post-imputation QC concluded with extracting MTBC clade-matched samples from the remaining samples which had passed all QC filters.

4.7. Covariable data

All available covariables were obtained including sex, and age at time of active TB and subsequent recruitment into the study. To correct for differences in ethnicity amongst participants, either ancestry proportions or principal components were calculated and included as covariables.

4.8. Multi-phenotype GWAS

For the association analysis, a multinomial logistic regression (MLR) analysis using an additive genetic model was performed using SNPTEST v2.5.2 (Marchini, 2010). Two discrete variables, namely sex and superclade, as well as continuous variables, namely age at TB onset and ancestry proportions or principal components were included in the analysis. The phenotype

tested was specified as the MTBC superclade. Thus, the MLR model specified was the
occurrence of the MTBC superclade as a function of the baseline covariables given, as well as
the host genotypes supplied.

SNPTEST is unable to include covariables when the variance in the values provided is "too
small" as indicated (Marchini, n.d.). At the time of this study, SNPTEST was still under
development, and it had not yet been established, or recorded in the software manual, to what
degree of variance covariable data would not be accepted for inclusion in the logistic
regression. For developing the method, it was established through trial and error that if the
variance was below 0.001, these covariables could not be included.

The standard genome-wide significance cut-off of alpha $= 5\mathrm{x}10^{-8}$ was used when reporting
significance of SNPs (Pe'er et al., 2008; The International HapMap Consortium, 2005). Odds
ratios (OR) for the multiple phenotypes tested were calculated against a baseline phenotype by
setting the odds of that phenotype occurring, given the genotype, to 1. For this study, the
baseline phenotype was specified as the dominant superclade in the cohort, or a common
superclade of intermediate frequency if more than one cohort was being studied. Thus, the
LAM- and LAM_CAM superclades were used as the baseline phenotype for the association
analyses of the South African, and Ghanaian cohorts, respectively.

SNPs with a Likelihood Ratio Threshold (LRT) p-value of less than $5\mathrm{x}10^{-4}$ were selected and
analysed in R (R Core Team, 2017) and odds ratios were calculated from the beta values
generated by SNPTEST. SNPs with a standard error greater than 1.5 for their odds ratios were
excluded and SNPs with an LRT p-value less than $1\mathrm{x}10^{-6}$ were prioritised for further
investigation. These thresholds were chosen pragmatically to facilitate the completion of
method development. Finally, the Variant Effect Predictor (VEP) Tool (McLaren et al. 2016)
was used to retrieve gene annotations for the SNPs of interest.

Acknowledgements: The authors would like to acknowledge and thank the study participants
for their contribution and participation. This research was partially funded by the South African
government through the South African Medical Research Council. This work was also
supported by the National Research Foundation of South Africa (grant number 93460) to E.H.
and by a Strategic Health Innovation Partnership grant from the South African Medical
Research Council and Department of Science and Innovation/South African Tuberculosis
Bioinformatics Initiative to G.T. The content is solely the responsibility of the authors and does
not necessarily represent the official views of the South African Medical Research Council.

Ethics approval and consent to participate: For the South African cohort, blood and sputum
samples were collected from study participants as approved by the Health Research Ethics
Committee of Stellenbosch University (Project numbers: S17/01/013 and 95/072). For the
Ghanaian cohort, ethics for the study protocol was granted by the Committee on Human
Research, Publications and Ethics, School of Medical Sciences, Kwame Nkrumah University
of Science and Technology, Kumasi, Ghana, and the Ethics Committee of the Ghana Health
Service, Accra, Ghana (Thye et al., 2012). Venous blood samples were taken only after a
detailed explanation of the aims of the study, and consent was obtained of individuals enrolled
or their parents/guardians by signature or by thumbprint in case of illiteracy.

Availability of data and materials: Summary statistics for the South African cohort's data
can be made available to researchers who meet the criteria for access to confidential data after
application to the Health Research Ethics Committee of Stellenbosch University. Requests may
be sent to: Prof Craig Kinnear, E-mail: gkin@sun.ac.za.

Competing interests: The authors declare that they have no competing interests.

Author contributions:

Stephanie J. Müller: Conceptualization, Methodology, Software, Investigation, Formal
analysis, Writing - Original Draft, Visualization

Haiko Schurz: Software, Validation, Writing - Review & Editing

Gerard Tromp: Conceptualization, Supervision, Visualization, Writing - Review & Editing
Gian D. van der Spuy: Conceptualization, Supervision, Data Curation, Resources, Writing
- Review & Editing

Eileen G. Hoal: Conceptualization, Supervision, Writing - Review & Editing

Paul D. van Helden: Conceptualization, Writing - Review & Editing

Ellis Owusu-Dabo: Data Curation, Writing - Review & Editing

Christian G. Meyer: Data Curation, Writing - Review & Editing

Thorsten Thye: Data Curation, Resources, Writing - Review & Editing

Stefan Niemann: Data Curation, Writing - Review & Editing

Robin M. Warren: Data Curation, Resources, Writing - Review & Editing

Elizabeth Streicher: Data Curation, Writing - Review & Editing

Marlo Möller: Conceptualization, Supervision, Writing - Review & Editing

Craig Kinnear: Conceptualization, Supervision, Writing - Review & Editing

References

Alexander, D.H., Novembre, J., Lange, K., 2009. Fast Model-Based Estimation of Ancestry in Unrelated Individuals. Genome Res. 1655–1664. https://doi.org/10.1101/gr.094052.109.vidual

Asante-Poku, A., Yeboah-Manu, D., Otchere, I.D., Aboagye, S.Y., Stucki, D., Hattendorf, J., Borrell, S., Feldmann, J., Danso, E., Gagneux, S., 2015. Mycobacterium africanum Is Associated with Patient Ethnicity in Ghana. PLoS Negl. Trop. Dis. 9, e3370. https://doi.org/10.1371/journal.pntd.0003370

Bellamy, R., 1998. Genetic susceptibility to tuberculosis in human populations. Thorax 53, 588–593.

Bergtold, J.S., Yeager, E.A., Featherstone, A., 2011. Sample Size and Robustness of Inferences from Logistic Regression in the Presence of Nonlinearity and Multicollinearity. Presented at the Agricultural & Applied Economics Association's 2011 AAEA & NAREA Joint Annual Meeting, Pittsburgh, Pennsylvania.

Brites, D., Gagneux, S., 2015. Co-evolution of Mycobacterium tuberculosis and Homo sapiens. Immunol. Rev. 264, 6–24. https://doi.org/10.1111/imr.12264

Caws, M., Thwaites, G., Dunstan, S., Hawn, T.R., Thi Ngoc Lan, N., Thuong, N.T.T., Stepniewska, K., Huyen, M.N.T., Bang, N.D., Huu Loc, T., Gagneux, S., van Soolingen, D., Kremer, K., van der Sande, M., Small, P., Thi Hoang Anh, P., Chinh, N.T., Thi Quy, H., Thi Hong Duyen, N., Quang Tho, D., Hieu, N.T., Torok, E., Hien, T.T., Dung, N.H., Thi Quynh Nhu, N., Duy, P.M., van Vinh Chau, N., Farrar, J., 2008. The Influence of Host and Bacterial Genotype on the Development of Disseminated Disease with Mycobacterium tuberculosis. PLoS Pathog. 4. https://doi.org/10.1371/journal.ppat.1000034

Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M., Lee, J.J., 2015. Second-generation PLINK: rising to the challenge of larger and richer datasets. GigaScience 4. https://doi.org/10.1186/s13742-015-0047-8

Cheng, M.P., Abou Chakra, C.N., Yansouni, C.P., Cnossen, S., Shrier, I., Menzies, D., Greenaway, C., 2016. Risk of Active Tuberculosis in Patients with Cancer: A Systematic Review and Meta-Analysis. Clin. Infect. Dis. ciw838. https://doi.org/10.1093/cid/ciw838

Chimusa, E.R., Zaitlen, N., Daya, M., Möller, M., Helden, P.D. van, Nicola, J.M., Price, A.L., Hoal, E.G., 2014. Genome-wide association study of ancestry-specific TB risk in the South African coloured population. Hum. Mol. Genet. 23, 796–809. https://doi.org/10.1093/hmg/ddt462

Das, S., Forer, L., Schönherr, S., Sidore, C., Locke, A.E., Kwong, A., Vrieze, S.I., Chew, E.Y., Levy, S., McGue, M., Schlessinger, D., Stambolian, D., Loh, P.-R., Iacono, W.G., Swaroop, A., Scott, L.J., Cucca, F., Kronenberg, F., Boehnke, M., Abecasis, G.R., Fuchsberger, C., 2016. Next-generation genotype imputation service and methods. Nat. Genet. 48, 1284–1287. https://doi.org/10.1038/ng.3656

Daya, M., Van Der Merwe, L., Galal, U., Möller, M., Salie, M., Chimusa, E.R., Galanter, J.M., Van Helden, P.D., Henn, B.M., Gignoux, C.R., Hoal, E., 2013. A panel of ancestry informative markers for the complex five-way admixed South African Coloured population. PLoS ONE 8, 12–12. https://doi.org/10.1371/journal.pone.0082224

De Wit, E., Delport, W., Rugamika, C.E., Meintjes, A., Moller, M., Van Helden, P.D., Seoighe, C., Hoal, E.G., 2010. Genome-wide analysis of the structure of the South African Coloured Population in the Western Cape. Hum. Genet. 128, 145–153. https://doi.org/10.1007/s00439-010-0836-1

Deelen, P., Bonder, M.J., van der Velde, K.J., Westra, H.-J., Winder, E., Hendriksen, D.,
Franke, L., Swertz, M.A., 2014. Genotype harmonizer: automatic strand alignment and
format conversion for genotype data integration. BMC Res. Notes 7, 901.
Delaneau, O., Coulonges, C., Zagury, J.-F., 2008. Shape-IT: new rapid and accurate algorithm
for haplotype inference. BMC Bioinformatics 9, 540. https://doi.org/10.1186/1471-
2105-9-540
Dippenaar, A., 2014. A phylogenomic-and proteomic investigation into the evolution and
biological characteristics of the members of the group 2 Latin-American Mediterranean
(LAM) genotype of Mycobacterium tuberculosis (PhD Thesis). Stellenbosch:
Stellenbosch University.
Durbin, R., 2014. Efficient haplotype matching and storage using the positional Burrows–
Wheeler   transform   (PBWT).   Bioinformatics   30,   1266–1272.
https://doi.org/10.1093/bioinformatics/btu014
Gagneux, S., 2012. Host-pathogen coevolution in human tuberculosis. Philos. Trans. R. Soc.
Lond. B. Biol. Sci. 367, 850–9. https://doi.org/10.1098/rstb.2011.0316
Gagneux, S., Small, P.M., 2007. Global phylogeography of Mycobacterium tuberculosis and
implications for tuberculosis product development. Lancet Infect. Dis. 7, 328–337.
https://doi.org/10.1016/S1473-3099(07)70108-1
Herb, F., Thye, T., Niemann, S., Browne, E.N.L., Chinbuah, M.A., Gyapong, J., Osei, I.,
Owusu-Dabo, E., Werz, O., Rusch-Gerdes, S., Horstmann, R.D., Meyer, C.G., 2007.
ALOX5 variants associated with susceptibility to human pulmonary tuberculosis. Hum.
Mol. Genet. 17, 1052–1060. https://doi.org/10.1093/hmg/ddm378
Hirschhorn, J.N., Daly, M.J., 2005. Genome-wide association studies for common diseases and
complex traits. Nat. Rev. Genet. 6, 95–108. https://doi.org/10.1038/nrg1521
Hoal, E.G., Dippenaar, A., Kinnear, C., van Helden, P.D., Möller, M., 2017. The arms race
between man and Mycobacterium tuberculosis : Time to regroup. Infect. Genet. Evol.
https://doi.org/10.1016/j.meegid.2017.08.021
Hong, E.P., Go, M.J., Kim, H.-L., Park, J.W., 2017. Risk prediction of pulmonary tuberculosis
using genetic and conventional risk factors in adult Korean population. PLOS ONE 12,
e0174642. https://doi.org/10.1371/journal.pone.0174642
Howie, B.N., Donnelly, P., Marchini, J., 2009. A Flexible and Accurate Genotype Imputation
Method for the Next Generation of Genome-Wide Association Studies. PLOS Genet.
5, e1000529. https://doi.org/10.1371/journal.pgen.1000529
Kinnear, C., Hoal, E.G., Schurz, H., Van Helden, P.D., Moller, M., 2017. The role of human
host genetics in tuberculosis resistance. Expert Rev. Respir. Med. 11, 721–737.
https://doi.org/10.1080/17476348.2017.1354700
Kritzinger, F.E., den Boon, S., Verver, S., Enarson, D.A., Lombard, C.J., Borgdorff, M.W.,
Gie, R.P., Beyers, N., 2009. No decrease in annual risk of tuberculosis infection in
endemic area in Cape Town, South Africa. Trop. Med. Int. Health 14, 136–142.
https://doi.org/10.1111/j.1365-3156.2008.02213.x
Lim, C.H., Chen, H.-H., Chen, Y.-H., Chen, D.-Y., Huang, W.-N., Tsai, J.-J., Hsieh, T.-Y.,
Hsieh, C.-W., Hung, W.-T., Lin, C.-T., Lai, K.-L., Tang, K.-T., Tseng, C.-W., Chen,
Y.-M., 2017. The risk of tuberculosis disease in rheumatoid arthritis patients on
biologics and targeted therapy: A 15-year real world experience in Taiwan. PLOS ONE
12, e0178035. https://doi.org/10.1371/journal.pone.0178035
Lin, M., Caberto, C., Wan, P., Li, Y., Lum-Jones, A., Tiirikainen, M., Pooler, L., Nakamura,
B., Sheng, X., Porcel, J., Lim, U., Setiawan, V.W., Le Marchand, L., Wilkens, L.R.,
Haiman, C.A., Cheng, I., Chiang, C.W.K., n.d. Population-specific reference panels are
crucial for genetic analyses: an example of the CREBRF locus in Native Hawaiians.
Hum. Mol. Genet. https://doi.org/10.1093/hmg/ddaa083

Marchini, J., 2010. SNPTEST v2 Technical Details 10.
Marchini, J., n.d. SNPTest.
Mathias, R.A., Taub, M.A., Gignoux, C.R., Fu, W., Musharoff, S., O'Connor, T.D., Vergara,
C., Torgerson, D.G., Pino-Yanes, M., Shringarpure, S.S., Huang, L., Rafaels, N.,
Boorgula, M.P., Johnston, H.R., Ortega, V.E., Levin, A.M., Song, W., Torres, R.,
Padhukasahasram, B., Eng, C., Mejia-Mejia, D.-A., Ferguson, T., Qin, Z.S., Scott, A.F.,
Yazdanbakhsh, M., Wilson, J.G., Marrugo, J., Lange, L.A., Kumar, R., Avila, P.C.,
Williams, L.K., Watson, H., Ware, L.B., Olopade, C., Olopade, O., Oliveira, R., Ober,
C., Nicolae, D.L., Meyers, D., Mayorga, A., Knight-Madden, J., Hartert, T., Hansel,
N.N., Foreman, M.G., Ford, J.G., Faruque, M.U., Dunston, G.M., Caraballo, L.,
Burchard, E.G., Bleecker, E., Araujo, M.I., Herrera-Paz, E.F., Gietzen, K., Grus, W.E.,
Bamshad, M., Bustamante, C.D., Kenny, E.E., Hernandez, R.D., Beaty, T.H.,
Ruczinski, I., Akey, J., Barnes, K.C., 2016. A continuum of admixture in the Western
Hemisphere revealed by the African Diaspora genome. Nat. Commun. 7.
https://doi.org/10.1038/ncomms12522
McCarthy, S., Das, S., Kretzschmar, W., Delaneau, O., Wood, A.R., Teumer, A., Kang, H.M.,
Fuchsberger, C., Danecek, P., Sharp, K., Luo, Y., Sidore, C., Kwong, A., Timpson, N.,
Koskinen, S., Vrieze, S., Scott, L.J., Zhang, H., Mahajan, A., Veldink, J., Peters, U.,
Pato, C., van Duijn, C.M., Gillies, C.E., Gandin, I., Mezzavilla, M., Gilly, A., Cocca,
M., Traglia, M., Angius, A., Barrett, J.C., Boomsma, D., Branham, K., Breen, G.,
Brummett, C.M., Busonero, F., Campbell, H., Chan, A., Chen, S., Chew, E., Collins,
F.S., Corbin, L.J., Smith, G.D., Dedoussis, G., Dorr, M., Farmaki, A.-E., Ferrucci, L.,
Forer, L., Fraser, R.M., Gabriel, S., Levy, S., Groop, L., Harrison, T., Hattersley, A.,
Holmen, O.L., Hveem, K., Kretzler, M., Lee, J.C., McGue, M., Meitinger, T., Melzer,
D., Min, J.L., Mohlke, K.L., Vincent, J.B., Nauck, M., Nickerson, D., Palotie, A., Pato,
M., Pirastu, N., McInnis, M., Richards, J.B., Sala, C., Salomaa, V., Schlessinger, D.,
Schoenherr, S., Slagboom, P.E., Small, K., Spector, T., Stambolian, D., Tuke, M.,
Tuomilehto, J., Van den Berg, L.H., Van Rheenen, W., Volker, U., Wijmenga, C.,
Toniolo, D., Zeggini, E., Gasparini, P., Sampson, M.G., Wilson, J.F., Frayling, T., de
Bakker, P.I.W., Swertz, M.A., McCarroll, S., Kooperberg, C., Dekker, A., Altshuler,
D., Willer, C., Iacono, W., Ripatti, S., Soranzo, N., Walter, K., Swaroop, A., Cucca, F.,
Anderson, C.A., Myers, R.M., Boehnke, M., McCarthy, M.I., Durbin, R., Abecasis, G.,
Marchini, J., the Haplotype Reference Consortium, 2016. A reference panel of 64,976
haplotypes   for   genotype   imputation.   Nat.   Genet.   48,   1279–1283.
https://doi.org/10.1038/ng.3643
McHenry, M.L., Bartlett, J., Igo, R.P., Wampande, E., Benchek, P., Mayanja-Kizza, H.,
Fluegge, K., Hall, N.B., Gagneux, S., Tishkoff, S.A., Wejse, C., Sirugo, G., Boom,
W.H., Joloba, M., Williams, S.M., Stein, C.M., 2019. Interaction between host genes
and M. tuberculosis lineage can affect tuberculosis severity: evidence for coevolution.
bioRxiv 769448. https://doi.org/10.1101/769448
Möller, M., Hoal, E.G., 2010. Current findings, challenges and novel approaches in human
genetic   susceptibility   to   tuberculosis.   Tuberculosis   90,   71–83.
https://doi.org/10.1016/j.tube.2010.02.002
Oki, N.O., Motsinger-Reif, A.A., Antas, P.R., Levy, S., Holland, S.M., Sterling, T.R., 2011.
Novel human genetic variants associated with extrapulmonary tuberculosis: a pilot
genome wide association study. BMC Res. Notes 4, 28. https://doi.org/10.1186/1756-
0500-4-28
Omae, Y., Toyo-oka, L., Yanai, H., Nedsuwan, S., Wattanapokayakit, S., Satproedprai, N.,
Smittipat, N., Palittapongarnpim, P., Sawanpanyalert, P., Inunchot, W., Pasomsub, E.,
Wichukchinda, N., Mushiroda, T., Kubo, M., Tokunaga, K., Mahasirimongkol, S.,

2017. Pathogen lineage-based genome-wide association study identified CD53 as susceptible locus in tuberculosis. J. Hum. Genet. https://doi.org/10.1038/jhg.2017.82

Owusu-Dabo, E., Adjei, O., Meyer, C.G., Horstmann, R.D., Enimil, A., Kruppa, T.F., Bonsu, F., Browne, E.N.L., Chinbuah, M.A., Osei, I., Gyapong, J., Berberich, C., Kubica, T., Niemann, S., Ruesch-Gerdes, S., 2006. Mycobacterium tuberculosis Drug Resistance, Ghana. Emerg. Infect. Dis. 12, 1170–1172. https://doi.org/10.3201/eid1207.051028

Pastinen, T., Liitsola, K., Niini, P., Salminen, M., SyväNen, A.-C., 1998. Contribution of the CCR5 and MBL Genes to Susceptibility to HIV Type 1 Infection in the Finnish Population. AIDS Res. Hum. Retroviruses 14, 695–698. https://doi.org/10.1089/aid.1998.14.695

Pe'er, I., Yelensky, R., Altshuler, D., Daly, M.J., 2008. Estimation of the multiple testing burden for genomewide association studies of nearly all common variants. Genet. Epidemiol. 32, 381–385. https://doi.org/10.1002/gepi.20303

Png, E., Alisjahbana, B., Sahiratmadja, E., Marzuki, S., Nelwan, R., Balabanova, Y., Nikolayevskyy, V., Drobniewski, F., Nejentsev, S., Adnan, I., van de Vosse, E., Hibberd, M.L., van Crevel, R., Ottenhoff, T.H., Seielstad, M., 2012. A genome wide association study of pulmonary tuberculosis susceptibility in Indonesians. BMC Med. Genet. 13. https://doi.org/10.1186/1471-2350-13-5

Pratiwi, R.D., 2016. Socio-economic and environmenys risk factors of Tuberculosis in Wonosobo, Central Java, Indonesia. Graduate Studies in Public Health, Graduate Program, Sebelas Maret University Jl. Ir Sutami 36A, Surakarta 57126. Telp/Fax: (0271) 632 450 ext.208 First website:http//:s2ikm.pasca.uns.ac.id Second website: www.theicph.com. Email: theicph2016@gmail.com, p. 89. https://doi.org/10.26911/theicph.2016.027

Purcell, S., Chang, C., n.d. PLINK 1.9.

R Core Team, 2017. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria.

Rockett, K.A., Clarke, G.M., Fitzpatrick, K., Hubbart, C., Jeffreys, A.E., Rowlands, K., Craik, R., Jallow, M., Conway, D.J., Bojang, K.A., Pinder, M., Usen, S., Sisay-Joof, F., Sirugo, G., Toure, O., Thera, M.A., Konate, S., Sissoko, S., Niangaly, A., Poudiougou, B., Mangano, V.D., Bougouma, E.C., Sirima, S.B., Modiano, D., Amenga-Etego, L.N., Ghansah, A., Koram, K.A., Wilson, M.D., Enimil, A., Evans, J., Amodu, O., Olaniyan, S., Apinjoh, T., Mugri, R., Ndi, A., Ndila, C.M., Uyoga, S., Macharia, A., Peshu, N., Williams, T.N., Manjurano, A., Riley, E., Drakeley, C., Reyburn, H., Nyirongo, V., Kachala, D., Molyneux, M., Dunstan, S.J., Phu, N.H., Ngoc Quyen, N.T., Thai, C.Q., Hien, T.T., Manning, L., Laman, M., Siba, P., Karunajeewa, H., Allen, S., Allen, A., Davis, T.M.E., Michon, P., Mueller, I., Green, A., Molloy, S., Johnson, K.J., Kerasidou, A., Cornelius, V., Hart, L., Vanderwal, A., SanJoaquin, M., Band, G., Le, S.Q., Pirinen, M., Sepúlveda, N., Spencer, C.C.A., Clark, T.G., Agbenyega, T., Achidi, E., Doumbo, O., Farrar, J., Marsh, K., Taylor, T., Kwiatkowski, D.P., 2014. Reappraisal of known malaria resistance loci in a large multi-centre study. Nat. Genet. 46, 1197–1204. https://doi.org/10.1038/ng.3107

Salie, M., Van Der Merwe, L., Moller, M., Daya, M., Van Der Spuy, G.D., Van Helden, P.D., Martin, M.P., Gao, X.J., Warren, R.M., Carrington, M., Hoal, E.G., 2014. Associations between human leukocyte antigen class i variants and the mycobacterium tuberculosis subtypes causing disease. J. Infect. Dis. 209, 216–223. https://doi.org/10.1093/infdis/jit443

Schurz, H., Kinnear, C.J., Gignoux, C., Wojcik, G., van Helden, P.D., Tromp, G., Henn, B., Hoal, E.G., Möller, M., 2019a. A Sex-Stratified Genome-Wide Association Study of

Tuberculosis Using a Multi-Ethnic Genotyping Array. Front. Genet. 9. https://doi.org/10.3389/fgene.2018.00678

Schurz, H., Müller, S.J., van Helden, P.D., Tromp, G., Hoal, E.G., Kinnear, C.J., Möller, M., 2019b. Evaluating the Accuracy of Imputation Methods in a Five-Way Admixed Population. Front. Genet. 10. https://doi.org/10.3389/fgene.2019.00034

Seddon, J.A., Hesseling, A.C., Godfrey-Faussett, P., Fielding, K., Schaaf, H.S., 2013. Risk factors for infection and disease in child contacts of multidrug-resistant tuberculosis: a cross-sectional study. BMC Infect. Dis. 13. https://doi.org/10.1186/1471-2334-13-392

Sherry, S.T., Ward, M.-H., Kholodov, M., Baker, J., Phan, L., Smigielski, E.M., Sirotkin, K., 2001. dbSNP: the NCBI database of genetic variation. Nucleic Acids Res. 29, 308–311.

Shisana, O., Rhele, T., Simbayi, L.C., Zuma, K., Jooste, S., Zungu, N., Labadarios, D., Onoya, D., 2012. South African National HIV Prevalence, Incidence and Behaviour Survey, 2012.

Søborg, C., Madsen, H.O., Andersen, Å.B., Lillebaek, T., Kok-Jensen, A., Garred, P., 2003. Mannose-Binding Lectin Polymorphisms in Clinical Tuberculosis. J. Infect. Dis. 188, 777–782. https://doi.org/10.1086/377183

Stucki, D., Brites, D., Jeljeli, L., Coscolla, M., Liu, Q., Trauner, A., Fenner, L., Rutaihwa, L., Borrell, S., Luo, T., Gao, Q., Kato-Maeda, M., Ballif, M., Egger, M., Macedo, R., Mardassi, H., Moreno, M., Vilanova, G.T., Fyfe, J., Globan, M., Thomas, J., Jamieson, F., Guthrie, J.L., Asante-Poku, A., Yeboah-Manu, D., Wampande, E., Ssengooba, W., Joloba, M., Boom, W.H., Basu, I., Bower, J., Saraiva, M., Vasconcellos, S.E.G., Suffys, P., Koch, A., Wilkinson, R., Gail-Bekker, L., Malla, B., Ley, S.D., Beck, H.-P., de Jong, B.C., Toit, K., Sanchez-Padilla, E., Bonnet, M., Gil-Brusola, A., Frank, M., Penlap Beng, V.N., Eisenach, K., Alani, I., Ndung'u, P.W., Revathi, G., Gehre, F., Akter, S., Ntoumi, F., Stewart-Isherwood, L., Ntinginya, N.E., Rachow, A., Hoelscher, M., Cirillo, D.M., Skenders, G., Hoffner, S., Bakonyte, D., Stakenas, P., Diel, R., Crudu, V., Moldovan, O., Al-Hajoj, S., Otero, L., Barletta, F., Carter, E.J., Diero, L., Supply, P., Comas, I., Niemann, S., Gagneux, S., 2016. Mycobacterium tuberculosis lineage 4 comprises globally distributed and geographically restricted sublineages. Nat. Genet. 48, 1535–1543. https://doi.org/10.1038/ng.3704

Sudmant, P.H., Gardner, E.J., Handsaker, R.E., Abyzov, A., Huddleston, J., Zhang, Y., Ye, K., Jun, G., Hsi-Yang Fritz, M., Konkel, M.K., Malhotra, A., Stütz, A.M., Shi, X., Paolo Casale, F., Chen, J., Hormozdiari, F., Dayama, G., Chen, K., Malig, M., Chaisson, M.J.P., Walter, K., Meiers, S., Kashin, S., Garrison, E., Auton, A., Lam, H.Y.K., Jasmine Mu, X., Alkan, C., Antaki, D., Bae, T., Cerveira, E., Chines, P., Chong, Z., Clarke, L., Dal, E., Ding, L., Emery, S., Fan, X., Gujral, M., Kahveci, F., Kidd, J.M., Kong, Y., Lameijer, E.-W., McCarthy, S., Flicek, P., Gibbs, R.A., Marth, G., Mason, C.E., Menelaou, A., Muzny, D.M., Nelson, B.J., Noor, A., Parrish, N.F., Pendleton, M., Quitadamo, A., Raeder, B., Schadt, E.E., Romanovitch, M., Schlattl, A., Sebra, R., Shabalin, A.A., Untergasser, A., Walker, J.A., Wang, M., Yu, F., Zhang, C., Zhang, J., Zheng-Bradley, X., Zhou, W., Zichner, T., Sebat, J., Batzer, M.A., McCarroll, S.A., Mills, R.E., Gerstein, M.B., Bashir, A., Stegle, O., Devine, S.E., Lee, C., Eichler, E.E., Korbel, J.O., 2015. An integrated map of structural variation in 2,504 human genomes. Nature 526, 75–81. https://doi.org/10.1038/nature15394

Supply, P., Allix, C., Lesjean, S., Cardoso-Oelemann, M., Rusch-Gerdes, S., Willery, E., Savine, E., de Haas, P., van Deutekom, H., Roring, S., Bifani, P., Kurepina, N., Kreiswirth, B., Sola, C., Rastogi, N., Vatin, V., Gutierrez, M.C., Fauville, M., Niemann, S., Skuce, R., Kremer, K., Locht, C., van Soolingen, D., 2006. Proposal for Standardization of Optimized Mycobacterial Interspersed Repetitive Unit-Variable-

920    Number Tandem Repeat Typing of Mycobacterium tuberculosis. J. Clin. Microbiol. 44,
921     4498–4510. https://doi.org/10.1128/JCM.01392-06
922 The International HapMap Consortium, 2005. A haplotype map of the human genome. Nature
923     437, 1299–1320. https://doi.org/10.1038/nature04226
924 Thye, T., Owusu-Dabo, E., Vannberg, F.O., Van Crevel, R., Curtis, J., Sahiratmadja, E.,
925     Balabanova, Y., Ehmen, C., Muntau, B., Ruge, G., Sievertsen, J., Gyapong, J.,
926     Nikolayevskyy, V., Hill, P.C., Sirugo, G., Drobniewski, F., Van De Vosse, E., Newport,
927     M., Alisjahbana, B., Nejentsev, S., Ottenhoff, T.H.M., Hill, A.V.S., Horstmann, R.D.,
928     Meyer, C.G., 2012. Common variants at 11p13 are associated with susceptibility to
929     tuberculosis. Nat. Genet. 44, 257–259. https://doi.org/10.1038/ng.1080
930 Thye, T., Vannberg, F.O., Wong, S.H., Owusu-Dabo, E., Osei, I., Gyapong, J., Sirugo, G.,
931     Sisay-Joof, F., Enimil, A., Chinbuah, M.A., Floyd, S., Warndorff, D.K., Sichali, L.,
932     Malema, S., Crampin, A.C., Ngwira, B., Teo, Y.Y., Small, K., Rockett, K.,
933     Kwiatkowski, D., Fine, P.E., Hill, P.C., Newport, M., Lienhardt, C., Adegbola, R.A.,
934     Corrah, T., Ziegler, A., Morris, A.P., Meyer, C.G., Horstmann, R.D., Hill, A.V.S.,
935     2010. Genome-wide association analyses identifies a susceptibility locus for
936     tuberculosis on chromosome 18q11.2. Nat. Genet. 42, 739–741.
937     https://doi.org/10.1038/ng.639
938 van der Spuy, G.D., Kremer, K., Ndabambi, S.L., Beyers, N., Dunbar, R., Marais, B.J., van
939     Helden, P.D., Warren, R.M., 2009. Changing Mycobacterium tuberculosis population
940     highlights clade-specific pathogenic characteristics. Tuberculosis 89, 120–125.
941     https://doi.org/10.1016/j.tube.2008.09.003
942 WHO, 2001. World Medical Association Declaration of Helsinki.
943 Yim, J.J., Selvaraj, P., 2010. Genetic susceptibility in tuberculosis. Respirology 15, 241–256.
944     https://doi.org/10.1111/j.1440-1843.2009.01690.x
945
946

Tables:

Table 1: Summary of patient recruitment for the South African and Ghanaian cohorts

Cohort	South Africa	Ghana
Cases	853	1 359
Male	516	2 087
Female	431	1 224
Cases + Male	469 (55%)	933 (69%)
Cases + Female	384 (45%)	426 (31%)
Total number of participants	947	3 311
Total number of variants	397 337	7 838

Table 2: Summary of data pre–processing on the Michigan Imputation Server

Cohort	South Africa		Ghana	
Reference panel	**1000G**	**CAAPA**	**1000G**	**CAAPA**
Chromosomes	1–23	1–22	1–22	1–22
Samples	**919**	**919**	**3 239**	**3 239**
Sex undefined	3	0	0	0
SNPs	**239 612**	**233 309**	**617 409**	**617 409**
Alternative allele frequency > 0.5	166	0	0	0
Reference Overlap	99.49	97.94	99.55	97.23
Match	164 643	153 863	420 805	412 495
Allele Switch	74 116	68 912	176 073	172 432
Strand flip	1	0	0	0
Strand flip and allele switch	0	0	1	0
A/T, C/G genotypes	6 090	5 723	15 683	15 330
Filtered sites				
Filter flag set	0	0	0	0
Invalid alleles	0	0	0	0
Duplicated sites	0	0	0	0
NonSNP sites	0	0	0	0
Monomorphic sites	0	0	0	0
Allele mismatch	54	8	2 051	25
SNPs call rate < 90%:	0	0	0	0
Excluded sites in total:	55	8	2 052	25
Sites remaining (before imputation)	**239 477**	**228 498**	**612 561**	**600 257**
Samples remaining	**916**	**916**	**3 239**	**3 239**

954 *Table 3: Percentage proportion of SNPs with a quality metric greater than 0.45*

Cohort	South Africa		Ghana	
Chromosome	Chr 1	Chr X	Chr 1	Chr 22
IH–1000G[1]	39	29	38	39
MIS–1000G[2]	32	18	49	45
MIS–CAAPA[3]	22	–	56	50
SIS–1000G[4]	36	32	41	40
SIS–AGR[5]	43	40	38	36

[1]IH–1000G: In–House workflow using 1000G reference panel

[2]MIS–1000G: Michigan Imputation Server workflow using 1000G reference panel

[3]MIS–CAAPA: Michigan Imputation Server workflow using CAAPA reference panel

[4]SIS–1000G: Sanger Imputation Server workflow using 1000G reference panel

[5]SIS–AGR: Sanger Imputation Server workflow using AGR reference panel

Table 4: Top 11 SNPs identified by MLR to be associated with strains of different MTBC superclades in the South African cohort

		Allele		Odd Ratio (95% C.I.)				
Chr	SNP ID	Ref[a]	Risk	BeijingCAS1	HaarlemsLCC	Quebec	Other	LRT p–val [b]
5	rs17458866	C	T	0.34 (0.19–0.61)	0.44 (0.26–0.76)	1.99 (1.07–3.68)	0.46 (0.23–0.95)	10e–07
5	rs13355101	G	A	0.31 (0.17–0.57)	0.46 (0.27–0.79)	2.00 (1.08–3.70)	0.47 (0.23–0.96)	6.43e–07
5	rs12518239	C	A	0.29 (0.15–0.56)	0.39 (0.22–0.70)	1.91 (1.01–3.62)	0.51 (0.25–1.05)	9.41e–07
5	rs28769614	C	T	0.27 (0.13–0.53)	0.37 (0.20–0.68)	1.92 (1.00–3.66)	0.48 (0.23–1.01)	3.03e–07
6	rs9389610	G	A	2.19 (1.35–3.55)	1.07 (0.64–1.76)	0.25 (0.08–0.73)	2.78 (1.52–5.08)	1.60e–07
17	rs78022196	G	A	1.04 (0.60–1.80)	0.59 (0.32–1.09)	5.31 (2.44–11.57)	1.96 (1.03–3.73)	5.13e–07
17	rs72843143	C	T	0.98 (0.57–1.69)	0.57 (0.31–1.04)	4.81 (2.26–10.25)	1.89 (1.00–3.58)	8.18e–07
17	rs8071332	A	G	0.94 (0.55–1.60)	0.59 (0.33–1.06)	4.77 (2.22–10.28)	2.03 (1.08–3.81)	6.54e–07
17	rs10438776	T	C	0.99 (0.58–1.71)	0.55 (0.30–1.01)	4.99 (2.30–10.85)	1.94 (1.02–3.69)	3.93e–07
17	rs17682747	G	A	0.98 (0.57–1.68)	0.56 (0.31–1.03)	4.85 (2.27–10.33)	1.90 (1.01–3.58)	5.93e–07
17	rs7208461	T	C	0.94 (0.55–1.61)	0.54 (0.30–0.98)	4.68 (2.17–10.10)	1.77 (0.94–3.34)	8.64e–07

962 [a] Reference allele

963 [b] LAM was used as the reference superclade
964
965
966
967

Table 5: Top 32 SNPs identified by MLR to be associated with strains of different MTBC superclades in the Ghanaian cohort

Chr	SNP ID	Ref[a]	Risk	BeijingCAS	EAI_afri	Ghana2	HaarlemX	T_U	LRT p–val[b]
		Allele				**Odd Ratio (95% C.I.)**			
6	rs529920[*]	A	G	0.40 (0.22–0.70)	0.69 (0.57–0.84)	1.04 (0.60–1.80)	1.10 (0.84–1.44)	1.24 (0.98–1.56)	1.86e–07
12	rs73418916	A	G	0.30 (0.11–0.81)	34.15 (20.00–58.33)	1.38 (0.64–2.96)	0.76 (0.51–1.12)	0.88 (0.63–1.22)	2.31e–97
12	rs138396290	T	C	0.32 (0.13–0.77)	4.59 (3.63–5.82)	1.31 (0.73–2.37)	0.87 (0.63–1.19)	0.91 (0.70–1.19)	3.32e–62
12	rs75717431	T	C	2.46 (1.40–4.35)	0.98 (0.78–1.24)	3.30 (1.88–5.82)	0.95 (0.69–1.31)	0.59 (0.44–0.80)	8.55e–09
12	rs77428482	G	A	2.56 (1.45–4.52)	1.01 (0.81–1.28)	3.20 (1.81–5.64)	0.95 (0.69–1.32)	0.60 (0.45–0.81)	1.93e–08
12	rs77562721	G	A	2.50 (1.42–4.41)	1.04 (0.82–1.30)	3.12 (1.77–5.50)	0.93 (0.68–1.29)	0.60 (0.45–0.81)	2.53e–08
12	rs41524146	C	G	2.38 (1.36–4.18)	1.00 (0.80–1.26)	3.19 (1.82–5.58)	0.97 (0.71–1.34)	0.61 (0.45–0.81)	2.62e–08
12	rs7299395	G	A	2.16 (1.22–3.82)	0.99 (0.80–1.24)	3.12 (1.77–5.52)	0.93 (0.69–1.27)	0.64 (0.48–0.84)	2.50e–07
12	rs74550821	G	A	2.59 (1.47–4.57)	1.05 (0.84–1.32)	3.20 (1.82–5.63)	1.01 (0.74–1.39)	0.63 (0.47–0.84)	2.89e–08
12	rs144335343	C	T	2.33 (1.32–4.11)	1.08 (0.87–1.35)	3.63 (2.05–6.41)	1.00 (0.73–1.36)	0.69 (0.52–0.92)	9.15e–08
12	rs6582329	A	T	2.58 (1.46–4.57)	1.07 (0.85–1.34)	3.42 (1.94–6.02)	1.04 (0.76–1.43)	0.67 (0.50–0.90)	6.00e–08
12	rs12296167	T	G	2.42 (1.25–4.70)	2.03 (1.60–2.59)	3.33 (1.71–6.48)	1.03 (0.74–1.42)	1.41 (1.07–1.85)	2.02e–09
12	rs544003050	A	G	0.5	4.72	1.42	0.75	0.74	4.17e–40

				(0.24–1.05)	(3.47–6.43)	(0.68–2.98)	(0.52–1.07)	(0.54–1.00)	
12	rs58262822	C	G	0.58	8.50	0.75	0.81	0.93	1.96e–124
				(0.34–0.99)	(6.29–11.48)	(0.47–1.20)	(0.65–1.01)	(0.78–1.11)	
12	rs11108508	T	C	1.09	6.61	1.33	0.71	1.11	1.04e–112
				(0.56–2.09)	(5.26–8.32)	(0.73–2.44)	(0.48–1.05)	(0.85–1.46)	
12	rs41472447*	A	G	2.56	0.97	2.94	0.92	0.59	5.70e–09
				(1.48–4.41)	(0.78–1.21)	(1.71–5.08)	(0.68–1.26)	(0.44–0.79)	
13	rs549053537	A	T	0.63	5.51	1.23	0.86	0.91	8.12e–79
				(0.41–0.96)	(4.12–7.39)	(0.83–1.82)	(0.71–1.04)	(0.78–1.07)	
13	rs73497904	C	G	1.34	65.39	1.20	0.96	1.23	2.23e–244
				(0.69–2.59)	(40.67–105.13)	(0.60–2.39)	(0.67–1.39)	(0.92–1.65)	
13	rs9524738	G	C	1.88	0.31	1.32	1.34	1.16	2.67e–52
				(0.86–4.14)	(0.26–0.38)	(0.68–2.55)	(0.97–1.84)	(0.90–1.50)	
15	rs551641937	G	A	0.52	275.89	0.52	0.39	1.11	3.61e–236
				(0.07–3.97)	(152.64–498.69)	(0.07–3.98)	(0.14–1.14)	(0.59–2.08)	
15	rs35799802	C	T	0.76	3.08	0.89	1.09	1.19	3.94e–15
				(0.36–1.62)	(2.29–4.13)	(0.42–1.88)	(0.75–1.59)	(0.87–1.64)	
15	rs55747528	C	T	0.53	28.58	1.47	0.72	0.79	1.49e–69
				(0.24–1.15)	(15.28–53.45)	(0.66–3.25)	(0.49–1.05)	(0.57–1.08)	
16	rs577800201	C	T	0.31	93.05	0.87	0.67	1.01	1.20e–167
				(0.07–1.33)	(54.17–159.85)	(0.32–2.36)	(0.4–1.13)	(0.68–1.50)	
16	rs187181146	C	T	0.64	8.65	0.94	0.83	0.92	9.77e–152
				(0.35–1.17)	(6.61–11.31)	(0.59–1.49)	(0.66–1.06)	(0.76–1.12)	
16	rs35868343	G	A	0.72	10.65	1.01	0.91	0.92	1.90e–93
				(0.46–1.11)	(7.14–15.87)	(0.66–1.55)	(0.74–1.12)	(0.77–1.10)	
17	rs143309838	G	A	0.57	9.38	0.93	0.84	0.99	2.08e–165
				(0.28–1.17)	(7.20–12.23)	(0.57–1.53)	(0.65–1.08)	(0.81–1.21)	
17	rs144224512	C	G	0.36	220.49	0.50	0.42	0.74	1.21e–215

Chr	SNP	a	b						p-value
				(0.05–2.69)	(118.29–411.01)	(0.08–3.13)	(0.17–1.01)	(0.40–1.36)	
17	rs374315920	C	T	3.52 (0.94–13.13)	548.96 (270.65–1113.45)	1.11 (0.14–8.79)	0.85 (0.27–2.63)	0.93 (0.37–2.36)	1.68e–255
17	rs77139740	A	G	0.34 (0.12–1.01)	44.94 (26.85–75.20)	0.97 (0.43–2.22)	0.73 (0.48–1.12)	0.89 (0.63–1.26)	3.74e–120
17	rs77641928	G	A	1.24 (0.37–4.17)	57.29 (33.69–97.42)	0.46 (0.06–3.62)	0.67 (0.30–1.49)	0.94 (0.53–1.67)	4.98e–282
22	rs553728019	A	C	0.51 (0.23–1.09)	23.66 (12.10–46.25)	1.05 (0.47–2.33)	0.76 (0.52–1.11)	0.79 (0.57–1.09)	2.83e–55
22	rs60153275	C	T	0.93 (0.32–2.71)	61.01 (36.66–101.54)	0.93 (0.32–2.71)	0.70 (0.38–1.28)	0.98 (0.63–1.51)	7.19e–281

969 [a] Reference allele

970 [b] LAM_CAM was used as the reference superclade

971 [*] Genotyped SNPs

 Figure Legends:

Fig. 1. Clustering of MTBC clades on the SNP–based phylogenetic tree (adapted from original

tree sourced with permission from Dippenaar 2014). MTBC clades were grouped into

superclades near a point of divergence on the phylogenetic tree. Clustering reduced 12 distinct

clades into seven closely related superclades. TB cases identified to be due to infection with

the East African Indian (EAI) and *M. africanum* clades (green bracket), were merged into the

"EAI_afri" superclade. Similarly, CAS and Beijing clades were merged into the "BeijingCAS"

superclade (red bracket). The LCC, Pre-Haarlem, Haarlem-like, and Haarlem clades merged

into one superclade designated as "HaarlemsLCC" (orange bracket), while the Quebec,

Latin-American Mediterranean (LAM), T, and Lineage 7 clades remained unchanged due to

the lack of a common progenitor on the red dotted line, were subsequently treated as individual

superclades and are indicated by black brackets. The clades denoted as LCC 1 bander (F110)

and 'unknown (F32)' were not clustered into superclades.

Fig. 2. Frequency distributions of MTBC Clades and Superclades. Frequency distributions of

A: MTBC clades and **B:** superclades for the South African cohort and **C:** clades and **D:**

superclades for the Ghanaian cohort. The South African cohort was dominated by the LAM

superclade, while the Ghanaian cohort was dominated by the EAI_afri and LAM_CAM

superclades.

Fig. 3. SNP density plots for Chromosome 1 of the Ghanaian cohort post-imputation using the

five workflows. **A:** IH with 1000G, **B:** MIS with 1000G, **C:** SIS with 1000G, **D:** MIS with

CAAPA, **E:** SIS with AGR. The MIS workflow using the CAAPA resource imputed the

greatest proportion of SNPs with a quality metric greater than 0.45.

Fig. 4. SNP density plots for Chromosome 22 of the Ghanaian cohort post-imputation using

the five workflows. **A:** IH with 1000G, **B:** MIS with 1000G, **C:** SIS with 1000G, **D:** MIS with

CAAPA, **E:** SIS with AGR.

1000

1001 Fig. 5. Median quality scores across MAF bins for the Ghanaian cohort, using the five

1002 protocols. **A:** Chromosome 1 **B:** Chromosome 22. For both these representative chromosomes,

1003 the IH imputation protocol using the 1000G reference panel outperformed the other four

1004 workflows.

1005

1006   Fig. 6. Standard errors of odds ratios calculated for each superclade against the reference LAM
1007   superclade. **A:** South African cohort **B:** Ghanaian cohort. For both cohorts, the smallest
1008   superclades had the largest variation in their standard errors.

Figure 1

Figure 3

Figure 4

Figure 5

Figure 6