Pulmonary function and risk of Alzheimer dementia: two-sample Mendelian randomization study

Tom C. Russ,[1,4] * Sarah E. Harris,[4] G. David Batty[1,5]

1. Alzheimer Scotland Dementia Research Centre, University of Edinburgh;

2. Edinburgh Dementia Prevention Group, University of Edinburgh;

3. Division of Psychiatry, Centre for Clinical Brain Sciences, University of Edinburgh;

4. Lothian Birth Cohorts, Department of Psychology, University of Edinburgh;

5. UCL Research Department of Epidemiology & Public Health, University College London

* Correspondence to: Dr Tom Russ, Alzheimer Scotland Dementia Research Centre,

University of Edinburgh, 7 George Square, Edinburgh, EH8 9JZ, UK

Telephone: +44 (0)131 650 4340; Email: T.C.Russ@ed.ac.uk

Manuscript statistics: 1000 words with one table

Funding: This article received no specific funding. The Alzheimer Scotland Dementia Research Centre is funded by Alzheimer Scotland. The Lothian Birth Cohorts are funded by Age UK. TCR is employed by the UK National Health Service and the Scottish Government. GDB is supported by the UK Medical Research Council (MR/P023444/1) and the US National Institute on Aging (1R56AG052519-01; 1R01AG052519-01A1). All researchers are independent of their funders.

ABSTRACT

Dementia is a major global public health concern and in addition to recognised risk factors there is emerging evidence that poorer pulmonary function is linked with subsequent dementia risk. However, it is unclear if this observed association is causal or whether it might result from confounding. Therefore, we present the first two-sample Mendelian randomisation study of the association between pulmonary function and Alzheimer dementia using the most recent genome-wide association studies to produce instrumental variables for both. We found no evidence of a causal effect of reduced Forced Expiratory Volume in 1 second (FEV_1) or Forced Vital Capacity (FVC) on Alzheimer dementia risk (both $P>0.35$). However, the FEV_1/FVC ratio was associated with Alzheimer dementia risk with, in fact, superior function predicting an increased dementia risk ($\beta=0.11$, $P=0.016$) which may result from survivor bias. While we can conclude that there is no causal link between impaired pulmonary function and Alzheimer dementia, our study sheds less light on potential links with other types of dementia.

It is well-documented that dementia is a major and growing public health problem — the number of cases are projected to increase, particularly in low-to-middle income countries. (1) To date, the risk of developing Alzheimer disease — the most common neurodegenerative disease culminating in the dementia syndrome — is thought to be raised in the presence of a relatively small series of environmental and genetic factors. These include lower educational attainment, hypertension, obesity, diabetes, cigarette smoking, and carriage of the ε4 allele of the *APOE* gene. (2)

Recent findings from observational studies also suggest that impaired pulmonary function is consistently associated with around a 40% evaluation in the risk of later dementia. While there is mechanistic evidence to support this association — including the hypoxic effect of extended sub-optimal ventilatory function (3) — crucially, given the observational nature of these studies, it is unclear if this relationship is causal. (4) The primary impediment to drawing causal inference from such observational studies is the perennial problem of confounding — the notion that, in the case of pulmonary function, the characteristics of people in the low functioning group differ from the unexposed in a variety of ways that may explain the association. Investigators attempt to include as many relevant covariates as possible but the possibility of confounding by factors which have not been measured, or imprecisely quantified, is universal. The technique of Mendelian randomisation (MR) has been seen as a possible remedy to this problem. (5) MR exploits natural genetic variation to create instrumental exposure variables which are less prone to confounding than the directly measured phenotype. This technique has been extended to a two-sample MR where both the exposure and outcome are instrumental genetic variables. (6) Accordingly, for the first time to our knowledge, we present here a two-sample MR study to clarify whether our original observed

association between poorer pulmonary function and subsequent Alzheimer dementia is

causal.

METHODS

We ran a two-sample MR using the using summary data from the UKBiobank/SpiroMeta

Consortium Genome-Wide Association Study (GWAS; 7) which comprised 400,102

individuals. We derived two genetic instruments for lung function: Forced Expiratory

Volume in one second (in litres; FEV_1) and Forced Vital Capacity (in litres; FVC). Of the

279 SNPs associated with lung function, only those related to the relevant trait with P<5x10$^-$

^{8} and with the same direction of effect in UKBiobank and SpiroMeta were used as genetic

instruments. In addition, we included a more exploratory measure of pulmonary function, the

FEV_1/FVC ratio, which has been used in the diagnosis of chronic obstructive pulmonary

disease whereby lower values are more suggestive of this disorder. (8) For the outcome,

Alzheimer dementia, we used summary data from the most recent GWAS for Alzheimer

dementia which included 21,982 people with Alzheimer dementia and 41,944 controls. (9)

The models were run in the R statistical computing environment using the TwoSampleMR

package. (10)

RESULTS

In **Table 1** we show the relationship between our markers of lung function with the

subsequent risk of Alzheimer dementia. There was no evidence of a causal effect of poorer

lung function — measured using FEV_1 or FVC — on Alzheimer dementia risk (both

P>0.35). However, the FEV_1/FVC ratio was associated with Alzheimer dementia risk with, in

fact, superior function predicting an increased dementia risk (β=0.11, P=0.016). The MR

Egger intercept for the latter indicates little horizontal pleiotropy ($\beta=0.0002$, P=0.96) and

the inverse-variance weighted Q-value (177.7, P=0.08) suggests no substantial heterogeneity.

DISCUSSION

We found that the association between lower pulmonary function and Alzheimer dementia

risk based on conventional epidemiological analyses was not supported as being causal in the

present MR analyses. Thus, it is possible that the original relationship resulted from

confounding by one or more unmeasured or poorly measured confounding factors. Multiple

candidates exist including an adverse intrauterine environment leading to reduced maximal

lung function, exposure to environmental factors (e.g., tobacco smoke, atmospheric

pollution) affecting lung function and development, and socioeconomic factors (poverty,

educational failure, and less-advantaged social class). In our systematic review and meta-

analysis, most included studies took account of smoking and cardiovascular disease risk

factors, and slightly fewer included height.(4) Socioeconomic position was variably accounted

for and there was little coverage of the whole life course in terms of all covariables included.

The FEV_1/FVC ratio has not been routinely examined in relation to dementia risk using

conventional epidemiology methods.(4) However, we found a link between high pulmonary

function as captured using this measure and increased Alzheimer dementia risk. This may

possibly be explained by survivor bias, with participants with poorer pulmonary function dying

before they reach late life.

While it is less susceptible to unmeasured confounding, a two-sample MR approach is not

without its limitations. It relies on the instrument being a strong predictor of the exposure

and the exposure being measured accurately in the GWAS from which the instrument is

derived. Pulmonary function was accurately measured with rigorous quality control in

UKBiobank (87.2% participants), as in the individual studies of the SpiroMeta

consortium.(7) The Alzheimer dementia GWAS included 46 case-control studies from four

consortia.(9) These studies used various methods of ascertaining dementia, with multiple

diagnostic criteria being applied and some diagnosing dementia clinically and some identifying

Alzheimer-type pathology post mortem. This variation is likely to affect the applicability of

the GWAS findings in our analysis.

In contrast to the instrumental Alzheimer dementia variable used here, most observational

studies use a more general category of 'dementia.'(4, 9) This lack of clarity is common and the

multiple diseases which cause the dementia syndrome — such as Alzheimer disease,

cerebrovascular disease, Lewy body disease, and the Fronto-Temporal Lobar Degenerative

syndromes — are frequently conflated. Depending on the methodology used to ascertain

dementia status, it can be more or less straightforward to clarify an individual's precise

diagnosis. For example, death certificates frequently only record the broad dementia

syndrome, rather than the specific sub-type. Thus, while we can conclude that there is no

causal link between impaired pulmonary function and Alzheimer dementia, our study sheds

less light on potential links with other types of dementia. It is plausible that there may be a

different relationship between pulmonary function and vascular dementia, for instance.

REFERENCES

1. Prince MJ, Wimo A, Guerchet M, Ali G-C, Wu Y-T, Prina AM. World Alzheimer Report 2015. The Global Impact of Dementia: an analysis of prevalence, incidence, cost and trends. London: Alzheimer's Disease International; 2015.

2. Livingston G, Sommerlad A, Orgeta V, Costafreda SG, Huntley J, Ames D, Ballard C, Banerjee S, Burns A, Cohen-Mansfield J, Cooper C, Fox N, Gitlin LN, Howard R, Kales HC, Larson EB, Ritchie K, Rockwood K, Sampson EL, Samus Q, Schneider LS, Selbaek G, Teri L, Mukadam N. Dementia prevention, intervention, and care. *Lancet* 2017; 390: 2673-2734.

3. Horsburgh K, Wardlaw JM, van Agtmael T, Allan SM, Ashford MLJ, Bath Philip M, Brown R, Berwick J, Cader MZ, Carare Roxana O, Davis John B, Duncombe J, Farr Tracy D, Fowler JH, Goense J, Granata A, Hall CN, Hainsworth AH, Harvey A, Hawkes CA, Joutel A, Kalaria RN, Kehoe Patrick G, Lawrence CB, Lockhart A, Love S, Macleod MR, Macrae IM, Markus Hugh S, McCabe C, McColl Barry W, Meakin Paul J, Miller A, Nedergaard M, O'Sullivan M, Quinn TJ, Rajani R, Saksida LM, Smith C, Smith Kenneth J, Touyz RM, Trueman Rebecca C, Wang T, Williams A, Williams SCR, Work LM. Small vessels, dementia and chronic diseases – molecular mechanisms and pathophysiology. *Clinical Science* 2018; 132: 851-868.

4. Russ TC, Kivimaki M, Batty GD. Respiratory Disease and Lower Pulmonary Function as Risk Factors for Dementia: A Systematic Review With Meta-analysis. *Chest* 2020.

5. Davey Smith G, Ebrahim S. 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? *International Journal of Epidemiology* 2003; 32: 1-22.

6. Hartwig FP, Davies NM, Hemani G, Davey Smith G. Two-sample Mendelian randomization: avoiding the downsides of a powerful, widely applicable but potentially fallible technique. *International Journal of Epidemiology* 2017; 45: 1717-1726.

7. Shrine N, Guyatt AL, Erzurumluoglu AM, Jackson VE, Hobbs BD, Melbourne CA, Batini C, Fawcett KA, Song K, Sakornsakolpat P. New genetic signals for lung function highlight pathways and chronic obstructive pulmonary disease associations across multiple ancestries. *Nature Genetics* 2019; 51: 481-493.

8. Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P, Fukuchi Y, Jenkins C, Rodriguez-Roisin R, van Weel C, Zielinski J. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 2007; 176: 532-555.

9. Kunkle BW, Grenier-Boley B, Sims R, Bis JC, Damotte V, Naj AC, Boland A, Vronskaya M, Van Der Lee SJ, Amlie-Wolf A. Genetic meta-analysis of diagnosed Alzheimer's disease identifies new risk loci and implicates A β, tau, immunity and lipid processing. *Nature Genetics* 2019; 51: 414-430.

10. Hemani G, Tilling K, Davey Smith G. Orienting the causal relationship between imprecisely measured traits using GWAS summary data. *PLOS Genetics* 2017; 13: e1007081.

Table 1. Inverse-variance weighted causal effect estimates of the association between pulmonary function (FEV_1, FVC, and FEV_1/FVC ratio) and Alzheimer dementia from a two-sample Mendelian randomization

	N SNPs	β	SE	P
FEV_1	179	0.0554	0.05989	0.354949
FVC	133	-0.01728	0.074022	0.815442
FEV_1/FVC ratio	154	0.113478	0.047106	0.015996