

1 **Article Summary Line:** Rapid increase in seroprevalence of antibodies against SARS-CoV-2  
2 was seen in Karachi, Pakistan from April to June 2020 with a high conditional risk of infection  
3 within the household

4 **Running Title:** SARS-CoV-2 serosurvey in Karachi, Pakistan

5 **Keywords:** Seroepidemiologic Studies; Coronavirus; Pakistan

6 **Title:** Serial population based household survey of antibodies to SARS-CoV-2 in low and high  
7 transmission neighborhoods of urban Pakistan

8 **Authors:** Muhammad Imran Nisar MBBS, MSc; Nadia Ansari MBBS, MSc; Mashal Amin  
9 MBBS, MSc; Aneeta Hotwani MSc; Farah Khalid MSc; Najeeb Rehman BSc; Arjumand Rizvi  
10 MPhil; Arslan Memon MBBS, MPH; Zahoor Ahmed MBBS; Ashfaque Ahmed MBBS; Junaid  
11 Iqbal PhD; Ali Faisal Saleem MBBS, MSc; Uzma Bashir Aamir MBBS, PhD; Daniel B  
12 Larremore PhD; Bailey Fosdick PhD; Fyezah Jehan MBBS, MSc

13 **Affiliations:**

14 The Aga Khan University, Karachi, Sind, Pakistan (MI. Nisar, N. Ansari, M. Amin, A. Hotwani,  
15 F. Khalid, N. Rahman, A. Rizvi, J. Iqbal, AF. Saleem, F. Jehan)

16 Health Department, Government of Sind, Karachi, Pakistan (A. Memon, Z. Ahmed, A. Ahmed)

17 World Health Organization Country Office, Karachi, Sind, Pakistan (UB. Aamir)

18 University of Colorado, Boulder, Colorado, USA (DB. Larremore)

19 Colorado State University, Fort Collins, Colorado, USA (B. Fosdick)

20  
1

21 **Abstract-** 146 words

22 Serial household antibody sero-surveys informs the pandemic where testing is non-  
23 uniform. Young populations with intergenerational co-residence may have different transmission  
24 dynamics. We conducted two serial cross-sectional surveys in April and June 2020 in low- and  
25 high-transmission neighborhoods of Karachi, Pakistan, using random sampling. Symptoms were  
26 assessed and blood tested for antibody using chemiluminescence. Seroprevalence was adjusted  
27 using Bayesian regression and post stratification. CRI with 95% confidence intervals was  
28 obtained. We enrolled 2004 participants from 406 households. In June 8.7% (95% CI 5.1-13.1)  
29 and 15.1% (95% CI 9.4 -21.7) were infected in low- and high-transmission-areas respectively  
30 compared with 0.2% (95% CI 0-0.7) and 0.4% (95% CI 0 - 1.3) in April. Conditional risk of  
31 infection was 0.31 (95% CI 0.16-0.47) and 0.41(95% CI 0.28-0.52) respectively with only 5.4%  
32 symptomatic. Rapid increase in seroprevalence from baseline is seen in Karachi, with a high  
33 probability of infection within household.

34

35

36

37

38

39

40

41 **Text-2541 words**

## 42 **INTRODUCTION**

43 The global COVID-19 pandemic has resulted in more than 15.5 million confirmed cases  
44 (until July 24) and more than 633000 deaths, with an estimated case fatality rate (CFR) of 4.1 %  
45 (1). Pakistan was among the first of low- and middle- income countries (LMICs) to be affected,  
46 and since then, there have been 269000 cases with 5709 deaths (CFR 2.1 %) (2). Karachi, a large  
47 metropolitan city became the epicenter of epidemic on 26<sup>th</sup> Feb 2020. Since then Karachi has  
48 seen the largest number of cases (~83000 or 31% of all cases) in Pakistan.

49 The population demographics of Pakistan are typical of most LMICs with over 65%  
50 younger than 30 years of age, nuclear families with intergenerational co-residence and average  
51 household sizes of greater than 6 in urban areas. In Karachi, crowded neighborhoods, urban slum  
52 dwellings and poor adherence to social distancing measures add to the dynamics of the epidemic.  
53 Symptom-based surveillance strategies are the mainstay of prevention but supply (equipment,  
54 reagents, and NP-swabs) and demand ( stigma and fear) side issues limit its effectiveness along  
55 with undetected pre-symptomatic and asymptomatic transmission (3).

56 In this milieu, population-based sero surveys can be ideal (4, 5). However, high quality  
57 reliable population based surveys may not be feasible in an ongoing pandemic because they  
58 mandate training on personal protective equipment (PPE) for data collectors and allaying fear or  
59 anxiety in potential participants to avoid non-response bias. Sampling entire households can ease  
60 procedures of data and blood collection and provide an opportunity to study transmission at  
61 household level especially when lockdown and social distancing measures are in place.

62 Household transmission is a concern in closed congested neighborhoods of metropolitan  
63 cities when lockdown measures are in place. It is also of importance because of intergenerational

64 co residence of young and old, especially in light of recent evidence indicating that children can  
65 be major drivers of the epidemic (6). Secondary transmission from index cases in households  
66 using prospective follow up and active symptom monitoring with NP-PCR has indicated  
67 household attack rates of 32.4% (95% CI 22.4%–44.4%) (7). However it is resource intensive  
68 and may differ between symptomatic and asymptomatic households, as symptomatic individuals  
69 are more likely to transmit the virus (8-11). In a seroprevalence data, household transmission is  
70 easily estimable through the conditional risk of infection (CRI), namely the probability that an  
71 individual in a household is infected given that another household member is infected.

72 With this approach, we estimated changes in seroprevalence in low- and high-  
73 transmission neighborhoods of Karachi between April and June 2020 and compared the  
74 seroprevalence to total reported positive tests in the area to understand the level of under-  
75 reporting from gold standard RT-PCR testing. We also determined age - gender stratified  
76 estimates of seroprevalence and assessed the role of household transmission through conditional  
77 risk of infection (CRI) estimation; using the adapted World Health Organization (WHO) Unity  
78 protocol(12).

## 79 **MATERIALS and METHODS**

### 80 **Study Participants and sample collection**

81 We conducted the study in two areas (Figure 1). Four sub-administrative units or union  
82 councils of District East were purposely selected as high-transmission area based on highest  
83 number of cases to date and one union council of District Malir which had reported only 4 out of  
84 164 PCR tests was designated as a low-transmission area (Table 1)

85 Two serial cross-sectional surveys were performed at the household level, between April  
86 15-25 and June 25- July 11, 2020. Four teams, comprising of one data collector and one

87 phlebotomist each conducted the survey. A detailed list of cases was available in District East  
88 which allowed households to be selected through systematic random sampling as follows.  
89 Reference point was identified randomly from the list of current positive cases and spinning a  
90 bottle or pen indicated direction. The reference household with the positive case did not take part  
91 in the survey, instead every *n*th household was sampled based on second to last digit of a bank  
92 note until the lane ended in which case the team moved to the next lane or until the day was  
93 completed. Same sampling strategy was repeated daily until the overall sample size of the survey  
94 was achieved. In case of refusal from a household, the household on the right was sampled.  
95 Household level approval and individual participant written informed consent or assent was  
96 obtained. A slightly different approach was adapted in District Malir where a list of active cases  
97 was not available but a line listing of households from ongoing work (13) allowed simple  
98 random sampling to be used for household identification daily until the sample size was  
99 achieved. In both areas, all household members were eligible to participate irrespective of their  
100 case status.

101 All team members were trained in use of PPE, hand hygiene, disinfection techniques and  
102 safe transportation of biological samples. A 3 ml sample of blood from infants and 5 ml from  
103 those older than 1 year was collected by a trained phlebotomist and transported to the Infectious  
104 Diseases Research Lab for centrifugation, serum separation and storage at -20<sup>0</sup>C. Demographic  
105 and clinical data were collected using the adapted survey instrument from the WHO Unity  
106 Protocol for sero-epidemiology survey of COVID-19 (4).

### 107 **Laboratory analysis**

108 A commercial Elecsys® Anti-SARS-CoV-2 immunoassay (Roche Diagnostics), targeting  
109 both IgG and IgM against SARS-CoV-2 was performed at the Nutritional Research Laboratory

110 (NRL) at Aga Khan University. The manufacturer reports a specificity greater than 99.8% and  
111 sensitivity of 100% for individuals with a positive PCR test at least two weeks prior, and 88.1%  
112 sensitivity for those 7-13 days post a PCR positive test (14). Optimization of assay at the NRL  
113 was done on a small number of samples; 20 stored sera from before 2019 and 25 newly collected  
114 sera from RT-PCR-confirmed COVID-19 cases. All 20 pre-pandemic sera were negative while  
115 20 of 25 sera from cases were positive.

## 116 **STATISTICAL ANALYSIS**

### 117 **Sample Size**

118 The sample size for each phase of the survey per site was 500, totaling 1000 participants  
119 per survey. This allowed us to estimate an age adjusted prevalence for each site from 20-30 % at  
120 95 % confidence with precision of  $\pm 5\%$  and a design effect of 1.5 for household level clustering.

### 121 **Data Management**

122 All data was double entered on an SQL database and checked for missingness and  
123 consistency. Age was categorized and also presented as mean with standard deviations and  
124 median with interquartile ranges. Occupation was recoded to create a category of those working  
125 outside home. Those reporting fever with or without respiratory symptoms in the previous two  
126 months were categorized as symptomatic.

### 127 **Estimation of overall, age and gender stratified seroprevalence**

128 Age and gender-stratified seroprevalence estimates were computed using a Bayesian  
129 hierarchical regression model. This approach, described in detail in the Supplementary Material,  
130 accounts for uncertainty due to finite lab validation data,(15) and produces estimates with

131 uncertainty across age and gender groups using typical choices of uninformative or weakly  
132 informative prior distributions(16, 17).

133 Seroprevalence estimates were computed for each district and each survey phase  
134 independently, and all seroprevalence estimates are expressed as posterior means and 95% equal-  
135 tailed credible intervals based on 20000 samples from the Bayesian posterior distribution. All  
136 calculations were performed in R and samples from the posterior distributions were obtained  
137 using Stan (18).

### 138 **Household conditional risk of infection (CRI) analysis**

139 CRI, the probability that an individual in a household is infected, given that another  
140 household member is infected(19) was estimated by a fraction whose numerator is the total  
141 number of ordered pairs among infected individuals in the same household and whose  
142 denominator is the total number of ordered pairs in the same household in which the first  
143 individual in the pair is infected. A 95% confidence interval was estimated via bootstrap for each  
144 area by resampling households with replacement.

### 145 **Results**

146 A total of 2004 participants were enrolled across two phases from District East and  
147 District Malir. Figure 2 Panel A and B describe the flow of participants. Among households who  
148 agreed to take part, individual participation rate was 82.3% (1000 out of 1215 household  
149 members eligible) in phase 1 and 76.5% (1004 out of 1312 household members eligible) in phase  
150 2. Table 2 describes the baseline sociodemographic and clinical characteristics of the enrolled  
151 participants.

152 In Phase 1 of the study, only 2 of 500 samples tested positive in District East and 0 of 500  
153 tested positive in District Malir. In Phase 2 of the study, 100 of 500 samples (20.0%) tested  
154 positive in District East and 64 of 504 samples (12.7%) tested positive in District Malir. Both  
155 districts showed marked and significant increase in seroprevalence between time points. Of the  
156 total 166 participants who tested positive, only 9 (5.4%) gave a history of any symptoms.

157 To measure whether individuals in the same household were more likely to have similar  
158 sero- status, we computed the conditional risk of infection (CRI) for phase two. CRI estimates  
159 were 0.41 (95% CI 0.28-0.52) in District East and 0.31 (95% CI 0.16-0.4) in District Malir.  
160 These estimates are consistent with reported within household transmission, but cannot be  
161 completed attributed to within household transmission in the absence of a prospective  
162 design(19).

163 Given the correlation among household seropositivity values, we estimated  
164 seroprevalence via a Bayesian regression model (see Methods) which took into account  
165 household membership, age, and gender for each individual, as well as the lab validation data  
166 reported by the test manufacturer(14). Seroprevalence estimates by age and gender were then  
167 post stratified to adjust for the demographic makeup of the respective district.

168 In Phase 1, post-stratified seroprevalence was estimated to be 0.4% (95% CI 0%-1.3%) in  
169 District East and 0.2% (95% CI 0%-0.7%) in District Malir. In Phase 2, post-stratified  
170 seroprevalence was estimated to be 15.1% (95% CI 9.4%-21.7%) in District East, and 8.7%  
171 (95% CI 5.1%-13.1%) in District Malir. Seropositivity rates were indistinguishable between men  
172 and women within each district. Point estimates of seroprevalence were lower across all age

173 groups in District Malir than in District East, however there was significant overlap between the  
174 credible intervals for the two locations (Figure 3)

## 175 **Discussion**

176 The two serial serosurveys conducted in a metropolitan city of Pakistan in low- and high-  
177 transmission neighborhoods indicate a sharp increase in seroprevalence rates between April and  
178 June. While increase in the high transmission area is slightly greater and anticipated based on  
179 their identification as current “hotspots” of infection, as per the local health authorities, there has  
180 been a notable increase in the low transmission areas as well. The survey did not identify any  
181 difference in patterns of seroprevalence rates across age and gender categories. In both areas  
182 infection was mostly seen in individuals without any symptoms and within household sero-  
183 concordance suggests household transmission rates were high.

184 This is the first report from Pakistan and a low- and middle-income country, of a  
185 longitudinal population representative household survey conducted simultaneously in areas of  
186 low and high transmission.

187 The sharp elevation in seroprevalence in an area of presumably low transmission  
188 indicates that the virus continues to spread unchecked in populations where test timing is  
189 irregular or test numbers are suboptimal. It will be critical to monitor such populations because  
190 of the possible risk of resurgence within such areas once lockdown and contact tracing in high  
191 transmission areas is eased, as is currently being seen in North America (20).

192 Reasons for heterogeneity in SARS-CoV-2 transmission are largely unclear. Factors such  
193 as low socioeconomic area with poor access to clean water and sanitation and recurrent  
194 pneumonia and diarrheal illness in low-transmission area may be in play (21, 22). Transmission

195 of COVID-19 in areas with poor hygienic conditions is possibly reduced due to environmental  
196 microbes and resulting trained immunity seen with other SARS infections (23). However, several  
197 of the District East neighborhoods that were part of the study had similar characteristics  
198 including household size. This heterogeneity and its underlying causes needs to be further  
199 explored.

200 Seroprevalence in children was not different from that seen in adults and elderly. Recent  
201 evidence suggests that children may be more likely to transmit the virus. However data from  
202 population-based studies in any countries suggest children may be less likely to be infected (24).  
203 Our study found that children might be infected at a rate similar to adults. This may be due to the  
204 household nature of our survey during school closure, where we were powered better for the  
205 pediatric population compared to other studies. However, the margins of errors were broad  
206 suggesting the need of better-powered studies to establish rates of infection in children relative to  
207 adults. Specifically, such studies could better elucidate the role of children in transmission  
208 blocking strategies such as lockdowns, school closures and vaccination measures, once available.

209 Spread of infection in the household is an important consideration in SARS-CoV2  
210 transmission, especially in areas where there is lockdown and large families are roomed-in in  
211 small poorly ventilated spaces, typical of the neighborhoods of Karachi (25). The probability of  
212 an individual to have an infection in the presence of another infected household member in our  
213 study, as measured by conditional risk of infection (CRI) , was found to be similar to the actual  
214 and modelled secondary attack rates reported in other studies.(19)CRI can function as a  
215 substitute in situations where comprehensive surveillance and disease notification strategy is  
216 absent and secondary attack rates are difficult to calculate.

217           The strength of our study is knowledge of baseline seropositivity absent in most  
218 serosurveys and the serial nature with a two month inter survey interval allowing for a study of  
219 change in seroprevalence. A third round of survey is planned in August 2020. About one third or  
220 more of our sample includes children less than 18 years of age focusing on an understudied age  
221 group in the pandemic. Probability sampling which is community based is less biased for  
222 estimating prevalence and results are more generalizable than those from self-selecting or  
223 specific groups of individuals. Adaptation of the UNITY protocol allows for pooling of our  
224 results with other reports. The Electro Chemi Luminescence (ECL) technology used to test  
225 antibodies is sensitive and precise as per manufactures report (14).

226           Limitations include the limited geographical area of the study and the slight lag in  
227 duration of conduct of the second serosurvey that may have led to a slight over-estimation in  
228 District Malir. In-house validation on local samples was also not performed due to a limited  
229 supply chain of testing kits in Pakistan, however this was somewhat compensated by modeling  
230 directly on the data reported by the manufacturer.

## 231 **Conclusion**

232           There is a large increase in seroprevalence to SARS-CoV-2 infection even in areas where  
233 transmission is reportedly low. Most seropositives are asymptomatic and a majority of the  
234 population is still seronegative. There is high probability of an individual to be infected given  
235 exposure to another infected in the household, irrespective of symptoms. Enhanced surveillance  
236 activities of COVID-19 are required especially in low-transmission sites in order to determine  
237 the real direction of the pandemic and the risks of household transmission in tightly knit  
238 neighborhoods in urban LMIC settings.

239 **Acknowledgments**

240 The authors would like to acknowledge all the data collectors, phlebotomists and  
241 laboratory personnel who made this happen in the most difficult of circumstances.

242 **Author Bio**

243 Dr M Imran Nisar is an Assistant Professor in the Department of Pediatrics and Child  
244 Health at The Aga Khan University. He is an epidemiologist by training and his areas of interest  
245 include maternal & child health and infectious diseases.

246 **Funding:** This study was funded by the Infectious Disease Research Laboratory(IDRL) at the  
247 Aga Khan University in Karachi, Pakistan. None of the authors have any financial interest to  
248 disclose.

249 **References**

- 250 1. University JH. COVID-19 Dashboard by the Center for Systems Science and Engineering  
251 (CSSE) at Johns Hopkins University (JHU). 2020 [cited 2020 24/7/2020]; Available from:  
252 <https://coronavirus.jhu.edu/map.html>
- 253 2. Pakistan Go. COVID-19 Health Advisory Platform by Ministry of National Health  
254 Services Regulations and Coordination. 2020 [cited 2020 24/7/2020]; Available from:  
255 <https://covid.gov.pk/stats/pakistan>
- 256 3. Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, et al. Estimates of the  
257 severity of coronavirus disease 2019: a model-based analysis. The Lancet infectious diseases.  
258 2020.

- 259 4. Organization WH. Population-based age-stratified seroepidemiological investigation  
260 protocol for coronavirus 2019 (COVID-19) infection, 26 May 2020: World Health Organization;  
261 2020.
- 262 5. Winter AK, Martinez ME, Cutts FT, Moss WJ, Ferrari MJ, McKee A, et al. Benefits and  
263 challenges in using seroprevalence data to inform models for measles and rubella elimination.  
264 *The Journal of infectious diseases*. 2018;218(3):355-64.
- 265 6. Heald-Sargent T, Muller WJ, Zheng X, Rippe J, Patel AB, Kociolek LK. Age-Related  
266 Differences in Nasopharyngeal Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-  
267 2) Levels in Patients With Mild to Moderate Coronavirus Disease 2019 (COVID-19). *JAMA*  
268 *Pediatrics*.
- 269 7. Wu J, Huang Y, Tu C, Bi C, Chen Z, Luo L, et al. Household Transmission of SARS-  
270 CoV-2, Zhuhai, China, 2020. *Clinical Infectious Diseases*. 2020.
- 271 8. Lavezzo E, Franchin E, Ciavarella C, Cuomo-Dannenburg G, Barzon L, Del Vecchio C,  
272 et al. Suppression of a SARS-CoV-2 outbreak in the Italian municipality of Vo'. *Nature*. 2020:1-.
- 273 9. Gudbjartsson DF, Helgason A, Jonsson H, Magnusson OT, Melsted P, Norddahl GL, et  
274 al. Spread of SARS-CoV-2 in the Icelandic population. *New England Journal of Medicine*. 2020.
- 275 10. Ferretti L, Wymant C, Kendall M, Zhao L, Nurtay A, Abeler-Dörner L, et al. Quantifying  
276 SARS-CoV-2 transmission suggests epidemic control with digital contact tracing. *Science*.  
277 2020;368(6491).
- 278 11. Streeck H, Schulte B, Kuemmerer B, Richter E, Höller T, Fuhrmann C, et al. Infection  
279 fatality rate of SARS-CoV-2 infection in a German community with a super-spreading event.  
280 medrxiv. 2020.

- 281 12. Organization WH. Coronavirus disease (COVID-19) technical guidance: The Unity  
282 Studies: Early Investigations Protocols. 2020 [cited 2020; Available from: Coronavirus disease  
283 (COVID-19) technical guidance: The Unity Studies: Early Investigations Protocols
- 284 13. Ilyas M, Naeem K, Fatima U, Nisar MI, Kazi AM, Jehan F, et al. Profile: health and  
285 demographic surveillance system in peri-urban areas of Karachi, Pakistan. Gates Open Research.  
286 2018;2(2):2.
- 287 14. Ltd. RDI. ElectroChemiLuminescence (ECL). 2020 [cited; Available from:  
288 [https://diagnostics.roche.com/global/en/article-listing/electroChemiLuminescence-unique-](https://diagnostics.roche.com/global/en/article-listing/electroChemiLuminescence-unique-immunoassay-technology.html)  
289 [immunoassay-technology.html](https://diagnostics.roche.com/global/en/article-listing/electroChemiLuminescence-unique-immunoassay-technology.html)
- 290 15. Larremore DB, Fosdick BK, Zhang S, Grad YH. Jointly modeling prevalence, sensitivity  
291 and specificity for optimal sample allocation. bioRxiv. 2020.
- 292 16. Larremore DB, Fosdick BK, Bubar KM, Zhang S, Kissler SM, Metcalf CJE, et al.  
293 Estimating SARS-CoV-2 seroprevalence and epidemiological parameters with uncertainty from  
294 serological surveys. medRxiv. 2020.
- 295 17. Gelman A, Carpenter B. Bayesian analysis of tests with unknown specificity and  
296 sensitivity. medRxiv. 2020.
- 297 18. Stan - Stan. 2020 [cited 2020 25/7/2020]; Available from: <https://mc-stan.org/about/>
- 298 19. Curmei M, Ilyas A, Evans O, Steinhardt J. Estimating Household Transmission of SARS-  
299 CoV-2. medRxiv. 2020.
- 300 20. Tuite AR, Greer AL, De Keninck S, Fisman DN. Risk for COVID-19 Resurgence  
301 Related to Duration and Effectiveness of Physical Distancing in Ontario, Canada. Annals of  
302 Internal Medicine. 2020.

- 303 21. Quadri F, Nasrin D, Khan A, Bokhari T, Tikmani SS, Nisar MI, et al. Health care use  
304 patterns for diarrhea in children in low-income periurban communities of Karachi, Pakistan. The  
305 American journal of tropical medicine and hygiene. 2013;89(1\_Suppl):49-55.
- 306 22. Kerai S, Nisar I, Muhammad I, Qaisar S, Feroz K, Raza A, et al. A Community-Based  
307 Survey on Health-Care Utilization for Pneumonia in Children in Peri-Urban Slums of Karachi,  
308 Pakistan. The American journal of tropical medicine and hygiene. 2019;101(5):1034-41.
- 309 23. Netea MG, Giamarellos-Bourboulis EJ, Domínguez-Andrés J, Curtis N, van Crevel R,  
310 van de Veerdonk FL, et al. Trained Immunity: a Tool for Reducing Susceptibility to and the  
311 Severity of SARS-CoV-2 Infection. Cell. 2020.
- 312 24. Li X, Xu W, Dozier M, He Y, Kirolos A, Theodoratou E. The role of children in  
313 transmission of SARS-CoV-2: A rapid review. Journal of global health. 2020;10(1).
- 314 25. Statistics GoPBo. Housing units by number of rooms and type [cited; Available from:  
315 [www.pbs.gov.pk](http://www.pbs.gov.pk)

316 **Address for correspondence:** Fyezah Jehan, Associate Professor, Department of Paediatrics  
317 and Child Health, Aga Khan University, Stadium Road, Karachi 74800, Pakistan; Email:  
318 [Fyezah.jehan@aku.edu](mailto:Fyezah.jehan@aku.edu); Tel: +92 21 3486 4981

319

320 Table 1- District size, population density and population of the selected union councils

District	Union councils	Population
East	Dalmia/ Shanti Nagar	50,692
	Pehlwan Goth	136,229
	Safoora Goth	223,771
	Faisal cantonment	69,576
Malir	Ibrahim Hyderi	85,000

321

322

323 Table 2. General characteristics of the study participants

Characteristics	District East		District Malir	
	Phase 1	Phase 2	Phase 1	Phase 2
	(n= 500)	(n= 500)	(n= 500)	(n= 504)
Sex, male n (%)	256 (51.2%)	225 (45.0%)	207 (41.4%)	199 (39.5%)
Age in years , mean(SD)	26.2 (17.9)	25.9(16.7)	28.5 (17.9)	24.32 (16.7)
Age, categories, n (%)				
0-4 year	35 (7.0%)	22 (4.4%)	26 (5.2%)	33 (6.6%)
5-9 year	57 (11.4%)	54 (10.8%)	52 (10.4%)	74 (14.7%)
10-18 year	107 (21.4%)	139 (27.8%)	86 (17.2%)	120 (23.9%)
19-39 year	185 (37.0%)	170 (34.0%)	203 (40.6%)	186 (37.0%)
40-59 year	83 (16.6%)	97 (19.4%)	103 (20.6%)	65 (12.9%)
60 and above	33 ( 6.6%)	18 ( 3.6%)	30 ( 6.0%)	25 ( 5.0%)
Household size, mean (SD)	6.1(4.17)	6.6(3.6)	6(2.8)	5.89(3.2)
Working outside home, n (%)	155 (31.0%)	145 (29.0%)	143 (28.6%)	124 (24.6%)
Comorbids, n (%)				
None	453 (92.3%)	463 (93.0%)	473 (94.6%)	479 (95.0%)
Diabetes	20 ( 4.0%)	14 ( 2.8%)	6 ( 1.2%)	10 ( 2.0%)
Hypertension	14 (2.8%)	13 (2.6%)	16 (3.2%)	11 (2.2%)
Asthma or Allergy	3 (0.6%)	3 (0.6%)	1 (0.2%)	0 (0.0%)
Chronic Hepatitis	2 (0.4%)	3 (0.6%)	3 (0.6%)	3 (0.6%)
Chronic Heart Disease	0 (0.0%)	3 (0.6%)	1 (0.2%)	1 (0.2%)
Asymptomatic	435 (87.0%)	453 (90.6%)	472 (94.4%)	487 (96.6%)

Sought care	9 ( 1.8%)	9 ( 1.8%)	3 ( 0.6%)	4 ( 0.8%)
Hospitalization	1 ( 0.2%)	5 ( 1.0%)	1 ( 0.2%)	0 ( 0.0%)
History of Travel	2 (0.4%)	11 (2.2%)	8 ( 1.6%)	1 ( 0.2%)
Contact with suspected or confirmed COVID-19 case	1 ( 0.2%)	1 (0.2%)	0 ( 0.0%)	1 ( 0.2%)

324

325

326 Figure 1. Study sites.

327 Figure 2 Panel A. Flow chart of participants in the phase 1 of study


328 Figure 2 Panel B. Flow chart of participants in the phase two of study

329 Figure 3: Prevalence estimates by age and gender based on the data from the second survey. The  
330 circle represents the posterior mean seroprevalence and the bar represents the 95% equal-tailed  
331 credible interval. Posterior mean estimates for District East are consistently greater than those  
332 for District Malir, although there is significant overlap in the credible intervals for all age and  
333 gender subpopulations. No consistent patterns exist between the prevalence rates for males and  
334 females.


335

336


## Pakistan





## Karachi



## Study Area







prevalence

