

1 EVALUATION OF THE DURABILITY OF LONG-LASTING INSECTICIDAL NETS IN  
2 GUATEMALA

3 María Eugenia Castellanos<sup>1,2\*</sup>, Soledad Rodas<sup>1\*</sup>, Jose Guillermo Juarez<sup>1</sup>, Juan Carlos Lol<sup>1</sup>,  
4 Sayra Chanquin<sup>3</sup>, Zoraida Morales<sup>4</sup>, Lucrecia Vizcaino<sup>5</sup>, Stephen C. Smith<sup>5</sup>, Jodi Vanden Eng<sup>5</sup>,  
5 Henok G. Woldu<sup>6</sup>, Audrey Lenhart<sup>5</sup> and Norma Padilla<sup>1</sup>

6 <sup>1</sup> Center for Health Studies, Universidad del Valle de Guatemala, Guatemala, Guatemala

7 <sup>2</sup> Global Health Institute, College of Public Health, University of Georgia, Athens, Georgia,  
8 United States

9 <sup>3</sup>Medical Entomology Section, Vector-borne Diseases Program, Ministry of Public Health and  
10 Social Welfare, Guatemala, Guatemala

11 <sup>4</sup>National Malaria Sub-program, Vector-borne Diseases Program, Ministry of Public Health and  
12 Social Welfare, Guatemala, Guatemala

13 <sup>5</sup>Division of Parasitic Diseases and Malaria, Centers for Disease Control and Prevention,  
14 Atlanta, United States

15 <sup>6</sup> The Center For Health Analytics For National and Global Equity (C.H.A.N.G.E.), Columbia,  
16 United States

17 \*Both authors contributed equally to this manuscript.

18 RUNNING TITLE. Durability of LLINs in Guatemala

19 CORRESPONDING AUTHOR: Norma Padilla, PhD., Center for Health Studies, Universidad  
20 del Valle de Guatemala, Guatemala. E-mail: npadilla@ces.uvg.edu.gt

21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

## ABSTRACT

Background. Insecticide-treated bednets (ITNs) are widely used for the prevention and control of malaria. In Guatemala, since 2006, ITNs have been distributed free of charge in the highest risk malaria-endemic areas and constitute one of the primary vector control measures in the country. Despite relying on ITNs for almost 15 years, there is a lack of data to inform the timely replacement of ITNs whose effectiveness becomes diminished by routine use.

Methods. We assessed the survivorship, physical integrity, insecticide content and bio-efficacy of ITNs through cross-sectional surveys conducted at 18, 24 and 32 months after a 2012 distribution of PermaNet® 2.0 in a malaria focus in Guatemala. A total of 988 ITNs were analyzed (290 at 18 months, 349 at 24 months and 349 at 32 months).

Results. The functional survivorship of bednets decreased over time, from 92% at 18 months, to 81% at 24 months and 69% at 32 months. Independent of the time of the survey, less than 80% of the bednets that were still present in the household were reported to have been used the night before. Most of the bednets had been washed at least once (88% at 18 months, 92% at 24 months and 96% at 32 months). The proportion of bednets categorized as “in good condition” per WHO guidelines of the total hole surface area, diminished from 77% at 18 months to 58% at 32 months. The portion of ITNs with deltamethrin concentration less than 10mg/m<sup>2</sup> increased over time (14% at 18 months, 23% at 24 months, and 35% at 32 months). Among the bednets for which bioassays were conducted, the percentage that met WHO criteria for efficacy dropped from 90% at 18 months to 52% at 32 months.

Conclusion. While our assessment demonstrated that nets were in relatively good physical condition over time, the combination of declining bio-efficacy over time and low use rates limited the overall effectiveness of the LLINs. Efforts to encourage the community to retain, use, and properly care for the LLINs may improve their impact. Durability assessments should be included in future campaigns.

46 KEYWORDS. insecticide treated bednets, malaria, long-lasting insecticide nets, durability,

47 deltamethrin, attrition, survivorship, bio-efficacy, insecticide content

48

49

50

## INTRODUCTION

51 Globally, insecticide-treated bednets (ITNs) are a standard intervention for the  
52 prevention and control of malaria. Thirteen of the seventeen malaria-endemic countries in the  
53 Americas have incorporated within their malaria control and prevention programs the universal  
54 distribution of ITNs, and report having a policy of distributing ITNs through mass campaigns  
55 (World Health Organization, 2018).

56 The use of ITNs has had a significant impact on personal protection and in reducing  
57 malaria transmission. If high ITN coverage is sustained (> 80%), nets can provide protection to  
58 the larger community, including those households that do not use them (Hawley et al., 2003).  
59 The Global Malaria Program of the World Health Organization (WHO) recommends universal  
60 access and use of ITNs as a primary intervention for malaria control (World Health  
61 Organization, 2017).

62 Previous assessments of ITNs show a high degree of variation in ITN durability (Fettene,  
63 Balkew, & Gimblet, 2009; Kilian et al., 2011; Rosas-Aguirre et al., 2011), with ITNs from the  
64 same manufacturer showing significantly different levels of durability. The factors that influence  
65 this variability include the frequency of washing (including the quality of the water), type of  
66 laundry soap, washing and drying techniques, and daily levels of “wear and tear”. Translating  
67 this variability into a single 3-5 year durability estimate for ITNs could result in an inaccurate  
68 presumption of ITN effectiveness in areas where durability is compromised considerably faster.  
69 As such, understanding durability is a key consideration for the timing of ITN replacement  
70 strategies, and should be a component of any distribution campaign.

71 While ITN durability has been much more broadly studied in Africa, limited data are  
72 available from the Americas. An ITN distribution took place in Guatemala in 2012-2013 as part  
73 of malaria control activities supported by the Global Fund. Approximately 929,000 PermaNet®  
74 2.0 long-lasting insecticidal nets (LLINs) were distributed in 14 Health Areas that comprise a  
75 total of 111 municipalities in 12 departments of Guatemala. The primary objective of this study

76 was to assess the durability of the PermaNet® 2.0 LLINs over 32 months of routine use in an  
77 area with high malaria burden in the Mesoamerican sub-region. The three elements of  
78 durability, survivorship, physical integrity and insecticidal activity were evaluated, following the  
79 WHO guidelines for monitoring the durability of ITNs under operational conditions (World Health  
80 Organization., 2011). This evaluation was designed to provide relevant data for future plans  
81 around ITN replacement strategies that could be relevant in both Guatemala and more broadly  
82 in Latin America.

83

84

## METHODS

85 Study site and description of the ITN distribution. The study was conducted in the municipality of  
86 La Gomera in the department of Escuintla in southern Guatemala (Figure 1). Escuintla was  
87 included in the 2012-2013 ITN mass distribution campaign. According to national surveillance  
88 data, 56% of all malaria cases in Guatemala are reported from Escuintla (2012). As of 2012, La  
89 Gomera is divided into 19 communities, had a population of 60,299 inhabitants and reported  
90 23% of the malaria cases in the country (PAHO, 2017). In Escuintla, malaria is mainly  
91 transmitted by *Anopheles albimanus* and *Plasmodium vivax* is responsible for over 95% of the  
92 cases. 69,214 PermaNet® 2.0 (Vestergaard Frandsen, Lausanne, Switzerland) deltamethrin-  
93 treated LLINs were distributed in Escuintla from April 2012 to September 2013. It was the first  
94 mass LLIN distribution campaign in the region.

95 Experimental design. The evaluation was carried out from November 2013 to March 2015  
96 through three prospective cross-sectional surveys in randomly selected households using a  
97 multistage sample design. The first stage of the sampling consisted of a non-probabilistic  
98 sampling during which one municipality within the LLINs distribution area was selected (La  
99 Gomera). The following stage was done using a cluster survey methodology with each cluster  
100 size proportional to its population. The evaluation included only those communities that received  
101 LLINs approximately 18 months (range=17-19 months) prior to the first survey. Twelve

102 communities met this criterion and were included in the 18-and 24-months survey. In this twelve  
103 communities, 4,076 had been delivered in the study area between April-June 2012. At the 32-  
104 month survey, only 8 of these 12 communities were included since in the other four  
105 communities a round of LLIN replacement occurred in 2014. The selection of households to be  
106 sampled within each stratum was done through a simple random sampling. Nets were not  
107 labelled prior to distribution and therefore users were not aware that the nets were to be  
108 evaluated, reducing the possibility of the Hawthorne effect (Earl-Slater, 2002).

109 Sample size for the cross-sectional surveys. To determine survivorship and physical integrity of  
110 the LLINs, sample size estimates were based on the confidence interval around the attrition rate  
111 at each time period, according to values reported for Kenya (11% for 18 months, 15% for 24  
112 months, 23% for 30 months) (Gimnig, 2013). We assumed a relative error of 5%, a confidence  
113 level of 95%, and a design effect of 2.0. We assumed a non-response proportion of 15% at 18  
114 months. For the surveys conducted at 24 months and 32 months we had to increase this  
115 proportion to 40% and 55%, respectively as when we visited the households many of them  
116 reported that they had not received any LLIN. Also, for the surveys at 24 months and 32  
117 months, we increased by 5% the number of households to survey because we had used Google  
118 Maps to select the households (more details in the next section) and it was possible that some  
119 structures that had been enumerated as households were another type of structure (Wampler,  
120 Rediske, & Molla, 2013). Thus, the numbers of households selected to survey at each time  
121 point were as follows: n=348 at 18 months, n=583 at 24 months, and n=887 at 32 months, for a  
122 total of 1,818 households.

123 Selection of households and nets evaluated. For the 18-month survey, households were  
124 identified using the Ministry of Health (MoH) 2012 LLINs campaign distribution list. We used the  
125 OpenEpi Random Program ([www.openepi.com](http://www.openepi.com)) to generate a random number list for the final  
126 selection of households. We then used the addresses of the households provided by the MoH to  
127 locate the household. In the 24-month and 32-month surveys, we used aerial Google Earth

128 imagery to identify houses for sampling (Map data: Google, DigitalGlobe). The high resolution  
129 of the 2013 images of the selected communities allowed for good accuracy when mapping  
130 individual houses. The Google Earth file was imported to ArcMap 9.3 (ESRI, Redlands, CA,  
131 USA) using a KML to shapefile extension (Parent, 2009). Longitude and latitude coordinates  
132 (WGS84 coordinate systems) were added to the attributes table which was saved as a dBASE  
133 file. This was imported to an Excel spreadsheet for the random household selection, using  
134 simple random sampling with the OpenEpi Random Program. The geographic information  
135 system (GIS) information about each selected community was uploaded to GPS Visualizer  
136 ([www.gpsvisualizer.com](http://www.gpsvisualizer.com)), and subsequently added to a Garmin global positioning system (GPS)  
137 map 62S (Garmin, USA) with a 20m proximity alarm. A field validation of the maps and GPS  
138 equipment was carried out prior to conducting the fieldwork.

139 Questionnaires. Program staff from the National Malaria Sub-program of the Vector-borne  
140 Diseases Program of the MoH and researchers from Universidad del Valle de Guatemala  
141 conducted the surveys. The survey teams were trained to understand the structure of the survey  
142 and how to deliver it using the personal digital assistants (PDAs) that included a GPS element.  
143 Survey questions were based on WHO Durability guidelines (World Health Organization., 2011),  
144 adapted to local settings, translated to Spanish and validated. The questionnaire was  
145 administered primarily to the adult woman in charge of the household in order to collect as much  
146 data as possible on the maintenance and washing frequencies of the LLINs. If an adult woman  
147 was not present, then the questionnaire was administered to any adult household member.  
148 Questions were included on a) demographic characteristics of the enrolled household; b) net  
149 survivorship, attrition and use; and c) physical integrity of the fabric measured *in situ*. Of the  
150 total number of LLINs that were delivered to the household, one was randomly selected for  
151 further analysis (1 enrolled household = 1 enrolled LLIN). Questions on net use and physical  
152 integrity were directed towards this randomly selected specific net if it was still present in the  
153 household. If the selected LLIN was no longer present, other questions related to reasons for its

154 absence were pursued. In the 24-month and 32-month surveys, if a previously surveyed  
155 household was selected, we still randomly selected a LLIN to evaluate. If the selected LLIN had  
156 already been evaluated, we would select another one if available.

157 Physical integrity. LLINs were assessed *in situ* with the net left hanging over the sleeping space  
158 if the position allowed it. Holes were classified according to the WHO 2011 categories: (1) holes  
159 smaller than a thumb (~0.5 to 2 cm diameter), (2) bigger than a thumb but smaller than a fist (~2  
160 to 10 cm diameter), (3) bigger than a fist but smaller than a head (~10 to 25 cm diameter) and  
161 (4) bigger than a head (> ~ 25 cm diameter) (World Health Organization., 2011). A standard  
162 ruler was available in case there was uncertainty on a measurement.

163 Insecticide content. Deltamethrin content, in mg/m<sup>2</sup>, was measured using a Tracer III-SD  
164 handheld X-ray fluorescence (XRF) analyzer (Bruker Nano Analytics, Inc., Kennewick, WA,  
165 USA). For this analysis, deltamethrin content was calculated from the intensity of 11.549-  
166 12.248 keV X-rays emitted by the bromine atoms of the deltamethrin in the sample (Smith et al.,  
167 2007). Calibration samples of polyester mosquito netting treated with six different levels of  
168 deltamethrin (0-114 mg/m<sup>2</sup>) were used to correlate intensity with deltamethrin content. The  
169 correlation was linear with R<sup>2</sup>=0.98, minimum. For whole bednet analysis, the net was folded in  
170 a way to create a 24-layer sampling point, comprising 8 points from the roof of the net, and 4  
171 points from each of the 4 sides. This multilayer sample was then analyzed to yield the average  
172 deltamethrin content over the 24 locations. Since folding the net and collecting the data takes  
173 only 2-3 minutes, the average deltamethrin content of many nets can be measured quickly and  
174 conveniently (Anshebo et al., 2014; Craig et al., 2015).

175 Sample size for cone bioassays. From the LLINs evaluated in the households in each  
176 community, a sub-sample of the nets reported to have been used to sleep at least once was  
177 randomly selected and collected for bio-efficacy testing. In order to detect the loss of at least  
178 5.6% of insecticide, with respect to the baseline level of 55 mg/m<sup>2</sup>, a power of 90%, alpha=0.05,  
179 and using a standard deviation of 5, a subset of 55 LLINs were randomly selected during the

180 18-month survey, 60 at 24 months, and 65 at 32 months, with increasing number of nets  
181 selected in order to compensate for an estimated non-response effect of 15%.

182 Collected nets were placed individually in plastic bags and grouped inside large black  
183 polyethylene bags and kept at  $\pm 20^{\circ}\text{C}$  until their arrival to the laboratory. Householders were  
184 given a new PermaNet<sup>®</sup> 2.0 bednet to replace the collected net.

185 WHO cone bioassays. Cone bioassays were performed on samples from each of the five sides  
186 of each bednet and following the sampling scheme recommended by WHO (World Health  
187 Organization., 2011). Until processed the net samples were stored at  $4^{\circ}\text{C}$  wrapped in aluminum  
188 foil and placed inside plastic bags. Briefly, we used 2-3-day old, non-blood fed *Anopheles*  
189 *albimanus* females from the laboratory insecticide susceptible SANARATE strain. Two  
190 replicates of 20 mosquitoes were exposed to two study PermaNet<sup>®</sup> net pieces and two  
191 replicates of 10 mosquitoes were exposed to an untreated polyester net as control. Mosquitoes  
192 were exposed for 3 minutes and the 60 minutes knock-down and 24-hour mortality recorded  
193 according to WHO criteria. Data were adjusted according to Abbot's formula. Bioassays were  
194 performed under laboratory conditions,  $27 \pm 2^{\circ}\text{C}$  and  $70 \pm 10\%$  relative humidity. Knock down  
195 at 60 minutes (KD60) and mortality at 24 hours were calculated as the number of mosquitoes  
196 knocked down at 60 min and the number of dead mosquitoes at 24 hours proportional to the  
197 total number exposed. Results were considered valid if the control mortality was lower than 5%;  
198 between 5-20% control mortality, results were corrected using Abbott's formula and if control  
199 mortality was greater than 20%, the bioassays for that net were repeated (World Health  
200 Organization., 2011).

201 Verification of baseline LLIN insecticidal content. From the PermaNet<sup>®</sup> 2.0 batch distributed in  
202 2012, 11 nets were retained and analyzed using XRF and WHO cone bioassays at 24 months  
203 post-distribution to assess the insecticidal content of the unused LLINs (considered time-point  
204 0).

205 Data analysis

206 The main outcomes of the study were overall and functional survivorship, physical integrity,  
207 insecticide content measured by XRF and bio-efficacy measured by the WHO cone bioassay.

208 Descriptive demographic characteristics of the households that were enrolled and details of their  
209 reported use and handling of the LLINs at the three-survey time-points (18 months, 24 months  
210 and 32 months) were estimated using proportions and measures of central tendency.

211 The overall survivorship at each time-point was defined as the number of enrolled LLINs still  
212 present in the households divided by the total number of enrolled LLINs. The functional  
213 survivorship was calculated at each time-point as follows: number of enrolled LLINs still present  
214 and “serviceable” (see definition below) in the households divided by the number of LLINs still  
215 present + LLINs not present owed to attrition reasons (damaged and thrown away or LLIN used  
216 for other purposes) (Committee & Secretariat, 2013; Tan et al., 2016; WHO Malaria Policy  
217 Advisory Committee, 2013). We then calculated the median survival time, defined as “the time  
218 point at which the estimate of functional LLIN survival crosses the 50% mark” (WHO Malaria  
219 Policy Advisory Committee, 2013) using the following formula:  $tm = t_1 + \frac{(t_2 - t_1) \times (p_1 - 50)}{(p_1 - p_2)}$ , where  
220  $tm$  is the median survival time,  $t_1$  and  $t_2$  were the 24 and 32 month survey points (in years) and  
221  $p_1$  and  $p_2$  were the functional survivorship proportions for the 24 and 32 month points,  
222 respectively.

223 The physical integrity of LLINs was measured by estimating the proportion of LLINs with at least  
224 one hole of any size, the total number of holes/LLIN, and the total hole area (cm<sup>2</sup>)/ LLIN in  
225 quartiles. The total hole surface area (THSA) was calculated by multiplying the number of holes  
226 by the area of each hole (1.2, 28.3, 240.5 or 706.9 depending on the size category of the holes)  
227 and then summing across the categories (Vanden Eng et al., 2015; World Health Organization,  
228 2013). Holes smaller than 0.5 cm were not included in the assessment. We classified the LLIN  
229 condition as good (<79 cm<sup>2</sup> THSA), damaged (79-788 cm<sup>2</sup> THSA) or severely torn (>=789 cm<sup>2</sup>  
230 THSA) following the categorization recommended by WHO, and an LLIN was considered

231 “serviceable” if it was classified as good or damaged (WHO Malaria Policy Advisory Committee,  
232 2013). The proportion of LLINs with evidence of repairs was also calculated.  
233 The median concentration and interquartile range (IQR) of deltamethrin ( $\text{mg}/\text{m}^2$ ) obtained  
234 through the XRF analyses was estimated at each time point, including unused nets. We  
235 calculated the proportions of LLINs with a concentration of less than  $10 \text{ mg}/\text{m}^2$  and  $25 \text{ mg}/\text{m}^2$ .  
236 These values were chosen because LLINs with less than  $10 \text{ mg}/\text{m}^2$  are considered to not have  
237 the minimum effective concentration of insecticide, and nets with less than  $25 \text{ mg}/\text{m}^2$  are  
238 considered to not contain an optimum insecticide concentration (Ashebo et al 2014).  
239 For the bio-efficacy results, we estimated the distribution and geometric mean (CI95) for the  
240 KD60 and mortality at 24 hours, for each survey time point including unused nets (Tan et al.,  
241 2016). We also calculated the proportion (and CI95) of LLINs that showed  $\geq 80\%$  mortality at 24  
242 hours, or  $\geq 95\%$  KD60 at each time point, including for unused nets (World Health Organization,  
243 2013).  
244 We assessed the association between the cone bioassay results and the deltamethrin  
245 concentration obtained by the XRF using Locally Weighted Scatterplot Smoothing (LOESS)  
246 regression analysis to estimate Kendall's correlation coefficient. We then developed a  
247 predictive model using piecewise (segmented) regression in which we explored if the use of the  
248 deltamethrin concentration measured by XRF and adjusted by other variables (More details in  
249 Appendix) can predict if results of a bioassay on the LLIN will exceed or not the WHO threshold  
250 of  $\geq 80\%$  mortality at 24 hours.  
251 We analyzed nets based on a series of sequential criteria: a) Distributed in the household, b)  
252 Still present in the household, c) Used at least once, d) Used the night before the survey, e) In  
253 serviceable condition and e) With a concentration of deltamethrin above  $10 \text{ mg}/\text{m}^2$  as measured  
254 by XRF (we didn't include this criterion in the 18-month survey as we only analyzed a small  
255 number of LLINs with XRF). For the 24- and 32-month surveys, we developed a working  
256 definition of 'LLIN providing adequate protection', for nets that met all of these criteria. For this

257 purpose, we estimated the proportion of LLINs that sequentially meet each criterion. In order to  
258 be analyzed in each criterion, the LLINs should have met the requirement of the previous  
259 criterion. We then estimated the overall proportion of LLINs providing adequate protection  
260 based on our study definition.

261 We used the Cochran-Armitage Trend Test to assess trends across the time-points and the  
262 different categorical outcomes. For continuous outcomes, we used the Kruskal-Wallis test  
263 (differences among all groups) and the Wilcoxon rank sum test (pairwise comparisons).

264 All analyses were conducted using SAS software v 9.4 (SAS Institute, Cary, NC, US) and R  
265 v3.6.0 (R Foundation for Statistical Computing, Vienna, Austria, 2016).

266

267

## RESULTS

268 Descriptive characteristics of the study population. A total of 1,013 LLINs were enrolled in the  
269 study, of which 988 (98%) of them were considered to have valid information. Of these, 290  
270 were evaluated at 18 months, 349 at 24 months and 349 at 32 months (Figure 2). The  
271 education level of the head of household and the proportion of households with electricity that  
272 participated in the surveys remained consistent throughout the duration of the study (Table 1).  
273 We observed a two-fold increment in the proportion of households that owned a flush toilet and  
274 that had piped water as their source of drinking water in the 32-month survey as compared with  
275 the previous time points. The median number of individuals that slept in the household the night  
276 before the survey was four (Supplemental material, Table S1). According to the census of  
277 LLINs distributed, each household received a median of three LLINs, which corresponded to the  
278 median number of sleeping places used in the household the night before of the survey.

279 Overall and functional survivorship. The proportion of LLINs still present in the household  
280 decreased over time ( $p < 0.001$ ), with an overall survivorship of 86% (CI95: 82-90) at 18 months,  
281 76% (CI95: 72-81) at 24 months and 66% (CI95: 61-71) at 32 months. The main reported cause  
282 of loss at 18 months (56% , CI95: 40-72) and 24 months (54%, CI95: 43-65) was that the net

283 was given away to others, whereas at the 32-month survey, the main cause was that the net  
284 was damaged and thrown away (43% CI95: 34-52) (Figure 3 and Table S2). The functional  
285 survivorship was 92% (CI95: 88-95) at 18 months, 81% (CI95: 76-86) at 24 months and 69%  
286 (CI95: 64-75) at 32 months. The median survival time was extrapolated to be 3.7 years (CI95:  
287 3.4-4.2).

288 LLIN use and handling. More than 90% of the LLINs still present in the households were  
289 reported to have been used at least once at any of the time points (93% at 18 months, 97% at  
290 24 months and 95% at 32 months). However, among the LLINs used at least once, less than  
291 80% were reported to have been used the night before the survey (74% at 18 months, 79% at  
292 24 months and 71% at 32 months) (Table 2). Almost a third of these LLINs were found not  
293 hanging (stored away or visible but not hung up) at each of the survey time-points.

294 A large proportion of the LLINs that were used at least once had been washed (88% at 18  
295 months, 92% at 24 months, 96% at 32 months) and in >55% of the cases, nets had been  
296 washed less than 1 month preceding the survey (Table 2). Among the washed LLINs, the  
297 majority were soaked (64% at 18 months, 62% at 24 months, 60% at 32 months), although in  
298 over 70% of these, the reported soaking time was less than one hour (80% at 18 months, 73%  
299 at 24 months, 74% at 32 months). The principal soap used to wash the LLINs was detergent  
300 powder, either alone or in combination with bar soap and/or bleach**Error! Reference source**  
301 **not found.** Over time, there was a reduction in the proportion of LLINs that were scrubbed hard  
302 or beaten on a hard surface during washing (62% at 18 months, 51% at 24 months and 46% at  
303 32 months,  $p = 0.0007$ ). In most cases, the washed LLINs were dried outside in the sun (70% at  
304 18 months, 69% at 24 months and 60% at 32 months).

305 Physical integrity. The median number of holes in the LLINs was 2 (IQR 0,7) at 18 months, 4  
306 (0,15) at 24 months and 5.5 (IQR 1,16) at 32 months (Table 3). The median total hole area  
307 (cm<sup>2</sup>) in the bednets was 2.4 (IQR 0,63.8) at 18 months, 8.4 (IQR 0-271.5) at 24 months and  
308 34.3 (IQR 1.2,327.8) at 32 months. The proportion of LLINs considered in good condition

309 decreased from 77% at 18 months to 70% at 24 months and to 58% at 32 months ( $p < 0.0001$ ).

310 At 32 months, 84% of the LLINs were considered still in 'serviceable' condition.

311 Respondents reported new holes in the month prior to the survey in 24% of bednets at 18  
312 months, in 15% of bednets at 24 months and in 27% at 32 months. The principal reported cause  
313 of these new holes was that the bednet tore or was split when caught on an object (45% at 18  
314 months, 31% at 24 months and 38% at 32 months). Other reported causes of holes were  
315 children (21% at 18 months, 10% at 24 months and 19% at 32 months) and animals (11% at 18  
316 months, 13% at 24 months and 10% at 32 months). However, a large proportion of these new  
317 holes did not have a known cause reported by the respondent (21% at 18 months, 36% at 24  
318 months and 24% at 32 months). No new holes were reported due to burns. A minority of the  
319 LLINs with holes had been repaired (23% at 18 and 24 months, 44% at 32 months), with  
320 stitches as the principal mode of repair (Table 3).

321 Insecticide content. The median deltamethrin concentrations ( $\text{mg}/\text{m}^2$ ) measured by XRF  
322 differed significantly by survey time-point ( $p < 0.0001$ ). For unused nets, the median was 49.8  
323  $\text{mg}/\text{m}^2$  (IQR 47.2, 53.0,  $n=11$ ), at 18 months it was 23.2  $\text{mg}/\text{m}^2$  (IQR 11.2, 31.2,  $n=51$ ), at 24  
324 months it was 14.6  $\text{mg}/\text{m}^2$  (IQR 8.8, 25.6,  $n=264$ ) and at 32 months it was 13.9  $\text{mg}/\text{m}^2$  (IQR  
325 6.6, 30.0,  $n=218$ ). The reductions in insecticide content were statistically significant between  
326 unused nets and values measured at 18 months ( $p < 0.001$ ) and between 24 months and 32  
327 months ( $p=0.027$ ), but not between 18 months to 24 months ( $p=0.21$ ) (Figure 4). The proportion  
328 of LLINs with a deltamethrin concentration of less than 10  $\text{mg}/\text{m}^2$  increased over time: 0% for  
329 unused nets, 14% (CI95: 4-24) at 18 months, 23% (CI95: 18-29) at 24 months and 35% (CI95:  
330 29-42) at 32 months ( $p < 0.0001$ ). When the cut-off was set at 25  $\text{mg}/\text{m}^2$ , the proportion of LLINs  
331 below the threshold was: 0% for unused nets, 57% (CI95: 43-71) at 18 months, 69% (CI95: 63-  
332 75) at 24 months and 73% (CI95: 67-79) at 32 months ( $p < 0.0001$ ).

333 Bio-efficacy. Among the sub-set of LLINs on which bioassays were conducted, the geometric  
334 mean mortality at 24 hours decreased over time: 100% (CI95: 100, 100) for unused nets ( $n=11$ ),

335 82% (CI95: 76-89) at 18 months (n=51), 52% (CI95: 43-61) at 24 months (n=60) and 45%  
336 (CI95: 35-55) at 32 months (n=63) (Figure S1). In the case of KD60, the geometric means were  
337 100% (CI95: 99-100) for unused nets (n=11), 96% (CI95: 94-99) at 18 months (n=51), 84%  
338 (CI95: 77-92) at 24 months (n=60) and 80% (CI95: 73-87) at 32 months (n=63) (Figure S2). On  
339 summary, the proportion of LLINs that passed the WHO  $\geq 80\%$  mortality threshold or  $\geq 95\%$   
340 KD60 for bio-efficacy were 100% for unused nets, 90% (CI95: 82-99) at 18 months, 68% (CI95:  
341 56-80) at 24 months and 52% (CI95: 40-65) at 32 months ( $p < 0.0001$ ).

342 Association between mortality at 24 hours and deltamethrin concentrations ( $\text{mg}/\text{m}^2$ ) measured  
343 by XRF. There was a positive linear relationship between deltamethrin concentration measured  
344 by XRF and the percent mortality at 24 hours up to approximately  $25 \text{ mg}/\text{m}^2$  of deltamethrin  
345 concentration ( $r=0.62$ ,  $p < 0.0001$ ) (Figure 5 & Figure S3). The summary statistics from the  
346 fitted segmented regression model confirmed the positive relationship between deltamethrin  
347 content and mosquito mortality (Table S3), with an estimated breakpoint at  $24.4 \text{ mg}/\text{m}^2$ . The age  
348 of the LLINs was also an independent predictor of mosquito mortality (%) after 24 hours. The  
349 mortality decreased with increasing age of the LLIN. This model using segmented regression  
350 was able to predict 89.7% of the time if a LLIN will achieve the WHO 80% of mortality threshold  
351 for bio-efficacy using only two predictors: insecticide content measurements obtained from  
352 bromine x-ray fluorescence testing and the age of the LLINs.

353 Overall proportion of LLINs providing adequate protection. The proportion of LLINs providing  
354 adequate protection was 38% (CI95: 33-44) at 24 months and 21% (CI95: 17-26) at 32 months.  
355 Three components contributed most to the observed decreases in adequate protection: 1) A  
356 decrease between the number of LLINs distributed and the proportion of LLINs still present in  
357 the household-which was particularly important at 32 months (drop of 34%); 2) A reduction  
358 between the proportion of LLINs present and used at least once to the proportion of LLINs  
359 present and used the night before of the survey (drop of 21%, 16% and 18% for 18, 24 and 32  
360 months, respectively); and 3) A drop between the proportion of LLINs in serviceable condition

361 and the proportion of LLINs in serviceable condition with a deltamethrin concentration of at least  
362 10 mg/m<sup>2</sup> (drop of 12% and 15% for 24 and 32 months, respectively) (Figure 6 and Table S4).

363

## 364 DISCUSSION

365 In this prospective multiple cross-sectional study, we assessed the durability of PermaNet<sup>®</sup> 2.0  
366 LLINs distributed in a region of persistent malaria transmission in Guatemala by measuring their  
367 overall and functional survivorship, physical integrity, insecticide content and bio-efficacy.

368 The proportion of LLINs present in the household and in ‘serviceable’ condition was 69% after  
369 32 months of routine use, with an extrapolated median survival time of 3.7 years. The median

370 survival time reported in our study is higher than similar studies conducted in Africa that have

371 also evaluated PermaNet<sup>®</sup> 2.0 LLINs. In Tanzania the estimated survival was 2.5 years, in

372 Zanzibar it was 3.1-3.3 years and in Zambia it was 2.5-3 years (Haji et al., 2020; Lorenz et al.,

373 2019; Tan et al., 2016). In our study, the physical integrity of the LLINs—a key component for

374 the functional survivorship—was high in each of the survey time-points (range: 84-94%),

375 whereas other studies have reported lower values (e.g. 68% at 36 months) (Haji et al., 2020).

376 These differences are further highlighted by the median total hole area, which in our study was

377 only 34 cm<sup>2</sup> at 32 months, whereas other studies evaluating PemaNet<sup>®</sup> 2.0 LLINs reported

378 values of 390 cm<sup>2</sup> and 106 cm<sup>2</sup> (Craig et al., 2015; Vanden Eng et al., 2015). In general, the

379 size of the holes found on the LLINs in Guatemala was smaller, resulting in a greater number of

380 LLINs categorized as being in ‘serviceable’ condition and a higher functional survivorship. The

381 low proportion of new holes caused by thermal damage or animals as compared to other

382 studies may also help explain these differences (Morgan et al., 2015; Mutuku et al., 2013).

383 Of the LLINs containing holes, only 44% showed evidence of repair.

384 From a programmatic perspective, we also estimated the overall survivorship of the LLINs,

385 which was 66% after 32 months. This was very similar to the functional survivorship, indicating

386 that two thirds of the campaign LLINs were still present and functioning well in the surveyed

387 households nearly 3 years after their distribution. However, reasons explaining the overall and  
388 functional survivorship were different in the 18- and 24-month surveys. At these time points, the  
389 main reason reported for not having a net present in the household was because it had been  
390 'given away to others', not because of wear and tear. These findings are similar to previous  
391 reports (Gnanguenon, Azondekon, Oke-Agbo, Beach, & Akogbeto, 2014; Mansiangi et al.,  
392 2020; Obi et al., 2020), highlighting the real-life challenges faced by national malaria programs,  
393 which reach beyond the quality of the fabric of the nets. In a large study evaluating 14  
394 household surveys from four African countries, 34% of the nets that were not present in the  
395 household had been given away, the majority to family members and within the first month after  
396 a distribution campaign (Koenker et al., 2014).

397 In terms of usage and handling, between 71-79% of the users reported to have used the LLINs  
398 the night preceding the survey, with almost a third of the nets observed not being in use as they  
399 were not hanging in the sleeping place. Previous studies that have evaluated the usage of  
400 LLINs in Latin America have shown a variety of results around LLIN usage. In Colombia, the  
401 usage was 51.1% after 8 months, whereas in Venezuela, it was over 90% after 6 months,  
402 highlighting intra-regional differences (Alvarado, García, Villarroel, & Rosas-Aguirre, 2011;  
403 Cabrera, Diaz, Pareja, & Santamaría, 2009).

404 We detected decreases in insecticide content over time, with a drop of 53% detected between  
405 unused nets ( $49.8 \text{ mg/m}^2$ ) and nets at the 18-month survey point ( $23.2 \text{ mg/m}^2$ ). These levels  
406 were lower than levels detected on PermaNet<sup>®</sup> 2.0 in Ethiopia, where the mean concentration of  
407 deltamethrin was  $44.1 \text{ mg/m}^2$  after 14-20 months of use (Anshebo et al., 2014). In our study,  
408 the proportion of nets with deltamethrin levels  $\geq 10 \text{ mg/m}^2$  after 32 months was 65%, a stark  
409 contrast to the 95% observed in the aforementioned study. The values obtained by XRF to  
410 estimate total insecticide content showed similar trends to the bioassays, where just 52% of the  
411 LLINs at 32 months met the WHO bioassay threshold of bio-efficacy. The frequent washing of  
412 the LLINs and the high proportion that were dried under direct sunlight are some of the factors

413 that could have contributed to this decline (Agossa et al., 2014; Atieli, Munga, Ofulla, & Vulule,  
414 2010). As other studies have detected, we also found that a measurement of  $\geq 25$  mg/m<sup>2</sup> by  
415 XRF was more likely to indicate a LLIN with a mortality at 24 hrs. of  $\geq 80\%$ , suggesting that this  
416 threshold is more closely associated with optimal bio-efficacy than 10 mg/m<sup>2</sup> (Anshebo et al.,  
417 2014).

418 When we integrated the results of survivorship, physical integrity, usage and insecticide content,  
419 the estimated that the level of protection provided by the LLINs was sub-optimal. The overall  
420 proportion of LLINs providing adequate protection was only 38% after 24 months and dropped  
421 further to 21% at 32 months. National malaria data showed that the number of cases in  
422 Escuintla remained constant during the period of 2012-2015, suggesting that the LLINs did not  
423 provide sufficient protection.

424 Several lessons can be gathered from our findings. First, we suggest that three key factors  
425 contributed to the sub-optimal level of protection: the loss of LLINs still present in the household,  
426 the degree of usage of the LLINs, and the low percentage of LLINs with adequate insecticide  
427 content. Given that these factors are potentially compromising the protective effectiveness of  
428 the LLINs more than the durability of the fabric, activities focusing on community engagement  
429 and the inclusion of local civil society around LLIN use could be particularly impactful. Second,  
430 our findings suggest that personal redistribution of nets is an important trend in this region,  
431 similar as in Africa. By giving nets away to friends and family, communities may be  
432 compensating for inadequate campaign planning. National malaria programs may consider  
433 including local leadership and shift toward a more decentralized approach to improve the design  
434 of LLIN campaign distributions (Gosling et al., 2020). Third, we were able to show that in areas  
435 with limited resources to carry out the WHO cone bioassay, the measurement of insecticide  
436 content by bromine x-ray fluorescence might be an accurate alternative methodology to  
437 estimate the bio-efficacy of LLINs.

438 A major strength of our study is that we concurrently measured survivorship, physical integrity,  
439 insecticide content and bio-efficacy. There are few data on these indicators reported from Latin  
440 America, and our study provides one of the most comprehensive evaluations to date for the  
441 Americas. In addition, by following standard WHO guidelines, our results can be compared to  
442 outcomes arising from similar studies conducted elsewhere, while at the same time including  
443 variables that were customized to the local context. The use of GIS in the 24- and 32-month  
444 surveys provided us with a more accurate identification of the households in each community,  
445 and our randomization process allowed us to limit bias associated with the selection of the  
446 surveyed households. However, our study also had several limitations. First, we were not able  
447 to conduct a household survey at the 6- or 12-month time-points. Second, we had to exclude  
448 four out of the twelve selected communities in the 32-month survey, which could introduce  
449 selection bias. Third, the answers for the questions related to usage, care and handling of the  
450 LLINs are prone to recall and social desirability bias (Althubaiti, 2016).

451

## 452 CONCLUSION

453 The lifetime of PermaNet<sup>®</sup> 2.0 is expected to be approximately 3 years. In our study, after 2  
454 years, over two thirds of the LLINs did not contain the optimal deltamethrin concentration of 25  
455 mg/m<sup>2</sup> and 32% of them did not meet the WHO criteria for bio-efficacy. Even though the nets  
456 were in good physical condition at 32 months, their diminished bio-efficacy and reported lack of  
457 consistent use compromised the protection provided to the users. The widespread provision of  
458 LLINs is a cornerstone of global efforts for the control and potential elimination of malaria, which  
459 is now a near-term goal in several Latin American countries (World Health Organization, 2018).  
460 We recommend that future evaluations of the durability of LLINs should always include  
461 measurements of survivorship, usage and handling of the LLIN, physical integrity, and  
462 insecticidal activity (insecticide content and/or bio-efficacy) in order to estimate the protective

463 effect of the nets in a community. The engagement of the National Malaria Program with the  
464 community is critical to the success of future LLIN distributions in Guatemala.

#### 465 LIST OF ABBREVIATIONS

466 GIS = geographic information system

467 GIS = geographic information system, GPS=Global positioning system, IQR= interquartile

468 range, ITN= Insecticide-treated bednet, LLIN=long-lasting insecticide-treated bednet, LOESS=

469 Locally Weighted Scatterplot Smoothing, MoH= Ministry of Health, PDA = personal digital

470 assistant, THSA=total hole surface area, tm=median survival time, WHO= World Health

471 Organization, XRF= x-ray fluorescence

#### 472 DECLARATIONS

##### 473 *Ethics approval and consent to participate*

474 Oral informed consent was obtained from all participants prior to study inclusion. This study  
475 was approved by the Ethics Committee of the Center for Health Studies at Universidad del Valle  
476 de Guatemala (Approval Number: 081-06-2013); CDC investigators were not considered to be  
477 engaged in human subjects research.

##### 478 *Consent for publication*

479 Not applicable

##### 480 *Availability of data and materials*

481 The data that support the findings of this study are available on request from the senior author,  
482 NP. The data are not publicly available due to containing information that could compromise the  
483 privacy of participants.

##### 484 *Competing interests*

485 The authors declare that they have no competing interests

##### 486 *Funding*

487 The funding for this study was provided by the United States Agency for International  
488 Development (USAID) via the Amazon Malaria Initiative (AMI), Centers for Disease Control and

489 Prevention (CDC) of the United States of America, Guatemalan Ministry of Public Health and  
490 Social Welfare and Center for Health Studies and Universidad del Valle de Guatemala. The  
491 funding bodies had no role in the design of the study and collection, analysis, and interpretation  
492 of the data and in writing the manuscript.

#### 493 *Author's contributions*

494 Conception and design: NP, AL, SC, ZM, JVE, SS, MEC; survey data collection, supervision  
495 and coordination: SC, NP, MEC, JCL, JGJ, insecticide content and bio-efficacy assays: SR, LV,  
496 SS; data analysis: MEC, JVE, HV, SR, draft of manuscript: MEC, SR, review and edit of the  
497 manuscript: MEC, SR, JCL, JGJ, ZM, SC, SS, LV, JVE, HW, AL, NP. All authors approved the  
498 final version of the manuscript.

#### 499 *Acknowledgements*

500 We thank the invaluable contribution of the survey team from the National Malaria Sub-program,  
501 Vector-borne Diseases Program and from Universidad del Valle de Guatemala, particularly  
502 Rodrigo Flores, Pedro Peralta, and Alfonso Salam. We thank Gerard Lopez for his help in the  
503 design of the survey and installation in the PDAs. We also thank all the participants in the  
504 survey and all the help provided by the community leaders at La Gomera, Escuintla. We thank  
505 Vestergaard for providing us with LLINs to replace the ones we used for the laboratory  
506 analyses. We are grateful to Barbara Marston and William Hawley for their critical suggestions  
507 and for their careful reading of the manuscript.

#### 508 *Disclaimer*

509 The findings and conclusions in this paper are those of the authors and do not necessarily  
510 represent the official position of the Centers for Disease Control and Prevention.

511

512

## REFERENCES

513

- 514 Agossa, F. R., et al. (2014). Laboratory and field evaluation of the impact of washings on the  
515 effectiveness of LifeNet(R), Olyset(R) and PermaNet(R) 2.0 in two areas, where there is  
516 a high level of resistance of *Anopheles gambiae* to pyrethroids, Benin, West Africa.  
517 *Malar J*, 13, 193. doi:10.1186/1475-2875-13-193
- 518 Althubaiti, A. (2016). Information bias in health research: definition, pitfalls, and adjustment  
519 methods. *J Multidiscip Healthc*, 9, 211-217. doi:10.2147/JMDH.S104807
- 520 Alvarado, G., et al. (2011). Monitoreo de la posesión y uso de mosquiteros tratados con  
521 insecticidas de larga duración en un área endémica de malaria de la Amazonia  
522 Venezolana. *Revista Pan-Amazônica de Saúde*, 2(1), 45-51.
- 523 Anshebo, G. Y., et al. (2014). Estimation of insecticide persistence, biological activity and  
524 mosquito resistance to PermaNet(R) 2 long-lasting insecticidal nets over three to 32  
525 months of use in Ethiopia. *Malar J*, 13, 80. doi:10.1186/1475-2875-13-80
- 526 Atieli, F. K., et al. (2010). Wash durability and optimal drying regimen of four brands of long-  
527 lasting insecticide-treated nets after repeated washing under tropical conditions. *Malar J*,  
528 9, 248. doi:10.1186/1475-2875-9-248
- 529 Cabrera, O. L., et al. (2009). Aceptabilidad y eficacia de mosquiteros tratados con insecticida de  
530 larga duración Olyset® en un área endémica para malaria en Colombia. *Bol Mal Salud*  
531 *Amb*, 49, 241-250.
- 532 Committee, W. H. O. M. P. A., et al. (2013). Malaria Policy Advisory Committee to the WHO:  
533 conclusions and recommendations of September 2013 meeting. *Malar J*, 12, 456.  
534 doi:10.1186/1475-2875-12-456
- 535 Craig, A. S., et al. (2015). Long-lasting insecticidal nets in Zambia: a cross-sectional analysis of  
536 net integrity and insecticide content. *Malar J*, 14, 239. doi:10.1186/s12936-015-0754-8
- 537 Earl-Slater, A. (2002). *The handbook of clinical trials and other research*: Radcliffe Publishing.
- 538 Fettene, M., et al. (2009). Utilization, retention and bio-efficacy studies of PermaNet in selected  
539 villages in Buie and Fentalie districts of Ethiopia. *Malar J*, 8, 114. doi:10.1186/1475-  
540 2875-8-114
- 541 Gimnig, J. (2013). *Assessing the Durability of Long Lasting Insecticidal Nets within PMI*.  
542 Retrieved from (<https://endmalaria.org/sites/default/files>)
- 543 Gnanguenon, V., et al. (2014). Durability assessment results suggest a serviceable life of two,  
544 rather than three, years for the current long-lasting insecticidal (mosquito) net (LLIN)  
545 intervention in Benin. *BMC Infect Dis*, 14, 69. doi:10.1186/1471-2334-14-69
- 546 Gosling, R., et al. (2020). District-level approach for tailoring and targeting interventions: a new  
547 path for malaria control and elimination. *Malar J*, 19(1), 125. doi:10.1186/s12936-020-  
548 03185-w
- 549 Haji, K. A., et al. (2020). Monitoring the durability of the long-lasting insecticidal nets Olyset((R))  
550 and PermaNet((R)) 2.0 in similar use environments in Zanzibar. *Malar J*, 19(1), 187.  
551 doi:10.1186/s12936-020-03258-w
- 552 Hawley, W. A., et al. (2003). Community-wide effects of permethrin-treated bed nets on child  
553 mortality and malaria morbidity in western Kenya. *Am J Trop Med Hyg*, 68(4 Suppl),  
554 121-127.
- 555 Kilian, A., et al. (2011). Evidence for a useful life of more than three years for a polyester-based  
556 long-lasting insecticidal mosquito net in Western Uganda. *Malar J*, 10, 299.  
557 doi:10.1186/1475-2875-10-299

- 558 Koenker, H., et al. (2014). What happens to lost nets: a multi-country analysis of reasons for  
559 LLIN attrition using 14 household surveys in four countries. *Malar J*, 13, 464.  
560 doi:10.1186/1475-2875-13-464
- 561 Lorenz, L. M., et al. (2019). Comparative Functional Survival and Equivalent Annual Cost of  
562 Three Long Lasting Insecticidal Net (LLIN) Products in Tanzania: A Three-Year  
563 Prospective Cohort Study of LLIN Attrition, Physical Integrity, and Insecticidal Activity.  
564 *Physical Integrity, and Insecticidal Activity (June 21, 2019)*.
- 565 Mansiangi, P., et al. (2020). Comparing the durability of the long-lasting insecticidal nets  
566 DawaPlus((R)) 2.0 and DuraNet(c) in northwest Democratic Republic of Congo. *Malar J*,  
567 19(1), 189. doi:10.1186/s12936-020-03262-0
- 568 Morgan, J., et al. (2015). Physical durability of two types of long-lasting insecticidal nets (LLINs)  
569 three years after a mass LLIN distribution campaign in Mozambique, 2008-2011. *Am J*  
570 *Trop Med Hyg*, 92(2), 286-293. doi:10.4269/ajtmh.14-0023
- 571 Mutuku, F. M., et al. (2013). Physical condition and maintenance of mosquito bed nets in Kwale  
572 County, coastal Kenya. *Malar J*, 12, 46. doi:10.1186/1475-2875-12-46
- 573 Obi, E., et al. (2020). Monitoring the physical and insecticidal durability of the long-lasting  
574 insecticidal net DawaPlus((R)) 2.0 in three States in Nigeria. *Malar J*, 19(1), 124.  
575 doi:10.1186/s12936-020-03194-9
- 576 Rosas-Aguirre, A., et al. (2011). [Long-lasting insecticide - treated bednet ownership, retention  
577 and usage one year after their distribution in Loreto, Peru]. *Rev Peru Med Exp Salud*  
578 *Publica*, 28(2), 228-236.
- 579 Smith, S. C., et al. (2007). Evaluation of bednets after 38 months of household use in northwest  
580 Ghana. *Am J Trop Med Hyg*, 77(6 Suppl), 243-248.
- 581 Tan, K. R., et al. (2016). A longitudinal study of the durability of long-lasting insecticidal nets in  
582 Zambia. *Malar J*, 15, 106. doi:10.1186/s12936-016-1154-4
- 583 Vanden Eng, J. L., et al. (2015). Bed Net Durability Assessments: Exploring a Composite  
584 Measure of Net Damage. *PLoS One*, 10(6), e0128499.  
585 doi:10.1371/journal.pone.0128499
- 586 Wampler, P. J., et al. (2013). Using ArcMap, Google Earth, and Global Positioning Systems to  
587 select and locate random households in rural Haiti. *Int J Health Geogr*, 12, 3.  
588 doi:10.1186/1476-072X-12-3
- 589 WHO Malaria Policy Advisory Committee. (2013). Malaria Policy Advisory Committee to the  
590 WHO: conclusions and recommendations of September 2013 meeting: Springer.
- 591 World Health Organization. (2013). *Guidelines for laboratory and field-testing of long-lasting*  
592 *insecticidal nets* (9241505273). Retrieved from
- 593 World Health Organization. (2017). *A framework for malaria elimination*: World Health  
594 Organization.
- 595 World Health Organization. (2018). World malaria report 2018. . *World Health Organization*:  
596 Geneva.
- 597 World Health Organization. (2011). *Guidelines for monitoring the durability of long-lasting*  
598 *insecticidal mosquito nets under operational conditions*. Geneva: World Health  
599 Organization.

600

601

602 Tables Legends

603

604 Table 1. Demographic characteristics of the surveyed households and head household at the  
605 three time-points.

606 Table 2. Long-lasting insecticide-treated bednets (LLINs) use and handling as reported by  
607 householders at each survey time-point.

608 Table 3. Physical integrity of long-lasting insecticide-treated bednets (LLINs), used at least once  
609 for sleeping and with complete data regarding holes.

610

611 Table 3. Demographic characteristics of the surveyed households and head of household at the  
 612 three time-points.

613

Characteristic	18 months n=290	24 months n=349	32 months n=349
<b>Education level head of household</b>			
None (n, %)	84 (29)	79 (23)	99 (28)
Primary school through 3rd grade (n, %)	65 (22)	93 (27)	102 (29)
Primary school through 6th grade (n, %)	88 (30)	99 (28)	92 (26)
Basic level (n, %) ±	24 (8)	36 (10)	29 (8)
Diversified level (n, %) π	16 (6)	30 (9)	20 (6)
Higher education/University (n, %)	4 (1)	1 (0)	6 (2)
Does not know/unclear (n, %)	9 (3)	11(3)	1 (0)
Mother as the respondent (n, %)	216 (74)	269 (77)	270 (77)
Electricity present (n, %)	265 (91)	338 (97)	343 (98)
Own flush toilet (n, %)	75 (26)	99 (28)	187 (54)
Piped water into home (n, %)	77 (27)	77 (22)	147 (42)

614

615 ± Secondary education, first three years

616 π Secondary education, last two or three years

617 Table 4. Long-lasting insecticide-treated bednets (LLINs) use and handling as reported by  
618 householders at each survey time-point.

	18 months n=249	24 months n=266	32 months n=229
LLIN used at least once to sleep (n, %) ±	231 (93)	257 (96)	217 (95)
LLIN used the night before the survey (n, %)	171 (74)	203 (79)	154 (71)
LLIN used every night during the week before the survey (n, %)	189 (82)	208 (81)	167 (77)
LLIN used year-round (n, %)	205 (89)	215 (84)	178 (82)
LLIN tucked under bed at night (n, %)	213 (92)	241 (94)	194 (89)
LLIN observed at time of survey, among LLIN used at least once to sleep			
Hanging loose over sleeping place (n, %)	66 (29)	87 (34)	69 (32)
Hanging tied in knot (n, %)	17 (7)	33 (13)	35 (16)
Hanging folded (n, %)	37 (16)	28 (11)	24 (11)
Visible but not hanging (n, %)	13 (6)	15 (6)	10 (5)
Stored away (n, %)	53 (23)	56 (22)	60 (28)
Hanging next to the wall (n, %)	45 (19)	38 (15)	19 (9)
Washing activities, among LLIN used at least once to sleep			
Ever washed <sup>π</sup> (n, %)	204 (88)	237 (92)	209 (96)
Bed net washed <1 month ago (n, %)	128 (63)	131 (55)	121(58)
When washed, soaked (n, %)	130 (64)	147 (62)	126 (60)
Detergent used (n, %)	148 (72)	189 (80)	163 (78)
When washed, scrubbed hard (n, %)	127(62)	122 (51)	97(46)
Dried outside in the sun (n, %)	143 (70)	163 (69)	126 (60)
Dried outside in the shade (n, %)	54 (26)	70 (30)	75 (36)
Dried inside (n, %)	5 (2)	3 (2)	4 (2)

619 Abbreviations: LLIN= long-lasting insecticide-treated bednets, IQR = Interquartile boundary values (p25,  
620 p75)

621 ± This 'n' serves as the denominator for the subsequent categories of use and position of LLIN observed  
622 at time of survey.

623 π This 'n' serves as the denominator for the subsequent washing categories.

624 Table 3. Physical integrity of long-lasting insecticide-treated bednets (LLINs), used at least once  
 625 for sleeping and with complete data regarding holes.

626

Characteristic	18 months n=222	24 months n=256	32 months n=214
Any holes (n, %)	131 (59)	181 (71)	162 (76)
Any holes repaired (n, %)±	30 (23)	42 (23)	71 (44)
By stitches (n, %)	23 (77)	27 (64)	54 (76)
By knots (n, %)	9 (30)	20(48)	33 (46)
By patches (n, %)	1 (3)	0 (0)	1 (1)
Median total number of holes [IQR]	2 [0,7]	4 [0,15]	5.5 [1,16]
Median total hole area (cm <sup>2</sup> ) [IQR]	2.4 [0,63.8]	8.4 [0,271.5]	34.3 [1.2,327.8]
Categorization based on total hole area (cm <sup>2</sup> )			
<i>Serviceable condition</i> <sup>††</sup> (n, %)	208 (94)	225 (88)	180 (84)
Good condition (n, %)	171 (77)	180 (70)	125 (58)
Damaged condition (n, %)	37 (17)	45 (18)	55 (26)
<i>Too torn condition</i> (n, %)	14 (6)	31 (12)	34 (16)

627 Abbreviations: IQR = Interquartile boundary values (p25, p75)

628 ± This 'n' serves as the denominator for the type of repair categories.

629 <sup>††</sup>Serviceable condition includes long-lasting insecticide-treated bednets with good or damaged condition.

630

631

632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656

## FIGURE LEGENDS

Figure 1. Location of the 12 study communities in La Gomera, Escuintla, Guatemala.

Figure 2. Flow chart of the data collection and number of long-lasting insecticide-treated bednets (LLINs) that were evaluated for each of the measurements. Invalid questionnaires represent questionnaires with incomplete or discrepant data or questionnaires carried out on nets that were not part of the PermaNet 2.0 2012 distribution.

Figure 3. Reasons for long-lasting insecticide-treated bednet (LLIN) loss.


Figure 4. Estimated total deltamethrin concentration ( $\text{mg}/\text{m}^2$ ) measured by X-ray fluorescence (XRF) at the survey time-points. The time-point of 0 represents values on unused long-lasting insecticide-treated bednets (LLINs) that were from the same batch as those distributed. The dashed horizontal line represents the threshold of  $25 \text{ mg}/\text{m}^2$  and the solid horizontal line represents the threshold of  $10 \text{ mg}/\text{m}^2$ .

Figure 5. Locally weighted regression (LOESS) analysis between results of the cone bioassays measuring percent mortality at 24 hours and concentration of deltamethrin ( $\text{mg}/\text{m}^2$ ) as measured by X-ray fluorescence (XRF) analysis for all survey time-points. Each dot represents one long-lasting insecticide-treated bednet (LLIN) plotted based corresponding cone bioassays and XRF measurements. The gray area represents the 95% confidence interval. The solid vertical line represents the threshold of  $10 \text{ mg}/\text{m}^2$  and the dashed vertical line represents the threshold of  $25 \text{ mg}/\text{m}^2$ . The black horizontal line represents the 80% mortality threshold.

657 Figure 6. Estimate of effective protection of long-lasting insecticide-treated bednets (LLINs) in  
658 La Gomera, Escuintla. The proportion of LLINs that met each of the criteria listed on the left is  
659 shown in sequential descending order, with each value shown as a subset of the criteria above.  
660 If the LLIN was able to fulfill all criteria, it was considered a LLIN that provided adequate  
661 protection. We did not include the stage of deltamethrin above 10 mg/m<sup>2</sup> as measured by XRF  
662 for the 18-month survey as we only analyzed a small number of LLINs with XRF at that time-  
663 point.


664 Figure 1.

665



666

667 Figure 2.




668

669

670 Figure 3.

671



672

673

674

675


676

677

678 Figure 4.

679

680




681

682

683 Figure 5.


684



685  
686

687 Figure 6.

688



689

690

691

692 SUPPLEMENTARY MATERIAL

693

694

Table Legends

695

696 Table S1. Additional descriptive characteristics of the households that were enrolled in the  
697 surveys at the three time-points.

698 Table S2. Reasons for LLIN loss.

699 Table S3. Association of deltamethrin ( $\text{mg}/\text{m}^2$ ) content level measured by X-ray fluorescence  
700 (XRF) and age of the long-lasting insecticide-treated bednets (LLINs) in predicting the  
701 percentage of mosquito mortality within 24 hrs.

702 Table S4. Estimate of effective protection of long-lasting insecticide-treated bednets (LLINs) in  
703 La Gomera, Escuintla. Results are shown for a) the overall study population, b) Data excluding  
704 LLINs with incomplete hole section and c) Data excluding LLINs with incomplete hole section  
705 and with no X-ray fluorescence (XRF) analysis. Results in bold are the ones presented in Figure  
706 6.

707

708

709

710 Table S1. Additional descriptive characteristics of the households that were enrolled in the  
711 surveys at the three time-points.

712

Characteristic	18 months n=290	24 months n=349	32 months n=349
Number of people slept in household last night			
Adults > 15 years (median, IQR)	2 [2,4]	2 [2,4]	3 [2,4]
5-15 years (median, IQR)	1 [0,2]	1 [0,2]	1 [0,2]
<5 years (median, IQR)	0 [0,1]	0 [0,1]	0 [0,1]
Total (median, IQR)	4 [3,6]	4 [3,6]	4 [3,5]
Number of sleeping places used last night in your household (median, IQR)	3 [2,4]	3 [2,4]	3 [2,4]
According to census, number of LLINs given by MoH in 2012 (median, IQR)	3 [2,4]	3 [2,5]	3 [2,4]
Number of any bed nets in household that can be used for sleeping (median, IQR)	4 [2,5]	4 [2,5]	3 [2,5]
Number of LLINs in household that can be used for sleeping (median, IQR)	3 [2,4]	3 [2,4]	2 [1,3]

713 Abbreviations: LLIN= long-lasting insecticide-treated bednets, IQR = Interquartile boundary values (p25,  
714 p75)

715

716 Table S2. Reasons for LLIN loss.

	18 months	24 months	32 months
Characteristic	N=290	N=349	N=349
LLINs still present (n, %)	249 (86)	266 (76)	229 (66)
LLINs not present (n, %)	41 (14)	83 (24)	120 (34)
Reason for loss (n, %)			
Net was given away to others (n, %)	23 (56)	45 (54)	39 (32)
Net was damaged and thrown away (n, %)	6 (15)	21 (25)	52 (43)
Net is being used in another location (n, %)	5 (12)	9 (11)	18 (15)
Net was stolen (n, %)	2 (5)	2 (2)	3 (2)
Do not know (n, %)	2 (5)	2 (2)	3 (2)
Net being used for another purpose (n, %)	1 (2)	2 (2)	0 (0)
Allergies (n, %)	1 (2)	0 (0)	3 (2)
Unclear reason (n, %)	1 (2)	2 (2)	2 (2)

717 Abbreviations: LLIN= long-lasting insecticide-treated bednets

718

719

720 Table S3. Association of deltamethrin ( $\text{mg}/\text{m}^2$ ) content level measured by X-ray fluorescence  
721 (XRF) and age of the long-lasting insecticide-treated bednets (LLINs) in predicting the  
722 percentage of mosquito mortality within 24 hrs.

Exposure variable (1st segment)	Estimate (se)	p-value
Deltamethrin ( $\text{mg}/\text{m}^2$ ) in first segment	2.8 (0.3)	<0.001
18 months	27.1 (5.6)	<0.001
24 months	-18.8 (4.3)	<0.001
32 months	-24.9 (4.3)	<0.001

723  
724 Association of Deltamethrin ( $\text{mg}/\text{m}^2$ ) in the 2nd segment was 1.1 (se = 0.5, CI95: 0.2, 2.0)

725

726 Table S4. Estimate of effective protection of long-lasting insecticide-treated bednets (LLINs) in  
 727 La Gomera, Escuintla. Results are shown for a) the overall study population, b) Data excluding  
 728 LLINs with incomplete hole section and c) Data excluding LLINs with incomplete hole section  
 729 and with no X-ray fluorescence (XRF) analysis. Results in bold are the ones presented in Figure  
 730 6.

Criterion	18 months	24 months	32 months
Overall data			
Evaluated $\pm$ (n, %)	<b>290 (100)</b>	<b>349 (100)</b>	<b>349 (100)</b>
Still present (n, %)	<b>249 (86)</b>	<b>266 (76)</b>	<b>229 (66)</b>
Used at least once (n, %)	<b>231 (80)</b>	<b>257 (74)</b>	<b>217 (62)</b>
Used last night (n, %)	<b>171 (59)</b>	<b>203 (58)</b>	<b>154 (44)</b>
Data excluding LLINs with incomplete hole section			
Evaluated $\pi$ (n, %)	281 (100)	348 (100)	346 (100)
Still present (n, %)	240 (85)	266 (76)	226 (65)
Used at least once (n, %)	222 (79)	256 (74)	214 (62)
Used last night (n, %)	164 (58)	203 (58)	152 (44)
Serviceable condition (based on total hole area) (n, %)	<b>151 (54)</b>	<b>174 (50)</b>	<b>126 (36)</b>
Data excluding LLINs with incomplete hole section and with no XRF analysis			
Evaluated $\pi\pi$ (n, %)	91 (100)	346 (100)	335 (100)
Still present (n, %)	NE	263 (76)	215 (64)
Used at least once (n, %)	NE	254 (73)	204 (61)
Used last night (n, %)	NE	201 (58)	145 (43)
Serviceable condition (based on total hole area) (n, %)	NE	172 (50)	121 (36)
Over 10 mg/m <sup>2</sup> (based on XRF analysis) (n, %)	NE	<b>133 (38)</b>	<b>72 (21)</b>

731 Abbreviations: LLIN= long-lasting insecticide-treated bednets, XRF = X-ray fluorescence

732  $\pm$  This 'n' serves as the denominator for the overall data analysis.

733  $\pi$  This 'n' serves as the denominator for the data excluding LLINs with incomplete hole section

734  $\pi\pi$  This 'n' serves as the denominator for the data excluding LLINs with incomplete hole section and with  
 735 no XRF measurement.

736 NE =Not estimated because only a small proportion of LLINs were analyzed by XRF analysis, so this  
 737 measurement is likely to be biased.

738 SUPPLEMENTARY MATERIAL

739 Figure Legends

740 Figure S1. Mortality at 24 hours of *An. albimanus* (Sanarate strain) mosquitoes exposed in cone  
741 bioassays to long-lasting insecticide-treated bednets (LLINs) at the surveys time-points. The  
742 time-point of 0 represents values on unused LLINs that were from the same batch as those  
743 distributed. Each box represents the interquartile distance, center line in each box indicates the  
744 median. The black dots indicate each of the values of a LLIN. The dashed horizontal line  
745 represents the 80% mortality threshold.

746 Figure S2. Knockdown after a 60-min (KD60) exposure of *An. albimanus* (Sanarate strain)  
747 mosquitoes exposed in cone bioassays to long-lasting insecticide-treated bednets (LLINs) at the  
748 surveys time-points. The time-point of 0 represents values on unused LLINs that were from the  
749 same batch as those distributed. Each box represents the interquartile distance, center line in  
750 each box indicates the median. The black dots indicate each of the values of a LLIN. The  
751 shaded horizontal line represents the 95% KD60 threshold.

752 Figure S3. Locally weighted regression (LOESS) analysis between results of the cone  
753 bioassays measuring percent mortality at 24 hours and concentration of deltamethrin ( $\text{mg}/\text{m}^2$ )  
754 as measured by X-ray fluorescence (XRF), stratified by surveys time-points (18 months, 24  
755 months an 32 months). Each dot represents one long-lasting insecticide-treated bednets  
756 (LLINs) with their corresponding cone bioasays and XRF measurements. The gray area  
757 represents 95% confidence interval. The solid vertical line represents the threshold of  $10 \text{ mg}/\text{m}^2$   
758 and the dashed vertical line represents the threshold of  $25 \text{ mg}/\text{m}^2$ . The black horizontal line  
759 represents the 80% mortality threshold.


760

761

762 Figure S1. Mortality at 24 hours of *An. albimanus* (Sanarate strain) mosquitoes exposed in cone  
763 bioassays to long-lasting insecticide-treated bednets (LLINs) at the surveys time-points. The  
764 time-point of 0 represents values on unused LLINs that were from the same batch as those  
765 distributed. Each box represents the interquartile distance, center line in each box indicates the  
766 median. The black dots indicate each of the values of a LLIN. The dashed horizontal line  
767 represents the 80% mortality threshold.

768

769



770


771

772

773

774 Figure S2. Knockdown after a 60-min (KD60) exposure of *An. albimanus* (Sanarate strain)  
775 mosquitoes exposed in cone bioassays to long-lasting insecticide-treated bednets (LLINs) at the  
776 surveys time-points. The time-point of 0 represents values on unused LLINs that were from the  
777 same batch as those distributed. Each box represents the interquartile distance, center line in  
778 each box indicates the median. The black dots indicate each of the values of a LLIN. The  
779 shaded horizontal line represents the 95% KD60 threshold.

780



781

782

783

784

785

786

787


788

789

790 Figure S3. Locally weighted regression (LOESS) analysis between results of the cone  
791 bioassays measuring percent mortality at 24 hours and concentration of deltamethrin ( $\text{mg}/\text{m}^2$ )  
792 as measured by X-ray fluorescence (XRF), stratified by surveys time-points (18 months, 24  
793 months and 32 months). Each dot represents one long-lasting insecticide-treated bednets  
794 (LLINs) with their corresponding cone bioassays and XRF measurements. The gray area  
795 represents 95% confidence interval. The solid vertical line represents the threshold of  $10 \text{ mg}/\text{m}^2$   
796 and the dashed vertical line represents the threshold of  $25 \text{ mg}/\text{m}^2$ . The black horizontal line  
797 represents the 80% mortality threshold.

798

799



800

801

802 APPENDIX FILE

803 **Methodology to assess if insecticide concent measurement obtained by XRF could serve**  
804 **as a predictor to estimate the mortality at 24 hours.**

805

- 806 • We developed a predictive model using piecewise (segmented) regression to  
807 accommodate the change in the magnitude and direction of the linear relationship  
808 between the outcome and exposure variables observed in our exploratory analysis.

809 We use selected responses from the surveys as exposure variables:

- 810 – Months after distribution of LLIN
- 811 – Highest level of education of the head of the household
- 812 – Net still present in the household
- 813 – Net used the night before of the survey to sleep under
- 814 – Frequency of the net used in the week before of the survey to sleep under
- 815 – Adult (>15 years) slept under the net the night before of the survey
- 816 – Children (5-15 years) slept under the net the night before of the survey
- 817 – Children (<15 years) slept under the net the night before of the survey
- 818 – Net has been washed
- 819 – Last time the net was washed
- 820 – Type of soap used to wash the net
- 821 – Net was soaked when washing
- 822 – Length of time a net was soaked
- 823 – Net was scrubbed when washing
- 824 – Location were net was dried
- 825 – Presence of open flame were the net was located
- 826 – Total area of holes of LLIN

827 – Condition of LLIN based on holes area (Good, damaged or too torn)

828 – Deltamethin content ( $\text{mg}/\text{m}^2$ ) measured by XRF

829

830 • Outcome variable: % Mortality at 24 hours

831 • The model was first trained using 80% of the data and validated in the remaining 20%.

832 We used random sampling to separate the data as training and test data. This process

833 was repeated 1000 times. To assess the accuracy of the fitted model, a confusion matrix

834 was built to calculate the proportion of correct classification when the model was applied

835 to the test data

836

837

838

839