Gait and Balance Function Improves After 10 Weeks of Using a Wearable Sensory Neuroprosthesis in Persons with Peripheral Neuropathy and High Fall Risk – the walk2Wellness Trial

Lars IE Oddsson[1,2,3], Teresa Bisson[2,4], Helen S Cohen[5], Laura Jacobs[1], Mohammad Khoshnoodi[11], Doris Kung[5], Lewis A Lipsitz[6,7,8], Brad Manor[6,7,8], Patricia McCracken[9], Yvonne Rumsey[1], Sara R Koehler-McNicholas[2,9], Diane M Wrisley[10].

[1] RxFunction Inc. MN, USA;
[2] University of Minnesota, MN, USA;
[3] Ben Gurion University of the Negev, Israel;
[4] M Health Fairview, Minneapolis, MN, USA;
[5] Baylor College of Medicine, Houston, TX, USA;
[6] Hinda and Arthur Marcus Institute for Aging Research, Hebrew SeniorLife, Roslindale, MA, USA.
[7] Division of Gerontology, Beth Israel Deaconess Medical Center, Boston, MA, USA;
[8] Harvard Medical School, Boston, MA, USA;
[9] Minneapolis Department of Veterans Affairs Health Care System, Minneapolis, MN, USA;
[10] College of St Mary, Omaha, NE, USA;
[11] Johns Hopkins University School of Medicine, Baltimore, MD, USA;

Abstract

Background: Patients with sensory peripheral neuropathy (PN) commonly present with gait and balance problems increasing their risk of falls. The multi-site walk2Wellness trial investigates effects of long-term, home-based daily use of a wearable lower limb sensory neuroprosthesis on gait function, balance, quality of life and fall rates in a cohort of PN patients. The device (Walkasins®, RxFunction Inc., MN, USA) is designed to replace lost nerve function related to foot pressure sensation by providing directional tactile cues around the ankle reflecting foot pressure measurements during standing and walking activities. We hypothesized that previously shown short-term in-clinic improvements in gait and balance would be sustained after 10 weeks of use.

Methods: Participants had a PN diagnosis with loss of plantar sensation associated with gait and balance problems, a Functional Gait Assessment (FGA) score <23 (cut-off for high fall risk), and ability to sense Leg Unit tactile stimuli. Clinical outcomes included FGA, Gait Speed, Timed Up&Go (TUG) and Four-Stage Balance Test. Patient-reported outcomes included Activities-Specific Balance Confidence (ABC) scale, Vestibular Disorders Activities of Daily Living Scale (VADL), PROMIS participation and satisfaction scores, pain rating, and falls. Evaluations were performed at baseline visit and after 2, 6, and 10 weeks. Subjects were not made aware of any changes in outcomes and no additional balance interventions were allowed.

Results: Forty-five participants of 52 enrolled across four sites completed all in-clinic assessments. FGA scores improved from 15.0 at baseline to 19.1 at 10 weeks (p<0.000001), normal and fast gait speed from 0.86m/s to 0.95m/s (p<0.00005) and 1.24m/s to 1.33m/s (p<0.002), respectively, and TUG from 13.8s to 12.5s (p<0.012). Four-Stage Balance Test did not improve significantly. Several patient-reported outcomes were in normal range at baseline and remained largely unchanged. Interestingly, while FGA scores improved similarly across the baseline range, subjects with baseline ABC scores lower than 67% (cut-off for high fall risk) showed an increase in their ABC scores (from 49.9% to 59.3%, p<0.01), whereas subjects with baseline ABC scores above 67% showed a decrease (from 76.6% to 71.8%, p<0.019). Subjects who reported falls in the prior six months (n=25) showed a decrease in the number of fall-risk factors (from 5.1 to 4.3, p<0.023) as well as a decrease in fall rate from 13.8 to 7.4 falls/1000 days (p<0.014). Four subjects in the pre-study non-faller group (n=20) fell during the 10 weeks of the study.

Conclusion: A wearable sensory neuroprosthesis may provide a new way to treat gait and balance problems and manage falls in high fall-risk patients with PN.

Trial registration: ClinicalTrials.gov (#NCT03538756)

Keywords: Peripheral Neuropathy, Falls, Neuroprosthesis, Balance, Gait Speed, Function, Neuromodulation, Clinical Trial

1   Introduction

Falls are a widely recognized problem in the elderly and commonly related to problems with balance (Ganz and Latham, 2020). About 29% of community-dwelling adults 65 years or older fall once annually and 10% fall at least twice annually (Ganz et al., 2007; Bergen et al., 2016). Data from the Centers for Disease Control (CDC) indicate that medical treatment was required by 37.5% of individuals who fell in 2014 (Bergen et al., 2016). Sterling et al. (Sterling et al., 2001) reported that 30% of falls in the elderly result in serious injury. In 2015, medical cost related to fatal and nonfatal falls was approximately $50.0 billion (Florence et al., 2018). Overall, falls are associated with poor health, shortened survival (Jónsdóttir and Ruthig, 2020), reduced quality of life, and a fear of falling (Lawrence et al., 1998; Scheffer et al., 2008). Sensory peripheral neuropathy (PN) is associated with poor balance and is an independent risk factor for falls (Richardson and Hurvitz, 1995), regardless if the etiology is idiopathic (Riskowski et al., 2012), due to diabetes (Mustapa et al., 2016; Vinik et al., 2017), or chemotherapy (Winters-Stone et al., 2017). The prevalence of PN in the US population for those over age 40 has been reported to be nearly 15% (Gregg et al., 2004). The importance of sensory information from plantar cutaneous mechanoreceptors for balance control has been shown in healthy individuals (Meyer et al., 2004a; b), with loss of such information in patients with PN likely leading to problems with gait and balance function and increased risk of falls (Menz et al., 2004; DeMott et al., 2007; Dixon et al., 2017; Lipsitz et al., 2018). The occurrence of fall-related injuries is up to 15 times higher in patients with diabetic PN than in healthy individuals (Cavanagh et al., 1992). Furthermore, the prevalence of polyneuropathy has been reported to be almost four times higher in persons older than 60 years and to independently contribute to functional impairments including difficulty walking and tendency to fall (Hoffman et al., 2015). Persons with polyneuropathy are more likely to fall and more often incur fall-related injuries (Hanewinckel et al., 2017). In a prospective study, 65% of older individuals with PN fell during a one-year period and 30% reported an injury from a fall (DeMott et al., 2007). In addition, low gait speed is a risk factor for falls (Studenski et al., 2003; Montero-Odasso et al., 2005), an important indicator of frailty (Kim et al., 2019) and a predictor of survival (Studenski et al., 2011). Although gait speed declines with healthy aging (Buracchio et al., 2010), the decline in individuals with progressive sensory loss may be four times as high (Lipsitz et al., 2018). Interestingly, interventions designed to improve gait speed may also increase survival (Hardy et al., 2007).

Clinical treatment of gait and balance problems related to PN is mainly limited to the use of canes, walkers, physical therapy interventions and balance exercises (Richardson et al., 2001; Ganz and Latham, 2020) including Tai-Chi (Li and Manor, 2010; Manor et al., 2014; Quigley et al., 2014). Long-term use of bilateral ankle foot orthoses in elderly individuals with a history of falls showed positive changes in certain in-clinic static sway measures (Wang et al., 2019a), although long-term benefits related to fall rates and gait function appear limited (Wang et al., 2019b).

Several review studies support the hypothesis that strength and balance training interventions can improve balance and reduce fall risk and falls in patients with PN (Ites et al., 2011; Tofthagen et al., 2012; Streckmann et al., 2014). The training, however, should be specific to balance (Bulat et al., 2007; Oddsson et al., 2007; Halvarsson et al., 2011; Akbari et al., 2012) because strength and/or endurance training in patients with PN appears to have less impact on balance (Streckmann et al., 2014). In addition, unless balance activities, including Tai Chi or balance therapies are conducted with sufficient intensity, frequency (Lipsitz et al., 2019) and specificity, benefits may be limited or absent (Kruse et al., 2010; Lipsitz et al., 2018; Lipsitz et al., 2019) leading to mixed outcomes. Furthermore, continued exercise is required to maintain benefits long-term (Wolf et al., 2001;

Halvarsson et al., 2013; Melzer and Oddsson, 2013), although some improvements last up to six
months (Allet et al., 2010). Guidelines regarding physical activity for older adults with mobility
problems recommend a minimum of activity at least twice a week (Chodzko-Zajko et al., 2009).
Some studies on patients with diabetic PN following a period of balance training 2-3 times /week
over 6-12 weeks did show improved balance and reduced fall risk (Morrison et al., 2010; Morrison et
al., 2018). However, there currently are no specific guidelines regarding frequency of balance
exercises and even three times a week may be insufficient to see an improvement in balance function
(Kruse et al., 2010). Consequently, there is a need for additional solutions to help improve gait and
balance function in patients with PN.
109
A growing body of literature on various sensory substitution and augmentation technologies suggest
novel ways of improving gait and balance function in different populations of patients. The concept
of sensory substitution related to brain plasticity was laid out by Bach-y-Rita and colleagues, initially
for vision and the vestibular system (Bach-y-Rita et al., 1969; Bach-y-Rita, 2004) and other sensory
systems (Tyler et al., 2003; Bach-y-Rita, 2004). Recent efforts include wearable systems showing
benefits to patients with vestibular loss (Hegeman et al., 2005; Wall et al., 2009; Basta et al., 2011;
Yamanaka et al., 2016), PN (Wall et al., 2012; Wrisley et al., 2018) and Parkinson's Disease (Rossi-
Izquierdo et al., 2013; Lee et al., 2015). Combining wearable neurostimulation with balance therapy
has shown benefits in patients with multiple sclerosis (Leonard et al., 2017), cerebellar ataxia (Cakrt
et al., 2012), stroke (Badke et al., 2011) traumatic brain injury (Ptito et al., 2020) and in-home
balance therapy (Bao et al., 2018)
121
In a randomized crossover trial, a recent study further supported findings from an earlier pilot study
(Wall et al., 2012) and demonstrated meaningful short-term, in-clinic improvements in Functional
Gait Assessment (FGA) scores and gait speed in subjects with PN using a wearable sensory
neuroprosthesis (Koehler-McNicholas et al., 2019). The device (Walkasins®, RxFunction Inc., MN,
USA, Figure 1) is an external lower limb sensory prosthesis designed to replace lost nerve function
used for detection and signaling of foot pressure sensation in patients with PN. It provides gentle
directional tactile stimuli (in the form of low-intensity vibrations) around the lower leg that reflect
changes in foot pressure distribution measured with an instrumented Foot Pad in the shoe. The
subject's nervous system senses these new tactile cues and incorporates them to improve gait and
balance. Currently, effects of long-term daily use of Walkasins on clinical outcomes are unknown.
The multi-site clinical trial, walk2Wellness, (NCT #03538756, www.clinicaltrials.gov) investigates
long-term, home-based use of Walkasins on clinical and patient-reported outcomes of balance and
gait function, quality of life, physical activity, social participation, pain and fall rates. Data from the
primary endpoint of the study at 10 weeks are reported here. We hypothesized that previously
demonstrated short-term in-clinic improvements in FGA score by at least four points would be
sustained long-term (Beninato et al., 2014; Koehler-McNicholas et al., 2019). Early data from the
trial were presented in abstract form (Oddsson et al., 2019; Oddsson et al., 2020).
139

2   Materials and Methods

2.1   Subject Recruitment

Human subject testing was approved according to the Declaration of Helsinki by Advarra IRB
(formerly Quorum Review IRB), serving as the Institutional Review Board (IRB) of record for three
of the participating sites under the study protocol for *walk2Wellness: Long-term Use Effects of*
Walkasins® Wearable Sensory Prosthesis on Gait Function, Balance-Confidence, and Social
Participation. The three sites include Baylor College of Medicine, Houston, TX; Hebrew SeniorLife,
a Harvard Medical School Affiliate, Boston, MA; and M Health Fairview, Minneapolis, MN.
Advarra IRB determined that Walkasins are a non-significant risk device because they do not meet
the criteria of a significant risk device according to U.S. Food and Drug Administration regulations.
The IRB Subcommittee, the Subcommittee on Research Safety, and the Research and Development
Committee of the Minneapolis VA Health Care System (MVAHCS) also approved the trial. The
study is registered on ClinicalTrials.gov (#NCT03538756). At the time this study began, Walkasins
were available only for research purposes.

2.2   Inclusion and Exclusion Criteria

Inclusion criteria for the study were similar to Koehler-McNicholas et al. (Koehler-McNicholas et al.,
2019): age 21-90 years; male or female; a formal diagnosis of sensory PN prior to participating in the
study as indicated by subject's medical record or a signed letter by a physician; self-reported
problems with balance; ability for transfers or ambulation on level surfaces at fixed cadence as
assessed by trained study personnel; an Functional Gait Assessment (FGA) score <23, the cut-off
score for high fall risk (Wrisley and Kumar, 2010); ability to understand and provide informed
consent; foot size to allow the Walkasins device to function properly, and ability to complete all
functional outcome measures without the use of an assistive device to ensure sufficient motor
function. Subjects could use an assistive device at their own discretion during the trial. Subjects were
excluded from participation if they were unable to perceive tactile stimuli from the Walkasins leg
unit or used an ankle-foot orthosis for ambulation that prevented donning of the device. Subjects with
any of the following conditions were also excluded: acute thrombophlebitis; deep vein thrombosis;
untreated lymphedema; a lesion of any kind, swelling, infection, inflamed area of skin, or eruptions
on the lower leg near placement of the device; foot or ankle fractures; or severe peripheral vascular
disease. In addition, subjects with any musculoskeletal or other neurological conditions that would
prohibit use of the device, as determined by a clinician, were excluded. Due to risk of overloading the
pressure sensor Foot Pad, subjects weighing over 136 kg (300lbs) were excluded from participation.
Furthermore, subjects were prohibited from initiating any balance training (e.g., Tai-Chi etc.) or
balance-related therapy during the ten weeks of the trial. Subjects were not systematically provided
information about changes in any outcomes scores or changes in their performance throughout the 10
weeks of the trial, nor did study personnel monitor outcomes during the study. Potential subjects
responded to announcements that specifically targeted individuals with PN and balance problems, or
they were referred by clinicians who were familiar with the study and believed them to be good
candidates for the trial.

2.3   Study Procedures

All participants signed IRB-approved consent forms prior to the initiation of study activities.
Following the informed consent process, a study team member tested the subjects on both legs to
determine whether they could feel the stimuli from the Walkasins Leg Unit (Figure 1). Subjects wore

Walkasins on both feet. A small number of subjects who were unable to perceive the stimuli from the
Walkasins Leg Unit (Figure 1) were excluded from participation in the study. Participants then
completed a demographics and health screening questionnaire to assess common health issues related
to neurological, musculoskeletal, cardiopulmonary disorders, and other systemic diseases along with
information on their history of falls over the past six and twelve months and regular use of an
assistive device (Koehler-McNicholas et al., 2019). Falls were defined according to the World Health
Organization: "an event which results in a person coming to rest inadvertently on the ground or floor
or other lower level". Subjects enrolled in the study were instructed to wear the device as much as
possible throughout their daily activities, indoors as well as outdoors. At each follow-up visit,
participants were asked about changes in their health status and any falls and adverse events they
experienced since the previous visit. During the baseline visit, participants also provided a list of
their medications (medication name, indication, dose, and frequency), which was updated over the
course of the study. Medications are a known risk factor for falling, based on side effects of
medication use or drug interactions (Woolcott et al., 2009).

2.4 Initial Assessments

Subjects then completed the Activities-Specific Balance Confidence (ABC) Questionnaire, which
measures levels of balance confidence in elderly persons. The ABC asks the question "How
confident are you that you will not lose your balance or become unsteady" when performing 16
different tasks (Powell and Myers, 1995). Subjects rated themselves on a scale from 0 to 100, and an
average score was calculated across the 16 responses. An ABC score below 67% has been associated
with high fall risk (Lajoie and Gallagher, 2004). In addition, subjects completed the Vestibular
Activities of Daily Living Scale (VADL) (Cohen et al., 2000), which evaluates self-reported effects
of vertigo and balance disorders on independence in everyday activities of daily living that are
relevant for individuals living in the community. Individuals rate their level of functional ability for
basic and instrumental activities of daily living on a scale from 1 (independent) to 10 (dependent),
which incorporates the use of assistive devices.
209
Following completion of the questionnaires, a study team member performed tactile and vibration
sensation testing to document loss of sensation. Loss of sensation was tested with the Weinstein
Enhanced Sensory Test (WEST) monofilament foot test (0.5g, 2g, 10g, 50g, and 200g) applied
perpendicular to the skin at four test sites on the plantar surface of the foot, including the first, third,
and fifth metatarsal heads as well as the great toe. Study personnel began testing with the 10g
filament and used a smaller filament if the subject was sensate and a larger filament if the subject was
insensate. The smallest filament the subject was able to feel was recorded (if none were felt this
result was recorded as "none"). Vibration sensation was assessed with a Rydel-Seiffer tuning fork,
which is a 128Hz tuning fork with end weights that convert the tuning fork from 128 to 64 Hz. The
weights are scaled allowing a score 0-8 (lower scores indicating less sensation), allowing reliable
quantitative vibratory testing. Scores were read from the black triangle and rounded to the nearest
whole number. Vibration values ≤ 4 are categorized as abnormal at the first metatarsal joint
(Kästenbauer et al., 2004). The tuning fork was applied firmly and perpendicular to the lateral aspect
of the first metatarsophalangeal, lateral malleolus, and patella testing sites. The monofilament and
vibration tests were repeated at the 10-week visit.

2.5 Clinical Outcome Measures

Upon completion of the monofilament and vibration sensation testing, subjects performed a series of
functional outcome measures while wearing the device turned off (baseline). Tests were repeated at

weeks 2, 6, and 10. Subjects could rest as needed during the clinical assessments. For study purposes
the clinical outcomes were standardized and performed by study personnel who were trained by one
of the investigators (DW). Observation visits were conducted periodically during the study to ensure
standardization among the sites.

Functional Gait Assessment (FGA): The FGA (Wrisley et al., 2004) is the recommended clinical
outcome measure for walking balance based on current physical therapy Clinical Practice Guidelines
for outcome measures for adults with neurologic conditions (Moore et al., 2018). It is a reliable and
valid measure of gait function related to postural stability and has been shown to be effective in
classifying fall risk in older adults and predicting unexplained falls in community-dwelling older
adults (scores ≤22/30) (Wrisley et al., 2004; Wrisley and Kumar, 2010). It has also been validated in
multiple neurological conditions (stroke, Parkinson's, vestibular conditions) (Lin et al., 2010; Leddy
et al., 2011) and has less floor and ceiling effects than the similar Dynamic Gait Index (Lin et al.,
2010). The FGA includes 10 different items that challenge gait balance where each item is scored
from 0 to 3 (3 = normal, 2 = mild impairment, 1 = moderate impairment, 0 = severe impairment) with
a maximum score of 30. An increase of ≥4 points is considered the Minimal Clinically Important
Difference (MCID) for community-dwelling elderly individuals (Beninato et al., 2014). Subjects
whose baseline FGA score was 23 or higher were excluded from further participation in the study.
Subjects completed the FGA in a large open area with a 6-m (20-ft) walkway marked with tape
according to published recommendations (Wrisley et al., 2004).

10-Meter Walk Test (10MWT): The 10m-walk (Perera et al., 2006) is the recommended clinical
outcome measure for walking speed based on current physical therapy Clinical Practice Guidelines
for outcome measures for adults with neurologic conditions (Moore et al., 2018). It is routinely used
in rehabilitation and has excellent reliability in multiple neurologic conditions (chronic stroke,
traumatic brain injury, Parkinson's) (Hiengkaew et al., 2012). Gait speed has been found to be an
important predictor of survival in older adults (Hardy et al., 2007), further emphasizing its
importance as a clinical outcomes measure. Gait speed was assessed during the middle 6 meters of a
10-meter-long pathway to allow for acceleration and deceleration, completed in one trial under two
conditions: 1) walk at normal speed and 2) walk as fast as possible. An increase by 0.05 m/s is
deemed "small meaningful" and 0.10 m/s as "substantial" (Perera et al., 2006). These are considered
the MCID in the geriatric population (Perera et al., 2006).

Timed Up and Go (TUG): The TUG (Mathias et al., 1986) is part of the CDC recommended
STEADI test protocol for balance function (CDC, 2017). It is commonly used in rehabilitation and
has excellent validity and reliability for elderly adults and has been shown to be effective in
classifying community dwelling adults at risk for falls (Podsiadlo and Richardson, 1991; Shumway-
Cook et al., 2000; Bischoff et al., 2003; CDC, 2017). From a seated position in a standard armchair,
the subject is asked to do the following: 1) stand up from the chair, 2) walk at normal pace around a
tape mark on the floor 10 feet from the chair, 3) turn, 4) walk back to the chair at a normal pace, and
5) sit down again. Subjects were provided one practice trial that was not recorded followed by the
recorded timed trial. The tester recorded the time from the command "Go" until the subject's
buttocks returned to the chair when sitting. We used >12s as a cut-off for high fall risk (Bischoff et
al., 2003; CDC, 2017). The Minimal Detectable Change (MDC) for older adults with type 2 diabetes
has been reported to be 1s (Alfonso-Rosa et al., 2014).

4-Stage Balance Test: The 4-Stage Balance Test is part of the Centers for Disease Control and
Prevention (CDC)-recommended STEADI test protocol for balance function (CDC, 2017). It

includes four gradually more challenging postures the subject is exposed to: 1) stand with feet side by
side, 2) stand with feet in semi-tandem stance, 3) stand with feet in tandem stance, and 4) stand on
one leg. Subjects were allowed upper extremity support to obtain the position and passed each level
if they were able to hold the stance unsupported for 10 seconds. The assessment ended when subjects
were unable to hold a stance for 10 seconds. The times for each position held was recorded and
summed as a measure of overall performance. A fail of stances 1, 2, or 3 (i.e., total time < 30s)
indicates high risk of falling (CDC, 2017).

2.6   Learning Protocol

As part of the baseline visit, subjects performed a standardized set of balance activities, once while
wearing the device turned off and once while wearing it turned on (Koehler-McNicholas et al., 2019).
Activities lasted approximately 10 minutes and included standing (two-leg standing, tandem
standing, and one-legged standing) and walking (walking straight, turning right and left) at both
normal and fast speeds. Activities were repeated with the eyes closed. During standing exercises
subjects were challenged to explore their base of support in both mediolateral and anteroposterior
directions and to notice the pattern of tactile stimuli when the device was turned on. During walking
activities subjects were instructed to notice the pattern of tactile stimuli when the foot was in contact
with the ground, how it matched their pace of walking, and the flow of tactile stimuli from step to
step. Subjects were not instructed how to respond to the tactile stimuli from Walkasins instead, the
activities focused on orientation and familiarization with the device.

2.7   Participant Reported Outcomes

At the baseline visit and at each follow-up visit at weeks 2, 6, and 10, subjects also completed the
five subject-reported outcome measures described below:
298
299 **Patient Health Questionnaire (PHQ-9):** The PHQ-9 (Kroenke et al., 2001) is a concise, self-
300 administered tool for assessing depression. Commonly used for screening and diagnosis of
301 depression, the PHQ-9 incorporates depression criteria according to the 4th edition of the Diagnostic
302 and Statistical Manual of Mental Disorders (DSM-IV) with other leading major depressive
303 symptoms.
304
305 **PROMIS Pain Interference Short Form 6b** (Askew et al., 2016): The PROMIS Pain Interference
306 instrument measures the self-reported impact of pain on relevant aspects of a person's life within the
307 past seven days. Items capture the extent to which pain hinders social, cognitive, emotional, physical,
308 and recreational activities. The Pain Interference short form is a global scale rather than disease
309 specific.
310
311 **PROMIS Numeric Rating Scale v1.0 - Pain Intensity Form 1a**: The PROMIS Pain Intensity
312 instrument assesses reported average pain intensity on a scale from 0-10 with higher scores indicating
313 greater levels of pains. The Pain Intensity short form is global rather than disease specific.
314
315 **PROMIS Ability to Participate Short Form 8a**: The PROMIS Ability to Participate in Social
316 Roles and Activities instrument (Hahn et al., 2016b) assesses the individual's perceived ability to
317 perform usual social roles and activities. The measure does not use a designated time frame (e.g. over
318 the past seven days), and higher scores represent fewer limitations (e.g., I have trouble doing all of
319 my regular leisure activities with others).
320

PROMIS Satisfaction with Participation in Social Roles Short Form 8a: The PROMIS
Satisfaction with Social Roles and Activities (Hahn et al., 2014; Hahn et al., 2016a) is a self-reported
instrument to assess satisfaction with the ability to perform usual social roles and activities (e.g., "I
am satisfied with my ability to do things for my family").
PROMIS scores are presented as T-scores, a standardized score with a mean of 50 (representing
average for the US population) and a standard deviation of 10.
327
User Experience Survey: At the 2 and 10-week visits, subjects completed a 10-question survey to
collect information concerning their experience with the device. Subjects rated aspects of Walkasins
use (e.g., donning and doffing, charging, etc.) on a 7-point Likert scale ranging from "Very Easy" to
"Very Hard". Subjects also rated their overall satisfaction with the device and were able to provide
additional comments and feedback regarding their experience with the device. Between visits,
subjects were asked to document their use of Walkasins on a calendar by marking the days they wore
their Walkasins and for how many hours. The subject calendar was also used to facilitate
documentation of falls that occurred over the course of the study (Hannan et al., 2010). Subjects were
asked to return their calendars at their next study visit.

2.8   Number of Subjects

Sample size estimation was based on data from the recent study of a similar population of subjects
(Koehler-McNicholas et al., 2019), showing a baseline average FGA score of 15.2 and standard
deviation 4.8. The data was normally distributed according to the Shapiro-Wilk's test. To detect a
mean difference in pre- and post-FGA score ≥4 points, the Minimal Clinically Important Difference
for community dwelling elderly individuals (Beninato et al., 2014), required at least 20 subjects using
a significance level of 0.01 and a power of 0.8. Accounting for an expected ~20% drop-out rate
(National Heart, 2020) target enrollment was set at 25 subjects (20/0.8=25). Multiple sites were
engaged in the trial to expand geographical, ethnical, and clinical variation in the data initially
allowing each site to recruit up to 25 subjects. Due to the COVID-19 pandemic, the trial was
interrupted and continued enrollment as well as in-clinic testing was halted. At this time, sufficient
overall statistical power based on the sample size calculation above has been achieved following
enrollment of 52 subjects, well over the 20 subjects required to achieve statistical significance.
Collection of participant-reported outcomes has continued through phone calls and the longer-term
outcomes assessments as originally planned (at 26 and 52 weeks) are expected to continue.

2.9   Statistical Analysis and Availability of Data

Descriptive statistics were calculated and presented as mean and standard deviation of the mean.
Variables were tested for normality using the Shapiro-Wilk's test. The two-proportion Z-test was
used to compare proportion-based measures. Subjects who reported falls in the previous six months
(Pre-Fallers, n=30) were analyzed separately from the remaining subjects (Pre-NonFallers, n=22).
Comparisons of baseline characteristics between Pre-Fallers and Pre-NonFallers were made with a t-
test for independent samples or a Mann-Whitney U test if data was not normally distributed based on
a Shapiro-Wilk's test. Repeated measures analysis of variance (ANOVA) was performed for
outcomes measured across the four assessment events, baseline, 2, 6 and 10 weeks. If the ANOVA
was significant ($p<0.05$), three pairwise comparisons were made using dependent t-tests between the
baseline assessments and each of the 2, 6 and 10-week assessments. If the ANOVA was non-
significant, no further comparisons were made. A Bonferroni's adjustment of significance levels for
correlated measures was applied, ranging from $p<0.0167$ (0.05/3 for three comparisons) for a full
correction (non-correlated measures, $r=0$) and $p<0.05$ for perfectly correlated measures ($r=1$)

(Uitenbroek, 1997). Effect sizes were calculated using Cohen's d (Lakens, 2013) and were
interpreted according to recommendations by Cohen (Cohen, 1988) with 0.2 representing a small
effect, 0.5 a medium effect, and 0.8 a large effect. Ninety-five percent confidence intervals of effect
sizes were estimated according to Algina (Algina et al., 2005). Statistical analysis was performed
using the Analysis-ToolPak module in Microsoft Excel 2016 and the Real Statistics Resource Pack
software, release 6.8 (Zaiontz, 2020). Table 1 shows baseline characteristics of subjects enrolled in
the study from the four different clinical sites. Subject data were pooled for the continued analysis
presented here.
Datasets from the current study are available upon request. The raw data supporting the conclusions
of this article will be made available to qualified researchers, without undue reservation.
376

3   Results

3.1   Enrollment and Allocation

The flow chart for the study is shown in Figure 2. Sixty-seven subjects were assessed for eligibility across the four participating sites. Their respective baseline characteristics are shown in Table 1. First enrollment occurred at the MVAHCS on 10/22/2018, followed by M Health Fairview/University of Minnesota on 10/23/2018, Baylor College of Medicine on 12/20/2018 and Marcus Institute/Harvard Medical School on 09/19/2019. The last enrollment occurred at Baylor College of Medicine on 01/10/2020. Subjects across the four sites were pooled for the continued analysis presented here. Fifteen subjects were excluded from participation of which five had an FGA score higher than 22; two were unable to sense tactile stimuli from the device; two had neurological conditions that prevented device use; two were planning to start physical therapy treatment, one declined to participate and three for other reasons. Fifty-two subjects were enrolled for baseline assessment and allocated for the intervention. A total of seven subjects discontinued participation, four subjects at the two-week assessment (two related to device use, one due to transportation issues, and one due to an unrelated adverse event, Figure 2), one ahead of the six-week assessment and two ahead of the 10-week assessment (due to device and study issues), respectively (Figure 2). A total of 45 subjects (87%) completed all in-clinic assessments from baseline to the 10-week endpoint.

3.2   Baseline Characteristics and Outcomes

Table 2 shows baseline characteristics of all enrolled participants (n=52), then separately for subjects who reported having fallen in the six months preceding the study (Pre-Faller, n=30) and those who did not report a fall (Pre-NonFaller, n=22). Overall, participants were elderly (74.4±8.7 yrs.), overweight (BMI >25) and mostly male (79%). A majority used an assistive device (54%) and had fallen in the previous year (71%) or in the past six months (58%). Furthermore, a majority showed high fall risk based on low 4-Stage Balance Test outcomes (63% of subjects < 30s) (CDC, 2017) or low ABC scores (56% of subjects scored < 67%) (Lajoie and Gallagher, 2004). Twenty-five percent of participants had a normal gait speed less than 0.7 m/s (Montero-Odasso et al., 2005) and half of the participants performed the TUG slower than 12 s (CDC, 2017), the commonly used thresholds for high fall risk. The Pre-Faller group had a higher number of fall-risk factors as compared to the Pre-NonFaller group (5.3±1.0, vs. 3.5±1.3, respectively, $p<0.000001$). The baseline FGA score was statistically significantly lower in Pre-Faller (13.5±3.7) as compared to Pre-NonFaller (16.7±3.6, $p<0.004$). There was no difference in normal gait speed between the two groups ($p=0.12$), although fast gait speed was higher in the Pre-NonFaller (1.41±0.35 m/s) as compared to the Pre-Faller subjects (1.13±0.34 m/s, $p<0.006$). Furthermore, TUG times were significantly slower in the Pre-Faller group compared to the Pre-NonFaller (14.7±6.3s and 12.0±2.9, respectively, $p<0.05$).

The ABC score was higher in the Pre-NonFaller compared to the Pre-Faller group, although the difference was not statistically significant (66.0±19.7 vs. 57.0±15.2, respectively, $p=0.07$). The VADL score was marginally lower in the Pre-NonFaller group (3.29±1.02, vs. 3.94±1.04, $p<0.03$). Pain scores were similar for the two groups (2.6±2.1 vs. 2.9±2.4). The PHQ-9 score was slightly higher in the Pre-Faller group, although the difference was not statistically significant ($p=0.11$). PROMIS outcome scores for "Pain Interference," "Satisfaction with Social Roles," and "Ability to Participate" (Table 2) showed mean values around 50 for both groups, which is considered average for the US population (Askew et al., 2016; Hahn et al., 2016a; Hahn et al., 2016b). Any differences were well within 10, which is one standard deviation of these measures in the US population (Table 2) (Askew et al., 2016; Hahn et al., 2016a; Hahn et al., 2016b).

3.3 Chronic Conditions and Medication Use

Table 3 shows self-reported chronic conditions and medication use for subjects enrolled in the study. On average, subjects reported having 8.2±3.3 chronic conditions. All subjects had a diagnosis of PN either in their medical chart or provided in a letter signed by their physician. Most subjects reported having neuropathic pain in their feet (73%) as well hypertension (63%) and half of participants reported having chronic back pain. All subjects reported having difficulty with walking and balance. The Pre-Faller group reported a higher incidence of cancer as a chronic condition than the Pre-NonFaller group (43% vs. 14%, respectively, p<0.03). Falls in the 12 months preceding study participation were reported by all the Pre-Faller participants (i.e. reporting falls over the past 12 and six months) versus by seven of the 22 in the Pre-NonFaller group (i.e. reporting falls over the past 12 but not six months) (p<0.00001). Ninety percent of the falls reported in the 7-12 months preceding the study were reported by the Pre-Faller participants (p<0.000001). Overall, participants reported taking one non-prescription medication and eight (median) prescription medications of which three are known to cause potential balance issues and increase the risk of falling (Woolcott et al., 2009). Medication use was similar between the Pre-Faller and Pre-NonFaller groups (Table 3).

3.4 Clinical Outcomes

Table 4 shows clinical outcomes for the 45 subjects who completed assessments at baseline, 2, 6 and 10 weeks. For all subjects, repeated measures ANOVA showed highly statistically significant differences across the assessment events for all subjects and all clinical outcomes ($0.00001 < p < 0.01$) except for the 4-Stage Balance Test, which was not statistically significant (p=0.23, Table 4 ANOVA column). Further pairwise comparisons following statistically significant ANOVA showed highly statistically significant differences between the baseline assessment and the 2, 6, and 10-week assessments, respectively, for the FGA score and normal gait speed and between baseline and 6 and 10 weeks, respectively, for fast gait speed and the TUG scores. Cohen's d effect size calculated between the baseline and 10-week endpoint was large for FGA (0.92, FGA change from 15.0 to 19.1) and small to medium for normal gait speed (0.42, 0.86m/s to 0.95m/s), fast gait speed (0.27, 1.24 m/s to 1.33 m/s) and the TUG (0.28, 13.8 s to 12.5 s, Table 4).

Both the Pre-Faller and Pre-NonFaller groups increased their FGA scores from baseline to the 2-, 6-, and 10-week assessments, respectively (p<0.00016). The Pre-Faller group had lower FGA scores both at baseline as compared to the Pre-NonFaller (13.5 vs. 16.9) and at the 10-week assessment (16.8 vs. 22.0). Effect sizes for FGA changes for both groups were large, although higher in the Pre-NonFaller group compared to the Pre-Faller group (Cohen's d 1.38 vs 0.82, respectively). Similarly, normal gait speed increased for both groups although the effect size was larger in the Pre-NonFaller group (Cohen's d 0.74 vs. 0.28, respectively). Interestingly, fast gait speed only improved in the Pre-Faller group from 1.09 m/s at baseline to 1.17 m/s at 10 weeks (p <0.015, Table 4) although effect size was small (Cohen's d 0.27). There was an overall statistically significant change in TUG for the Pre-Faller group (ANOVA, p<0.025) although none of the pairwise comparisons reached statistical significance required after Bonferroni correction (0.047 < p <0.14). The overall ANOVA for TUG in the Pre-NonFaller group was not statistically significant (p=0.49, Table 4).

3.5 Participant Reported Outcomes, Sensation Tests, and Device Use

Patient reported outcomes for the 45 subjects who completed all assessments are shown in Table 5 as well as for the Pre-Faller and Pre-NonFaller groups. For all subjects, there was an overall significant ANOVA for the VADL score (p<0.044) although none of the pairwise comparisons reached

466 statistical significance following Bonferroni correction (0.053 < p <0.99). There were no other
467 statistically significant differences across all subjects during the 10-week period.
468
Table 5 shows and data from vibration sensation testing using the Rydel-Seiffer graduated tuning
fork for all subjects. There were no differences in vibration sensation between Pre-Faller and Pre-
NonFaller and no differences between baseline and 10-week assessments. Vibration sensitivity
showed bilateral symmetry across the three sites tested, right and left patella, lateral malleolus and
first metatarsophalangeal joints, respectively. There was a gradual proximal to distal decrease in
vibration sensitivity across the three sites from 4.3-4.4 proximally and 2.0 distally based on the 0-8
Rydel-Seiffer scale (Table 5). Similarly, there was bilateral symmetry for the WEST monofilament
sensitivity test. At baseline, the median for the monofilament test was 50g for both feet and across all
four anatomical test sites, the first, third, and fifth metatarsal heads as well as the great toe. This was
similar at the 10-week assessment with the exception for the first metatarsals of both feet where the
median monofilament increased to 200g.
480
In the Pre-Faller subgroup, there were no overall statistically significant differences for any of the
participant reported outcomes across the 10-week period (Table 5). For the Pre-NonFaller group,
there was a statistically significant difference in the PROMIS "Ability to Participate" score at two
weeks (ANOVA, p<0.04, 49.7±7.1 to 53.7±7.3, p<0.02, Table 5). The improvement appeared
maintained at six (52.8±7.6) and 10 weeks (53.7±8.4) although it did not fully reach statistical
significance (p<0.06). None of the other participant reported outcomes for the Pre-NonFaller group
were statistically different across the 10-week period (Table 5). Subjects reported using the device on
average 5.5±1.4 days/week (range 3.5-7.0) for a total of 36.1±14.9 hours/week (range 5.3-56.0).

3.6   Falls Assessment

Parameters related to falls and fall risk assessed at baseline and throughout the 10 weeks are shown
in Table 6. Overall, after 10 weeks, 13 of the 45 subjects had achieved FGA scores higher than 22,
the cut-off for normal fall risk (Wrisley and Kumar, 2010). Four of these subjects were part of the
Pre-Faller subgroup and nine were from the Pre-NonFaller group. Subjects reported a total of 62 falls
in the six months prior to the study. During the 10-week study period, 17 falls were reported, 13 of
which occurred in the Pre-Faller group and four in the Pre-NonFaller group (Table 6). Sixteen of the
25 Pre-Fallers did not fall during the 10-week period of the trial. There was a non-significant
decrease in fall rate across all subjects (from 7.7 to 5.4 falls/1000 patient days, p<0.27). In the Pre-
Faller group, there was a 46% statistically significant decrease in fall rate (from 13.8 to 7.4 falls per
1000 patient days, p<0.014). The Pre-NonFaller group showed a smaller non-significant increase in
fall rate (from 0 to 2.9 falls per 1000 patient days, p=0.125). This increase was based on one fall each
by four subjects who had not fallen in the prior six months (Table 6). There was a statistically
significant decrease in the number of fall risk factors (Table 6) from baseline to 10 weeks across all
subjects (from 4.2 to 3.8 fall risk factors, p<0.047). This decrease was larger and statistically
significant in the Pre-Faller group (from 5.1 to 4.3 fall risk factors, p<0.023). There was no change in
the number of fall risk factors in the Pre-NonFaller group (p=0.76, Table 6).
506

4   Discussion

4.1   Key Findings

Results from this multi-site clinical trial supported our *a priori* hypothesis that patients with gait and
balance problems due to sensory PN and high risk of falls would show clinically meaningful
improvements of gait and dynamic balance function after 10 weeks of using a wearable sensory
neuroprosthesis, confirming previously demonstrated in-clinic findings in a randomized controlled
cross-over trial (Koehler-McNicholas et al., 2019). Overall, the mean FGA score improved from 15.0
at baseline to 19.1 at 10 weeks across all subjects (Table 4), beyond the MCID for the FGA
(Beninato et al., 2014). Thirteen subjects reached normal fall risk status showing an FGA score
higher than 22 (Wrisley and Kumar, 2010) after 10 weeks of device use. Both normal and fast gait
speed improved overall by 0.09m/s, near what is considered substantial change (0.10m/s) for older
adults (Perera et al., 2006). Across both groups, TUG improved from 13.8 s to 12.5 s, which is
beyond the MDC for older adults with type 2 diabetes (Alfonso-Rosa et al., 2014). Effect sizes
ranged from large for FGA scores (Cohen's d, 0.92) to small for fast gait speed and TUG (0.27 and
0.28, respectively).
522
As expected, vibratory sensation did not improve during the trial suggesting new sensory information
from the device provided relevant input to improve function. Although both Pre-NonFallers and Pre-
Fallers showed improvements, effect sizes were larger in the Pre-NonFaller group for FGA and
normal gait speed while only the Pre-Faller group showed a statistically significant improvement in
fast gait speed with an effect size of 0.27. The increase in fast gait speed from 1.41 m/s to 1.52 m/s in
the Pre-NonFaller group did not reach statistical significance (ANOVA main effect, p=0.10),
although such gait speeds are considered in the high range of elderly community ambulators
(Middleton et al., 2015). Interestingly, the 4-Stage Balance measure, an indicator of static balance,
did not improve significantly over the 10-week period in either of the two subgroups although our
recent in-clinic study (Koehler-McNicholas et al., 2019) found a statistically significant improvement
in this measure in a group of PN patients. One important difference between these two groups may be
a slightly lower baseline static balance performance in the first cohort, a mean of 22.2 s versus 26.2 s
in the current one. The first group improved their performance in clinic to a mean of 27.6 s (p<0.001)
while they improved to a nearly identical 27.8 s (n.s.) in the current study. It may be that further
improvement of static balance performance would require some additional training challenge in
addition to device use (e.g. Romberg and sharpened Romberg types of activities) and simply wearing
the device daily does not sufficiently challenge the static balance ability assessed by the 4-Stage
Balance Test.

4.2   Physical Intervention versus Sensory Substitution

To our knowledge, this is the first trial where a cohort of subjects with gait and balance problems
related to PN have worn a sensory prosthetic device of this kind in the community and on a regular
basis. Other wearable neuromodulation technologies have been used as a treatment modality in a
home setting, although typically only worn in conjunction with a home therapy program for balance
and mobility in different categories of subjects, e.g., multiple sclerosis (Leonard et al., 2017) and
healthy elderly (Bao et al., 2018). A recent case study of a patient with PN wearing Walkasins for a
year (Wrisley et al., 2018), found dramatic improvements in gait and balance outcomes when daily
continuous device use was combined with balance therapy. The patient had received balance therapy
twice a week for over five months prior to using Walkasins and no longer noticed any further
improvements (Wrisley et al., 2018). In the current study, in an attempt to isolate the effect of long-

term device use on clinical outcomes, subjects were instructed to wear the device as much as possible
and were not allowed to participate in any additional balance training or therapy. Furthermore, they
were not systematically informed of any changes in their outcomes or performance, which they
would normally receive in regular clinical care. Still, improvements in clinical outcomes were similar
to those reported after various balance intervention programs conducted over comparable time
periods up to 12 weeks in similar patient populations (Shumway-Cook et al., 1997; Wolf et al., 2001;
Li et al., 2005; Manor et al., 2014), or even longer up to six months and nearly a year (Wolf et al.,
2003; Li et al., 2005; Li and Manor, 2010).
560
Interestingly, two systematic reviews of interventions specifically for patients with diabetic PN gave
lower extremity strengthening only a fair recommendation while other interventions showed
insufficient evidence to increase function (Ites et al., 2011; Tofthagen et al., 2012). More recently,
improvement in function in patients with PN following specific task-oriented training has been
shown (Salsabili et al., 2015). Consequently, it may be particularly difficult to enhance gait and
balance function in individuals with PN using physical therapy or balance training activities alone,
possibly because these interventions do not replace lost somatosensory input. Furthermore, effects of
a training or therapy program on gait and balance function are likely due to different mechanisms
than the use of a sensory substitution device.
570
For training/therapy programs to be effective, well known principles of training and exercise
physiology must be adhered to (Oddsson et al., 2007) ensuring that sensorimotor systems are
sufficiently challenged to adapt and improve their capabilities leading to improved muscle function
and neuromotor coordination. However, patients with PN have lost important cutaneous afferent
systems that typically would not be affected by such interventions. Instead, improvements related to
use of Walkasins seen in the current study are most likely due to participants receiving new tactile
balance stimuli that are relevant for gait and balance and therefore become integrated into their
neuromotor control and movement repertoire. Subjects in the current study received hundreds of such
stimuli per hour from the device during their regular standing and walking activities throughout the
day.
581
Consequently, we hypothesize that any balance-related therapy or training activity in conjunction
with wearing the device would provide an additive effect to overall function and balance outcomes.
This hypothesis is supported by our recent case study (Wrisley et al., 2018) as well as our previous
in-clinic study where subjects following a baseline assessment were randomized to either wearing the
device turned on or turned off while performing a brief 10-15 min standardized balance activity
session with a physical therapist (Koehler-McNicholas et al., 2019). Ten of 15 subjects in the on
group increased their FGA scores by at least 4 points compared to five of 16 in the off group
(p<0.05). Furthermore, seven of 15 subjects in the on group increased gait speed by >0.13m/s
compared to 3 of 16 in the off group (p<0.05) (Koehler-McNicholas et al., 2019).

4.3   The Importance of Gait Speed

Gait speed is a powerful indicator of overall health and survival in the elderly population and
improving gait speed is an important therapeutic goal. Based on a large population study, Studenski
et al. (Studenski et al., 2011) found that gait speed, age, and gender predicted survival as well as
factors related to chronic conditions, smoking history, blood pressure, and hospitalization. In fact,
improvement in gait speed by 0.10 m/s was found to predict better survival in older adults (Hardy et
al., 2007). Furthermore, while each decrease in gait speed by 0.10m/s has been associated with longer
hospital stays and higher healthcare costs, each 0.10m/s per year increase in gait speed has been

shown to be predictive of shorter hospital stays and healthcare cost reduction (Purser et al., 2005)
further emphasizing the importance of gait speed as a relevant health indicator and vital sign
(Middleton et al., 2015).
602
Healthy aging is associated with an annual decrease in gait speed by 0.013 m/s (Buracchio et al.,
2010). However, it is well know that individuals with PN walk slower than their healthy counterparts
(Menz et al., 2004; Lipsitz et al., 2018), likely as an adapted strategy to maintain balance (Dingwell
et al., 2000). In fact, subjects with peripheral sensory loss (Lipsitz et al., 2018) who were consistently
impaired over five years showed a decline in gait speed of 0.23 m/s over that time period, i.e., 0.046
m/s/yr., more than 3.5 times higher than reported by Buracchio in healthy aging (Buracchio et al.,
2010). Our current population of subjects with PN, who appear similar to the "impaired" category of
community-dwelling older individuals in Lipsitz et al. (Lipsitz et al., 2018), showed an increase in
gait speed of 0.09 m/s following 10 weeks of use of the Walkasins device, corresponding to an
annual rate of 0.47 m/s. Interestingly, the Pre-Faller group increased both their normal and fast gait
speed although the effect size was smaller compared to the Pre-NonFaller group who improved only
their normal gait speed.

4.4   Plantar Sensation and Balance Control

The improvement in gait speed and function from wearing Walkasins may be interpreted from our
understanding of the sensorimotor control of human locomotion and balance. When subjects with PN
perform gait and balance activities, sensory information related to foot pressure is either completely
absent or at least distorted and, therefore, likely non-veridical; and it is unlikely that remaining
balance-related somatosensory information can sufficiently compensate, leading to decreased
stability and increased risk of balance loss. Plantar cutaneous sensory information is important for
standing balance (Meyer et al., 2004b; Strzalkowski et al., 2018) gait stability (Zehr et al., 2014)
signaling stance limb placement and withdrawal to facilitate phase-dependent modulation of
controlling reflex responses (Zehr and Stein, 1999) as well as when responding to balance
perturbations (Meyer et al., 2004a). Hlavacka et al. (Hlavacka et al., 1995) stated that an internal
representation of the body vertical requires integration of somatosensory and vestibular inputs, later
emphasized by Bronstein who concluded that somatosensory information has a "prominent role" in
verticality perception, which is crucial for optimal balance control (Bronstein, 1999). Marsden et al.
(Marsden et al., 2003) concluded that the processing of vestibular information is influenced by load-
related afferent feedback for control of balance.
631
The integration of somatosensory and vestibular information appears of particular importance for gait
function during the double support phase following foot placement during walking (Bent et al.,
2004), which is near the events when Walkasins provides tactile stimuli. Although of lesser fidelity
than intact plantar sensation, the Walkasins device may provide sufficient and relevant sensory
information that is veridical both during standing and walking activities by signaling out of balance
events during standing as well as indicating stance and swing phases of gait, which can help improve
gait and balance function. Furthermore, these tactile stimuli are provided just proximally to the
original sensory loss and mainly along the same dermatomes representing the plantar surface of the
foot possibly making it intuitive to integrate into functional behavior (Koehler-McNicholas et al.,
2019) by providing relevant sensory information to spinal central pattern generators for locomotion
(Guertin, 2012).
643
644

4.5 Falls Data

Although we noticed an encouraging decrease in fall rate as well as in the number of fall-risk factors
in the Pre-faller subgroup (Table 6), a 10-week time period with a fairly small number of subjects is
too short to draw any final conclusions related to prevention of falls. Continued evaluation of longer-
term data from the walk2Wellness trial during continued use of the device is currently ongoing (26
and 52 weeks) and will be reported at a future date. Interestingly, studies have found that effects of
improved clinical gait and balance outcomes may lead to a delayed effect of fall reduction, which
was reported following six months of Tai Chi (Li et al., 2004; Li et al., 2005) and also observed by
Wolf et al. (Wolf et al., 1996; Wolf et al., 2003).

4.6 Participant Reported Outcomes

There appeared to be none or only minimal changes in the self-report measures throughout the 10
weeks of the trial (Table 5). Although this may seem counterintuitive, Richardson et al. (Richardson
et al., 2001) reported significant improvements in clinical balance outcomes, but non-significant
improvement in the ABC score following a strength and balance intervention for patients with PN.
Similarly, an exercise intervention study for older adults found discrepancies between balance and
ABC score improvements suggesting that the relationship between balance confidence and functional
performance may not be well understood (Cyarto et al., 2008). In the current study, a lack of overall
improvement in the ABC score may also be influenced by subject expectations, the short duration of
study, or the time of the year as some subjects were enrolled in the winter months during snowy and
icy conditions. Furthermore, results of the ABC balance confidence scores may also be viewed in the
context of the subjects not being systematically informed of changes in clinical outcomes during the
study, nor receiving any organized encouragement reflecting their performance.
667
Interestingly, upon further investigation we observed differences between the overall pattern of
improvement in clinical outcomes versus self-reported measures of balance confidence as illustrated
in Figure 3. Clinical outcomes at 10 weeks showed similar improvement across the full range of
baseline scores, indicated by a regression line between baseline and 10-week scores being near
parallel with the line of unity as illustrated with the FGA scores in Figure 3 A. However, this was not
the case for the balance confidence ABC-score as seen in Figure 3 B (similar observations were made
for the VADL scores, not shown here). As can be seen, the regression line between baseline and 10-
week ABC scores intersects the line of unity and it has a slope of 0.47 (Figure 3 B). Interestingly, the
two lines intersect at the baseline value 67%, the published cut-off value for high fall risk (Lajoie and
Gallagher, 2004).
678
Subjects enrolled in the current study were all at high fall risk based on their FGA score ≤22 (Wrisley
and Kumar, 2010) as well as a diagnosis of PN (Richardson and Hurvitz, 1995). Based on additional
outcomes and baseline characteristics, subjects had on average more than four fall risk factors (Table
1, Table 2, Table 6). Consequently, it could be argued that study participants with multiple
impairments related to balance maybe "should not" have a balance confidence ABC score above 67%
and those who report such scores are either overly confident and/or simply unaware of their true
balance capabilities. Compellingly, subjects with greater balance confidence at baseline actually
showed a decrease in their balance confidence during the study (from 76.5±8.1 to 71.8±9.9, p<0.02)
while subjects with a baseline ABC score <67% (in the range of high-fall risk) increased their
balance confidence scores (49.9±12.5% to 59.3±15.1%, p<0.01).
689

690 If we consider a simple ratio between ABC and FGA scores representing "amount" of balance
691 confidence per FGA point, someone with 100% balance confidence and the maximum FGA score of
692 30 would have a ratio of 3.3. A ratio between the established cut-offs for high fall risk for ABC and
693 FGA scores, 67% and 22, respectively, represents a ratio of 3.0. At baseline in the current trial, the
694 overly confident subjects had a ratio of 5.1 ± 1.3, versus 3.6 ± 1.1 for the low confidence subjects. After
695 10 weeks of device use, with increased FGA scores across the board, the ratio was 3.8 ± 1.4 for the
696 overly confident subjects versus 3.4 ± 1.0 for the lower confidence subjects. It appears the overly
697 confident subjects may have "normalized" their self-perception of their balance ability while the
698 lower confidence subjects increased their ABC score proportionally to their improved FGA score and
699 maintained a similar self-confidence to FGA ratio. Although the FGA and ABC capture different
700 constructs related to balance, it may be of importance for clinicians to be aware of the potential
701 discrepancy between a patient's self-perception and actual functional performance when developing a
702 plan of care targeting gait, balance function and fall prevention.

704 The T-scores for baseline PROMIS measures (Pain Interference, Satisfaction Social Roles, Ability to
705 Participate) were all close to 50, which was unexpected since it is considered the average for the US
706 population (Hahn et al., 2014; Askew et al., 2016; Hahn et al., 2016a; Hahn et al., 2016b). We had
707 expected these outcomes to deviate significantly in this complex clinical population. Consequently,
708 any major changes in these measures should not be expected although a small statistically significant
709 improvement in PROMIS Ability to Participate score, which remained through 10 weeks, was seen
710 for the Pre-NonFaller sub-group from baseline to the 2-week assessment.

712 Some additional trends in the patient reported outcomes of interest for further research were
713 observed, especially some differences between the Pre-Faller and Pre-NonFaller group. The Pre-
714 Faller group had a PHQ-9 score of 5.3 at baseline, >5 considered mild depression (Kroenke et al.,
715 2001), which decreased to 3.9 at 6 weeks and reached 4.5 at 10 weeks, changes that nearly reached
716 overall statistical significance (p=0.08), and of minimal clinical relevance (Kroenke et al., 2001). It
717 may be of interest to further investigate individuals with higher initial PHQ-9 scores to better
718 understand this observation. In the Pre-NonFaller group, a non-significant (p=0.12), increasing trend
719 in PROMIS Satisfaction Social Roles scores and a near significant improvement in VADL score
720 (p=0.08) were seen. Finally, while VAS Pain scores were overall in the range of mild pain ($\leq$3) and
721 remained steady throughout 10 weeks, pain scores appeared to trend slightly lower in the Pre-
722 NonFaller group, although not statistically significant (p=0.10) and not clinically meaningful. Here it
723 may be of interest to further investigate individuals with higher initial pain levels, especially those
724 with neuropathy-related foot pain by using a more disease-specific pain rating scale.

725 **4.7 Study Limitations**

726 There are several limitations to this trial. It is not blinded, lacks a control group and a placebo
727 treatment. Unfortunately, it is not feasible to blind subjects from treatment in the current study since
728 being able to feel the tactile stimuli from the device is an inclusion criterion. Using some form of
729 random pattern stimuli as a sham may seem possible (Basta et al., 2011), although it is not known if
730 such stimuli may have an effect of their own and it would not help address the question whether
731 using the device as currently designed, according to principles of sensorimotor control of balance and
732 gait, has an effect on gait and balance function. Consequently, the best placebo treatment would
733 likely be wearing a device that is turned off. However, without using some form of deceit claiming
734 the device is working although it cannot be felt, it would likely be difficult to recruit participants for
735 such research and/or to ensure long-term compliance. In addition, incorporating a minimal
736 stimulation amplitude as a sham, assuming it has no effect may be incorrect since studies

implementing stochastic resonance using subsensory mechanical noise have demonstrated
improvement in balance (Lipsitz et al., 2015). Furthermore, using a randomized control cross-over
design, we recently demonstrated in-clinic improvements in clinical outcomes when the Walkasins
device was worn and turned on as compared to turned off (Koehler-McNicholas et al., 2019).
Consequently, we felt comfortable incorporating a single treatment arm design knowing the in-clinic
effects. Furthermore, any placebo effects were likely decreased by not systematically informing
subjects about any changes in outcomes and minimizing encouragement during interactions with
subjects that could affect expectation and beliefs in the treatment (Finniss et al., 2010; Enck et al.,
2013; Coste and Montel, 2017), and prohibiting any additional balance training/therapy intervention
during the 10 weeks of the trial.
747
If the effects in this study were placebo our findings should align with research findings on the
placebo arm of randomized, placebo-controlled trials (Wartolowska et al., 2016). A systematic
review of temporal changes in the placebo arm across 47 surgical randomized control trials found
that effects size of subjective outcomes was large (0.64), while effect size of objective clinical
outcomes was small (0.11) (Wartolowska et al., 2016). Furthermore, major differences in placebo-
effect sizes have been reported with subject-reported self-perception effects being larger than
observer-based ratings (Rief et al., 2009). On the contrary, effect sizes in the current study were large
for the clinical outcomes and small for the self-reported outcomes, supporting the interpretation that
effects were due to device use and not placebo. Further support of this view includes relatively high
subject compliance and reported device use and, a low subject dropout rate as well as the sustained
duration and continued gradual improvement in clinical outcomes throughout the 10-week period. An
additional weakness includes enrollment of mostly male subjects, partly due to nearly half of the
subjects being Veterans, who especially in this older generation are predominantly male. Strengths of
this trial include involvement of multiple sites across different geographies with different assessors at
different clinics limiting confounding balance interventions, and the use of standardized objective
clinical outcome measures.

5   Conclusion

765 A wearable sensory neuroprosthesis may provide a new way to treat gait and balance problems and
766 manage falls in high fall-risk patients with PN. Longer term data would be required to further
767 investigate actual decreases in falls.

6   Acknowledgments

We thank Lori Danzl, Alexandria Lloyd, Christine Olney (Minneapolis Department of Veterans
Affairs Health Care System, Minneapolis, MN), Tien Dat Nguyen (M Health Fairview, Minneapolis,
MN), Nathan Silver (Baylor College of Medicine, Houston, TX) and research staff at Hinda and
Arthur Marcus Institute for Aging Research, Hebrew SeniorLife, Roslindale, MA, for providing
study coordination, subject recruitment and data gathering activities, and Annette Xenopoulos-
Oddsson (University of Minnesota, MN,) for reviewing early drafts of the manuscript.
775

7   Contribution to the Field Statement

Peripheral neuropathy is a condition affecting over 20 Million individuals in the US alone that causes
damage to nerves in the peripheral nervous system. Peripheral neuropathy is often related to diabetes,
chemotherapy treatment or has unknown causes. Symptoms of the disease that involve numbness and

loss of sensation in the feet often lead to problems with gait and balance and a high risk of falls. We
report results from the multi-site clinical trial, walk2Wellness, that has investigated long-term effects
of daily use of a new wearable sensory neuroprosthesis on gait function, balance, fall rates, and
quality of life in a group of peripheral neuropathy patients. The device (Walkasins®, RxFunction
Inc., MN, USA) provides directional tactile stimuli around the ankle during standing and walking
activities reflecting changes in foot pressure related to balance. Patients can then "feel" their feet in
contact with the ground, notice body sway during standing replacing their lost afferent nerve
function. Our results after 10 weeks of daily use show improved walking balance, speed of walking,
ability to rise from a chair move around and sit down and an encouraging trend in lower fall rates.
Also, participants with initial low self-confidence in their balance showed higher self-confidence.
790

791
792 **Figure 1.** Picture of the Walkasins sensory neuroprosthesis device showing the pressure sensor embedded Foot Pad that
793 is placed in the subject's shoe and connected to the Leg Unit that houses an embedded microprocessor with software,
794 supporting electronics, a rechargeable battery, and four tactile stimulators placed around the lower leg. The system is
795 worn bilaterally. Leg Unit and left Foot Pad shown

796
797
798

Figure 2. Flowchart of the study.

799
800
801

802
803

	n	Age (yrs)	Height (m)	Weight (kg)	#ChrD	FGA Score	Gait Speed Normal (m/s)	Gait Speed Fast (m/s)	TUG (s)	4-Stage Balance Test (s)
VAMC	20	74.7 (6.0)	1.76 (0.06)	96.9 (13.5)	9.8 (2.4)	14.8 (4.4)	0.80 (0.16)	1.2 (0.23)	14.7 (6.3)	25.0 (7.3)
Baylor	18	75.1 (11.7)	1.74 (0.08)	80.9 (18.1)	8.2 (3.0)	15.1 (4.2)	0.87 (0.25)	1.28 (0.39)	13.2 (5.2)	28.1 (8.2)
M Health Fairview	9	70.1 (7.8)	1.79 (0.12)	87.6 (10.5)	6.6 (4.1)	15.0 (3.2)	1.02 (0.28)	1.32 (0.27)	10.8 (2.3)	23.6 (5.2)
Harvard	5	78.2 (5.1)	1.71 (0.1)	95.0 (22.0)	5.2 (2.9)	14.0 (3.4)	0.77 (0.25)	1.07 (0.45)	15.0 (3.4)	28.3 (6.6)

804 **Table 1.** Baseline characteristics of subjects from the four different clinical sites enrolled in the study. Values represent
805 Mean (Standard Deviation). #ChrD - Number of Chronic Diseases.
806

807
808

Baseline Assessment	All n=52	Pre-F n=30	Pre-NF n=22	p-level
Gender Female (n)	11 of 52 (21%)	9 of 30 (30%)	2 of 22 (10%)	0.069
Use of Assistive Device (n)	28 of 52 (54%)	19 of 30 (63%)	9 of 22 (41%)	0.11
Gait Speed Normal <0.7m/s (n)	13 of 52 (25%)	10 of 30 (33%)	3 of 22 (14%)	0.10
Timed Up and Go >12s (n)	26 of 52 (50%)	18 of 30 (60%)	8 of 22 (36%)	0.09
4-Stage Balance Test <30s (n)	33 of 52 (63%)	22 of 30 (73%)	11 of 22 (50%)	0.08
ABC Score <67% (n)	30 of 52 (56%)	20 of 30 (67%)	10 of 22 (45%)	0.13
Fallen in Last 6 Months (n)	30 of 52 (58%)	30 of 30 (100%)	0	n/a
Fallen in Last 12 Months (n)	37 of 52 (71%)	30 of 30 (100%)	7 of 22 (32%)	**<0.00001**
Number of Falls 6 Months	65	65	0	n/a
Number of Falls 12 Months	121	109 of 121 (90%)	12 of 121 (10%)	**<0.000001**
	Mean (SD) n=52	Mean (SD) n=30	Mean (SD) n=22	p-level
Age (yrs)	74.4 (8.7)	73.8 (9.1)	75.2 (8.2)	0.57
Height (m)	1.75 (0.08)	1.76 (0.09)	1.75 (0.07)	0.54
Weight (kg)	89.6 (16.8)	89.7 (17.8)	89.3 (15.8)	0.93
BMI (kg/m^2)	29.1 (5.2)	29.0 (5.8)	29.2 (5.2)	0.92
FGA Score	14.9 (4.0)	13.5 (3.7)	16.7 (3.6)	**<0.004**
Gait Speed, Normal (m/s)	0.86 (0.23)	0.81 (0.26)	0.92 (0.18)	0.12
Gait Speed, Fast (m/s)	1.25 (0.37)	1.13 (0.34)	1.41 (0.35)	**<0.006**
TUG (s)	13.5 (5.3)	14.7 (6.3)	12.0 (2.9)	**<0.05**
4-Stage Balance Test (s)	26.2 (7.3)	24.8 (6.3)	28.0 (8.3)	0.055
Fall-Risk Factors* (n of 7)	4.5 (1.5)	5.3 (1.0)	3.5 (1.3)	**<0.000001**
ABC Score (%)	60.8 (17.6)	57.0 (15.2)	66.0 (19.7)	0.07
VADL Mean Score	3.66 (1.07)	3.94 (1.04)	3.29 (1.02)	**<0.03**
VAS Pain Score (0-10)	2.7 (2.2)	2.6 (2.1)	2.9 (2.4)	0.67
PHQ-9	4.4 (3.8)	5.2 (4.3)	3.4 (3.8)	0.11
Pain Interference PROMIS® 6b	51.1 (8.0)	52.5 (8.1)	49.0 (8.1)	0.10
Satisfaction Social Roles PROMIS® 8a	50.4 (7.8)	49.1 (7.7)	52.1 (7.8)	0.13
Ability to Participate PROMIS® 8a	50.0 (7.2)	49.8 (7.1)	50.2 (7.2)	0.88

Table 2. Baseline characteristics of subjects enrolled in the study (n=52), then separately for subjects who reported
having fallen in the past six months (Pre-F) and those who did not (Pre-NF). Values shown are Mean (Standard
Deviation). Column p-level shows significance level for comparison between the Pre-F and Pre-NF groups. In bold if
p<0.05.

*Fall-risk factors assessed in the current study included, recent history of falls (Tinetti and Kumar, 2010), PN diagnosis
(Richardson and Hurvitz, 1995), FGA score<23 (Wrisley and Kumar, 2010), TUG >12 s (CDC, 2017), 4-Stage Balance
Test<30s (CDC, 2017), Gait Speed <0.7m/s (Studenski et al., 2003; Montero-Odasso et al., 2005), ABC score <67%
(Lajoie and Gallagher, 2004).

817
818

819
820

Baseline Assessment	All n=52	Pre-F n=30	Pre-NF n=22	p-level
Number of Chronic Conditions (n)	8.2 (3.3)	8.6 (3.2)	7.8 (3.4)	0.39
Peripheral Neuropathy (n)	52 (100%)	30 (100%)	22 (100%)	n/a
Numbness in Feet (n)	49 (94%)	30 (100%)	19 (86%)	**<0.05**
Neuropathic Pain in Feet (n)	38 (73%)	22 (73%)	16 (73%)	0.96
Hypertension (n)	33 (63%)	19 (63%)	14 (64%)	0.98
Back Pain (n)	26 (50%)	16 (53%)	10 (45%)	0.57
Arthritis (n)	24 (46%)	15 (50%)	9 (41%)	0.52
Knee Dysfunction (n)	23 (44%)	14 (47%)	9 (41%)	0.68
Diabetes Diagnosis (n)	19 (37%)	9 (30%)	10 (45%)	0.25
Inner Ear Problems (n)	17 (33%)	9 (30%)	8 (36%)	0.63
Heart Disease (n)	16 (31%)	10 (33%)	6 (27%)	0.64
Neck Pain (n)	16 (31%)	9 (30%)	7 (32%)	0.89
Cancer (n)	16 (31%)	13 (43%)	3 (14%)	**<0.03**
Lung Disease (n)	10 (19%)	7 (23%)	3 (14%)	0.38
Stroke (n)	9 (17%)	4 (13%)	5 (23%)	0.38
Osteoporosis(n)	10 (19%)	7 (23%)	3 (14%)	0.38
Seizures (n)	5 (10%)	3 (10%)	2 (9%)	0.91
Ankle Dysfunction (n)	5 (10%)	4 (13%)	1 (5%)	0.29
TMJ/Jaw Pain (n)	5 (10%)	2 (7%)	3 (14%)	0.40
Fainting (n)	5 (10%)	2 (7%)	3 (14%)	0.40
Migraines (n)	4 (8%)	2 (7%)	2 (9%)	0.75
Meningitis (n)	0	0	0	n/a
Other Conditions (n)	14 (27%)	11 (37%)	3 (14%)	0.06
Difficulty Walking/balance (n)	52 (100%)	30 (100%)	22 (100%)	n/a
	Mean (SD) Median	Mean (SD)	Mean (SD)	p-level
Prescription Medications (n)	8	8.5	7	0.50
Non-Prescription Medications (n)	1	1	1	n/a
Medications Associated with Falls (n)	3	3	3	n/a

821 **Table 3.** Self-reported chronic conditions as well as medication use for subjects enrolled in the study. Column p-level
822 shows significance level for comparison between the Pre-F and Pre-NF groups. In bold if p<0.05.

823
824

825
826

ALL	Baseline Mean (SD) n=45	2-week Mean (SD) n=45	6-week Mean (SD) n=45	10-week Mean (SD) n=45	ANOVA p-level	Cohen's d (95% CI's)
FGA Score	15.0 (4.0)	18.3 (4.4) **<0.0000001**	18.5(4.4) **<0.0000001**	19.1 (5.2) **<0.0000001**	**<0.00001**	0.92 (0.49, 1.35)
Gait Speed Normal (m/s)	0.86 (0.24)	0.92 (0.26) **<0.002**	0.94 (0.25) **<0.0001**	0.95 (0.24) **<0.00005**	**<0.00001**	0.42 (0.01, 0.82)
Gait Speed Fast (m/s)	1.24 (0.37)	1.27 (0.33) 0.30	1.30 (0.37) **<0.016**	1.33 (0.38) **<0.00168**	**<0.013**	0.27 (-0.14,0.67)
TUG (s)	13.8 (5.5)	12.7 (4.2) 0.06	12.3 (4.2) **<0.034** (*0.69*) 0.0397*	12.5 (3.7) **<0.012**	**<0.01**	0.28 (-0.12, 0.68)
4-Stage Balance Test (s)	26.2 (7.1)	27.1 (7.7)	27.7 (7.7)	27.8 (6.8)	0.23	n/a
Pre-Fallers	**Baseline Mean (SD) n=25**	**2-week Mean (SD) n=25**	**6-week Mean (SD) n=25**	**10-week Mean (SD) n=25**	**p-level ANOVA**	**Cohen's d (95% CI's)**
FGA Score	13.5 (3.6)	16.6 (3.7) **<0.0000005**	17.3 (4.1) **<0.00002**	16.8 (4.9) **<0.00016**	**<0.00001**	0.82 (0.26, 1.39)
Gait Speed Normal (m/s)	0.79 (0.27)	0.85 (0.26) **<0.022** (*0.90*) 0.047*	0.89 (0.25) **<0.0021**	0.86 (0.24) **<0.016**	**<0.0014**	0.28 (-0.27, 0.83)
Gait Speed Fast (m/s)	1.09 (0.33)	1.16 (0.30) **<0.035** (*0.88*) 0.046*	1.17 (0.32) **<0.0165**	1.17 (0.35) **<0.015**	**<0.028**	0.27 (-0.28, 0.82)
TUG (s)	15.3 (6.6)	14.0 (4.6) 0.14	13.3 (3.8) 0.047	13.7 (4.2) 0.056	**<0.025**	n/a
4-Stage Balance Test (s)	24.7 (5.5)	25.4 (7.8)	26.0 (7.9)	27.5 (6.1)	0.15	n/a
Pre-NonFallers	**Baseline Mean (SD) n=20**	**2-week Mean (SD) n=20**	**6-week Mean (SD) n=20**	**10-week Mean (SD) n=20**	**p-level ANOVA**	**Cohen's d (95% CI's)**
FGA Score	16.9 (3.6)	20.5 (4.4) **<0.00009**	20.1 (4.3) **<0.00004**	22.0 (4.0) **<0.000004**	**<0.0000001**	1.38 (0.70, 2.1)
Gait Speed Normal (m/s)	0.93 (0.17)	1.01 (0.22) **<0.031** (*0.78*) 0.043*	1.01 (0.24) **<0.019** (*0.84*) 0.045*	1.06 (0.20) **<0.0004**	**<0.0011**	0.74 (0.10, 1.37)
Gait Speed Fast (m/s)	1.41 (0.35)	1.42 (0.33)	1.43 (0.41)	1.52 (0.34)	0.10	n/a
TUG (s)	11.9 (2.9)	11.2 (3.2)	11.7 (5.0)	10.9 (2.2)	0.49	n/a
4-Stage Balance Test (s)	27.6 (8.5)	29.4 (7.1)	29.9 (7.1)	28.1 (7.7)	0.45	n/a

Table 4. Clinical outcomes for the 45 subjects completing all assessments for baseline, 2, 6 and 10 weeks as well as the
subgroups of Pre-Fallers and Pre-NonFallers. Values in **bold** indicate statistical significance. ANOVA column shows
significance levels of main effect from initial repeated measures ANOVA test. If significant (p<0.05), pairwise
comparisons were made using three dependent t-tests between baseline and 2, 6 and 10-week assessments, respectively.
Bonferroni's adjustment of significance levels for correlated measures was applied. As noted, values in *(italics)* indicate
Pearson's correlation coefficient followed by the adjusted significance level required for an overall significance of 0.05 as
marked with *. If ANOVA was non-significant, no further comparisons were made. Cohen's d indicates effect size where
0.2 is represents a small effect, 0.5 a medium effect, and 0.8 a large effect. Values in parenthesis show 95% confidence
interval.
836

837
838

ALL (n=45)	Baseline Mean (SD) n=45	2-week Mean (SD) n=45	6-week Mean (SD) n=45	10-week Mean (SD) n=45	ANOVA p-level
ABC-Score (%)	61.4 (17.9)	65.0 (13.1)	64.2 (15.1)	65.1 (14.1)	0.36
VADL Mean Score	3.70 (1.09)	3.37 (0.86) 0.053	3.52 (0.99) 0.16	3.63 (0.93) 0.99	**<0.044**
VAS Score (0-10)	2.8 (2.2)	2.5 (2.1)	2.5 (2.2)	2.6 (2.3)	0.31
PHQ-9	4.5 (3.9)	3.8 (3.5)	3.6 (3.8)	3.9 (4.5)	0.25
Pain Interference PROMIS® 6b	50.8 (7.9)	51.2 (8.1)	50.5 (8.9)	51.8 (8.5)	0.85
Satisfaction Social Roles PROMIS® 8a	50.2 (7.8)	52.1 (7.2)	51.7 (7.3)	52.1 (7.8)	0.10
Ability to Participate PROMIS® 8a	49.8 (7.3)	50.7 (6.4)	51.0 (7.1)	51.3 (8.1)	0.38
R 1st MTP Joint	2.0 (2.5)	n/a	n/a	2.0 (2.5)	n/a
R Lateral Malleolus	3.6 (2.7)	n/a	n/a	3.4 (2.6)	n/a
R Patella	4.5 (2.0)	n/a	n/a	4.5 (1.8)	n/a
L 1st MTP Joint	2.3 (2.6)	n/a	n/a	2.2 (2.7)	n/a
L Lateral Malleolus	3.7 (2.7)	n/a	n/a	3.6 (2.5)	n/a
L Patella	4.3 (2.1)	n/a	n/a	4.4 (1.6)	n/a
Pre-Fallers (n=25)	**Baseline Mean (SD)**	**2-week Mean (SD)**	**6-week Mean (SD)**	**10-week Mean (SD)**	**p-level**
ABC-Score (%)	57.7 (17.9)	61.5 (14.0)	61.8 (15.1)	61.2 (13.9)	0.29
VADL Mean Score	4.00 (1.13)	3.68 (0.90)	3.83 (1.01)	3.91 (0.97)	0.42
VAS Score (0-10)	2.5 (2.4)	2.5 (2.5)	2.8 (2.6)	2.6 (2.5)	0.55
PHQ-9	5.3 (3.9)	4.7 (3.3)	3.9 (3.3)	4.5. (3.5)	0.08
Pain Interference PROMIS® 6b	51.9 (8.3)	52.2 (8.5)	51.6 (9.9)	53.6 (8.0)	0.90
Satisfaction Social Roles PROMIS® 8a	49.1 (8.0)	49.9 (7.9)	49.2 (7.5)	49.6 (7.7)	0.68
Ability to Participate PROMIS® 8a	49.9 (7.5)	48.6 (5.8)	49.5 (6.8)	49.4 8.0)	0.58
Pre-NonFallers (n=20)	**Baseline Mean (SD)**	**2-week Mean (SD)**	**6-week Mean (SD)**	**10-week Mean (SD)**	**p-level**
ABC-Score (%)	66.2 (18.0)	69.8 (11.9)	67.2 (15.1)	70.2 (14.6)	0.96
VADL Mean Score	3.32 (1.06)	2.94 (0.80)	3.11 (0.99)	3.28 (0.89)	0.08
VAS Score (0-10)	3.0 (2.0)	2.4 (1.5)	2.2 (1.7)	2.7 (2.0)	0.10
PHQ-9	3.5 (3.9)	2.6.(3.7)	3.2 (4.4)	3.2 (5.6)	0.66
Pain Interference PROMIS® 6b	49.4 (7.6)	49.8 (7.6)	49.1 (7.6)	49.3 (9.1)	0.91
Satisfaction Social Roles PROMIS® 8a	51.7 (7.8)	55 .1 (6.4)	54.8 (7.0)	55.4 (8.0)	0.12
Ability to Participate PROMIS® 8a	49.7 (7.1)	53.7 (7.3) <**0.02** (0.64) 0.038*	52.8 (7.6) 0.06	53.7 (8.4) 0.05	**<0.041**

Table 5. Results from participant-reported outcomes and Rydel-Seiffer vibration sensation screening for the 45 subjects
who completed all assessments. Data is shown separately for the group as a whole and for Pre-Fallers (having reported
fallen in the previous 6 months) and Pre-NonFallers (no falls reported in the previous 6 months). Values in *(italics)*
indicate Pearson's correlation coefficient followed by the Bonferroni adjusted significance level required for an overall
significance of 0.05 as marked with *. There were no differences in vibration sensation between Pre-Fallers and Pre-
NonFallers and no differences between baseline and 10-week assessments.
845

846
847

All (n=45)	Baseline	10 Weeks	p-level
Number of Subjects FGA>22	0 of 45	13 of 45	n/a
#Falls (pre-6 mo & in study)	62	17	n/a
Fall Rate (pre-6mo & in study)	7.7	5.4	0.27*
#Fallers (pre-6mo & in study)	25	22	n/a
# Fall Risk Factors	4.2 (1.5)	3.8 (1.6)	**<0.047**
Pre-Fallers (n=25)			
Number of Subjects FGA>22	0 of 25	4 of 25	n/a
#Falls (pre-6 mo & in study)	62	13	n/a
Fall Rate (pre-6mo & in study)	13.8	7.4	**<0.014***
#Fallers (pre-6mo & in study)	25	9	n/a
# Fall Risk Factors	5.1 (1.3)	4.3 (1.7)	**<0.023**
Pre-NonFallers (n=20)			
Number of Subjects FGA>22	0 of 20	9 of 20	n/a
#Falls (pre-6 mo & in study)	0	4	n/a
Fall Rate (pre-6mo & in study)	0	2.9	0.125*
#Fallers (pre-6mo & in study)	0	4	n/a
# Fall Risk Factors	3.2 (1.8)	3.1 (1.7)	0.76

848   **Table 6.** Parameters related to falls and fall risk assessed at baseline and at 10 weeks. Fall rates are reported in number of
849   falls per 1000 patient days. Fall risk factors are identified in Table 2. *Wilcoxon Signed Rank test
850

851
852

853
854
855 **Figure 3.** Graphs showing baseline vs. 10-week FGA (A) and ABC scores (B). Markers above line of identity indicate
856 higher scores at 10-week assessment. Notice line of regression for FGA scores is near parallel to line of identity
857 indicating a similar increase across all baseline FGA scores. For ABC scores the line of regression intersects the line of
858 identify near 67% indicating an increase for lower baseline ABC scores and a decrease for higher baseline ABC scores.

859
860

8   References

Akbari, M., Jafari, H., Moshashaee, A., and Forugh, B. (2012). Do diabetic neuropathy patients benefit from balance training? *J Rehabil Res Dev* 49(2)**,** 333-338. doi: 10.1682/jrrd.2010.10.0197.

Alfonso-Rosa, R.M., Del Pozo-Cruz, B., Del Pozo-Cruz, J., Sañudo, B., and Rogers, M.E. (2014). Test-retest reliability and minimal detectable change scores for fitness assessment in older adults with type 2 diabetes. *Rehabil Nurs* 39(5)**,** 260-268. doi: 10.1002/rnj.111.

Algina, J, Keselman, H.J., and Penfield, R.D. (2005). Effect Sizes and their Intervals: The Two-Level Repeated Measures Case. *Edu Psych Meas* 65(April)**,** 241–258. doi: 10.1177/0013164404268675.

Allet, L., Armand, S., de Bie, R.A., Golay, A., Monnin, D., Aminian, K., et al. (2010). The gait and balance of patients with diabetes can be improved: a randomised controlled trial. *Diabetologia* 53(3)**,** 458-466. doi: 10.1007/s00125-009-1592-4.

Askew, R.L., Cook, K.F., Revicki, D.A., Cella, D., and Amtmann, D. (2016). Evidence from diverse clinical populations supported clinical validity of PROMIS pain interference and pain behavior. *J Clin Epidemiol* 73**,** 103-111. doi: 10.1016/j.jclinepi.2015.08.035.

Bach-y-Rita, P. (2004). Tactile sensory substitution studies. *Ann N Y Acad Sci* 1013**,** 83-91.

Bach-y-Rita, P., Collins, C.C., Saunders, F.A., White, B., and Scadden, L. (1969). Vision substitution by tactile image projection. *Trans Pac Coast Otoophthalmol Soc Annu Meet* 50**,** 83-91.

Badke, M.B., Sherman, J., Boyne, P., Page, S., and Dunning, K. (2011). Tongue-based biofeedback for balance in stroke: results of an 8-week pilot study. *Arch Phys Med Rehabil* 92(9)**,** 1364-1370. doi: 10.1016/j.apmr.2011.03.030.

Bao, T., Carender, W.J., Kinnaird, C., Barone, V.J., Peethambaran, G., Whitney, S.L., et al. (2018). Effects of long-term balance training with vibrotactile sensory augmentation among community-dwelling healthy older adults: a randomized preliminary study. *J Neuroeng Rehabil* 15(1)**,** 5. doi: 10.1186/s12984-017-0339-6.

Basta, D., Rossi-Izquierdo, M., Soto-Varela, A., Greters, M.E., Bittar, R.S., Steinhagen-Thiessen, E., et al. (2011). Efficacy of a vibrotactile neurofeedback training in stance and gait conditions for the treatment of balance deficits: a double-blind, placebo-controlled multicenter study. *Otol Neurotol* 32(9)**,** 1492-1499. doi: 10.1097/MAO.0b013e31823827ec.

Beninato, M., Fernandes, A., and Plummer, L.S. (2014). Minimal clinically important difference of the functional gait assessment in older adults. *Phys Ther* 94(11)**,** 1594-1603. doi: 10.2522/ptj.20130596.

Bent, L.R., Inglis, J.T., and McFadyen, B.J. (2004). When is vestibular information important during walking? *J Neurophysiol* 92(3)**,** 1269-1275. doi: 10.1152/jn.01260.2003.

Bergen, G., Stevens, M.R., and Burns, E.R. (2016). Falls and Fall Injuries Among Adults Aged ≥65 Years - United States, 2014. *MMWR Morb Mortal Wkly Rep* 65(37)**,** 993-998. doi: 10.15585/mmwr.mm6537a2.

Bischoff, H.A., Stähelin, H.B., Monsch, A.U., Iversen, M.D., Weyh, A., von Dechend, M., et al. (2003). Identifying a cut-off point for normal mobility: a comparison of the timed 'up and go' test in community-dwelling and institutionalised elderly women. *Age Ageing* 32(3)**,** 315-320. doi: 10.1093/ageing/32.3.315.

Bronstein, A.M. (1999). The interaction of otolith and proprioceptive information in the perception of verticality. The effects of labyrinthine and CNS disease. *Ann N Y Acad Sci* 871**,** 324-333. doi: 10.1111/j.1749-6632.1999.tb09195.x.

Bulat, T., Hart-Hughes, S., Ahmed, S., Quigley, P., Palacios, P., Werner, D.C., et al. (2007). Effect of
a group-based exercise program on balance in elderly. *Clin Interv Aging* 2(4)**,** 655-660. doi:
10.2147/cia.s204.
Buracchio, T., Dodge, H.H., Howieson, D., Wasserman, D., and Kaye, J. (2010). The trajectory of
gait speed preceding mild cognitive impairment. *Arch Neurol* 67(8)**,** 980-986. doi:
10.1001/archneurol.2010.159.
Cakrt, O., Vyhnálek, M., Slabý, K., Funda, T., Vuillerme, N., Kolář, P., et al. (2012). Balance
rehabilitation therapy by tongue electrotactile biofeedback in patients with degenerative
cerebellar disease. *NeuroRehabilitation* 31(4)**,** 429-434. doi: 10.3233/NRE-2012-00813.
Cavanagh, P.R., Derr, J.A., Ulbrecht, J.S., Maser, R.E., and Orchard, T.J. (1992). Problems with gait
and posture in neuropathic patients with insulin-dependent diabetes mellitus. *Diabet Med*
9(5)**,** 469-474. doi: 10.1111/j.1464-5491.1992.tb01819.x.
CDC (2017). *STEADI - Older Adult Fall Prevention* [Online]. On line: Centers for Disease Control
and Prevention. Available: https://www.cdc.gov/steadi/materials.html [Accessed May 5, 2018
2018].
Chodzko-Zajko, W.J., Proctor, D.N., Fiatarone Singh, M.A., Minson, C.T., Nigg, C.R., Salem, G.J.,
et al. (2009). American College of Sports Medicine position stand. Exercise and physical
activity for older adults. *Med Sci Sports Exerc* 41(7)**,** 1510-1530. doi:
10.1249/MSS.0b013e3181a0c95c.
Cohen, H.S., Kimball, K.T., and Adams, A.S. (2000). Application of the vestibular disorders
activities of daily living scale. *Laryngoscope* 110(7)**,** 1204-1209. doi: 10.1097/00005537-
200007000-00026.
Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences.* New York, NY: Routledge
Academic.
Coste, J., and Montel, S. (2017). Placebo-related effects: a meta-narrative review of
conceptualization, mechanisms and their relevance in rheumatology. *Rheumatology (Oxford)*
56(3)**,** 334-343. doi: 10.1093/rheumatology/kew274.
Cyarto, E.V., Brown, W.J., Marshall, A.L., and Trost, S.G. (2008). Comparative effects of home- and
group-based exercise on balance confidence and balance ability in older adults: cluster
randomized trial. *Gerontology* 54(5)**,** 272-280. doi: 10.1159/000155653.
DeMott, T.K., Richardson, J.K., Thies, S.B., and Ashton-Miller, J.A. (2007). Falls and gait
characteristics among older persons with peripheral neuropathy. *Am J Phys Med Rehabil*
86(2)**,** 125-132. doi: 10.1097/PHM.0b013e31802ee1d1.
Dingwell, J.B., Cusumano, J.P., Sternad, D., and Cavanagh, P.R. (2000). Slower speeds in patients
with diabetic neuropathy lead to improved local dynamic stability of continuous overground
walking. *J Biomech* 33(10)**,** 1269-1277. doi: 10.1016/s0021-9290(00)00092-0.
Dixon, C.J., Knight, T., Binns, E., Ihaka, B., and O'Brien, D. (2017). Clinical measures of balance in
people with type two diabetes: A systematic literature review. *Gait Posture* 58**,** 325-332. doi:
10.1016/j.gaitpost.2017.08.022.
Enck, P., Bingel, U., Schedlowski, M., and Rief, W. (2013). The placebo response in medicine:
minimize, maximize or personalize? *Nat Rev Drug Discov* 12(3)**,** 191-204. doi:
10.1038/nrd3923.
Finniss, D.G., Kaptchuk, T.J., Miller, F., and Benedetti, F. (2010). Biological, clinical, and ethical
advances of placebo effects. *Lancet* 375(9715)**,** 686-695. doi: 10.1016/S0140-6736(09)61706-
2.
Florence, C.S., Bergen, G., Atherly, A., Burns, E., Stevens, J., and Drake, C. (2018). Medical Costs
of Fatal and Nonfatal Falls in Older Adults. *J Am Geriatr Soc* 66(4)**,** 693-698. doi:
10.1111/jgs.15304.

Ganz, D.A., Bao, Y., Shekelle, P.G., and Rubenstein, L.Z. (2007). Will my patient fall? *JAMA*
297(1)**,** 77-86. doi: 10.1001/jama.297.1.77.

Ganz, D.A., and Latham, N.K. (2020). Prevention of Falls in Community-Dwelling Older Adults. *N*
Engl J Med 382(8)**,** 734-743. doi: 10.1056/NEJMcp1903252.

Gregg, E.W., Sorlie, P., Paulose-Ram, R., Gu, Q., Eberhardt, M.S., Wolz, M., et al. (2004).
Prevalence of lower-extremity disease in the US adult population >=40 years of age with and
without diabetes: 1999-2000 national health and nutrition examination survey. *Diabetes Care*
27(7)**,** 1591-1597. doi: 10.2337/diacare.27.7.1591.

Guertin, P.A. (2012). Central pattern generator for locomotion: anatomical, physiological, and
pathophysiological considerations. *Front Neurol* 3**,** 183. doi: 10.3389/fneur.2012.00183.

Hahn, E.A., Beaumont, J.L., Pilkonis, P.A., Garcia, S.F., Magasi, S., DeWalt, D.A., et al. (2016a).
The PROMIS satisfaction with social participation measures demonstrated responsiveness in
diverse clinical populations. *J Clin Epidemiol* 73**,** 135-141. doi:
10.1016/j.jclinepi.2015.08.034.

Hahn, E.A., DeWalt, D.A., Bode, R.K., Garcia, S.F., DeVellis, R.F., Correia, H., et al. (2014). New
English and Spanish social health measures will facilitate evaluating health determinants.
Health Psychol 33(5)**,** 490-499. doi: 10.1037/hea0000055.

Hahn, E.A., Kallen, M.A., Jensen, R.E., Potosky, A.L., Moinpour, C.M., Ramirez, M., et al. (2016b).
Measuring social function in diverse cancer populations: Evaluation of measurement
equivalence of the Patient Reported Outcomes Measurement Information System. *Psychol*
Test Assess Model 58(2)**,** 403-421.

Halvarsson, A., Franzén, E., Farén, E., Olsson, E., Oddsson, L., and Ståhle, A. (2013). Long-term
effects of new progressive group balance training for elderly people with increased risk of
falling - a randomized controlled trial. *Clin Rehabil* 27(5)**,** 450-458. doi:
10.1177/0269215512462908.

Halvarsson, A., Oddsson, L., Olsson, E., Farén, E., Pettersson, A., and Ståhle, A. (2011). Effects of
new, individually adjusted, progressive balance group training for elderly people with fear of
falling and tend to fall: a randomized controlled trial. *Clin Rehabil* 25(11)**,** 1021-1031. doi:
10.1177/0269215511411937.

Hanewinckel, R., Drenthen, J., Verlinden, V.J.A., Darweesh, S.K.L., van der Geest, J.N., Hofman,
A., et al. (2017). Polyneuropathy relates to impairment in daily activities, worse gait, and fall-
related injuries. *Neurology* 89(1)**,** 76-83. doi: 10.1212/WNL.0000000000004067.

Hannan, M.T., Gagnon, M.M., Aneja, J., Jones, R.N., Cupples, L.A., Lipsitz, L.A., et al. (2010).
Optimizing the tracking of falls in studies of older participants: comparison of quarterly
telephone recall with monthly falls calendars in the MOBILIZE Boston Study. *Am J*
Epidemiol 171(9)**,** 1031-1036. doi: 10.1093/aje/kwq024.

Hardy, S.E., Perera, S., Roumani, Y.F., Chandler, J.M., and Studenski, S.A. (2007). Improvement in
usual gait speed predicts better survival in older adults. *J Am Geriatr Soc* 55(11)**,** 1727-1734.
doi: 10.1111/j.1532-5415.2007.01413.x.

Hegeman, J., Honegger, F., Kupper, M., and Allum, J.H. (2005). The balance control of bilateral
peripheral vestibular loss subjects and its improvement with auditory prosthetic feedback. *J*
Vestib Res 15(2)**,** 109-117.

Hiengkaew, V., Jitaree, K., and Chaiyawat, P. (2012). Minimal detectable changes of the Berg
Balance Scale, Fugl-Meyer Assessment Scale, Timed "Up & Go" Test, gait speeds, and 2-
minute walk test in individuals with chronic stroke with different degrees of ankle
plantarflexor tone. *Arch Phys Med Rehabil* 93(7)**,** 1201-1208. doi:
10.1016/j.apmr.2012.01.014.

Hlavacka, F., Krizková, M., and Horak, F.B. (1995). Modification of human postural response to leg
muscle vibration by electrical vestibular stimulation. *Neurosci Lett* 189(1), 9-12. doi:
10.1016/0304-3940(95)11436-z.
Hoffman, E.M., Staff, N.P., Robb, J.M., St Sauver, J.L., Dyck, P.J., and Klein, C.J. (2015).
Impairments and comorbidities of polyneuropathy revealed by population-based analyses.
Neurology 84(16), 1644-1651. doi: 10.1212/WNL.0000000000001492.
Ites, K.I., Anderson, E.J., Cahill, M.L., Kearney, J.A., Post, E.C., and Gilchrist, L.S. (2011). Balance
interventions for diabetic peripheral neuropathy: a systematic review. *J Geriatr Phys Ther*
34(3), 109-116. doi: 10.1519/JPT.0b013e318212659a.
Jónsdóttir, H.L., and Ruthig, J.C. (2020). A longitudinal study of the negative impact of falls on
health, well-being, and survival in later life: the protective role of perceived control. *Aging*
Ment Health, 1-7. doi: 10.1080/13607863.2020.1725736.
Kim, D.H., Glynn, R.J., Avorn, J., Lipsitz, L.A., Rockwood, K., Pawar, A., et al. (2019). Validation
of a Claims-Based Frailty Index Against Physical Performance and Adverse Health Outcomes
in the Health and Retirement Study. *J Gerontol A Biol Sci Med Sci* 74(8), 1271-1276. doi:
10.1093/gerona/gly197.
Koehler-McNicholas, S.R., Danzl, L., Cataldo, A.Y., and Oddsson, L.I.E. (2019). Neuromodulation
to improve gait and balance function using a sensory neuroprosthesis in people who report
insensate feet - A randomized control cross-over study. *PLoS One* 14(4), e0216212. doi:
10.1371/journal.pone.0216212.
Kroenke, K., Spitzer, R.L., and Williams, J.B. (2001). The PHQ-9: validity of a brief depression
severity measure. *J Gen Intern Med* 16(9), 606-613. doi: 10.1046/j.1525-
1497.2001.016009606.x.
Kruse, R.L., Lemaster, J.W., and Madsen, R.W. (2010). Fall and balance outcomes after an
intervention to promote leg strength, balance, and walking in people with diabetic peripheral
neuropathy: "feet first" randomized controlled trial. *Phys Ther* 90(11), 1568-1579. doi:
10.2522/ptj.20090362.
Kästenbauer, T., Sauseng, S., Brath, H., Abrahamian, H., and Irsigler, K. (2004). The value of the
Rydel-Seiffer tuning fork as a predictor of diabetic polyneuropathy compared with a
neurothesiometer. *Diabet Med* 21(6), 563-567. doi: 10.1111/j.1464-5491.2004.01205.x.
Lajoie, Y., and Gallagher, S.P. (2004). Predicting falls within the elderly community: comparison of
postural sway, reaction time, the Berg balance scale and the Activities-specific Balance
Confidence (ABC) scale for comparing fallers and non-fallers. *Arch Gerontol Geriatr* 38(1),
11-26.
Lakens, D. (2013). Calculating and reporting effect sizes to facilitate cumulative science: a practical
primer for t-tests and ANOVAs. *Front Psychol* 4, 863. doi: 10.3389/fpsyg.2013.00863.
Lawrence, R.H., Tennstedt, S.L., Kasten, L.E., Shih, J., Howland, J., and Jette, A.M. (1998).
Intensity and correlates of fear of falling and hurting oneself in the next year: baseline
findings from a Roybal Center fear of falling intervention. *J Aging Health* 10(3), 267-286.
doi: 10.1177/089826439801000301.
Leddy, A.L., Crowner, B.E., and Earhart, G.M. (2011). Functional gait assessment and balance
evaluation system test: reliability, validity, sensitivity, and specificity for identifying
individuals with Parkinson disease who fall. *Phys Ther* 91(1), 102-113. doi:
10.2522/ptj.20100113.
Lee, B.C., Thrasher, T.A., Fisher, S.P., and Layne, C.S. (2015). The effects of different sensory
augmentation on weight-shifting balance exercises in Parkinson's disease and healthy elderly
people: a proof-of-concept study. *J Neuroeng Rehabil* 12, 75. doi: 10.1186/s12984-015-0064-
y.

Leonard, G., Lapierre, Y., Chen, J.K., Wardini, R., Crane, J., and Ptito, A. (2017). Noninvasive
tongue stimulation combined with intensive cognitive and physical rehabilitation induces
neuroplastic changes in patients with multiple sclerosis: A multimodal neuroimaging study.
Mult Scler J Exp Transl Clin 3(1)**,** 2055217317690561. doi: 10.1177/2055217317690561.
Li, F., Harmer, P., Fisher, K.J., and McAuley, E. (2004). Tai Chi: improving functional balance and
predicting subsequent falls in older persons. *Med Sci Sports Exerc* 36(12)**,** 2046-2052.
Li, F., Harmer, P., Fisher, K.J., McAuley, E., Chaumeton, N., Eckstrom, E., et al. (2005). Tai Chi and
fall reductions in older adults: a randomized controlled trial. *J Gerontol A Biol Sci Med Sci*
60(2)**,** 187-194. doi: 10.1093/gerona/60.2.187.
Li, L., and Manor, B. (2010). Long term Tai Chi exercise improves physical performance among
people with peripheral neuropathy. *Am J Chin Med* 38(3)**,** 449-459. doi:
10.1142/S0192415X1000797X.
Lin, J.H., Hsu, M.J., Hsu, H.W., Wu, H.C., and Hsieh, C.L. (2010). Psychometric comparisons of 3
functional ambulation measures for patients with stroke. *Stroke* 41(9)**,** 2021-2025. doi:
10.1161/STROKEAHA.110.589739.
Lipsitz, L.A., Lough, M., Niemi, J., Travison, T., Howlett, H., and Manor, B. (2015). A shoe insole
delivering subsensory vibratory noise improves balance and gait in healthy elderly people.
Arch Phys Med Rehabil 96(3)**,** 432-439. doi: 10.1016/j.apmr.2014.10.004.
Lipsitz, L.A., Macklin, E.A., Travison, T.G., Manor, B., Gagnon, P., Tsai, T., et al. (2019). A Cluster
Randomized Trial of Tai Chi vs Health Education in Subsidized Housing: The MI-WiSH
Study. *J Am Geriatr Soc* 67(9)**,** 1812-1819. doi: 10.1111/jgs.15986.
Lipsitz, L.A., Manor, B., Habtemariam, D., Iloputaife, I., Zhou, J., and Travison, T.G. (2018). The
pace and prognosis of peripheral sensory loss in advanced age: association with gait speed
and falls. *BMC Geriatr* 18(1)**,** 274. doi: 10.1186/s12877-018-0970-5.
Manor, B., Lough, M., Gagnon, M.M., Cupples, A., Wayne, P.M., and Lipsitz, L.A. (2014).
Functional benefits of tai chi training in senior housing facilities. *J Am Geriatr Soc* 62(8)**,**
1484-1489. doi: 10.1111/jgs.12946.
Marsden, J.F., Blakey, G., and Day, B.L. (2003). Modulation of human vestibular-evoked postural
responses by alterations in load. *J Physiol* 548(Pt 3)**,** 949-953. doi:
10.1113/jphysiol.2002.029991.
Mathias, S., Nayak, U.S., and Isaacs, B. (1986). Balance in elderly patients: the "get-up and go" test.
Arch Phys Med Rehabil 67(6)**,** 387-389.
Melzer, I., and Oddsson, L.I.e. (2013). Improving balance control and self-reported lower extremity
function in community-dwelling older adults: a randomized control trial. *Clin Rehabil* 27(3)**,**
195-206. doi: 10.1177/0269215512450295.
Menz, H.B., Lord, S.R., St George, R., and Fitzpatrick, R.C. (2004). Walking stability and
sensorimotor function in older people with diabetic peripheral neuropathy. *Arch Phys Med*
Rehabil 85(2)**,** 245-252. doi: 10.1016/j.apmr.2003.06.015.
Meyer, P.F., Oddsson, L.I., and De Luca, C.J. (2004a). Reduced plantar sensitivity alters postural
responses to lateral perturbations of balance. *Exp Brain Res* 157(4)**,** 526-536. doi:
10.1007/s00221-004-1868-3.
Meyer, P.F., Oddsson, L.I., and De Luca, C.J. (2004b). The role of plantar cutaneous sensation in
unperturbed stance. *Exp Brain Res* 156(4)**,** 505-512. doi: 10.1007/s00221-003-1804-y.
Middleton, A., Fritz, S.L., and Lusardi, M. (2015). Walking speed: the functional vital sign. *J Aging*
Phys Act 23(2)**,** 314-322. doi: 10.1123/japa.2013-0236.
Montero-Odasso, M., Schapira, M., Soriano, E.R., Varela, M., Kaplan, R., Camera, L.A., et al.
(2005). Gait velocity as a single predictor of adverse events in healthy seniors aged 75 years
and older. *J Gerontol A Biol Sci Med Sci* 60(10)**,** 1304-1309. doi: 10.1093/gerona/60.10.1304.

Moore, J.L., Potter, K., Blankshain, K., Kaplan, S.L., O'Dwyer, L.C., and Sullivan, J.E. (2018). A
Core Set of Outcome Measures for Adults With Neurologic Conditions Undergoing
Rehabilitation: A CLINICAL PRACTICE GUIDELINE. *J Neurol Phys Ther* 42(3)**,** 174-220.
doi: 10.1097/NPT.0000000000000229.
Morrison, S., Colberg, S.R., Mariano, M., Parson, H.K., and Vinik, A.I. (2010). Balance training
reduces falls risk in older individuals with type 2 diabetes. *Diabetes Care* 33(4)**,** 748-750. doi:
10.2337/dc09-1699.
Morrison, S., Simmons, R., Colberg, S.R., Parson, H.K., and Vinik, A.I. (2018). Supervised Balance
Training and Wii Fit-Based Exercises Lower Falls Risk in Older Adults With Type 2
Diabetes. *J Am Med Dir Assoc* 19(2)**,** 185.e187-185.e113. doi: 10.1016/j.jamda.2017.11.004.
Mustapa, A., Justine, M., Mohd Mustafah, N., Jamil, N., and Manaf, H. (2016). Postural Control and
Gait Performance in the Diabetic Peripheral Neuropathy: A Systematic Review. *Biomed Res*
Int 2016**,** 9305025. doi: 10.1155/2016/9305025.
National Heart, L.a.B.I. (2020). *Study Quality Assessment Tools* [Online]. Available:
https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools [Accessed 2020].
Oddsson, L., Bisson, T., Cohen, H., Iloputaife, I., Koehler-McNicholas, S., Kung, D., et al. (2020). A
Wearable Sensory Neuroprosthesis to Improve Gait and Balance Function in Persons with
Sensory Peripheral Neuropathy – the walk2Wellness Trial (1170). *Neurology* 94(15
Supplement)**,** 1170.
Oddsson, L., Bisson, T., Cohen, H., Koehler-McNicholas, S., Kung, D., and Wrisley, D. (2019).
"Slips and Falls and Record Snow in Minnesota: walk2wellness Clinical Trial Update", in:
AAPM&R Annual Assembly. (San Antonio, Texas).
Oddsson, L.I.E., Boissy, P., and Melzer, I. (2007). How to improve gait and balance function in
elderly individuals—compliance with principles of training. *Eur Rev Aging Phys Act* Volume
4(Issue 1)**,** 15-23. doi: https://doi.org/10.1007/s11556-007-0019-9.
Perera, S., Mody, S.H., Woodman, R.C., and Studenski, S.A. (2006). Meaningful change and
responsiveness in common physical performance measures in older adults. *J Am Geriatr Soc*
54(5)**,** 743-749. doi: 10.1111/j.1532-5415.2006.00701.x.
Podsiadlo, D., and Richardson, S. (1991). The timed "Up & Go": a test of basic functional mobility
for frail elderly persons. *J Am Geriatr Soc* 39(2)**,** 142-148. doi: 10.1111/j.1532-
5415.1991.tb01616.x.
Powell, L.E., and Myers, A.M. (1995). The Activities-specific Balance Confidence (ABC) Scale. *J*
Gerontol A Biol Sci Med Sci 50A(1)**,** M28-34.
Ptito, A., Papa, L., Gregory, K., Folmer, R.L., Walker, W.C., Prabhakaran, V., et al. (2020). A
Prospective, Multicenter Study to Assess the Safety and Efficacy of Translingual
Neurostimulation Plus Physical Therapy for the Treatment of a Chronic Balance Deficit Due
to Mild-to-Moderate Traumatic Brain Injury. *Neuromodulation*. doi: 10.1111/ner.13159.
Purser, J.L., Weinberger, M., Cohen, H.J., Pieper, C.F., Morey, M.C., Li, T., et al. (2005). Walking
speed predicts health status and hospital costs for frail elderly male veterans. *J Rehabil Res*
Dev 42(4)**,** 535-546. doi: 10.1682/jrrd.2004.07.0087.
Quigley, P.A., Bulat, T., Schulz, B., Friedman, Y., Hart-Hughes, S., Richardson, J.K., et al. (2014).
Exercise interventions, gait, and balance in older subjects with distal symmetric
polyneuropathy: a three-group randomized clinical trial. *Am J Phys Med Rehabil* 93(1)**,** 1-12;
quiz 13-16. doi: 10.1097/PHM.0000000000000052.
Richardson, J.K., and Hurvitz, E.A. (1995). Peripheral neuropathy: a true risk factor for falls. *J*
Gerontol A Biol Sci Med Sci 50(4)**,** M211-215.

Richardson, J.K., Sandman, D., and Vela, S. (2001). A focused exercise regimen improves clinical measures of balance in patients with peripheral neuropathy. *Arch Phys Med Rehabil* 82(2)**,** 205-209. doi: 10.1053/apmr.2001.19742.

Rief, W., Nestoriuc, Y., Weiss, S., Welzel, E., Barsky, A.J., and Hofmann, S.G. (2009). Meta-analysis of the placebo response in antidepressant trials. *J Affect Disord* 118(1-3)**,** 1-8. doi: 10.1016/j.jad.2009.01.029.

Riskowski, J.L., Quach, L., Manor, B., Menz, H.B., Lipsitz, L.A., and Hannan, M.T. (2012). Idiopathic peripheral neuropathy increases fall risk in a population-based cohort study of older adults. *Journal of Foot and Ankle Research* 5(1)**,** P19. doi: 10.1186/1757-1146-5-S1-P19.

Rossi-Izquierdo, M., Ernst, A., Soto-Varela, A., Santos-Pérez, S., Faraldo-García, A., Sesar-Ignacio, A., et al. (2013). Vibrotactile neurofeedback balance training in patients with Parkinson's disease: reducing the number of falls. *Gait Posture* 37(2)**,** 195-200. doi: 10.1016/j.gaitpost.2012.07.002.

Salsabili, H., Bahrpeyma, F., and Esteki, A. (2015). The effects of Task-Oriented Motor Training on gait characteristics of patients with type 2 diabetes neuropathy. *J Diabetes Metab Disord* 15**,** 14. doi: 10.1186/s40200-016-0236-8.

Scheffer, A.C., Schuurmans, M.J., van Dijk, N., van der Hooft, T., and de Rooij, S.E. (2008). Fear of falling: measurement strategy, prevalence, risk factors and consequences among older persons. *Age Ageing* 37(1)**,** 19-24. doi: 10.1093/ageing/afm169.

Shumway-Cook, A., Brauer, S., and Woollacott, M. (2000). Predicting the probability for falls in community-dwelling older adults using the Timed Up & Go Test. *Phys Ther* 80(9)**,** 896-903.

Shumway-Cook, A., Gruber, W., Baldwin, M., and Liao, S. (1997). The effect of multidimensional exercises on balance, mobility, and fall risk in community-dwelling older adults. *Phys Ther* 77(1)**,** 46-57. doi: 10.1093/ptj/77.1.46.

Sterling, D.A., O'Connor, J.A., and Bonadies, J. (2001). Geriatric falls: injury severity is high and disproportionate to mechanism. *J Trauma* 50(1)**,** 116-119. doi: 10.1097/00005373-200101000-00021.

Streckmann, F., Zopf, E.M., Lehmann, H.C., May, K., Rizza, J., Zimmer, P., et al. (2014). Exercise intervention studies in patients with peripheral neuropathy: a systematic review. *Sports Med* 44(9)**,** 1289-1304. doi: 10.1007/s40279-014-0207-5.

Strzalkowski, N.D.J., Peters, R.M., Inglis, J.T., and Bent, L.R. (2018). Cutaneous afferent innervation of the human foot sole: what can we learn from single-unit recordings? *J Neurophysiol* 120(3)**,** 1233-1246. doi: 10.1152/jn.00848.2017.

Studenski, S., Perera, S., Patel, K., Rosano, C., Faulkner, K., Inzitari, M., et al. (2011). Gait speed and survival in older adults. *JAMA* 305(1)**,** 50-58. doi: 10.1001/jama.2010.1923.

Studenski, S., Perera, S., Wallace, D., Chandler, J.M., Duncan, P.W., Rooney, E., et al. (2003). Physical performance measures in the clinical setting. *J Am Geriatr Soc* 51(3)**,** 314-322. doi: 10.1046/j.1532-5415.2003.51104.x.

Tinetti, M.E., and Kumar, C. (2010). The patient who falls: "It's always a trade-off". *JAMA* 303(3)**,** 258-266. doi: 10.1001/jama.2009.2024.

Tofthagen, C., Visovsky, C., and Berry, D.L. (2012). Strength and balance training for adults with peripheral neuropathy and high risk of fall: current evidence and implications for future research. *Oncol Nurs Forum* 39(5)**,** E416-424. doi: 10.1188/12.ONF.E416-E424.

Tyler, M., Danilov, Y., and Bach-Y-Rita, P. (2003). Closing an open-loop control system: vestibular substitution through the tongue. *J Integr Neurosci* 2(2)**,** 159-164.

Uitenbroek, D.G. (1997). *SISA-Binomial* [Online]. Available: https://www.quantitativeskills.com/sisa/calculations/bonfer.htm [Accessed May 2020].

Vinik, A.I., Camacho, P., Reddy, S., Valencia, W.M., Trence, D., Matsumoto, A.M., et al. (2017). AGING, DIABETES, AND FALLS. *Endocr Pract* 23(9)**,** 1117-1139. doi: 10.4158/EP171794.RA.

Wall, C., Wrisley, D., and Oddsson, L. (2012). Vibrotactile feedback of mediolateral trunk tilt or foot pressure increases locomotor performance in healthy older adults--a pilot study. *Conf Proc IEEE Eng Med Biol Soc* 2012**,** 6145-6148. doi: 10.1109/EMBC.2012.6347396.

Wall, C., Wrisley, D.M., and Statler, K.D. (2009). Vibrotactile tilt feedback improves dynamic gait index: a fall risk indicator in older adults. *Gait Posture* 30(1)**,** 16-21. doi: 10.1016/j.gaitpost.2009.02.019.

Wang, C., Goel, R., Rahemi, H., Zhang, Q., Lepow, B., and Najafi, B. (2019a). Effectiveness of Daily Use of Bilateral Custom-Made Ankle-Foot Orthoses on Balance, Fear of Falling, and Physical Activity in Older Adults: A Randomized Controlled Trial. *Gerontology* 65(3)**,** 299-307. doi: 10.1159/000494114.

Wang, C., Goel, R., Zhang, Q., Lepow, B., and Najafi, B. (2019b). Daily Use of Bilateral Custom-Made Ankle-Foot Orthoses for Fall Prevention in Older Adults: A Randomized Controlled Trial. *J Am Geriatr Soc* 67(8)**,** 1656-1661. doi: 10.1111/jgs.15929.

Wartolowska, K.A., Feakins, B.G., Collins, G.S., Cook, J., Judge, A., Rombach, I., et al. (2016). The magnitude and temporal changes of response in the placebo arm of surgical randomized controlled trials: a systematic review and meta-analysis. *Trials* 17(1)**,** 589. doi: 10.1186/s13063-016-1720-7.

Winters-Stone, K.M., Horak, F., Jacobs, P.G., Trubowitz, P., Dieckmann, N.F., Stoyles, S., et al. (2017). Falls, Functioning, and Disability Among Women With Persistent Symptoms of Chemotherapy-Induced Peripheral Neuropathy. *J Clin Oncol* 35(23)**,** 2604-2612. doi: 10.1200/JCO.2016.71.3552.

Wolf, B., Feys, H., De Weerdt, van der Meer, J., Noom, M., and Aufdemkampe, G. (2001). Effect of a physical therapeutic intervention for balance problems in the elderly: a single-blind, randomized, controlled multicentre trial. *Clin Rehabil* 15(6)**,** 624-636. doi: 10.1191/0269215501cr456oa.

Wolf, S.L., Barnhart, H.X., Kutner, N.G., McNeely, E., Coogler, C., and Xu, T. (1996). Reducing frailty and falls in older persons: an investigation of Tai Chi and computerized balance training. Atlanta FICSIT Group. Frailty and Injuries: Cooperative Studies of Intervention Techniques. *J Am Geriatr Soc* 44(5)**,** 489-497. doi: 10.1111/j.1532-5415.1996.tb01432.x.

Wolf, S.L., Sattin, R.W., Kutner, M., O'Grady, M., Greenspan, A.I., and Gregor, R.J. (2003). Intense tai chi exercise training and fall occurrences in older, transitionally frail adults: a randomized, controlled trial. *J Am Geriatr Soc* 51(12)**,** 1693-1701. doi: 10.1046/j.1532-5415.2003.51552.x.

Woolcott, J.C., Richardson, K.J., Wiens, M.O., Patel, B., Marin, J., Khan, K.M., et al. (2009). Meta-analysis of the impact of 9 medication classes on falls in elderly persons. *Arch Intern Med* 169(21)**,** 1952-1960. doi: 10.1001/archinternmed.2009.357.

Wrisley, D., McLean, G., and Lars, O. (Year). "The Use of a Wearable Sensory Prosthesis to Improve Gait and Balance in a Patient with Peripheral Neuropathy", in: *APTA Combined Sections Meeting*), #3196.

Wrisley, D.M., and Kumar, N.A. (2010). Functional gait assessment: concurrent, discriminative, and predictive validity in community-dwelling older adults. *Phys Ther* 90(5)**,** 761-773. doi: 10.2522/ptj.20090069.

Wrisley, D.M., Marchetti, G.F., Kuharsky, D.K., and Whitney, S.L. (2004). Reliability, internal consistency, and validity of data obtained with the functional gait assessment. *Phys Ther* 84(10)**,** 906-918.

Yamanaka, T., Sawai, Y., Murai, T., Nishimura, T., and Kitahara, T. (2016). Long-term effects of
electrotactile sensory substitution therapy on balance disorders. *Neuroreport* 27(10)**,** 744-748.
doi: 10.1097/WNR.0000000000000606.
Zaiontz, C. (2020). *Real Statistics Resource Pack (Release 6.8)* [Online]. Available:
http://www.real-statistics.com/ [Accessed].
Zehr, E.P., Nakajima, T., Barss, T., Klarner, T., Miklosovic, S., Mezzarane, R.A., et al. (2014).
Cutaneous stimulation of discrete regions of the sole during locomotion produces "sensory
steering" of the foot. *BMC Sports Sci Med Rehabil* 6**,** 33. doi: 10.1186/2052-1847-6-33.
Zehr, E.P., and Stein, R.B. (1999). What functions do reflexes serve during human locomotion? *Prog*
Neurobiol 58(2)**,** 185-205. doi: 10.1016/s0301-0082(98)00081-1.
1249