Gender disparities in international COVID-19 clinical trial leadership

Authors: Muge Cevik[1,2], Syed Arefinul Haque[3], Jennifer Manne-Goehler[4], Krutika Kuppalli[5], Paul

E. Sax[4,6], Maimuna S. Majumder[7], Chloe Orkin[8,9]

Affiliations:

1. Division of Infection and Global Health Research, School of Medicine, University of St Andrews, United

Kingdom

2. NHS Lothian Infection Service, Regional Infectious Diseases Unit, Western General Hospital,

Edinburgh, United Kingdom

3. Network Science Institute, Northeastern University, Boston, Massachusetts, United States

4. Division of Infectious Diseases, Brigham and Women's Hospital, Boston, Massachusetts, United States

5. Division of Infectious Diseases, Medical University of South Carolina, Charleston, South Carolina,

United States

6. Harvard Medical School, Boston, Massachusetts, United States

7. Boston Children's Hospital, Harvard Medical School, Boston, MA, United States

8. Blizzard Institute, Queen Mary University of London, London, United Kingdom

9. Royal London Hospital, Barts Health NHS Trust, London, United Kingdom

Author of correspondence:

Name: Dr Muge Cevik

Address: Division of Infection and Global Health Research, School of Medicine, University of St Andrews,

Fife, KY16 9TF

Telephone number: +447732800814

Email address: mc349@st-andrews.ac.uk

Abstract

The COVID-19 pandemic offers considerable possibilities for research and leadership that might equalize opportunity in a new field; however, our study finds instead that more than two-thirds of investigators leading COVID-19-related clinical studies are predicted to be men. These gender disparities in trial leadership during the pandemic suggest that the structural reproduction of inequalities in research has taken place once again in this new academic field. This indicates that policies are needed to facilitate the identification and implementation of strategies to correct gender bias. The active participation of women, trans and gender-nonconforming individuals are needed in research to drive scientific discovery and innovation as well as to better address health disparities.

Introduction

In addition to the human and financial loss associated with the novel Coronavirus Disease 2019 (COVID-19) pandemic, COVID-19 has also had a significant impact on both the personal and professional life of the global workforce, including that of the scientific research community [1-3]. Before COVID-19, women occupied fewer leadership positions, led a fewer funded studies, and applied for and received less grant funding than men when they did apply [4-7]. The employment gap that occurs when women take parental leave impacts the rate of academic advancement and in turn the receipt of institutional support to apply for and secure funding [6, 7]. These imbalances contribute to systemic inequalities that hamper women's access to and progress in science [2, 7, 8]. A review of the gender distribution of 24 COVID-19 national task forces suggests that many committees are comprised of less than a quarter women, indicating that women's voices and expertise have been excluded from decision making during this unprecedented public health emergency [9].

For example, emerging data suggest that across all disciplines, despite an increased number of peer-reviewed articles submitted to journals during the pandemic, women have published fewer papers than men thus far this year [10]. This may indicate a similarly reduced involvement of women in research leadership positions and an imbalanced distribution of grants and funding -- important indicators of advancement in a scientist's academic career [4-7, 10, 11]. Being principal investigator (PI) on a clinical trial is strongly associated with advancement to full professor among women academics in infectious diseases [8].

The COVID-19 pandemic offers numerous opportunities in clinical research. These include trials to assess the safety and efficacy of medical interventions, with protocols in various stages of implementation. Here, we provide a timely analysis to compare the gender distribution of clinical trial leadership in COVID-19 clinical trials.

Materials and Methods

We systematically searched https://clinicaltrials.gov/ and retrieved all clinical trials on COVID-19 registered from January 1, 2020 to June 26, 2020 using "COVID" as a keyword. As a comparator group, we have chosen two fields that are not related to emerging infections: breast cancer and type 2 diabetes mellitus (T2DM). We retrieved all clinical trials related to these comparator conditions registered at

https://clinicaltrials.gov/ within the aforementioned study period as well as those registered in the

preceding year (pre-study period: January 1, 2019 and December 31, 2019). Gender of the investigator

was predicted using the genderize.io API (application programming interface). This tool has been used to

predict the gender of first names in studies regarding gender bias [12, 13] and achieves a minimum

accuracy of 82%, with an F1 score of 90% for women and 86% for men [14]. Clinical trials were excluded

if i) investigator information was not provided; ii) the genderize.io API could not predict any of the

investigators' gender from their first name; or iii) organization or company names were provided as the

investigator. The number of studies that were excluded for the above reasons are reported in the

supplementary flow diagram. An exploratory temporal analysis was conducted with the available data.

Categorical variables were summarized by frequencies and percentages. We compared groups using Chi-

square testing for equality of proportions with continuity correction.[15] The analysis was performed

using R (Version 4.0.2). The repository of the datasets used to collect and analyse the data available at

https://osf.io/k2r57/.

Results

We identified 2 345 COVID-19-related clinical trials. Of those, 1 448 had at least one investigator (i.e.,

principal investigator, study director, or study chair) identified whose gender could be predicted. In the

comparator group, we identified 449 trials on breast cancer and 272 on T2DM that were registered. Of

those, 274 breast cancer studies and 139 T2DM studies had at least one investigator whose gender

could be predicted.

Overall 27.8% of PIs among COVID-19-related studies were predicted to be women, which is significantly

different compared to 54.4% and 42.1% for breast cancer ($p<0.005$) and T2DM ($p<0.005$) trials over the

same period, respectively (Table 1). While there has been a small increase in the proportion of PIs who

were predicted to be women in May 2020, clinical research leadership for COVID-19 among this group

was below 25% for the remainder of the study period (Supplementary Material). While 31.4% of study

chairs were predicted to be women in COVID-19-related studies, 32.1% ($p=0.7$) and 63.6% ($p<0.01$) were

predicted to be women in breast cancer and T2DM trials, respectively. Proportion of study chairs were

not significantly different across the three fields.

We also reviewed comparator group studies registered before January 1, 2020 to determine whether

the pandemic might have affected gender distribution of trial leadership. We identified 839 clinical trials

related to breast cancer and 533 on T2DM over a 12-month period prior to January 1, 2020. Of those,

573 breast cancer studies and 359 T2DM studies yielded at least one investigator whose gender could

be predicted. During this "pre-study" period, the proportion of PIs who were predicted to be women

were 49.7% and 44.4% for breast cancer and T2DM trials, respectively and the difference was not

statistically significant when compared to results from the study period (p>0.05).

Discussion

In this study, we demonstrate that less than one-third of COVID-19-related clinical trials are led by PIs

who were predicted to be women, half the proportion observed in non-COVID-19 (breast cancer and

T2DM) trials over the same period. The proportion of PIs in breast cancer and T2DM studies also

remained similar to the pre-study period. These gender disparities during the pandemic may indicate

not only a lack of women's leadership in international clinical trials and involvement in new projects, but

also may reveal imbalances in women's access to research activities and funding during health

emergencies [2, 16].

The COVID-19 pandemic offers numerous opportunities for research and leadership that could equalize

opportunity in a new field, but our results suggest the opposite. The pandemic has reinforced the

prevailing gender norms in which men continue to be allocated a disproportionate share of the funding,

as well as leadership and authorship roles [9, 10, 16]. One potential contributor for this discrepancy is

the speed demanded by the research agenda during the pandemic. The sense of urgency in starting

clinical trials may lead to an abandonment of any checks and balances around equality and inclusion

that would have otherwise encouraged the involvement of women scientists. Many women scientists

have already raised concerns about institutional funding distribution lacking gender balance or being left

out of research activities despite their expertise [2, 16]. During COVID-19 pandemic, a UK study showed

that women were more than twice as likely to take on childcare and schooling responsibilities of

children than men, while male academic counterparts leverage professional relationships and networks

more effectively [1, 2, 16].

As a community, we must recognise that there is a tendency to "turn to men" in times of crisis both for

leadership and scientific expertise [2, 3, 16, 17], highlighting the need to challenge this culture. Research

and academia are already competitive; being in the central decision-making group is often challenging

due to gender norms, along with roles and rules on how these groups are established and maintained;

during health emergencies, these same authoritative circles become more difficult for women scientists

to join [2, 16]. Our findings suggest that there is a need for transparency in opportunities and funding

that requires actively identifying and addressing the structurally implicit and unconscious biases that

favour men.

Our analysis has some limitations. We could include only ~50-75% of trials for which an investigator's

gender could be algorithmically predicted. Furthermore, while such algorithms allow for the rapid

analysis of gender disparities such as those conducted here, they can also be exclusionary to gender

non-conforming, non-binary, and trans individuals. Because of this, we have taken care to use the term

"predicted gender" throughout the text whenever discussing algorithm-based gender assignment in

acknowledgement of gender non-conforming, non-binary, and trans investigators. Beyond these

limitations, we did not consider studies that received private funding, which may not have been

registered on clinicaltrials.gov; however, it is worth noting that clinicaltrials.gov is an international

database with widespread international representation.

In summary, while the COVID-19 pandemic has thus far provided many new opportunities for research,

with numerous clinical trials initiated worldwide, a disproportionate proportion of PIs leading COVID-19

related studies are predicted to be men, despite women accounting for 70% of the global health

workforce [16]. Our demonstration of gender differences in trial leadership and grant allocation argue

for revised policies and strategies that encourage the participation of women in pandemic research. Not

only can these women drive discovery and innovation, but they can act to address health disparities and

provide role models for the next generation of women scientists [2, 16, 18, 19]. Ensuring gender

representation would also reflect the commitment of the global community to promoting gender

equality in academic medicine and research: inclusion, diversity, representation, progression, and

success for all. A health emergency is not an excuse for disregarding the importance of equitable

representation and given the disproportionate burden of such outbreaks for communities who are

marginalized due to their gender, sexuality, class, ethnicity, and ability an even greater reason we must

advocate for it [20-22].

Authors contributions

MC: conceptualisation, methodology, investigation, literature review, data curation, writing – original draft. SH and MM: investigation, data curation, formal analysis, writing – review and editing; JS, KK, PS: methodology, writing – review and editing, supervision. CO: conceptualisation, methodology, investigation, literature review, writing – original draft, supervision.

Conflict of interests

MC, SH, JS, KK, MM have none to disclose. CO has received honoraria, fees for lectures, and advisory boards from Gilead, MSD, Viiv, and Janssen. She has also received research grants to her institution from the above-mentioned companies. PES has received honoraria, fees for lectures, and advisory boards from Gilead, Merck, Janssen, and ViiV; he has also received research grants to his institution from Gilead and ViiV.

Table 1: Proportion of women leadership in clinical trials between January 1, 2020 and June 26, 2020 and before January 1, 2020

	Jan 1, 2020 - June 26, 2020					before Jan 1, 2020		
	COVID -19	Breast Cancer	p value	T2DM	p value	Breast Cancer	T2DM	p value
PI	27.8%	54.4%	<0.001	42.1%	< 0.001	49.7%	44.4%	0.158
Study Director	28.7%	48.9%	0.0103	22.2%	0.751	30.5%	47.6%	0.028
Study Chair	31.4%	32.1%	1	36.36%	0.98	33.3%	40.4%	0.54

Supplementary material:

Flow diagram of process of selection

Gender distribution over time (months)

References:

1.   Villadsen A, Conti, G. and Fitzsimons, E. Parental involvement in home schooling and developmental play during lockdown - Initial findings from the COVID-19 Survey in Five National Longitudinal Studies. . London: UCL Centre for Longitudinal Studies. 2020.

2.   Education TH. Women in science are battling both Covid-19 and the patriarchy 2020 [cited 2020 June 6]. Available from: https://www.timeshighereducation.com/blog/women-science-are-battling-both-covid-19-and-patriarchy#survey-answer.

3.   Brubaker L. Women Physicians and the COVID-19 Pandemic. JAMA. 2020. doi: 10.1001/jama.2020.14797.

4.   Head MG, Fitchett JR, Cooke MK, Wurie FB, Atun R. Sex discrepancies in infectious disease research funding 1997–2010: a systematic analysis. The Lancet. 2013;382:S44. doi: 10.1016/S0140-6736(13)62469-1.

5.   Hoa NT, Thuong NTT, Clapham HE, Thu TTA, Kestelyn E, Thwaites CL. Increasing women's leadership in science in Ho Chi Minh City. The Lancet. 2019;393(10171):523-4. doi: 10.1016/S0140-6736(18)32090-7.

6.   Blake M LVI. Who Applies for Research Funding?

7.   Jefferson L, Bloor K, Maynard A. Women in medicine: historical perspectives and recent
trends. British Medical Bulletin. 2015;114(1):5-15. doi: 10.1093/bmb/ldv007.
8.   Manne-Goehler J, Kapoor N, Blumenthal DM, Stead W. Sex Differences in Achievement
and Faculty Rank in Academic Infectious Diseases. Clinical Infectious Diseases.
2019;70(2):290-6. doi: 10.1093/cid/ciz200.
9.   Rajan D, Koch K, Rohrer K, Bajnoczki C, Socha A, Voss M, et al. Governance of the
Covid-19 response: a call for more inclusive and transparent decision-making. BMJ Global
Health. 2020;5(5):e002655. doi: 10.1136/bmjgh-2020-002655.
10.   Andersen JP, Nielsen MW, Simone NL, Lewiss RE, Jagsi R. COVID-19 medical papers
have fewer women first authors than expected. eLife. 2020;9:e58807. doi: 10.7554/eLife.58807.
11.   D'Armiento J, Witte SS, Dutt K, Wall M, McAllister G. Achieving women's equity in
academic medicine: challenging the standards. The Lancet. 2019;393(10171):e15-e6. doi:
10.1016/S0140-6736(19)30234-X.
12.   Holman L, Stuart-Fox D, Hauser CE. The gender gap in science: How long until women
are equally represented? PLoS Biol. 2018;16(4):e2004956. Epub 2018/04/20. doi:
10.1371/journal.pbio.2004956. PubMed PMID: 29672508; PubMed Central PMCID:
PMCPMC5908072.
13.   Huang J, Gates AJ, Sinatra R, Barabasi AL. Historical comparison of gender inequality in
scientific careers across countries and disciplines. Proc Natl Acad Sci U S A. 2020;117(9):4609-
16. Epub 2020/02/20. doi: 10.1073/pnas.1914221117. PubMed PMID: 32071248; PubMed
Central PMCID: PMCPMC7060730.
14.   Karimi F WC, Lemmerich F, Jadidi M, Strohmaier M. . Inferring gender from names on
the web: a comparative evaluation of gender detection meth- ods. Proceedings of the 25th
international conference companion on world wide web, WWW'16 companion. 2016.
15.   Newcombe RG. Interval estimation for the difference between independent proportions:
comparison of eleven methods. Stat Med. 1998;17(8):873-90. Epub 1998/05/22. doi:
10.1002/(sici)1097-0258(19980430)17:8<873::aid-sim779>3.0.co;2-i. PubMed PMID: 9595617.
16.   Gabster BP, van Daalen K, Dhatt R, Barry M. Challenges for the female academic during
the COVID-19 pandemic. Lancet. 2020;395(10242):1968-70. Epub 2020/06/22. doi:
10.1016/S0140-6736(20)31412-4. PubMed PMID: 32563275; PubMed Central PMCID:
PMCPMC7302767.
17.   WEP. Women Expert Project 2020. Available from: https://expertwomenproject.com.
18.   Nielsen MW, Bloch CW, Schiebinger L. Making gender diversity work for scientific
discovery and innovation. Nature Human Behaviour. 2018;2(10):726-34. doi: 10.1038/s41562-
018-0433-1.
19.   A call for more inclusive Covid-19 research: Tackling disparities during a pandemic
2020. Available from: https://www.statnews.com/sponsor/2020/05/22/a-call-for-more-inclusive-
covid-19-research-tackling-disparities-during-a-pandemic/.
20.   Koma W, Artiga, S., Neuman, T., Claxton, G.,Rae,M.,Kates,J.,Michaud, J. . Low-Income
and Communities of Color at Higher Risk of Serious Illness if Infected with Coronavirus
2020 [cited 2020 July 21]. Available from: https://www.kff.org/coronavirus-covid-19/issue-
brief/low-income-and-communities-of-color-at-higher-risk-of-serious-illness-if-infected-with-
coronavirus/.
21.   Finch WH, Hernández Finch ME. Poverty and Covid-19: Rates of Incidence and Deaths
in the United States During the First 10 Weeks of the Pandemic. Frontiers in Sociology.
2020;5(47). doi: 10.3389/fsoc.2020.00047.

22.   Chen J, Krieger, N. Revealing the unequal burden of COVID-19 by income,
race/ethnicity, and household crowding: US county vs. ZIP code analyses: Harvard Center for
Population and Development Studies; 2020 [cited 2020 July 26]. Available from:
https://cdn1.sph.harvard.edu/wp-content/uploads/sites/1266/2020/04/HCPDS_Volume-
19_No_1_20_covid19_RevealingUnequalBurden_HCPDSWorkingPaper_04212020-1.pdf.
300