Regional ventilation characteristics during non-invasive respiratory support in preterm infants

Thomson J[1,2], Ru□egger CM[3,4], Perkins, EJ[1], Pereira-Fantini PM[1], Farrell O[1,2], Owen LS[1,3,5], Tingay DG[1,2,3,6]

[1]Neonatal Research, Murdoch Children's Research Institute, Parkville, Australia

[2]Department of Paediatrics, The University of Melbourne, Parkville, Australia

[3]Neonatal Research, The Royal Women's Hospital, Parkville, Australia

[4]Newborn Research, Department of Neonatology, University Hospital and University of Zurich, Zurich, Switzerland

[5]Department of Obstetrics and Gynaecology, The University of Melbourne, Parkville, Australia

[6]Department of Neonatology, The Royal Children's Hospital, Parkville, Australia

Corresponding Author:

Jessica Thomson

Neonatal Research

Murdoch Children's Research Institute

50 Flemington Rd.

Parkville 3052 Victoria

jthomson2@student.unimelb.edu.au

Running title: Regional ventilation during NIV

Keywords: Preterm; Infant; Regional ventilation; non-invasive ventilation

Manuscript word count: 2500 words

ABSTRACT

Objective: To determine the regional ventilation characteristics during non-invasive ventilation in stable preterm infants. The secondary aims were to explore the relationship between indicators of ventilation homogeneity and other clinical measures of respiratory status.

Design: Prospective observational study.

Setting: Two tertiary neonatal intensive care units.

Patients: Forty stable preterm infants born <30 weeks gestation receiving either CPAP (n=32) or nHF(n=8) if <36 weeks corrected gestation, and extubated for at least 24 hours at time of study.

Interventions: Continuous electrical impedance tomography imaging of regional ventilation during 60-minutes of quiet breathing on clinician-determined non-invasive settings.

Main outcome measures: Gravity-dependent and right-left centre of ventilation (CoV), percentage of whole lung tidal volume by lung region, and percentage of lung unventilated were determined for 120 artefact-free breaths/infant (4770 breaths included). Oxygen saturation, heart and respiratory rates were also measured.

Results: Ventilation was greater in the right lung (mean (SD) CoV_{RL} 69.1 (14.9)%) and the gravity non-dependent lung; ideal-actual CoV 1.4 (4.5). The central third of the lung received the most tidal volume, followed by the non-dependent and dependent regions (p<0.0001 repeated measure ANOVA). Ventilation inhomogeneity was associated with worse SpO_2/FiO_2, (p=0.031, r^2 0.12; linear regression). In those infants that later developed bronchopulmonary dysplasia (n=25) SpO_2/FiO_2 was worse and non-dependent ventilation inhomogeneity greater than in those that did not (both p<0.05; t test Welch correction).

Conclusions: There is high breath-by-breath variability in regional ventilation patterns during NIV in preterm infants. Ventilation favoured the gravity non-dependent lung, with ventilation inhomogeneity associated with worse oxygenation.

Abstract word count: 249 words

INTRODUCTION

Most clinical guidelines now recommend non-invasive ventilation (NIV) as initial respiratory support for preterm infants with respiratory failure.[1] NIV is simpler to use than support requiring an endotracheal tube (invasive ventilation), is well tolerated and perceived to confer short and long-term lung protective benefits.[2] In meta-analysis of clinical trials, the use of NIV results in a reduction in bronchopulmonary dysplasia (BPD), albeit small.[2] Unfortunately, population follow-up studies have not demonstrated a reduction in BPD, with poorer late-childhood respiratory outcomes reported in recent cohorts, compared with earlier cohorts where NIV was less commonly used.[3-5] Beyond the delivery room, NIV is usually provided with either continuous positive airway pressure (CPAP) or high-flow nasal cannualae (nHF). The indications for use, and specific CPAP and nHF strategies vary widely.[1,6] This may also explain some of the discrepancy between trial and clinical respiratory outcomes.

BPD is a multi-factorial progressive process that results from injurious respiratory support delivered to an immature lung with reduced alveolar capacity.[7] Injurious ventilation states can be caused by inadequate lung volume (atelectasis) or excessive tidal ventilation (volutrauma), both of which can generate damaging mechanotransductive forces resulting in increased oxygen needs and a biotrauma response.[8-10] Understanding where these events are occurring in the lung requires methods of monitoring volume state, ideally on a breath-by-breath basis.[10,11] However, continuous monitoring during NIV is limited to peripheral measures of gas exchange and clinical observation of work of breathing.

Studies in adults and preterm animals have consistently shown that ventilation is

rarely uniform within the diseased lung.[10,12,13] In preclinical studies heterogeneous (non-uniform) ventilation worsens oxygenation and creates complex injury events, potentially allowing different volume states to occur in different lung regions simultaneously.[9,10,13] In animal studies, improving ventilation uniformity reduces acute markers of lung injury.[10,13-15] Currently, clinicians have no method of assessing ventilation uniformity at the bedside in preterm infants, limiting the ability to determine whether NIV settings are lung protective or injurious. This hampers the ability of respiratory treatments to be modified before the cascade of events causing BPD develops.[4]

Recently, electrical impedance tomography (EIT) systems for infants have become available.[11,16] EIT is a method of breath-by-breath imaging of regional ventilation homogeneity and relative tidal volume (V_T) that is ideally suited to preterm infants, being non-invasive, radiation-free and continuous during all forms of support.[11,17] Limited EIT data exists in preterm infants receiving NIV,[11,18,19] especially with regards to associations with clinical status. The aim of this study was to determine the regional ventilation characteristics during non-invasive ventilation in stable preterm infants. The secondary aims were to explore the relationship between indicators of ventilation homogeneity and other clinical measures of respiratory status, including BPD. This is the first step in understanding the potential clinical utility of measures of ventilation homogeneity to guide lung-protective management in the NICU.

METHODS

The study was performed in the Neonatal Intensive Care Units (NICUs) of the Royal Children's Hospital and Royal Women's Hospital, Melbourne, Australia between November 2016 to August 2017. Prospective written informed parental consent was provided.

Infants receiving either CPAP or nHF were eligible if they were born <30 weeks of gestation, were <36 weeks corrected gestation at time of study, and had been extubated for at least 24 hours. Infants were not studied if they had at least one period of >4 hours without any respiratory support, NIV was being used for a primary cardiac disease or to support central respiratory depression, had suppressed respiratory drive due to analgesia on day of study, had fragile skin or were deemed too medically unstable to handle. The mode and settings of NIV and infant body position were determined by the clinician and were not altered during study.

An ultrasound gel-coated, non-adhesive, 32-electrode neonatal EIT belt (SenTec AG, Landquart, Switzerland) was placed around the infant's chest at nipple level during a period of clinical care using our previously described methods.[16,20-23] The belt was connected to the SenTec Pioneer EIT system and electrode signal quality confirmed. After allowing the infant to settle, four 10-minute EIT recordings were performed at 48 frames per second during 60-minutes of quiet breathing. Peripheral capillary oxygen saturation (SpO_2), heart rate (HR) and respiratory rate (RR) were measured continuously (Intellivue MD70 monitor, Phillips Healthcare, Eindhoven, Netherlands) and values recorded minutely.

Raw EIT data were reconstructed with an anatomically correct custom-made algorithm of the human chest to exclude impedance change outside the right and left lung boundaries using the manufacturer's software package (ibeX, SenTec AG), in accordance with standardised guidelines.[11,24] Once reconstructed, each of the 10-minute recordings were manually reviewed using a standardised approach to identify the first 30 consecutive artefact-free inflations (total 120 inflations/infant) included for analysis.[17,19,22] Functional EIT images of the distribution of V_T for each selected inflation were created, and data extracted from the whole lung, right and left lung, and three gravity dependent regions of interest (ROIs); gravity-nondependent (ND), central (C) and gravity-dependent (D).[11] The following measures of ventilation homogeneity were generated: 1) the relative V_T within each ROI as a percentage of total V_T delivered to the lung and as a ratio of anatomical size of each ROI (%V_T in the gravity-nondependent, central and gravity-dependent ROI only analysed for prone and supine infants); 2) the centre of ventilation (geometric mean of V_T distribution along a single plane) along the right-left (CoV$_{RL}$), and gravity non-dependent to dependent plane,[11,25] and 3) the percentage of unventilated lung regions.[11] To adjust for various body positions the gravity-dependent CoV was referenced to the CoV value representing homogenous (ideal) ventilation for that body position (calculated by ibeX), generating the ideal CoV to actual CoV difference (CoV$_{I-A}$). CoV$_{I-A}$ 0 = homogenous ventilation; >0 ventilation inhomogeneity favouring the gravity non-dependent lung; <0 ventilation inhomogeneity favouring the dependent lung (Online Supplementary Material [OSM] Figure 1).

The true pattern of ventilation in the preterm lung is unknown. Based on previous EIT studies of preterm infants, 18 infants were required to detect a clinically meaningful

2% difference in gravity-dependent CoV (assuming a 3% SD, power 80% and alpha error 0.05).[17] A convenience sample size of 40 infants was chosen to allow exploration of pre-allocated variables that may impact ventilation pattern; type of NIV, oxygen deficit at study (measure of acute disease severity), and indicators of developmental lung state (gestational age (GA) at birth, corrected GA, age at time of study). Finally, all these variables were compared with subsequent bronchopulmonary dysplasia (BPD) status at 36-weeks corrected gestation (modified Walsh criteria).[26] The relationship between these variables and CoV measures were compared with linear regression. All data were tested for normality, and additional analysis performed with appropriate categorical or continuous test. Welch correction or robust error cluster analysis (by infant) were applied as required. Statistical analysis was performed using Prism (v7.0, GraphPad, CA, USA) and a p value <0.05 considered significant.

RESULTS

During the study period, 109 infants were eligible. Forty were studied and provided complete data sets (4770 breaths; OSM Figure 2). The characteristics of the infants studied are described in Table 1. At time of study 32 infants were receiving CPAP, and 8 nHF. The range of CPAP distending pressure was 5-10 (median 7) cmH$_2$O, and nHF flow rate 4-6 L/min. Infants receiving CPAP were more immature at birth, more likely to have been intubated and received exogenous surfactant, and had greater oxygen deficit and respiratory rate at time of study. Thirty-one infants were studied prone (30 CPAP), three supine (0 CPAP) and six were lateral (2 CPAP). The mean (SD) inspiratory time was 0.40 (0.14) s, with a mean (95% CI) difference between

CPAP and nHF of 0.07 (-0.01, 0.15) s (t-test).

Table 1. Infant characteristics.

	All; n=40	**CPAP; n=32**	**nHF; n=8**
Demographic Characteristics			
GA at birth (completed weeks)	27 (2)	26 (2)	28 (2)*
Corrected GA (completed weeks)	31 (2)	31 (2)	31 (1)
Postnatal age (days)	28 (6, 67)	33 (8, 67)	12 (6, 66)
Birth weight (g)	947 (226)	903 (178)	1123 (317)
Weight at study (g)	1332 (311)	1336 (339)	1316 (168)
Female, n (%)	19 (48%)	15 (47%)	4 (50%)
Twins, n (%)	10 (25%)	8 (25%)	2 (20%)
Respiratory Characteristics			
Intubated prior to study, n (%)	35 (88%)	30 (94%)	5 (63%)[†]
Exogenous surfactant therapy, n (%)	34 (85%)	30 (94%)[#]	4 (50%)[††]
Caffeine on day of study, n (%)	37 (93%)	29 (91%)	8 (100%)
Bronchopulmonary Dysplasia, n (%)	25 (63%)	22 (69%)	3 (38%)
Cardiorespiratory status at study commencement			
FiO_2	0.30 (0.21, 0.45)	0.30 (0.21, 0.45)	0.21 (0.21, 0.28)[‡]
SpO_2 (%)	93 (4)	93 (3)	96 (5)
SpO_2/FiO_2	345 (83)	321 (69)	443 (55)**
Heart rate (bpm)	163 (12)	163 (13)	165 (10)
Respiratory rate (breaths per min)	65 (30, 92)	68 (30, 92)	51 (37, 70)

All data mean (SD), median (range) or number (%).

Abbreviations: GA, gestational age; FiO_2, Fraction of inspired Oxygen; SpO_2, Peripheral oxygen saturation; bpm, beats per minute.

\# 10 infants received 2 doses of surfactant

CPAP vs nHF: *p=0.011, **p=0.0001 (t test with Welch correction), ‡p=0.0001 (Mann-Whitney test), and †p=0.046, ††p=0.0095 (Fisher exact test)

Distribution of ventilation

Table 2 describes the CoV data. Ventilation was greater in the right lung, with a mean

(SD) 69.1 (14.9)% of V_T during each inflation occurred in the right lung. There was

no difference between CPAP and nHF; 3.8 (-3.1, 10.7)% of V_T.

Overall, the relative V_T delivered to the gravity-nondependent, central and gravity-

dependent ROI of the chest differed (Figure 1; p<0.0001, repeated-measure ANOVA), even when adjusted for the different anatomical sizes of each ROI (p<0.0001). Ventilation was relatively greater in the non-dependent lung in infants receiving CPAP, and favoured the dependent lung in the eight infants receiving nHF, but the difference was not significant (p=0.058). CoV_{I-A} by each nursing position are shown in OSM Table 1.

Table 2. Uniformity of ventilation

	All (n=40)	CPAP (n=32)	nHF (n=8)	Difference
CoV_{RL} (%)	41.2 (7.4)%	40.7 (8.1)%	43.2 (3.5)%	2.5 (-1.4, 6.4)%
Ideal (uniform) ventilation occurs at 46% of the distance from the most right-hand (0%) to left-hand (100%) lung regions.				
CoV_{I-A}	1.4 (4.5)	2.0 (4.5)	-1.1 (3.6)	3.1 (-0.1, 6.3)
Ideal (uniform) ventilation occurs at 0, values >0 indicate relatively greater V_T in the gravity non-dependent lung, and <0 the dependent lung				
Percentage of lung regions without any apparent tidal ventilation				
Whole lung	12.5 (5.7)%	12.3 (5.9)%	13.4 (4.9)%	1.1 (-3.3, 5.5)%
Non-dependent lung	6.9 (3.9)%	7.1 (4.2)%	6.2 (2.1)%	0.8 (-1.3, 3.0)%
Dependent lung	5.6 (3.7)%	5.2 (3.7)%	7.1 (3.8)%	2.0 (-1.3, 5.3)%

All data mean (SD) and differences mean (95% CI); t test with robust error. n = number of infants (4770 breaths total).

Abbreviations: CoV_{RL}, Centre of ventilation along the right to left plane; CoV_{I-A}, ratio of actual to ideal centre of ventilation along the gravity dependent plane adjusted for body position at time of study.

Relationship between uniformity of ventilation, oxygenation and developmental lung state

There was no relationship between CoV_{RL} and oxygenation (SpO_2/FiO_2 ratio), GA at birth and age at study (data not shown). Inhomogeneity of ventilation favouring the non-dependent lung was associated with worse oxygenation (Figure 2A). There was

no relationship between CoV_{I-A} and GA, age at study, corrected GA, weight, birth weight and PEEP (Figure 2B and C, OSM Table 2).

BPD status

Twenty-five infants (62.5%) were assessed as later having BPD. Infants who later met the criteria for BPD were more immature and smaller at birth, but were older, heavier and had worse oxygenation at time of EIT study (Table 3). At the time of study, ventilation was more heterogenous, being greater in the non-dependent lung, and expressing greater variability in those infants who later developed BPD compared with those who did not.

Table 3. Outcome measures at time of study and later bronchopulmonary dysplasia status.

	No BPD (n=15)	BPD (n=25)	Difference
CoV_{I-A}	-0.5 (2.2)	2.6 (5.1)	3.1 (0.7, 5.4)
SpO_2/FiO_2 ratio	402 (73)	311 (69)	92 (44, 139)
GA at birth (completed weeks)	28 (1)	26 (2)	2 (1, 3)
Age at study (days)	11 (6, 33)	38 (10, 67)	p<0.0001*
Corrected GA (completed weeks)	31 (1)	30 (1)	1.5 (0.6, 2.3)
Birthweight (g)	1084 (212)	865 (191)	219 (79, 360)
Weight at study (g)	1193 (253)	1415 (317)	221 (36, 406)
PEEP $(cmH_2O)^{†}$	6.8 (1.5)	7.1 (1.2)	0.3 (-1.5, 0.8)

*All data mean (SD) except median (range). All differences mean (95% CI) t test with Welch's correction, except * Mann-Whitney test. [†] CPAP group only (10 no BPD, 22 BPD).*

DISCUSSION

Non-invasive ventilation is the most commonly used type of respiratory support for preterm infants, but lacks reliable methods of measuring the impact on lung function.

In our observational study we demonstrated that readily available EIT measures of ventilation homogeneity can be used during NIV, and were related to disease and developmental state. Importantly, preterm infants exhibited ventilation inhomogeneity during NIV, with ventilation favouring the gravity non-dependent lung regions, a pattern acknowledged to be injurious in other groups.[10,27,28] There was an association between worse oxygenation and relatively greater non-dependent ventilation inhomogeneity, suggesting that EIT may have merit in monitoring respiratory status and guiding care in the NICU.

The primary purpose of this study was to describe the regional ventilation patterns in a representative group of stable preterm infants receiving NIV. We found that ventilation was greater in the right lung and the non-dependent lung, with high breath-by-breath variability. This suggests that ventilation inhomogeneity favouring the non-dependent lung, a well-described characteristic of acute parenchymal lung disease,[19,29-33] persists beyond the period of initial surfactant-deficient RDS, and differs from the predominantly dependent ventilation pattern described in spontaneously breathing term infants.[23,34] Ventilation inhomogeneity creates the potential for opposing mechanisms of ventilator-induced lung injury to occur simultaneously; atelectasis in poorly ventilated regions and volutrauma in over-ventilated, even if the delivered overall V_T is appropriate for gas exchange.[9,10,28] Preterm infants often receive NIV for prolonged periods,[5] and ventilation homogeneity is rarely a treatment goal. Our study suggests the risk of injury to the under-developed preterm lung goes well beyond the period of acute RDS, and may have a role in the increasing rates of BPD.[3,5]

As expected, the study population did not have high oxygen needs. Thus, the finding that ventilation distribution along the gravity-dependent plane was associated with oxygenation, but not other parameters, is interesting. Oxygenation is the most commonly used parameter to guide respiratory support in the NICU,[1] but is influenced by non-pulmonary factors, and rarely discriminates beyond identifying treatment failure or suitability of ceasing NIV. EIT measures of ventilation and aeration have been shown to be more accurate at differentiating suitability of pressure settings during high-frequency ventilation and NIV,[18,19,30,35] Recently, EIT-guided PEEP titration, based on achieving ventilation homogeneity, has been demonstrated to improve oxygenation compared to existing approaches in acute respiratory distress syndrome.[33] EIT may have an important role in optimising NIV, especially as setting PEEP/flow levels during NIV in preterm infants still lacks evidence-based approaches[1] Although the goodness of fit was low ($r^2<0.5$), there is a biological plausibility between increased ventilation heterogeneity and worse oxygenation that supports the use of EIT in the NICU and should be validated in larger studies.

The majority of infants were receiving CPAP at the time of study, but 20% were being managed with nHF. Although CoV_{RL} and COV_{I-A} were more homogeneous in the infants studied during nHF these data likely reflect the lack of matching based on type of NIV, and the small number receiving nHF. Consistent with clinical practice broadly in Australia, infants receiving nHF had less severe lung disease before and at the time of study.[5] CPAP and nHF deliver NIV via different mechanisms, but EIT ventilation patterns were the same in a cross-over study of nHF and bubble CPAP (24 infants).[36] We postulate that the role of spontaneous breathing and disease state will be more important to regional ventilation than type of NIV if applied appropriately.[22]

To date EIT studies in infants have been observational.[11] In part this is due to the need to first describe the respiratory patterns of this poorly understood population in which diseases, lung development, interventions and clinical settings vary.[4,7] Clinicians will require evidence that EIT offers the potential to modulate outcomes before widespread use,[11] as has been demonstrated in animal studies.[10,31,37] Current neonatal EIT systems now have real-time processing and simple non-adhesive belts that can be applied during multiple scenarios, and over prolonged periods of time.[16,38] EIT has already been proposed as a simple radiation-free method of determining endotracheal tube location and air-leaks.[11,39,40] The observation that gravity-dependent V_T inhomogeneity differed between infants that developed BPD and those that did not is intriguing. This is especially so as inclusion in our study was based on feasibility, and a single measurement made at a non-standardised time point. An observation between ventilation inhomogeneity and later BPD status is purely exploratory but provides a justification to further investigate the ability of repeated EIT-measures of ventilation inhomogeneity from birth to predict BPD. The lack of an early biomarker of BPD risk when disease-modifying therapies may still be effective represents the largest single hurdle to meaningful impact on the long-term respiratory consequences of preterm birth.[4,7]

This study has limitations not already highlighted. The eligibility criteria were broad, sample size based on feasibility and sub-group analysis intentionally exploratory. The clinical status of infants was not mandated at entry, and a diverse population was studied. This limits interpretation but, as this is the largest study of NIV ventilation patterns in preterm infants, does not diminish the key finding that highly variable

ventilation inhomogeneity is a feature of NIV. Our study will inform the design of larger studies that can account for the multi-factorial processes influencing BPD.[7] We did not standardise infant position during the study, but did adapt our analysis to account for it. The majority of infants were studied prone, limiting the interpretation of the differences in COV_{I-A} seen in the small numbers nursed in other positions. If nursing position does alter ventilation patterns this is important and requires further investigation.

In conclusion, in our observational study, preterm infants generate high breath-by-breath variability in regional ventilation during NIV even when clinically stable. Ventilation favoured the gravity non-dependent lung, with ventilation inhomogeneity associated with worse oxygenation. The degree of ventilation inhomogeneity increases lung injury potential in other populations, and may have use in guiding clinical care and identifying those infants at greatest risk of BPD when disease risk is modifiable.

Competing interests: There are no competing interests to declare.

Ethics approval: This study was approved by the Royal Children's Hospital Human Research Ethics Committee (HREC 36159A) and the Royal Women's Hospital Human Research Ethics Committee (HREC 16-33), in accordance with the National Health and Medical Research Council Statement of Ethical Conduct in Human Research (2007).

Financial Support: This study is supported by the Victorian Government Operational Infrastructure Support Program (Melbourne, Australia). DGT is supported by a National Health and Medical Research Council Clinical Career Development Fellowship (Grant ID 1053889). CMR is supported by a Swiss National Science Foundation Early Postdoctoral Mobility fellowship (P2ZHP3_161749) and the Milupa Fellowship Award of the Swiss Society of Neonatology. SenTec AG (Landquart, Switzerland) manufactured custom-built EIT belts for infants. All EIT hardware was purchased by MCRI without any restrictions.

Author contributions: DGT and LSO developed the concept and experimental design. JT, CMR, EJP and DGT were involved in data acquisition and JT, OF and CMR analysed the data. DGT, PP-F, LSO and EP supervised the study. All authors participated in data interpretation. JT and DGT drafted the first manuscript and all authors contributed to editing.

Data sharing: Deidentified individual participant data, study protocols and statistical analysis codes are available from three months to 23 years following article publication to researchers who provide a methodologically sound proposal, with approval by an independent review committee ("learned intermediary"). Proposals should be directed to david.tingay@mcri.edu.au to gain access. Data requestors will need to sign a data access or material transfer agreement approved by MCRI.

References

1. Sweet DG, Carnielli V, Greisen G, et al. European Consensus Guidelines on the Management of Respiratory Distress Syndrome - 2016 Update. *Neonatology* 2017;111(2):107-25. doi: 10.1159/000448985

2. Subramaniam P, Ho JJ, Davis PG. Prophylactic nasal continuous positive airway pressure for preventing morbidity and mortality in very preterm infants. *Cochrane Database Syst Rev* 2016(6):CD001243. doi: 10.1002/14651858.CD001243.pub3

3. Doyle LW, Carse E, Adams AM, et al. Ventilation in Extremely Preterm Infants and Respiratory Function at 8 Years. *N Engl J Med* 2017;377(4):329-37. doi: 10.1056/NEJMoa1700827

4. Ciuffini F, Robertson CF, Tingay DG. How best to capture the respiratory consequences of prematurity? *Eur Respir Rev* 2018;27(147) doi: 10.1183/16000617.0108-2017

5. Chow SSW, Le Marsney, R., Creighton, P., Kander, V., Haslam, R., Lui, K. Report of the Australian and New Zealand Neonatal Network 2015. *Sydney ANZNN* 2017

6. Roehr CC, Schmalisch G, Khakban A, et al. Use of continuous positive airway pressure (CPAP) in neonatal units--a survey of current preferences and practice in Germany. *Eur J Med Res* 2007;12(4):139-44. [published Online First: 2007/05/19]

7. Thebaud B, Goss KN, Laughon M, et al. Bronchopulmonary dysplasia. *Nat Rev Dis Primers* 2019;5(1):78. doi: 10.1038/s41572-019-0127-7 [published Online First: 2019/11/16]

8. Dreyfuss D, Saumon G. Ventilator-induced lung injury: lessons from experimental studies. *Am J Respir Crit Care Med* 1998;157(1):294-323.

9. Pereira-Fantini PM, Pang B, Byars SG, et al. Preterm Lung Exhibits Distinct Spatiotemporal Proteome Expression at Initiation of Lung Injury. *Am J Respir Cell Mol Biol* 2019;61:631-42. doi: 10.1165/rcmb.2019-0084OC [published Online First: 2019/04/18]

10. Tingay DG, Pereira-Fantini PM, Oakley R, et al. Gradual Aeration at Birth is More Lung Protective than a Sustained Inflation in Preterm Lambs. *Am J Respir Crit Care Med* 2019;200:609-16. doi: 10.1164/rccm.201807-1397OC [published Online First: 2019/02/08]

11. Frerichs I, Amato MB, van Kaam AH, et al. Chest electrical impedance tomography examination, data analysis, terminology, clinical use and recommendations: consensus statement of the TRanslational EIT developmeNt stuDy group. *Thorax* 2017;72(1):83-93. doi: 10.1136/thoraxjnl-2016-208357

12. Gattinoni L, Caironi P, Pelosi P, et al. What has computed tomography taught us about the acute respiratory distress syndrome? *Am J Respir Crit Care Med* 2001;164(9):1701-11.

13. Tingay DG, Rajapaksa A, Zonneveld CE, et al. Spatiotemporal Aeration and Lung Injury Patterns Are Influenced by the First Inflation Strategy at Birth. *Am J Respir Cell Mol Biol* 2016;54(2):263-72. doi: 10.1165/rcmb.2015-0127OC

14. Tingay DG, Rajapaksa A, Zannin E, et al. Effectiveness of individualized lung recruitment strategies at birth: an experimental study in preterm lambs. *Am J Physiol Lung Cell Mol Physiol* 2017;312(1):L32-L41. doi: 10.1152/ajplung.00416.2016

15. Tingay DG, Polglase GR, Bhatia R, et al. Pressure-limited sustained inflation vs. gradual tidal inflations for resuscitation in preterm lambs. *J Appl Physiol* 2015;118(7):890-7. doi: 10.1152/japplphysiol.00985.2014

16. Sophocleous L, Frerichs I, Miedema M, et al. Clinical performance of a novel textile interface for neonatal chest electrical impedance tomography. *Physiol Meas* 2018;39(4):044004. doi: 10.1088/1361-6579/aab513 [published Online First: 2018/03/09]

17. Armstrong RK, Carlisle HR, Davis PG, et al. Distribution of tidal ventilation during volume-targeted ventilation is variable and influenced by age in the preterm lung. *Intensive Care Med* 2011;37(5):839-46. doi: 10.1007/s00134-011-2157-9 [published Online First: 2011/02/26]

18. Miedema M, van der Burg PS, Beuger S, et al. Effect of Nasal Continuous and Biphasic Positive Airway Pressure on Lung Volume in Preterm Infants. *J Pediatr* 2012 doi: 10.1016/j.jpeds.2012.09.027 [published Online First: 2012/10/30]

19. Bhatia R, Davis PG, Tingay DG. Regional Volume Characteristics of the Preterm Infant Receiving First Intention Continuous Positive Airway Pressure. *J Pediatr* 2017;187:80-88 e2. doi: 10.1016/j.jpeds.2017.04.046

20. Tingay DG, Waldmann AD, Frerichs I, et al. Electrical Impedance Tomography Can Identify Ventilation and Perfusion Defects: A Neonatal Case. *Am J Respir Crit Care Med* 2019;199(3):384-86. doi: 10.1164/rccm.201808-1551LE [published Online First: 2018/10/27]

21. Miedema M, Waldmann A, McCall KE, et al. Individualized Multiplanar Electrical Impedance Tomography in Infants to Optimize Lung Monitoring. *Am J Respir Crit Care Med* 2017;195(4):536-38. doi: 10.1164/rccm.201607-1370LE

22. Dowse GP, E.; Thomson, J.; Schinckel, N.; Pereira-Fantini, P.; Tingay, D.G. Synchronised inflations generate greater gravity dependent lung ventilation in neonates [preprint]. *medRxiv* 2020;2020.06.12.20129379 doi: doi.org/10.1101/2020.06.12.20129379

23. Schinckel NF, Hickey L, Perkins EJ, et al. Skin-to-skin care alters regional ventilation in stable neonates. *Arch Dis Child Fetal Neonatal Ed* 2020 doi: 10.1136/archdischild-2020-319136 [published Online First: 2020/08/01]

24. Adler A, Arnold JH, Bayford R, et al. GREIT: a unified approach to 2D linear EIT reconstruction of lung images. *Physiol Meas* 2009;30(6):S35.

25. Frerichs I, Dargaville PA, van Genderingen H, et al. Lung volume recruitment after surfactant administration modifies spatial distribution of ventilation. *Am J Respir Crit Care Med* 2006;174(7):772-79.

26. Walsh MC, Yao Q, Gettner P, et al. Impact of a physiologic definition on bronchopulmonary dysplasia rates. *Pediatrics* 2004;114(5):1305-11. doi: 10.1542/peds.2004-0204

27. Steinberg JM, Schiller HJ, Halter JM, et al. Alveolar instability causes early ventilator-induced lung injury independent of neutrophils. *Am J Respir Crit Care Med* 2004;169(1):57-63.

28. Gattinoni L, Carlesso E, Cadringher P, et al. Physical and biological triggers of ventilator-induced lung injury and its prevention. *Eur Respir J Suppl* 2003;47:15s-25s.

29. Tingay DG, Mills JF, Morley CJ, et al. The deflation limb of the pressure-volume relationship in infants during high-frequency ventilation. *Am J Respir Crit Care Med* 2006;173(4):414-20. doi: 10.1164/rccm.200502-299OC

30. Miedema M, de Jongh FH, Frerichs I, et al. Changes in lung volume and ventilation during lung recruitment in high-frequency ventilated preterm infants with respiratory distress syndrome. *J Pediatr* 2011;159(2):199-205 e2. doi: 10.1016/j.jpeds.2011.01.066 [published Online First: 2011/03/19]

31. Wolf GK, Gomez-Laberge C, Rettig JS, et al. Mechanical ventilation guided by electrical impedance tomography in experimental acute lung injury. *Critical Care Med* 2013;41(5):1296-304. doi: 10.1097/CCM.0b013e3182771516 [published Online First: 2013/03/12]

32. Gomez-Laberge C, Rettig JS, Smallwood CD, et al. Interaction of dependent and non-dependent regions of the acutely injured lung during a stepwise recruitment manoeuvre. *Physiol Meas* 2013;34(2):163-77. doi: 10.1088/0967-3334/34/2/163

33. Zhao Z, Chang MY, Chang MY, et al. Positive end-expiratory pressure titration with electrical impedance tomography and pressure-volume curve in severe acute respiratory distress syndrome. *Ann Intensive Care* 2019;9(1):7. doi: 10.1186/s13613-019-0484-0 [published Online First: 2019/01/19]

34. Pham TM, Yuill M, Dakin C, et al. Regional ventilation distribution in the first 6 months of life. *Eur Respir J* 2011;37(4):919-24. doi: 10.1183/09031936.00034310 [published Online First: 2010/07/24]

35. Miedema M, de Jongh FH, Frerichs I, et al. The effect of airway pressure and oscillation amplitude on ventilation in pre-term infants. *Eur Respir J* 2012;40(2):479-84. doi: 10.1183/09031936.00138311 [published Online First: 2012/03/01]

36. Hough JL, Shearman AD, Jardine L, et al. Effect of randomization of nasal high flow rate in preterm infants. *Pediatr Pulmonol* 2019;54(9):1410-16. doi: 10.1002/ppul.24418 [published Online First: 2019/07/10]

37. Tingay DG, Lavizzari A, Zonneveld CE, et al. An individualized approach to sustained inflation duration at birth improves outcomes in newborn preterm lambs. *Am J Physiol Lung Cell Mol Physiol* 2015;309(10):L1138-49. doi: 10.1152/ajplung.00277.2015

38. Sophocleous L, Waldmann AD, Becher T, et al. Effect of sternal electrode gap and belt rotation on the robustness of pulmonary electrical impedance tomography parameters. *Physiol Meas* 2020;41(3):035003. doi: 10.1088/1361-6579/ab7b42 [published Online First: 2020/02/29]

39. Schmolzer GM, Bhatia R, Davis PG, et al. A comparison of different bedside techniques to determine endotracheal tube position in a neonatal piglet model. *Pediatric Pulmonology* 2013;48(2):138-45. doi: 10.1002/ppul.22580 [published Online First: 2012/05/23]

40. Bhatia R, Schmolzer GM, Davis PG, et al. Electrical impedance tomography can rapidly detect small pneumothoraces in surfactant-depleted piglets. *Intensive Care Med* 2012;38(2):308-15. doi: 10.1007/s00134-011-2421-z [published Online First: 2011/11/29]

Figure Legends

Figure 1. A. Proportion (%) of V_T occurring in the gravity-nondependent (ND; yellow), central (C; blue) and gravity-dependent (D; pink) ROI thirds of the lungs. **B.** Proportion of V_T occurring in the thirds of the chest (using same symbols as Panel A) expressed as a ratio of the anatomical contribution (ventilation homogeneity ratio) of the gravity-nondependent (26.5%), central (40.7%) and gravity-dependent (32.8%) ROI thirds to the entire lung regions imaged. A value >1 indicates that V_T was relatively greater in that region compared to the size of the region, <1 V_T relatively less than the size of the region, and 0 = uniformity. **C.** Schematic of the shape of the lungs within the field of EIT imaging demonstrating the gravity-nondependent, central and gravity-dependent regions and the absolute number of pixels (and percentage of lung) that each region contains. Infants only included if in supine or prone position, and adjusted for position (n=34). All boxes SD and line mean, error bars range. * $p<0.0001$ Tukey post test (repeated-measure ANOVA)

Figure 2. Relationship between the gravity-dependent uniformity of ventilation (CoV_{I-A}) SpO_2/FiO_2 at start of study (**A**; p=0.031, R^2=0.12, linear regression), gestational age (GA, **B**; p=0.33, R^2=0.02) and age at study (**C**; p=0.56, R^2=0.01) for all 40 infants. A CoV_{I-A} of 0 (dashed line) represents uniform ventilation, and a value >0 relatively greater V_T in the gravity nondependent lung. Greater SpO_2/FiO_2 values represents better oxygenation. Solid line represents line of best fit (equation in panel; linear regression) and dashed lines the 95% CI.

SUPPLEMENTARY MATERIAL

Ventilation homogeneity during non-invasive ventilation in preterm infants

Thomson J[1,2], Ru☐egger CM[3,4], Perkins, EJ[1], Pereira-Fantini PM[1], Farrell O[1,2], Owen LS[1,3,5], Tingay DG[1,2,3,6]

[1]Neonatal Research, Murdoch Children's Research Institute, Parkville, Australia

[2]Department of Paediatrics, The University of Melbourne, Parkville, Australia

[3]Neonatal Research, The Royal Women's Hospital, Parkville, Australia.

[4]Newborn Research, Department of Neonatology, University Hospital and University of Zurich, Zurich, Switzerland

[5]Department of Obstetrics and Gynaecology, The University of Melbourne, Parkville, Australia

[6]Department of Neonatology, The Royal Children's Hospital, Parkville, Australia

Online Supplementary Material

1. **Supplementary Results**

OSM Table 1. Role of body position at study on CoV_{I-A}.

	Prone (n=31)	Supine (n=3)	Lateral	
			Left side down (n=4)	Right side down (n=2)
CoV_{I-A} (%)	0.6 (-3.5, 10.1)	-2.8 (-2.9, 1.4)	4.2 (0.1, 17.3)	-8.4, 1.9

All data median (range) or values (right lateral). No statistical comparison was performed due to small group numbers in supine and lateral positions.

OSM Table 2. Relationship between CoV_{I-A} and corrected gestational age, weights and Positive End-Expiratory Pressure (PEEP).

	Slope	y intercept	R^2	p value
Corrected A	0.16 (-0.81, 1.13)	-6.3 (-36.0, 23.4)	0.003	0.74
Birthweight	0.004 (-0.002, 0.010)	-5.3 (-11.5, 0.9)	0.043	0.20
Weight at study	-0.002 (-0.007, 0.003)	1.4 (-5.0, 7.7)	0.021	0.37
PEEP*	-0.77 (-2.02, 0.54)	3.4 (-5.9, 12.7)	0.046	0.24

All data linear regression and error 95% CI.
* infants only receiving CPAP (n=32)

OSM Figure 1. Study recruitment flow chart.

OSM Figure 2. Role of body position on location of ideal gravity-dependent centre of ventilation (CoV). To account for the differences in ideal gravity-dependent CoV all CoV data along the non-dependent (0%) to dependent (100%) plane were expressed against the ideal CoV representing uniform ventilation in that position.

Position	Schematic	Non-dependant (0%) to dependant (100%) hemithorax	Ideal CoV (non-dependant to dependant)
Supine		Ventral lung to Dorsal lung	55 %
Prone		Dorsal lung to ventral lung	45 %
Right side down		Left lung to right lung	54 %
Left side down		Right lung to left lung	46 %