SARS-CoV-2 and Stroke Characteristics:

A Report from the Multinational COVID-19 Stroke Study Group

Authors

Shima Shahjouei, MD, MPH,[1] Georgios Tsivgoulis, MD, PhD, MSc,[2] Ghasem Farahmand, MD,[3,4] Eric Koza, MD Candidate,[5] Ashkan Mowla, MD,[1,6] Alireza Vafaei Sadr, PhD,[7] Arash Kia, MD,[8] Alaleh Vaghefi Far, MD,[4] Stefania Mondello, MD, PhD, MPH,[9] Achille Cernigliaro, PhD, MPH,[10] Annemarei Ranta, MD, PhD, FRACP,[11] Martin Punter, PhD, MBChB,[11] Faezeh Khodadadi, Pharm.D,[12] Mirna Sabra, PhD,[13] Mahtab Ramezani, MD,[14] Soheil Naderi, MD,[15] Oluwaseyi Olulana, MS,[5] Durgesh Chaudhary, MBBS,[1] Aicha Lyoubi, MD,[16] Bruce Campbell, MD,[17,18] Juan F. Arenillas, MD,[19] Daniel Bock, MD,[20] Joan Montaner, MD,[21] Saeideh Aghayari Sheikh Neshin, MD,[22] Diana Aguiar de Sousa, MD, PhD,[23] Matthew S. Tenser, MD,[6] Ana Aires, MD,[24,25] Merccedes De Lera Alfonso, MD,[19] Orkhan Alizada,[26] MD, Elsa Azevedo, MD,[24,25] Nitin Goyal, MD,[27] Zabihollah Babaeepour, MD,[28] Gelareh Banihashemi, MD,[29] Leo H. Bonati, MD,[30] Carlo Cereda, MD,[31] Jason J. Chang, MD,[32] Miljenko Crnjakovic, MD,[33] GianMarco De Marchis, MD,[34] Massimo Del Sette, MD,[35] Seyed Amir Ebrahimzadeh, MD, MPH,[36] Mehdi Farhoudi, MD,[37] Ilaria Gandoglia, MD,[35] Bruno Gonçalves, MD,[38] Christoph Griessenauer, MD,[1] Mehmet Murat Hancı, MD,[26] Aristeidis H. Katsanos, MD,[2,39] Christos Krogias, MD,[40] Ronen Leker, MD,[41] Lev Lotman, MD,[42] Jeffrey Mai, MD,[43] Shailesh Male, MD,[44] Konark Malhotra, MD,[45] Branko Malojcic, MD, PhD,[46] Teresa Mesquita, MD,[47] Asadollah Mirghasemi, MD,[48] Hany Mohamed Aref, MD,[49] Zeinab Mohseni Afshar, MD,[50] Jusun Moon, MD,[51] Mika Niemelä, MD, PhD,[52] Behnam Rezaei Jahromi, MD,[52] Lawrence Nolan, MD,[41] Abhi Pandhi, MD,[27] Jong-Ho Park, MD,[53] João Pedro Marto, MD,[46] Francisco Purroy, MD, PhD[54] Sakineh Ranji-Burachaloo, MD,[3] Nuno Reis Carreira, MD,[55] Manuel Requena, MD,[56] Marta Rubiera, MD,[56] Seyed Aidin Sajedi, MD,[57] João Sargento-Freitas, MD,[58] Vijay Sharma, MD,[59] Thorsten Steiner, MD,[60,61] Kristi Tempro, MD,[41] Guillaume Turc,[38] MD, PhD, FESO, Yassaman Ahmadzadeh, MD,[62] Mostafa Almasi-Dooghaee, MD,[63,64] Farhad Assarzadegan, MD,[14] Arefeh Babazadeh, MD, MPH,[65] Humain Baharvahdat, MD,[66] Fabricio Cardoso, MD, MPH,[67] Apoorva Dev, PhD,[12] Mohammad Ghorbani, MD,[68] Ava Hamidi, MD,[69] Zeynab Sadat Hasheminejad, MD,[70] Sahar Hojjat-Anasri Komachali, MD,[71] Fariborz Khorvash, MD,[72] Firas Kobeissy, PhD,[73] Hamidreza Mirkarimi, MD,[74] Elahe Mohammadi-Vosough, MD,[74] Debdipto Misra, MS,[75] Ali Reza Noorian, MD,[76] Peyman Nowrouzi-Sohrabi, PhD,[77] Sepideh Paybast, MD,[78] Leila Poorsaadat, MD,[79] Mehrdad Roozbeh, MD,[80] Behnam Sabayan, MD, PhD,[81] Saeideh Salehizadeh,[82] Alia Saberi, MD,[22] Mercedeh Sepehrnia, MD,[70] Fahimeh Vahabizad, MD,[29] Thomas Yasuda, MD,[67] Ahmadreza Hojati Marvast, MD,[4] Mojdeh Ghabaee, MD[3,4] Nasrin Rahimian, MD, MPH,[83] Mohammad Hossein Harirchian, MD,[3] Afshin Borhani-Haghighi, MD,[84] Rohan Arora, MD,[85] Saeed Ansari, MD,[27] Venkatesh Avula, MS,[86] Jiang Li, MD,[86] Vida Abedi, MD,[86,87] Ramin Zand, MD, MPH.[1]

[1] Neurology Department, Neuroscience Institute, Geisinger Health System, Pennsylvania, USA;

[2] Second Department of Neurology, National and Kapodistrian University of Athens, School of Medicine, "Attikon" University Hospital, Athens, Greece;

[3] Iranian Center of Neurological Research, Neuroscience Institute, Tehran University of Medical Sciences, Tehran, Iran;

[4] Neurology Department, Tehran University of Medical Sciences, Tehran, Iran;

[5] Geisinger Commonwealth School of Medicine, Scranton, Pennsylvania, USA;

[6]Division of Stroke and Endovascular Neurosurgery, Department of Neurological Surgery, Keck School of Medicine, University of Southern California, California, USA;

[7]Department de Physique Theorique and Center for Astroparticle Physics, University Geneva, Switzerland;

[8]Icahn school of medicine at Mount Sinai, Department of Population Health Science and Policy, Institute for Healthcare Delivery Science, New York, USA;

[9]Department of Biomedical and Dental Sciences and Morphofunctional Imaging, University of Messina, Messina, Italy;

[10]Regional Health Authority of Sicily, Palermo, Italy;

[11]Department of Neurology, Wellington Hospital, Wellington, New Zealand;

[12]PES University, Bangaluru, Karnataka, India;

[13]Neurosciences Research Center (NRC), Lebanese University/ Medical School, Beirut, Lebanon;

[14]Neurology Department, Shahid Beheshti University of Medical Sciences, Tehran, Iran;

[15]Department of Neurosurgery, Tehran University of Medical Sciences, Tehran, Iran;

[16]Neurology Department, Delafontaine Hospital, Saint-Denis, France

[17]Department of Neurology, Royal Melbourne Hospital, Parkville, Australia;

[18]University of Melbourne, Parkville, Australia;

[19]Department of Neurology, University of Valladolid, Valladolid, Spain;

[20]Department of Cardiology, Klinikum Frankfurt Höchst, Frankfurt Germany;

[21]Department of Neurology, Hospital Universitario Virgen Macarena, Sevilla, Spain;

[22]Neurology Department, Poursina Hospital, Rasht, Guilan, Iran;

[23]Department of Neurology, Hospital de Santa Maria, University of Lisbon, Lisbon, Portugal;

[24]Department of Neurology, Centro Hospitalar Universitário de São João, Porto, Portugal;

[25]Faculty of Medicine, University of Porto, Porto, Portugal;

[26]Neurosurgery Department, Istanbul University-Cerrahpasa, Cerrahpasa Medical Faculty, Istanbul, Turkey;

[27]Department of Neurology, University of Tennessee, Tennessee, USA;

[28]Neurology Ward, Valiasr Hospital, Borujen, Iran;

[29]Neurology Department, Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran;

[30]Department of Neurology and Stroke Unit, University Hospital Basel, Switzerland;

[31]Stroke Center, Neurocenter of Southern Switzerland, Lugano, Switzerland;

[32]Department of Critical Care Medicine, MedStar Washington Hospital Center, Washington, DC, USA;

[33]Intensive Care Unit, Department of Neurology, Clinical Hospital Dubrava, Zagreb, Croatia;

[34]Neurorehabilitation Unit, University Center for Medicine of Aging and Rehabilitation Basel, Felix Platter Hospital, University of Basel, Switzerland;

[35]Neurology Unit, Galliera Hospital, Genova, Italy;

[36]Department of Radiology, Yasrebi Hospital, Kashan, Iran;

[37]Neurosciences Research Center (NSRC), Tabriz University of Medical Sciences, Tabriz, Iran;

[38]Department of Neurology, GHU Paris Psychiatrie et Neurosciences, Université de Paris, INSERM U1266, Paris, France;

[39]Division of Neurology, McMaster University/ Population Health Research Institute, Hamilton, ON, Canada

[40]Department of Neurology, St. Josef-Hospital, Ruhr University Bochum, Bochum, Germany;

[41]Department of Neurology, Hadassah-Hebrew University Medical Center, Jerusalem, Israel;

[42]Department of Neurology, Albany Medical Center, Albany, NY;

[43]Department of Neurosurgery, Georgetown University and MedStar Washington Hospital Center, Washington, DC, USA;

[44]Neurology Department, Vidant Medical Center, Greenville, North Carolina, USA;

[45]Department of Neurology, Allegheny Health Network (AHN), Pittsburgh, Pennsylvania, USA;

[46]TIA Clinic, Department of Neurology, University Hospital Centre Zagreb, Zagreb School of Medicine, University of Zagreb, Zagreb, Croatia;

[47]Department of Neurology, Hospital de Egas Moniz, Centro Hospitalar Lisboa Ocidental, Lisbon, Portugal;

[48]Department of Anesthesiology, University of Ottawa, Canada;

[49]Department of Neurology, Ain Shams University, Cairo, Egypt;

[50]Infection Disease Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran;

[51]Department of Neurology, National Medical Center, Seoul, South Korea;

[52]Department of Neurosurgery, Helsinki University Hospital, Helsinki, Finland;

[53]Department of Neurology, Myongji Hospital, Hanyang University College of Medicine, South Korea;

[54]Department of Neurology, Hospital Arnau de Vilanova, Lleida, Spain;

[55]Department of Neurology, Hospital Arnau de Vilanova, Institut de Recerca Biomèdica de Lleida (IRBLLeida), Universitat de Lleida UdL Lleida, Spain

[56]Stroke Unit, Department of Neurology, Hospital Vall d'Hebron, Department de Medicina, Universitat Autònoma de Barcelona, Barcelona, Spain;

[57]Neuroscience Research Center, Department of Neurology, Golestan University of Medical Sciences, Golestan, Iran;

[58]Department of Neurology, Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal;

[59]Division of Neurology, University Medicine Cluster, National University Health System, Singapore, Singapore;

[60]Department of Neurology, Klinikum Frankfurt Höchst, Frankfurt, Germany;

[61]Department of Neurology Heidelberg University Hospital, Germany

[62] Hospital for Special Surgery, New York, USA;

[63]Divisions of Vascular and Endovascular Neurosurgery and Neurology, Firoozgar Hospital, Iran University of Medical Sciences, Tehran, Iran;

[64]Divisions of Vascular and Endovascular Neurosurgery and Neurology, Rasoul-Akram Hospital, Iran University of Medical Sciences, Tehran, Iran;

[65]Infectious Diseases and Tropical Medicine Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran;

[66]Neurosurgical Department, Ghaem Hospital, Mashhad University of Medical Sciences, Mashhad, Iran;

[67]Neurology Department, Centro Médico de Campinas, São Paulo, Brazil;

[68]Division of Vascular and Endovascular Neurosurgery, Firoozgar Hospital, Iran University of Medical Sciences, Tehran, Iran;

[69]Neurology Ward, Gheshm Hospital, Gheshm, Iran;

[70]Department of Neurology, Imam Hosein Hospital, Shahid Beheshti Medical University, Tehran, Iran;

[71]Department of Neurology, Pirooz hospital, Gilan University of Medical Sciences, Lahijan, Iran;

[72]Professor Of Neurology, Isfahan University of Neurology, Isfahan, Iran;

[73]Program of Neurotrauma, Neuroproteomics and Biomarker Research (NNBR), University of Florida, Florida, USA;

[74]Neurology Ward, Modarres Hospital, Kashmar, Iran;

[75]Steele Institute of Health & Innovation, Geisinger Health System, Pennsylvania, USA;

[76]Department of Neurology, Southern California Permanente Medical Group, Irvine, California, USA;

[77]Student Research Committee, Shiraz University of Medical Sciences, Shiraz, Iran;

[78]Department of Neurology, Bou Ali hospital, Qazvin University of Medical Sciences, Qazvin, Iran;

[79]Department of Neurology, Arak University of Medical Sciences, Arak, Iran;

[80]Brain Mapping Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran;

[81]Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA;

[82]Salahadin Ayubi Hospital, Baneh, Iran;

[83]Department of Neurology, Yasrebi Hospital, Kashan, Iran;

[84]Clinical Neurology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran;

[85]Department of Neurology, Long Island Jewish Forest Hills, Queens, New York, USA;

[86]Department of Molecular and Functional Genomics, Geisinger Health System, Danville, Pennsylvania, USA;

[87]Biocomplexity Institute, Virginia Tech, Blacksburg, Virginia, USA.

Corresponding Author:

Ramin Zand M.D., M.P.H.

Neuroscience Institute, Geisinger,

100 North Academy Ave. Danville, PA, 17822

phone number: 570-214-4101

Fax number: (570) 808-3208

E-mail address: rzand@geisinger.edu; ramin.zand@gmail.com

Laboratory: www.thedecodelab.com

Abstract

Background: Stroke is reported as a consequence of SARS-CoV-2 infection. However, there is a lack of regarding

comprehensive stroke phenotype and characteristics

Methods: We conducted a multinational observational study on features of consecutive acute ischemic stroke (AIS),

intracranial hemorrhage (ICH), and cerebral venous or sinus thrombosis (CVST) among SARS-CoV-2 infected

patients. We further investigated the association of demographics, clinical data, geographical regions, and countries'

health expenditure among AIS patients with the risk of large vessel occlusion (LVO), stroke severity as measured by

National Institute of Health stroke scale (NIHSS), and stroke subtype as measured by the TOAST criteria.

Additionally, we applied unsupervised machine learning algorithms to uncover possible similarities among stroke

patients.

Results: Among the 136 tertiary centers of 32 countries who participated in this study, 71 centers from 17 countries

had at least one eligible stroke patient. Out of 432 patients included, 323(74.8%) had AIS, 91(21.1%) ICH, and

18(4.2%) CVST. Among 23 patients with subarachnoid hemorrhage, 16(69.5%) had no evidence of aneurysm. A total

of 183(42.4%) patients were women, 104(24.1%) patients were younger than 55 years, and 105(24.4%) patients had

no identifiable vascular risk factors. Among 380 patients who had known interval onset of the SARS-CoV-2 and

stroke, 144(37.8%) presented to the hospital with chief complaints of stroke-related symptoms, with asymptomatic or

undiagnosed SARS-CoV-2 infection. Among AIS patients 44.5% had LVO; 10% had small artery occlusion according

to the TOAST criteria. We observed a lower median NIHSS (8[3-17], versus 11[5-17]; p=0.02) and higher rate of

mechanical thrombectomy (12.4% versus 2%; p<0.001) in countries with middle to high-health expenditure when

compared to countries with lower health expenditure. The unsupervised machine learning identified 4 subgroups, with

a relatively large group with no or limited comorbidities.

Conclusions: We observed a relatively high number of young, and asymptomatic SARS-CoV-2 infections among

stroke patients. Traditional vascular risk factors were absent among a relatively large cohort of patients. Among

hospitalized patients, the stroke severity was lower and rate of mechanical thrombectomy was higher among countries

with middle to high-health expenditure.

Introduction:

Since the emergence of the Coronavirus disease 2019 (COVID-19) pandemic, several cases of cerebrovascular events were reported among patients with SARS-CoV-2.[1,2] Studies presented the incidence and prevalence of acute ischemic stroke (AIS), intracranial hemorrhage (ICH), and cerebral venous or sinus thrombosis (CVST) in SARS-CoV-2 infected patients.[3–11] Majority of these reports focus on AIS patients. Some reports highlighted AIS in critically ill and older patients with a higher number of comorbidities,[1,6,12] some suggested a higher risk in younger and healthy individuals with male predilection,[3,7,13] and other reports compared these patients with the cohort of stroke patients who were not infected with the virus.[14,15]

Many studies proposed coagulopathy as the underlying pathophysiological mechanism for the cerebrovascular events.[15,16] Accordingly, studies suggested that stroke patients with SARS-CoV-2 present multiple cerebral infarcts,[8,12,13] systemic coagulopathies in multiple organs,[17] uncommon thrombotic events such as aortic[18] or common carotid artery thrombosis,[19] and simultaneous arterial and venous thrombus formation.[20] Small case series demonstrated a higher proportion of large vessel occlusions,[3,6,13], or cryptogenic strokes,[5,21] with elevated D-dimer level, liver enzymes, and inflammatory or renal failure biomarkers among the patients who experienced SARS-CoV-2 infection.[5,7,12,22] Besides, most of the studies noted a higher severity and mortality rate among stroke patients diagnosed with SARS-CoV-2 compared with others.[2,5,6,23,24]

We devised a multinational multiple phase study to evaluate the stroke risk[25] and characteristics among the SARS-CoV-2 infected patients. We investigated the association among demographics, clinical data, geographical regions, and countries health expenditure among AIS patients with the risk of large vessel occlusion (LVO), stroke severity as measured by National Institute of Health stroke scale (NIHSS),[26] and stroke subtype (documented using TOAST criteria – the Trial of Org 10172 in Acute Stroke Treatment).[27] We further applied unsupervised machine learning algorithms to uncover the possible similarities among these patients.

Methods:

Study Design:

The details of the study design are available in Supplemental Document 1. This multicenter, multinational

observational study was conducted and reported according to the Strengthening the Reporting of Observational Studies

in Epidemiology (STROBE),[28] and Enhancing the QUAlity and Transparency Of health Research (EQUATOR)

guidelines.[29] The study protocol was designed by the investigators at the Neuroscience Institute of Geisinger Health

System, Pennsylvania, USA, and received approval by the Institutional Review Board of Geisinger Health System

and other participating institutions. Investigators from North America (Canada and six states of the United States),

South America (Brazil and Mexico), Europe (Belgium, Croatia, Czech Republic, Finland, France, Germany, Greece,

Ireland, Italy, Norway, Portugal, Spain, Sweden, and Switzerland), Asia and the Middle East (India, Iran, Iraq, Israel,

Lebanon, Singapore, South Korea, Turkey, and the United Arab Emirates), Oceania (Australia and New Zealand), and

Africa (Egypt, Nigeria, and Uganda) responded to our invitation. The centers were included by non-probability

sampling and data were recruited until June 10th, 2020.

<u>Participants</u>

We included consecutive SARS-CoV-2 infected adult patients who had imaging confirmed subsequent stroke[30]—

AIS, intracerebral hemorrhage, subarachnoid hemorrhage (SAH), and CVST. The preferred diagnostic criteria for

SARS-CoV-2 was defined according to the World Health Organization (WHO) interim guidance.[31] Ischemic or

hemorrhagic strokes were defined in the presence of a rapid onset of a neurological deficit with evidence of acute

ischemic or hemorrhagic lesions on Computed Tomography (CT) or Magnetic Resonance Imaging (MRI). Patients

who had transient stroke-like symptoms (transient ischemic attack, TIA) without acute lesions on CT or MRI were

not included in this study due to the high diagnostic error. [32–34]

Patients who initially presented to the hospital with stroke-related chief complaints and asymptomatic SARS-CoV-2

infection, those who had a stroke while being hospitalized for SARS-CoV-2 infection, or patients with stroke-related

admission who had confirmed prior diagnosis of SARS-CoV-2 were included in this study. We excluded patients with

stroke incidents prior to the SARS-CoV-2 infection onset. The onset of SARS-CoV-2 was considered as either the

symptoms onset or positive test, whichever was first.

Data Element and Processing

Collaborators were asked to provide data according to a core protocol. The age, sex, vascular risk factors and comorbidities (i.e., hypertension, diabetes mellitus, ischemic heart disease, atrial fibrillation, carotid stenosis, chronic kidney disease, cardiac ejection fraction <40%, active neoplasms, rheumatological diseases, smoking status, and history of TIA or stroke), and laboratory findings (i.e., the count for white blood cells, neutrophils, lymphocytes, and platelets, C-Reactive Protein, blood urea nitrogen, creatinine, alanine transaminase, aspartate transaminase, lactic acid dehydrogenase, fibrinogen, and D-dimer) were requested for the stroke patients. We also obtained additional data including the onset of the stroke and SARS-CoV-2 infection, the initiation of mechanical ventilation (if applicable), length of hospital stays, and patient disposition—still in the hospital, in-hospital death, being discharged to home, acute rehabilitation service, or nursing home. The details of neurological symptoms and investigations, imaging-based localization of the lesion(s), use of antiplatelets or anticoagulants prior to the stroke, and the National Institute of Health stroke scale (NIHSS) and the intracranial hemorrhage (ICH) score were also requested. In patients with AIS, beside TOAST criteria (the Trial of Org 10172 in Acute Stroke Treatment; defined as large artery atherosclerosis, cardioembolism, small artery occlusion, other determined etiology, and undetermined etiology)[27], the lesion(s) on diffusion-weighted imaging (DWI) or CT images were categorized as lacunar,[35] embolic/large vessel athero-thromboembolism,[36,37] vasculitis pattern,[38] or other phenotypes (borderzone or equivocal lesions). In this study, the AIS due to large-vessel occlusions (LVOs) are referred as occlusion of the internal carotid artery (ICA), middle cerebral artery (MCA) at M1 and M2, anterior cerebral artery (ACA) at A1, posterior cerebral artery (PCA) at P1, intracranial vertebral artery (VA), or basilar artery (BA).[39] Brain imaging findings were evaluated by local radiologists with expertise in neuroimaging.

We grouped the patients according to the age (younger versus older than 55-year-old, and younger versus older than 65-year-old),[40] comorbidities, laboratory findings, geographical regions (Middle East, Asia, Europe, and America), and countries' health expenditure (low versus middle and high income based on WHO reports).[41] We considered the countries' annual health expenditure of above US$1,000 (2015 to 2017, Supplemental Table 1) and total health expenditure of above US$10,000 (2010 to 2017, Supplemental Figure 1) as the cut-off. For the interval between SARS-CoV-2 infection and stroke, we grouped the patients as visiting the hospital with the chief complaint of stroke-

like symptoms and obscure viral infection versus those who presented the symptoms or were diagnosed by SARS-

CoV-2 infection prior to the stroke.

<u>Outcome measures</u>

The outcome measures in this study were the presence versus absence of LVO, stroke severity as measured by NIHSS,

and stroke subtype as measured by TOAST criteria. The severity of the stroke, according to NIHSS, was defined as

no stroke symptoms (NIHSS=0), minor (NIHSS of 1 to 4), moderate (NIHSS of 5 to 15), moderate to severe (NIHSS

of 16 to 20), and severe stroke (NIHSS 21-42).[26] We studied the outcome measures only among the patients with

AIS. We did not analyze the disposition and length of stay as outcome measures since many patients were still in the

acute phase or admitted in long-term acute care hospitals at the closure of our study.

<u>Clustering and Subgroup Analyses</u>

We performed clustering and subgroup analyses (Figure 1) to uncover the possible similarities among the patients in

the entire cohort. Patients with AIS and intraparenchymal hemorrhage (IPH, hemorrhage exclusively within brain

parenchyma) were grouped independently based on the reported comorbidities and laboratory findings with the aid of

unsupervised machine learning algorithms (ML) or expert opinion (EX) (details available in Supplemental Document

2). In ML models, we used hierarchical and K-means clustering (ML-K) to group the patients into 2 to 5 subgroups

(models $ML\text{-}K_2$ to $ML\text{-}K_5$). We also used the spectral clustering (ML-S) and clustered the patients into 2 to 5

subgroups ($ML\text{-}S_2$ to $ML\text{-}S_5$). The subgroup analysis considering the expert opinion (EX) defined the risk score as the

sum of either all the 11 collected comorbidities (Expert-All, EX-A) or the 8 selected comorbidities (Expert-Selected,

EX-S). We first divided the EX-A and EX-S into two subgroups ($EX\text{-}A_2$ and $EX\text{-}S_2$); subgroups "a" included the

patients with zero or one comorbidity, and subgroup "b" included the patients with >1 comorbidity. We repeated the

subgroup analysis by dividing the EX-A and EX-S into three subgroups ($EX\text{-}A_3$ and $EX\text{-}S_3$); we considered subgroup

"a" as patients without any known comorbidity, subgroup "b" with one or two comorbidities, and subgroup "c" as the

patients with >2 comorbidities. The above processes were repeated for the laboratory findings (data not shown);

however, no patterns or significant differences among the subgroups were observed. The clustering was not performed

for patients with SAH and CVST due to limited sample size.

<u>Statistical Analysis and Modeling</u>

We used descriptive statistics to summarize the data. Demographic data, comorbidities, laboratory findings, and neurological investigations were reported as medians and interquartile range (IQR), mean and standard deviations (SD), and under stratified categories when possible. The equality of the variances was assessed by Leven's test. Categorical variables were reported as absolute frequencies and percentages. The comparisons between categorical variables were conducted with the Pearson chi-square test, while the differences among continuous variables were assessed by independent t-test and analysis of variance (ANOVA). A post-hoc z-test on the adjusted residuals, and Cramér's phi, Tukey, or Dunnett's tests were used to demonstrate the degree and direction of the associations in comparison of means, while post-hoc comparison of medians was conducted by Dunn-Bonferroni approach to compare subgroups. All tests were performed using IBM SPSS Statistics version 26[42], and $p<0.05$ was considered statistically significant. Bonferroni correction was used for adjusting all p values in multiple comparisons.

We used unsupervised ML algorithms to cluster the patients based on the comorbidities and laboratory findings. The laboratory values were scaled to 1-100 range and underwent Log_{10} transformation prior to the clustering. We applied hierarchical (complete linkage method) and k-means (Hartigan-Wong algorithm) clustering, and spectral clustering,[43] to produce 2 to 5 clusters. We used the contingency matrix (a.k.a contingency table)[44] to present the clusters of each model versus other models. The similarity of the models was calculated as follows:

$Sim = \frac{\sum_1^i Maximum\ Value\ in\ Column\ i}{\sum_1^k Value\ in\ Cell\ k}$; where i is the number of columns and k is the total number of cells in the contingency matrix. Similarities among the models were considered to be mild (50%-65%), moderate (65%-80%), and strong (80%-100%). Packages *stat*[45] and *gplots*[46] in R version 3.6.3, and scikit-learn package[47] in Python version 3.7, were used for machine learning algorithms and visualization.

Results:

Collaborators from 136 tertiary centers of 32 countries participated in this study. Among them, 71 centers from 17 countries had at least one stroke patient eligible for this study. One center in the Middle East could not provide data by the deadline. The rest of the centers did not have stroke patients who met our inclusion criteria (Supplemental Document 1). We received data on 432 patients—America:114(26.4%), Europe: 82(19.0%), Middle East: 228(52.8%), and Asia: 8(1.9%). Among them, 203(47.0%) patients were from countries with middle- to high- health expenditure. Overall, 323(74.8%) patients had AIS, 91(21.1%) ICH, and 18(4.2%) CVST.

The mean age for the entire cohort was 65.7±15.7 years. Out of 432 patients, majority were men--249(57.6%), p<0.001. Among 380 patients who had known interval onset of the SARS-CoV-2 and stroke, 144(37.8%) presented to the hospital with chief complaints of stroke-related symptoms, with asymptomatic or undiagnosed SARS-CoV-2 infection. In total, 105(24.4%) out of the 430 patients with complete comorbidity profiles had no identifiable vascular risk factor at the time of stroke incidence. Details of patient phenotype and demographic characteristics under each stroke subtype are presented in Table 1.

AIS was diagnosed in 323(74.8%) patients (Table 2). AIS patients were grouped based on gender and age – ≤55 versus >55 years-old, ≤65 versus >65 years-old (Supplemental Table 2, 3 and 4). Out of 323 patients with AIS, 36.2% were younger than 55 years. Patients above 55 years had higher proportion of comorbidities—hypertension, 69.3% versus 39.4%, p<0.001; ischemic heart disease, 27.3% versus 13.8%, p=0.03; atrial fibrillation, 16.1% versus 6.1%, p=0.04; and carotid stenosis, 15.2% versus 4.6%, p=0.03. In addition, older stroke patients had higher cardioembolic origin according to the TOAST criteria (31.2% versus 15.6%, p=0.05). There were differences among men and women in smoking status (9.4% versus 21.2% in men, p=0.01), chronic kidney disease (7.9% versus 16.6% in men, p=0.01), NIHSS (12.0 [5.0 – 19.0] versus 8.0 [4.0 – 16.0] in men, p<0.001), and TOAST classification (p=0.05).

The distribution of AIS subtypes according to TOAST classification was the following: large-artery atherosclerosis (33%), cardioembolism (27%), small vessel occlusion (10%), ischemic stroke of other determined etiology (8%) and ischemic stroke of undetermined etiology (22%). The subgroups of the patients according to the TOAST classification were different in terms of age, gender, the prevalence of LVO, need for mechanical ventilation, hypertension, ischemic heart disease, atrial fibrillation, carotid stenosis, chronic kidney disease, and neoplasm, the median of laboratory results (lymphocytes counts, creatinine), and imaging patterns. We observed lower median of D-dimer in patients with large artery atherosclerosis compared to those with cardio-embolism strokes (486.5 [371.5 – 1422.5] ng/ml versus 1100.0 [955.0 – 2355.0] ng/ml, p=0.04). There were no differences in terms of LDH or fibrinogen among TOAST subgroups (Table 2, Supplemental Table 5). Patients grouped by NIHSS strata were different in terms of the prevalence of LVO, intravenous thrombolysis and mechanical thrombectomy, imaging patterns, TOAST classifications, need for mechanical ventilation, ischemic heart disease, atrial fibrillation, and mean of laboratory results (white blood cell counts, alanine transaminase, aspartate transaminase, creatinine, fibrinogen, and D-dimer) (Table 3, Supplemental Table 6). Patients with LVO were different from those without LVO in terms of NIHSS, imaging patterns, TOAST

criteria, and prevalence of intravenous thrombolysis, mechanical thrombectomy, and ischemic heart disease. (Table 3, Supplemental Table 7). Supplemental Tables 8 and 9 also present the difference among subgroups in terms of neuroimaging findings and the interval between SARS-CoV-2 infection and stroke. AIS patients were additionally compared based on geographical regions (Middle East, Asia, Europe, and America, Supplemental Table 10), and countries' health expenditures (low, or middle to high, Supplemental Table 11). We observed differences among the patients in terms of comorbidities: ischemic heart disease, chronic kidney disease, and congestive heart failure with ejection fraction <40%. We also observed differences among patients when considering the median of laboratory results (white blood cell count, platelet count, lactate dehydrogenase, fibrinogen, and D-dimer), the proportion of patients who underwent mechanical thrombectomy, and neuro-imaging findings. We observed a significant difference regarding the TOAST criteria in both regional and health-expenditure subgroups; small artery occlusion in the Middle East: 18.3%, America: 5.3%, and Europe: 4.1%, p<0.001; 4% versus 18.3% in countries with lower health-expenditure, p<0.001. We further detected higher NIHSS in countries with lower health-expenditure—11.0 [5.0–17.0] versus 8.0 [3.0–17.0], p=0.02. There was no significant difference among regions and health-expenditure subgroups in terms of the prevalence of LVO.

A total of 91(21.1%) patients presented with the ICH (Table 4). Among them, 3(3.3%) had simultaneous SAH and IPH without any evidence of aneurysm, and 4(4.4%) were presented with simultaneous intraventricular hemorrhage and IPH. Isolated SAH occurred in 23(25.3%), and isolated IPH in 61(67%) of the patients with hemorrhagic stroke. Eighteen (4.2%) patients experienced cerebral sinus or cortical venous thrombosis (Table 3); among them, 5(27.8%) had multiple vascular involvements.

Clustering and Subgroup Analysis

The structure of the models and proportion of the comorbidities under each model are available in Supplemental Tables 12-14 (AIS patients) and Supplemental Tables 15-17(IPH patients). Clustering the patients into 4 or 5 subgroups (ML-K$_4$, ML-S$_4$, ML-K$_5$, and ML-S$_5$; Supplemental Tables 14A, 14B, 17A, and 17B) were not informative and not included for further comparisons. Supplemental Figures 2 (AIS) and 3 (IPH) demonstrate contingency matrices. Similarity among different models was assessed to be strong (Sim>80) in all matrices except comparisons of unsupervised algorithms (ML-K$_3$ and ML-S$_3$) and expert opinions (EX-A$_3$ and EX-S$_3$) among AIS patients— moderate similarity, Sim=66%-73%. Figure 2A visualizes the comorbidities among AIS patients. The hierarchical

cluster demonstrates 4 subgroups, with a relatively large group with no or limited comorbidities. Figure 2B demonstrates comorbidities among IPH patients. Hypertension and diabetes were the most repeated and overlapped comorbidities among both AIS and IPH cohorts.

When the AIS patients were divided into two subgroups based on the existing comorbidities, all models (EX-A$_2$, EX-S$_2$, ML-K$_2$, and ML-S$_2$) presented differences among the patients according to the TOAST criteria and median of creatinine. EX-A$_2$, EX-S$_2$, and ML-K$_2$ also showed differences in the mean of age among the subgroups (Supplemental Table 18). When patients were divided into 3 subgroups, all models demonstrate a difference in mean of age among the patients. Additionally, subclasses were different in terms of TOAST criteria (EX-A$_3$, EX-S$_3$, ML-S$_3$), intravenous thrombolysis (ML-S$_3$ and EX-A$_3$), and LVO (ML-S$_3$) (Supplemental Table 19).

Dividing the IPH patients into two subgroups presented differences in white blood cells' count (ML-K$_2$ and EX-A$_2$) and neutrophils' count (ML-K$_2$, EX-A$_2$, and EX-S$_2$). We observed differences among subgroups of IPH patients in terms of age (ML-K$_3$, ML-S$_3$, EX-A$_3$, and EX-S$_3$), and ICH score (EX-A$_3$ and EX-S$_3$). Patients with no known comorbidities had younger age and higher ICH score in comparison with those having 1 or 2 comorbidities, or >2 comorbidities: EX-A3 model: mean age of 55±19 years compared to 69±15 years and 64±17 years, p=0.03; ICH score of 4[2-5] compared to 3[2-3] and 3[2-4], p=0.02. EX-S3 model: mean age of 53±19 years compared to 70±13 years and 63±17 years, p=0.001; ICH score of 4 [2-4] compared to 3 [2-3] and 3[2-5], p=0.02 (Supplemental Tables 20 and 21).

Discussion

To our knowledge, this is to date the largest study that comprehensively presents the characteristics and stroke subtypes of stroke in SARS-CoV-2 infected patients at a multinational level. The results of our work indicated a relatively high number of young AIS patients, male predominance, asymptomatic SARS-CoV-2 infection in more than one-third of the AIS patients, absence of traditional stroke risk factors in about one-fourth of the AIS patients, and a low rate of small artery occlusion and lacunar infarcts. We also noted significant differences regarding the TOAST criteria in both regional and health-expenditure subgroups as well as higher NIHSS among countries with lower health-expenditure. About one third of IPH patients did not have vascular risk factors but presented with higher ICH scores.

Although the definition of young stroke is debatable, the majority of the studies considered 50 or 55 years as the cut-

off.[40] We considered both 55 and 65 years-old as the cut-off and realized that younger patients had fewer

comorbidity rates (Supplemental Tables 2 and 3). Out study results showed that 36% of our AIS patients were younger

than 55 years old. This proportion was considerably higher than the reported proportions (12.9% to 20.7%) of young

stroke patients.[48,49] In our study, 64.4% of the patients under 55 years had AIS. The median age of AIS patients in

our study was 68[58–78] years, with no substantial differences across geographical regions (Supplemental Table 10).

A case series from New York on 32 AIS patients with SARS-CoV-2 showed a median of 63 years for these patients

which was significantly lower than AIS patients without SARS-CoV-2 in the same study and interval (median: 70

years), or historical cohort of the AIS patients presented to the center in 2019 (median: 68.5 years).[50] A

multinational study on 174 AIS patients with SARS-CoV-2 infection reported a median age of 71 years.[24] A national

surveillance study from the United Kingdom presented 77 SARS-CoV-2 infected patients with cerebrovascular events

and reported a median age of 74 years.[14] The authors stated that the overall median age of 71 in 125 SARS-CoV-2

infected patients with cerebrovascular disorders, altered mental status, and peripheral neuropathy in this study is

comparable to national data from the UK Government public health bodies over the same period. However, caution

should be made when interpreting the summary results from patients with different cerebrovascular disorders.

In our clustering, all models showed differences among the subgroups in terms of the TOAST criteria. In addition, we

observed the absence of traditional stroke risk factors in about one-fourth of the AIS patients (subgroup a of EX-A$_3$).

Comparing these patients with our stroke registry of 8,929 AIS patients (GNSIS, Geisinger NeuroScience Ischemic

Stroke database) indicated a higher prevalence of patients with no known risk factor (22.0% in this study versus 11.5%

in GNSIS, p<0.001). We also observed that the patients in this study were younger (60.0±18.0 years versus 62.9±17.6

years, p=0.004) and had more severe strokes (NIHSS of 8[4-22] versus 4[2-7], p<0.001). Further investigation is

needed to present the etiology and possible risk factors among these patients as well as to define the underlying

predispositions and pathogenic pathways occurring in SARS-CoV-2 infected individuals.

Among the AIS patients, 13.6% of patients received intravenous thrombolysis, while 7.4% underwent mechanical

thrombectomy. These rates are similar to the multinational study on 174 AIS SARS-CoV-2 infected patients (12.7%

thrombolysis, 6.9% thrombolysis, and thrombectomy, and 5.2% mechanical thrombectomy).[24] Although the rate of

LVOs and IVT were almost the same in various regions, we observed a significant difference among the regions in

the rate of mechanical thrombectomy—12.4% versus 2% in countries with lower health expenditure; Middle East: 2.6%, Asia: zero, America: 4.5%, and Europe: 21.1%. Overall, 44.5% of ischemic strokes were due to LVOs, without any age or sex predominance. This rate is comparable to a similar report from New York.[50] LVOs accounted for 24% to 46% of acute ischemic strokes.[51] When the definition of the LVOs is limited to ICA, MCA (M1 and M2), ACA (A1), PCA (P1), VA, and BA, similar to our study, the prevalence drops to 24-38%,[52,53] which was lower than our findings. We could not identify any significant difference in the risk of LVOs among various regions in our study (Supplemental Tables 10).

Stroke subtyping based on the TOAST criteria revealed a 33% large artery atherosclerosis (LAA) in our study, which is higher than reports from worldwide population-based studies (19-23%).[54,55] In the Middle East, our study presented 53.5% LAA, compared with the previously reported rate of 8-31%.[56–59] Small-vessel occlusion (SVO) accounted for stroke etiology in 10% of our patients, and analysis of neuro-imaging patterns showed 10.2% lacunar infarcts (lacune). These rates are lower than worldwide population-based studies—21-44% SVO,[55,60] and 21-30% lacune [55,61,62] (Supplemental Table 22). Likewise, we observed lower regional rates of SVO and lacune: in Europe, 4.1% SVO and 9.3% lacune, versus previous reports of 12-31% SVO [63–69]and 14-31% lacune;[65,70–72] in the North and South America, 5.3% SVO and 9.1% lacune, versus reported SVO rates of 15-18%[55,73] and 13-18% lacune;[55,62] and in the Middle Eastern countries, 18.3% SVO and 11.3% lacune, versus 20-25% SVO[56–60,74] and 19-26% lacune.[59,75,76] We could not identify any difference in imaging findings or subclasses of the TOAST criteria among SARS-CoV-2 infected patients who presented with chief complaints of stroke-related symptoms compared to patients with a prior diagnosis of SARS-CoV-2 infection. Similar to our findings, other reports on SARS-CoV-2 infected stroke patients suggested the lower rate of SVO and lacune, and a predilection of SARS-CoV-2 for inducing LVO.[77–79] Lacunar infarctions and SVO are more likely to produce milder deficits.[61,80–82] During the COVID-19 pandemic, patients with mild to moderate stroke symptoms were less likely to present at medical centers.[5,78,79,83] In addition, less severe stroke symptoms, mostly in critically ill patients or overwhelmed health centers, were more likely to be under-diagnosed. We observed a lower median NIHSS score in countries with middle to high-health expenditure—8[3-17] versus 11[5-17] in countries with lower health expenditure. The regional difference in the severity of stroke in our study—NIHSS of 12[6-17] in the Middle East, versus 7[0-16] in America and 8[4-18] in Europe— may either indicate that the patients with less severe strokes did not present to the hospital or reflect the under-diagnosis in some countries. Future studies such as CASCADE (Call to Action: SARS-CoV-2 and

Cerebrovascular DisordErs) are required to shed light on changes in stroke care protocols and hospitalization rate during the pandemic.[84]

Our study reported 91 patients with intracranial hemorrhage. Among the patients with SAH—23, no aneurysm was detected in 69.5% patients, which is higher than reported 15%(5%-34%) spontaneous SAH among patients without SARS-CoV-2 infection.[85] We observed that 27.9% of IPH patients had no vascular risk factor or comorbidities (subgroup 'a' of the EX-A3 model). These patients had higher ICH score and younger age in comparison with other patients with IPH.

In this study we observed 18 stroke patients with CVST; The average age of patients was 49 years, 78% were younger than 55 years old, and more than 60% were women. CVST was reported as a consequence of various viral infections such as the Varicella Zoster virus,[86–89] Influenza virus,[90] Herpes Simplex virus,[91] SARS,[92] and SARS-CoV-2.[11,93] Classically CVST is considered to occur in young adults, with the predilection of women.[94] However, the sex ratio varies widely – 44.7% to 83.4% in women.[95] A systematic review on the sex ratio of 23,638 patients with CVST demonstrated an increasing trend of women proportion (54.8% before 1981 to 69.8% after 2001), likely due to increased use of oral contraceptives.[96] Even though CVST patients in our study were younger than patients with other stroke subtypes, they were older than previously reported CVST patients without SARS-CoV-2 infection.[95–100] In addition, only 27.8% of the patients in our study had multiple sinus or venous involvement, which is considerably lower than previous reports in non-SARS-CoV-2 infected patients.[95,99,101,102] One reason might be the severe condition of the patients with multiple CVSTs that prevent the proper diagnosis of these patients.

To our knowledge, this is the largest study on stroke in patients infected with SARS-CoV-2. This work has several limitations. Despite that we included centers from multiple countries and presented a comprehensive panel of patients' characteristics, some of the specific laboratory parameters related to rare stroke causes (e.g., antiphospholipid antibodies) were not included in this study. The collaborators tried to identify SARS-CoV-2 patients who presented with stroke as the first and only symptom, but difficulty in measuring all symptoms related to COVID-19 (such as fatigue, anosmia, and ageusia) should be taken into consideration. Due to the small sample and heterogeneity of the patients with subclasses of SAH or CVST, we did not apply the machine learning on this subgroup for further exploration. Although attempts were made to minimize the selection bias by including patients from different

ethnicities, ecological conditions, and health care systems, this study may suffer from selection bias and low power in some subgroups. Further studies that include a control population are warranted.

In conclusion, we observed a relatively high number of young, and asymptomatic SARS-CoV-2 infections among stroke patients. Traditional vascular risk factors were absent among a relatively large cohort of patients. Among hospitalized patients, the stroke severity was lower and rate of mechanical thrombectomy was higher among countries with middle to high-health expenditure

Acknowledgement: We appreciate the efforts of all other health care providers and administrators who contributed data for this study.

Sources of funding: None

Disclosure: None

References:

[1]   Mao L, Jin H, Wang M, Hu Y, Chen S, He Q, et al. Neurologic Manifestations of Hospitalized Patients With Coronavirus Disease 2019 in Wuhan, China. JAMA Neurol 2020. doi:10.1001/jamaneurol.2020.1127.

[2]   Li Y, Wang M, Zhou Y, Chang J, Xian Y, Mao L, et al. Acute Cerebrovascular Disease Following COVID-19: A Single Center, Retrospective, Observational Study. SSRN Electron J 2020;19. doi:10.2139/ssrn.3550025.

[3]   Oxley TJ, Mocco J, Majidi S, Kellner CP, Shoirah H, Singh IP, et al. Large-Vessel Stroke as a Presenting Feature of Covid-19 in the Young We. N Engl J Med 2020. doi:10.1056/NEJMc2001272.

[4]   Sweid A, Hammoud B, Weinberg JH, Oneissi M, Raz E, Shapiro M, et al. Letter: Thrombotic Neurovascular Disease in COVID-19 Patients. Neurosurgery 2020;0. doi:10.1093/neuros/nyaa254.

[5]   Yaghi S, Ishida K, Torres J, Mac Grory B, Raz E, Humbert K, et al. SARS2-CoV-2 and Stroke in a New York Healthcare System. Stroke 2020:STROKEAHA120030335. doi:10.1161/STROKEAHA.120.030335.

[6]   Mehrpour M, Shuaib A, Farahani M, Hatamabadi HR, Fatehi Z, Ghaffari M, et al. EXPRESS: COVID-19 and stroke in Iran; a case series, and effects on stroke admissions. Int J Stroke 2020:174749302093739. doi:10.1177/1747493020937397.

[7]   Morassi M, Bagatto D, Cobelli M, D'Agostini S, Gigli GL, Bnà C, et al. Stroke in patients with SARS-CoV-2 infection: case series. J Neurol 2020:1–8. doi:10.1007/s00415-020-09885-2.

[8]   D'Anna L, Kwan J, Brown Z, Halse O, Jamil S, Kalladka D, et al. Characteristics and clinical course of Covid-19 patients admitted with acute stroke. J Neurol 2020. doi:10.1007/s00415-020-10012-4.

[9]   Craen A, Logan G, Ganti L. Novel Coronavirus Disease 2019 and Subarachnoid Hemorrhage: A Case Report. Cureus 2020;12. doi:10.7759/cureus.7846.

[10]   Reddy ST, Garg T, Shah C, Nascimento FA, Imran R, Kan P, et al. Cerebrovascular Disease in Patients with COVID-19: A Review of the Literature and Case Series. Case Rep Neurol 2020;77030:199–209. doi:10.1159/000508958.

[11]   Poillon G, Obadia M, Perrin M, Savatovsky J, Lecler A. Cerebral venous thrombosis associated with COVID-19 infection: Causality or coincidence? J Neuroradiol 2020. doi:10.1016/j.neurad.2020.05.003.

[12]   Zhang Y, Xiao M, Zhang S, Xia P, Cao W, Jiang W, et al. Coagulopathy and Antiphospholipid Antibodies in Patients with Covid-19. N Engl J Med 2020;382:e38. doi:10.1056/nejmc2007575.

[13]   Escalard S, Maïer B, Redjem H, Delvoye F, Hébert S, Smajda S, et al. Treatment of Acute Ischemic Stroke due to Large Vessel Occlusion With COVID-19. Stroke 2020:1–4. doi:10.1161/strokeaha.120.030574.

[14]   Varatharaj A, Thomas N, Ellul MA, Davies NWS, Pollak TA, Tenorio EL, et al. Neurological and neuropsychiatric complications of COVID-19 in 153 patients: a UK-wide surveillance study. The Lancet Psychiatry 2020;2:1–8. doi:10.1016/S2215-0366(20)30287-X.

[15]   Barrios-López JM, Rego-García I, Muñoz Martínez C, Romero-Fábrega JC, Rivero Rodríguez M, Ruiz Giménez JA, et al. Ischaemic stroke and SARS-CoV-2 infection: a causal or incidental association? Neurol (English Ed 2020. doi:10.1016/j.nrleng.2020.05.008.

[16]   Klok FA, Kruip MJHA, van der Meer NJM, Arbous MS, Gommers DAMPJ, Kant KM, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. Thromb Res 2020. doi:10.1016/j.thromres.2020.04.013.

[17]   García Espinosa J, Moya Sánchez E, Martínez Martínez A. Severe COVID-19 with debut as bilateral pneumonia, ischemic stroke, and acute myocardial infarction. Med Clin (Barc) 2020. doi:10.1016/j.medcli.2020.04.012.

[18]   Gomez-Arbelaez D, Ibarra-Sanchez G, Garcia-Gutierrez A, Comanges-Yeboles A, Ansuategui-Vicente M, Gonzalez-Fajardo JA. Covid-19-Related Aortic Thrombosis: a Report of Four Cases. Ann Vasc Surg 2020. doi:10.1016/j.avsg.2020.05.031.

[19]   Viguier A, Delamarre L, Duplantier J, Olivot JM, Bonneville F. Acute ischemic stroke complicating common carotid artery thrombosis during a severe COVID-19 infection. J Neuroradiol 2020. doi:10.1016/j.neurad.2020.04.003.

[20]   Malentacchi M, Gned D, Angelino V, Demichelis S, Perboni A, Veltri A, et al. Concomitant brain arterial

and venous thrombosis in a COVID-19 patient. Eur J Neurol 2020:0–3. doi:10.1111/ene.14380.

[21]   Valderrama EV, Humbert K, Lord A, Frontera J, Yaghi S. Severe Acute Respiratory Syndrome Coronavirus 2 Infection and Ischemic Stroke. Stroke 2020:STROKEAHA120030153. doi:10.1161/STROKEAHA.120.030153.

[22]   Tunç A, Ünlübaş Y, Alemdar M, Akyüz E. Coexistence of COVID-19 and acute ischemic stroke report of four cases. J Clin Neurosci Off J Neurosurg Soc Australas 2020. doi:10.1016/j.jocn.2020.05.018.

[23]   Avula A, Nalleballe K, Narula N, Sapozhnikov S, Dandu V, Toom S, et al. COVID-19 presenting as stroke. Brain Behav Immun 2020. doi:10.1016/j.bbi.2020.04.077.

[24]   Ntaios G, Michel P, Georgiopoulos G, Guo Y, Li W, Xiong J, et al. Characteristics and Outcomes in Patients With COVID-19 and Acute Ischemic Stroke The Global COVID-19 Stroke Registry. Stroke 2020. doi:10.1016/j.amjcard.2019.02.021.

[25]   Shahjouei S, Naderi S, Li J, Khan A, Chaudhary D, Farahmand G, et al. Risk of Cerebrovascular Events in Hospitalized Patients with SARS-CoV-2 Infection. Ebiomedicine 2020. doi:http://dx.doi.org/10.2139/ssrn.3605289.

[26]   Ver Hage A. The NIH stroke scale: a window into neurological status. Com Nurs Spectr 2011;24:44–49.

[27]   Adams H., Bendixen B., Kappelle L., Biller J, Love B., Gordon D., et al. Classification of Subtype of Acute Ischemic Stroke. Stroke 1993;23:35–41. doi:10.1161/01.STR.24.1.35.

[28]   von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: Guidelines for reporting observational studies. Int J Surg 2014;12:1495–1499. doi:10.1016/j.ijsu.2014.07.013.

[29]   Equator Network. Enhancing the QUAlity and Transparency Of health Research. Univ Oxford 2019. https://www.equator-network.org.

[30]   Sacco RL, Kasner SE, Broderick JP, Caplan LR, Connors JJB, Culebras A, et al. An updated definition of stroke for the 21st century: a statement for healthcare professionals from the American Heart Association/American Stroke Association. Stroke 2013;44:2064–2089.

doi:https://dx.doi.org/10.1161/STR.0b013e318296aeca.

[31] WHO. Laboratory testing for coronavirus disease 2019 (COVID-19) in suspected human cases. Interim

Guid 2020:1–7.

[32] Prabhakaran S, Silver AJ, Warrior L, McClenathan B, Lee VH. Misdiagnosis of Transient Ischemic Attacks

in the Emergency Room. Cerebrovasc Dis 2008;26:630–635. doi:10.1159/000166839.

[33] Sadighi A, Stanciu A, Banciu M, Abedi V, Andary N El, Holland N, et al. Rate and associated factors of

transient ischemic attack misdiagnosis. ENeurologicalSci 2019;15:100193.

doi:10.1016/j.ensci.2019.100193.

[34] Nadarajan V, Perry RJ, Johnson J, Werring DJ. Transient ischaemic attacks: mimics and chameleons. Pract

Neurol 2014;14:23–31. doi:https://dx.doi.org/10.1136/practneurol-2013-000782.

[35] Potter GM, Marlborough FJ, Wardlaw JM. Wide variation in definition, detection, and description of

lacunar lesions on imaging. Stroke 2011. doi:10.1161/STROKEAHA.110.594754.

[36] Wessels T, Röttger C, Jauss M, Kaps M, Traupe H, Stol E. Identification of embolic stroke patterns by

diffusion-weighted MRI in clinically defined lacunar stroke syndromes. Stroke 2005.

doi:10.1161/01.STR.0000158908.48022.d7.

[37] Bang OY, Ovbiagele B, Liebeskind DS, Restrepo L, Yoon SR, Saver JL. Clinical determinants of infarct

pattern subtypes in large vessel atherosclerotic stroke. J Neurol 2009;256:591–599. doi:10.1007/s00415-

009-0125-x.

[38] Abdel Razek AAK, Alvarez H, Bagg S, Refaat S, Castillo M. Imaging spectrum of CNS vasculitis.

Radiographics 2014;34:873–894. doi:10.1148/rg.344135028.

[39] Waqas M, Mokin M, Primiani CT, Gong AD, Rai HH, Chin F, et al. Large Vessel Occlusion in Acute

Ischemic Stroke Patients: A Dual-Center Estimate Based on a Broad Definition of Occlusion Site. J Stroke

Cerebrovasc Dis 2020;29:104504.

[40] F. C. Defining Young Stroke. YoungStroke, Inc 2017:1–15.

[41]   Global Health Expenditure Database n.d. https://apps.who.int/nha/database/Select/Indicators/en (accessed June 30, 2020).

[42]   IBM. Downloading IBM SPSS Statistics 26. Ibm 2020.

[43]   Von Luxburg U. A tutorial on spectral clustering. Stat Comput 2007;17:395–416. doi:10.1007/s11222-007-9033-z.

[44]   Palacio-Niño J-O, Berzal F. Evaluation Metrics for Unsupervised Learning Algorithms. ArXiv 2019:arXiv-1905.

[45]   Andy Bunn MK. An Introduction to dplR. Ind Commer Train 2008;10:11–18. doi:10.1108/eb003648.

[46]   Warnes GR, Bolker B, Bonebakker L, Gentleman R, Huber W, Liaw A, et al. gplots: Various R programming tools for plotting data. R Packag Version 2009;2:1.

[47]   Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn: Machine learning in Python. J Mach Learn Res 2011;12:2825–2830.

[48]   Cabral NL, Freire AT, Conforto AB, Dos Santos N, Reis FI, Nagel V, et al. Increase of stroke incidence in young adults in a middle-income country a 10-year population-based study. Stroke 2017;48:2925–2930. doi:10.1161/STROKEAHA.117.018531.

[49]   Kissela BM, J.C. K, Alwell K, Moomaw JC, Woo D, Adeoye O, et al. Age at stroke. Neurol 1781-1787 2012;79:1781–1787.

[50]   Yaghi S, Ishida K, Torres J, Grory B Mac, Raz E, Humbert K, et al. SARS2-CoV-2 and Stroke in a New York Healthcare System 2020:1–10. doi:10.1161/STROKEAHA.120.030335.

[51]   Rennert RC, Wali AR, Steinberg JA, Santiago-Dieppa DR, Olson SE, Pannell JS, et al. Epidemiology, Natural History, and Clinical Presentation of Large Vessel Ischemic Stroke. Clin Neurosurg 2019;85:S4–S8. doi:10.1093/neuros/nyz042.

[52]   Malhotra K, Gornbein J, Saver JL. Ischemic strokes due to large-vessel occlusions contribute disproportionately to stroke-related dependence and death: A review. Front Neurol 2017;8:1–5.

doi:10.3389/fneur.2017.00651.

[53]   Dozois A, Hampton L, Kingston CW, Lambert G, Porcelli TJ, Sorenson D, et al. Plumber study (prevalence of large vessel occlusion strokes in mecklenburg county emergency response). Stroke 2017;48:3397–3399. doi:10.1161/STROKEAHA.117.018925.

[54]   Ornello R, Degan D, Tiseo C, Di Carmine C, Perciballi L, Pistoia F, et al. Distribution and temporal trends from 1993 to 2015 of ischemic stroke subtypes a systematic review and meta-analysis. Stroke 2018;49:814–819. doi:10.1161/STROKEAHA.117.020031.

[55]   O'Donnell MJ, Denis X, Liu L, Zhang H, Chin SL, Rao-Melacini P, et al. Risk factors for ischaemic and intracerebral haemorrhagic stroke in 22 countries (the INTERSTROKE study): A case-control study. Lancet 2010;376:112–123. doi:10.1016/S0140-6736(10)60834-3.

[56]   Saber H, Thrift AG, Kapral MK, Shoamanesh A, Amiri A, Farzadfard MT, et al. Incidence, recurrence, and long-term survival of ischemic stroke subtypes: A population-based study in the Middle East. Int J Stroke 2017;12:835–843. doi:10.1177/1747493016684843.

[57]   Khealani BA, Khan M, Tariq M, Malik A, Siddiqi AI, Awan S, et al. Ischemic strokes in Pakistan: Observations from the national acute ischemic stroke database. J Stroke Cerebrovasc Dis 2014;23:1640–1647. doi:10.1016/j.jstrokecerebrovasdis.2014.01.009.

[58]   Khorvash F, Khalili M, Rezvani Habibabadi R, Sarafzadegan N, Givi M, Roohafza H, et al. Comparison of acute ischemic stroke evaluation and the etiologic subtypes between university and nonuniversity hospitals in Isfahan, Iran. Int J Stroke 2019;14:613–619. doi:10.1177/1747493019828648.

[59]   Lutski M, Zucker I, Shohat T, Tanne D. Characteristics and outcomes of young patients with first-ever ischemic stroke compared to older patients: The national acute stroke Israeli registry. Front Neurol 2017;8. doi:10.3389/fneur.2017.00421.

[60]   Rukn S Al, Mazya M V., Hentati F, Sassi S Ben, Nabli F, Said Z, et al. Stroke in the Middle-East and North Africa: A 2-year prospective observational study of stroke characteristics in the region—Results from the Safe Implementation of Treatments in Stroke (SITS)–Middle-East and North African (MENA). Int J Stroke

2019;14:715–722. doi:10.1177/1747493019830331.

[61]   M. P, L. N, P. D, B. L. Outcomes from ischemic stroke subtypes classified by the Oxfordshire Community Stroke Project: A systematic review. Eur J Phys Rehabil Med 2011;47:19–23.

[62]   O'Donnell MJ, Chin SL, Rangarajan S, Xavier D, Liu L, Zhang H, et al. Global and regional effects of potentially modifiable risk factors associated with acute stroke in 32 countries (INTERSTROKE): a case-control study. Lancet 2016;388:761–775. doi:10.1016/S0140-6736(16)30506-2.

[63]   Li L, Yiin GS, Geraghty OC, Schulz UG, Kuker W, Mehta Z, et al. Incidence, outcome, risk factors, and long-term prognosis of cryptogenic transient ischaemic attack and ischaemic stroke: A population-based study. Lancet Neurol 2015;14:903–913. doi:10.1016/S1474-4422(15)00132-5.

[64]   Yesilot Barlas N, Putaala J, Waje-Andreassen U, Vassilopoulou S, Nardi K, Odier C, et al. Etiology of first-ever ischaemic stroke in European young adults: The 15 cities young stroke study. Eur J Neurol 2013;20:1431–1439. doi:10.1111/ene.12228.

[65]   Ihle-Hansen H, Thommessen B, Wyller TB, Engedal K, Fure B. Risk factors for and incidence of subtypes of ischemic stroke. Funct Neurol 2012;27:35–40.

[66]   Hauer AJ, Ruigrok YM, Algra A, van Dijk EJ, Koudstaal PJ, Luijckx G-J, et al. Age-Specific Vascular Risk Factor Profiles According to Stroke Subtype. J Am Heart Assoc 2017;6:e005090. doi:10.1161/JAHA.116.005090.

[67]   Carrera E, Maeder-Ingvar M, Rossetti AO, Devuyst G, Bogousslavsky J. Trends in Risk Factors, Patterns and Causes in Hospitalized Strokes over 25 Years: The Lausanne Stroke Registry. Cerebrovasc Dis 2007;24:97–103. doi:10.1159/000103123.

[68]   Grau AJ, Weimar C, Buggle F, Heinrich A, Goertler M, Neumaier S, et al. Risk factors, outcome, and treatment in subtypes of ischemic stroke: The German stroke data bank. Stroke 2001;32:2559–2566. doi:10.1161/hs1101.098524.

[69]   Wafa HA, Wolfe CDA, Rudd A, Wang Y. Long-term trends in incidence and risk factors for ischaemic stroke subtypes: Prospective population study of the South London Stroke Register. PLoS Med 2018;15:1–

16. doi:10.1371/journal.pmed.1002669.

[70]   Czlonkowska A, Ryglewicz D, Weissbein T, Baranska-Gieruszczak M, Hier DB. A prospective community-based study of stroke in warsaw, poland. Stroke 1994;25:547–551. doi:10.1161/01.STR.25.3.547.

[71]   Alzamora MT, Sorribes M, Heras A, Vila N, Vicheto M, Forés R, et al. Ischemic stroke incidence in Santa Coloma de Gramenet (ISISCOG), Spain. A community-based study. BMC Neurol 2008;8:5. doi:10.1186/1471-2377-8-5.

[72]   Di Carlo A, Lamassa M, Baldereschi M, Pracucci G, Consoli D, Wolfe CDA, et al. Risk factors and outcome of subtypes of ischemic stroke. Data from a multicenter multinational hospital-based registry. The European Community Stroke Project. J Neurol Sci 2006;244:143–150. doi:10.1016/j.jns.2006.01.016.

[73]   Porcello Marrone LC, Diogo LP, De Oliveira FM, Trentin S, Scalco RS, De Almeida AG, et al. Risk factors among stroke subtypes in Brazil. J Stroke Cerebrovasc Dis 2013;22:32–35. doi:10.1016/j.jstrokecerebrovasdis.2011.05.022.

[74]   Senel GB, Elmali AD, Mehrvar K, Farhoudi M, Aboutalebi M, Rezaei M, et al. A survey from Turkey and Iran on comparison of risk factors and etiology in ischemic stroke. Iran J Neurol 2020;18:176–178. doi:10.18502/ijnl.v18i4.2189.

[75]   Kumral E, Özkaya B, Sagduyu A, Şirin H, Vardarli E, Pehliva M. The Ege Stroke Registry: A hospital-based study in the Aegean Region, Izmir, Turkey. Cerebrovasc Dis 1998;8:278–288. doi:10.1159/000015866.

[76]   Ghandehari K, Izadi-Mood Z. Khorasan stroke registry: Analysis of 1392 stroke patients. Arch Iran Med 2007;10:327–334. doi:07103/AIM.009.

[77]   Oxley TJ, Mocco J, Majidi S, Kellner CP, Shoirah H, Singh IP, et al. Large-Vessel Stroke as a Presenting Feature of Covid-19 in the Young. N Engl J Med 2020;382:e60. doi:10.1056/NEJMc2009787.

[78]   Jain R. Evolving Neuroimaging Findings during COVID-19. Am J Neuroradiol 2020;41:1–2. doi:10.3174/ajnr.a6658.

[79]   Siegler JE, Heslin ME, Thau L, Smith A, Jovin TG. Falling stroke rates during COVID-19 pandemic at a

comprehensive stroke center: Cover title: Falling stroke rates during COVID-19. J Stroke Cerebrovasc Dis 2020;29:104953. doi:10.1016/j.jstrokecerebrovasdis.2020.104953.

[80] Favate AS, Younger DS. Epidemiology of Ischemic Stroke. Neurol Clin 2016;34:967–980. doi:10.1016/j.ncl.2016.06.013.

[81] Wilterdink JL, Bendixen B, Adams HP, Woolson RF, Clarke WR, Hansen MD. Effect of prior aspirin use on stroke severity in the Trial of Org 10172 in Acute Stroke Treatment (TOAST). Stroke 2001;32:2836–2840. doi:10.1161/hs1201.099384.

[82] Pittock SJ, Meldrum D, Hardiman O, Thornton J, Brennan P, Moroney JT. The Oxfordshire Community Stroke Project Classification: Correlation with imaging, associated complications, and prediction of outcome in acute ischemic stroke. J Stroke Cerebrovasc Dis 2003;12:1–7. doi:10.1053/jscd.2003.7.

[83] Hoyer C, Ebert A, Huttner HB, Puetz V, Kallmünzer B, Barlinn K, et al. Acute Stroke in Times of the COVID-19 Pandemic: A Multicenter Study. Stroke 2020:STROKEAHA120030395. doi:10.1161/STROKEAHA.120.030395.

[84] Abootalebi S, Aertker BM, Andalibi MS, Asdaghi N, Aykac O, Azarpazhooh MR, et al. Call to Action: SARS-CoV-2 and CerebrovAscular DisordErs (CASCADE). J Stroke Cerebrovasc Dis 2020;29. doi:10.1016/j.jstrokecerebrovasdis.2020.104938.

[85] Kim YW, Lawson MF, Hoh BL. Nonaneurysmal subarachnoid hemorrhage: An update. Curr Atheroscler Rep 2012;14:328–334. doi:10.1007/s11883-012-0256-x.

[86] Sujay Sada, Anjaneyulu Kammineni, Meena A. Kanikannan JA. Cerebral sinus venous thrombosis: A rare complication of primary Varicella zoster virus. Neurol India 2012;60:645–646.

[87] Imam SF, Lodhi O ul haq, Fatima Z, Nasim S, Malik WT, Saleem MS. A Unique Case of Acute Cerebral Venous Sinus Thrombosis Secondary to Primary Varicella Zoster Virus Infection. Cureus 2017;9. doi:10.7759/cureus.1693.

[88] Siddiqi SA, Nishat S, Kanwar D, Ali F, Azeemuddin M, Wasay M. Cerebral venous sinus thrombosis: Association with primary varicella zoster virus infection. J Stroke Cerebrovasc Dis 2012;21:917.e1-917.e4.

1   doi:10.1016/j.jstrokecerebrovasdis.2012.04.013.

2   [89]   Sudhaker B, Dnyaneshwar MP, Jaidip CR, Rao SM. Cerebral venous sinus thrombosis (CVST) secondary to

3   varicella induced hypercoagulable state in a adult. Intern Med Insid 2014;2:1. doi:10.7243/2052-6954-2-1.

4   [90]   Taniguchi D, Nakajima S, Hayashida A, Kuroki T, Eguchi H, Machida Y, et al. Deep cerebral venous

5   thrombosis mimicking influenza-associated acute necrotizing encephalopathy: A case report. J Med Case

6   Rep 2017;11:10–15. doi:10.1186/s13256-017-1444-7.

7   [91]   Leite J, Ribeiro A, Gonçalves D, Sargento-Freitas J, Trindade L, Duque V. Cerebral Venous Thrombosis as

8   Rare Presentation of Herpes Simplex Virus Encephalitis. Case Rep Infect Dis 2019;2019:1–5.

9   doi:10.1155/2019/7835420.

10   [92]   KRAYENBUEHL H. Thrombosis of the cerebral veins and sinus. Acta Neurochir Suppl (Wien) 1961;Suppl

11   7:248–254.

12   [93]   Dahl-Cruz F, Guevara-Dalrymple N, López-Hernández N. [Cerebral venous thrombosis and SARS-CoV-2

13   infection]. Rev Neurol 2020;70:391–392. doi:10.33588/rn.7010.2020204.

14   [94]   Luo Y, Tian X, Wang X. Diagnosis and Treatment of Cerebral Venous Thrombosis: A Review. Front Aging

15   Neurosci 2018;10. doi:10.3389/fnagi.2018.00002.

16   [95]   Gunes HN, Cokal BG, Guler SK, Yoldas TK, Malkan UY, Demircan CS, et al. Clinical associations,

17   biological risk factors and outcomes of cerebral venous sinus thrombosis. J Int Med Res 2016;44:1454–

18   1461. doi:10.1177/0300060516664807.

19   [96]   Zuurbier SM, Middeldorp S, Stam J, Coutinho JM. Sex differences in cerebral venous thrombosis: A

20   systematic analysis of a shift over time. Int J Stroke 2016;11:164–170. doi:10.1177/1747493015620708.

21   [97]   Dentali F, Poli D, Scoditti U, di Minno MND, Stefano VD, Siragusa S, et al. Long-term outcomes of

22   patients with cerebral vein thrombosis: A multicenter study. J Thromb Haemost 2012;10:1297–1302.

23   doi:10.1111/j.1538-7836.2012.04774.x.

24   [98]   Coutinho JM, Zuurbier SM, Aramideh M, Stam J. The incidence of cerebral venous thrombosis: A cross-

25   sectional study. Stroke 2012;43:3375–3377. doi:10.1161/STROKEAHA.112.671453.

[99]   Sidhom Y, Mansour M, Messelmani M, Derbali H, Fekih-Mrissa N, Zaouali J, et al. Cerebral venous thrombosis: clinical features, risk factors, and long-term outcome in a Tunisian cohort. J Stroke Cerebrovasc Dis 2014;23:1291–1295.

[100]   Ferro JM, Aguiar de Sousa D. Cerebral Venous Thrombosis: an Update. Curr Neurol Neurosci Rep 2019;19:1–9. doi:10.1007/s11910-019-0988-x.

[101]   Sassi S Ben, Touati N, Baccouche H, Drissi C, Romdhane N Ben, Hentati F. Cerebral venous thrombosis: a tunisian monocenter study on 160 patients. Clin Appl Thromb 2017;23:1005–1009.

[102]   Janghorbani M, Zare M, Saadatnia M, Mousavi SA, Mojarrad M, Asgari E. Cerebral vein and dural sinus thrombosis in adults in Isfahan, Iran: Frequency and seasonal variation. Acta Neurol Scand 2008;117:117–121. doi:10.1111/j.1600-0404.2007.00915.x.

Figure 1. Flow chart. Overview of the included patients, approaches, and modeling.

3   Figure 2. The image demonstrates the hierarchical clustering of the patients based on the presence (blue line) or
4   absence of the comorbidities. A. SARS-CoV-2 infected patients with acute ischemic stroke. B. SARS-CoV-2
5   infected patients with intraparenchymal hemorrhage.

Table 1. The baseline characteristics, comorbidities, and laboratory findings among SARS-CoV-2 infected stroke.

Parameter	Acute Ischemic Stroke N = 323 (74.8%)	Intracerebral Hemorrhage N = 68 (15.7%)	Subarachnoid Hemorrhage N = 23 (5.3%)	Cerebral Venous/Sinus Thrombosis N = 18 (4.2%)
Age; Mean (SD); Years	67.2 ± 15.2	63 ± 16	62.6 ± 14.1	48.9 ± 11.5
Age; Median [IQR]; Years	68 [58 – 78]	65 [54 – 75]	62 [54 – 74]	51 [39.5 – 55.2]
<40; N (%)	23 (7.1)	8 (12.9)	1 (4.3)	5 (27.8)
41-64; N (%)	110 (34.1)	20 (32.3)	12 (52.2)	11 (61.1)
65-74; N (%)	77 (23.8)	13 (21)	5 (21.7)	2 (11.1)
≥75; N (%)	113 (53)	21 (33.9)	5 (21.7)	0 (0)
Sex; Women; N (%)	130 (40.2)	30 (44.1)	12 (52.2)	11 (61.1)
Stroke Symptoms as the Chief Complaint; N (%)	104 (36.1)	31 (53.4)	8 (36.4)	4 (22.2)
SARS-CoV-2 to Stroke Interval; Mean (SD); Days	5 ± 7	5 ± 8	3 ± 5	3 ± 4
SARS-CoV-2 to Stroke Interval; Median [IQR]; Days	3 [0-9]	0 [0-10.3]	1.5 [0-5.3]	4.5 [0.7-10.3]
National Institutes of Health Stroke Scale (NIHSS); Median [IQR]	9 [4-17]	10 [0-19]	6 [0-16]	14 [7-17]
Intracerebral Hemorrhage (ICH) Score; Median [IQR]	-	3 [2 - 4]	-	-
Mechanical Ventilation; N (%)	238 (73.7)	31 (50)	12 (52.2)	10 (55.6)
Initiation of Mechanical Ventilation; Median [IQR]; Days	2 [1-3]	2 [1-4]	5 [1-15]	1 [1-2]
Length of Hospital Stay; Mean (SD); Days	12 ± 12	15 ± 18	18 ± 20	18 ± 14
Length of Hospital Stay; Median [IQR]; Days	7 [4-16]	8 [3-21]	11 [4.2-31]	14 [7-30]
Disposition*				
Discharged Home; N (%)	127 (42.8)	10 (17.9)	3 (15)	12 (66.7)
In-Hospital Mortality; N (%)	82 (27.6)	35 (62.5)	11 (55)	3 (16.7)
Still in Hospital, or Disposioned to Subacute Care; N (%)	88 (29.6)	11 (22)	6 (30)	3 (16.7)
Medications				
Prior Anti-Platelet Therapy; N (%)	87 (28.0)	16 (28.1)	3 (14.3)	0 (0.0)
Prior Anti-Coagulant Therapy; N (%)	28 (9.0)	6 (10.5)	2 (9.5)	4 (23.5)
Region				
Middle East; N (%)	153 (47.4)	43 (63.2)	17 (73.9)	15 (83.3)
America; N (%)	6 (1.9)	0 (0.0)	0 (0.0)	2 (11.1)
Europe; N (%)	88 (27.2)	19 (27.9)	6 (26.1)	1 (5.6)
Asia; N (%)	76 (23.5)	6 (8.8)	0 (0.0)	0 (0.0)
Countries' Health Expenditure				
Middle to High; N (%)	170 (52.6)	26 (38.2)	5 (21.7)	2 (11.1)
Low; N (%)	153 (47.4)	42 (61.8)	18 (78.3)	16 (88.9)
Comorbidities				
Hypertension; N (%)	202 (63.1)	43 (63.2)	14 (60.9)	1 (5.6)
Diabetes Mellitus; N (%)	111 (34.6)	21 (30.9)	4 (17.4)	2 (11.1)
Ischemic Heart Disease; N (%)	72 (24.3)	12 (17.6)	4 (21.1)	0 (0)
Atrial Fibrillation; N (%)	45 (14.1)	3 (4.4)	2 (8.7)	0 (0)
Carotid Stenosis; N (%)	38 (12.8)	2 (2.9)	0 (0)	0 (0)
Chronic Kidney Disease; N (%)	42 (13.1)	9 (13.2)	3 (13)	2 (11.1)
Cardiac Ejection Fraction <40%; N (%)	24 (8.1)	7 (10.3)	0 (0)	0 (0)
Active Neoplasm; N (%)	21 (7.1)	4 (5.9)	1 (4.3)	0 (0)
Rheumatological Disease; N (%)	5 (1.7)	0 (0)	1 (4.3)	0 (0)
Prior Stroke or Transient Ischemic Attack; N (%)	5 (1.7)	0 (0)	0 (0)	0 (0)
Smoking; N (%)	53 (16.6)	11 (16.2)	3 (13)	0 (0)

Parameter	Acute Ischemic Stroke N = 323 (74.8%)	Intracerebral Hemorrhage N = 68 (15.7%)	Subarachnoid Hemorrhage N = 23 (5.3%)	Cerebral Venous/Sinus Thrombosis N = 18 (4.2%)
Laboratory findings				
White Blood Cell Count; Mean (SD); x109/L	9.8 ± 4.8	11.4 ± 10.0	13.5 ± 12.9	11.1 ± 6.0
White Blood Cell Count; Median [IQR]; x109/L	9 [6.8 – 11.2]	9.5 [7.7 – 12.1]	10 [8.8 – 12.2]	11.1 [6.7 – 14.8]
<4 x109/L; N (%)	7 (2.3)	3 (5.7)	0 (0)	2 (22.2)
4-10 x109/L; N (%)	184 (61.1)	29 (54.7)	10 (50)	4 (44.4)
10-20 x109/L; N (%)	96 (31.9)	18 (34)	9 (45)	3 (33.3)
≥20 x109/L; N (%)	14 (4.7)	5 (10)	1 (5)	0 (0)
Neutrophil Count; Mean (SD); x109/L	7.7 ± 4.5	9.3 ± 8.5	9.6 ± 6.1	9.2 ± 5.1
Neutrophil Count; Median [IQR]; x109/L	6.8 [4.8 – 9.2]	7.2 [5.2 – 10.6]	7.3 [5.4 – 11]	9.4 [4.7 – 14.2]
<4 x109/L; N (%)	43 (15.2)	5 (10)	1 (5)	1 (12.5)
4-10 x109/L; N (%)	182 (64.3)	33 (66)	12 (60)	6 (75)
10-20 x109/L; N (%)	49 (17.3)	10 (20)	6 (30)	1 (12.5)
≥20 x109/L; N (%)	9 (3.2)	2 (4)	1 (5)	0 (0)
Lymphocyte Count; Mean (SD); x109/L	1.5 ± 1.5	2.0 ± 3.6	2.8 ± 6.3	2.0 ± 1.3
Lymphocyte; Median [IQR]; x109/L	1.3 [0.9 – 1.9]	1.3 [0.9– 2.0]	1.3 [0.9-1.9]	1.9 [1.2 – 2.3]
<1 x109/L; N (%)	93 (32.5)	21 (39.6)	6 (28.6)	1 (12.5)
1-2 x109/L; N (%)	130 (45.5)	19 (35.8)	10 (47.6)	4 (50)
2-3 x109/L; N (%)	41 (14.3)	9 (35.8)	3 (14.3)	2 (25)
3-4 x109/L; N (%)	11 (3.8)	1 (1.9)	0 (0)	1 (12.5)
≥4 x109/L; N (%)	11 (3.8)	3 (5.7)	2 (9.5)	0 (0)
Platelet Count; Mean (SD); x109/L	314.5 ± 440.7	197.1 ± 89.1	191.45 ± 60.9	254.2 ± 170.1
Platelet Count; Median [IQR]; x109/L	229 [161 – 333.7]	178.5 [142.5 – 254]	183 [141 – 246]	237 [139 – 337]
<350 x109/L; N (%)	228 (78.1)	50 (94.3)	20 (100)	8 (88.9)
350-500 x109/L; N (%)	64 (21.9)	3 (5.7)	0 (0)	1 (11.1)
Alanine Transaminase (ALT); Mean (SD); U/L	63.3 ± 86.3	50.4 ± 66.0	67.2 ± 78.1	120.9 ± 231.6
Alanine Transaminase (ALT); Median [IQR]; U/L	29.8 [10.1 – 90.2]	22.9 [8.0– 67.7]	53 [11.6 – 75.2]	39 [29 – 107]
Aspartate Transaminase (AST); Mean (SD); U/L	32.1 ± 26.8	35.7 ± 24.5	43.4 ± 27.9	121.5 ± 265.1
Aspartate Transaminase (AST); Median [IQR]; U/L	23.9 [14-40.5]	28.9 [20 – 43]	35 [21 – 69.8]	44 [30.5 –63]
Blood Urea Nitrogen (BUN); Mean (SD); mg/dl	53.1± 104.2	99.9 ± 290.8	83.1 ± 160.6	31.9 ± 16.3
Blood Urea Nitrogen (BUN) mg/dl; Median [IQR]	32 [21 – 50]	33.5 [21.5 – 51]	36 [26 – 60]	30 [23 – 37]
Creatinine; Mean (SD); mg/dl	1.5 ± 1.7	2.2 ± 2.6	2.57 ± 3.12	1.46 ± 1.21
Creatinine; Median [IQR]; mg/dl	1.1 [1.1 – 1.5]	1.1 [0.9 – 1.8]	1 [0.9 – 2.8]	1.1 [0.9 – 1.6]
C-Reactive Protein (CRP); Mean (SD); mg/L	61 ± 131	79 ± 192	85 ± 155	54 ± 37
C-Reactive Protein (CRP); Median [IQR]; mg/L	36 [24 – 57]	37 [24 – 54]	41 [29.5 – 62.5]	30 [23.5 – 37]
Lactate Dehydrogenase (LDH); Mean (SD); U/L	604.8 ± 1536.7	450.8 ± 443.3	443.0 ± 268.5	766 ± 812.1
Lactate Dehydrogenase (LDH); Median [IQR]; U/L	377 [245 - 524]	366.5 [254.5 - 571]	252 [230 - 664]	453 [250 -1115]
Fibrinogen; Mean (SD); mg/dl	463.6 ± 989.6	706.1 ± 418.1	-	-
Fibrinogen; Median [IQR]; mg/dl	223 [3.9 – 490.5]	693 [303 - 1246]	-	-
D- Dimer; Mean (SD); ng/ml	2654.8 ± 6429.4	8668.4 ± 16318.1	-	1080.8 ± 1376.5
D- Dimer; Median [IQR]; ng/ml	1027 [551 - 2200]	2303 [584.5 - 6875]	-	-

* Data on patients' disposition were sparse.

Table 2. Neurological findings among patients with acute ischemic stroke.

Parameters	N (%)	Parameters		N (%)
Imaging Pattern of Ischemia		Vascular Territory based on Imaging*		
Embolic/Large Vessel Athero-thromboembolism	206 (80.5)	Anterior Cerebral Artery	Unilateral	7 (2.2)
Lacunar	26 (10.2)		Bilateral	1 (0.3)
Borderzone	23 (9.0)	Middle Cerebral Artery	Unilateral	181 (56)
Vasculitis	1 (0.4)		Bilateral	5 (1.5)
		Posterior Cerebral Artery	Unilateral	33 (10.2)
TOAST Criteria			Bilateral	4 (1.2)
Large Artery Atherosclerosis	56 (32.9)	Posterior Inferior Cerebellar Artery		9 (2.8)
Cardioembolism	46 (27.1)	Missing		64 (19.8)
Small Artery Occlusion	17 (10.0)	Symptoms at Onset*		
Other Determined Etiology	13 (7.6)	Hemineglect		1 (0.4)
Undetermined Etiology	38 (22.4)	Limb Paresis	Unilateral	168 (72.4)
			Bilateral	2 (0.9)
National Institute of Health Stroke Scale (NIHSS)		Sensory Loss	Unilateral	26 (11.2)
No Stroke Symptoms (NIHSS=0)	20 (7.5)		Bilateral	1 (0.4)
Minor Stroke (NIHSS of 1 to 4)	49 (18.4)	Visual Field Loss	Unilateral	14 (6.0)
Moderate Stroke (NIHSS of 5 to 15)	117 (43.8)		Bilateral	2 (0.9)
Moderate to Severe Stroke (NIHSS of 16 to 20)	35 (13.1)	Gaze Preference		13 (5.6)
Severe Stroke (NIHSS 21-42)	46 (17.2)	Facial Paresis		46 (19.8)
		Aphasia		56 (24.1)
Large vessel Occlusion	126 (44.5)	Dysarthria		41 (17.7)
		Altered Level of Consciousness		57 (24.6)
Mechanical Thrombectomy	24 (7.4)	Ataxia		21 (9.1)
		Seizure		7 (3.0)
Intravenous Thrombolysis	44 (13.6)	Missing		91 (28.2)

*Due to multiple infarcts in some patients, the summation may exceed 100%.

Table 3. Baseline characteristics and neuroimaging findings under each outcome measures .

Parameters	TOAST Criteria						Large Vessel Occlusion			National Institutes of Health Stroke Scale (NIHSS)					
	A: Large Artery Atherosclerosis N = 56 (32.9%)	B: Cardio-embolism N = 46 (27.1%)	C: Small Artery Occlusion N = 17 (10.0%)	D: Other Determined N = 13 (7.6%)	E: Undetermined N = 38 (22.4%)	P value	A: Large Vessel Occlusion N = 126 (44.5%)	B: Other Strokes N = 157 (55.5%)	P value	A: No Stroke Symptoms (NIHSS = 0) N = 20 (7.5%)	B: Minor Stroke (NIHSS = 1 to 4) N = 49 (18.4%)	C: Moderate Stroke (NIHSS 5 to 15) N = 117 (43.8%)	D: Moderate to Severe Stroke (NIHSS = 16 to 20) N = 35 (13.1%)	E: Severe Stroke (NIHSS = 21 to 42) N = 46 (17.2%)	P-value
Age; Mean (SD); Years	63 ± 15	72 ± 14 D (0.021)	67 ± 18	57 ± 16	65 ± 18	0.01	65.7 ± 14.	68 ± 16.	0.22	67 ± 13	65 ± 17	66 ± 15	66 ± 16	69 ± 15	0.77
Sex; Female; N (%)	19 (33.9)	21 (45.7)	3 (17.6)	9 (69.2) C (0.043)	15 (39.5)	0.05	46 (36.5	71 (45.2	0.14	6 (30.0)	20 (40.8)	44 (37.6)	16 (45.7)	22 (47.8	0.60
Large Vessel Occlusion; N (%)	46 (85.2) B (0.003) D (<0.001) E (<0.001)	24 (53.3)	0 (0.0)	2 (18.2)	7 (20.0)	<0.001	-	-	-	1 (8.3)	8 (17.0) B (0.01)	49 (44.5) A (0.004) B (<0.001)	23 (67.6) A (<0.001) B (<0.001) C (0.01)	31 (75.6)	<0.001
Intravenous Thrombolysis; N (%)	14 (25.0)	8 (17.4)	0 (0.0)	2 (15.4)	4 (10.5)	0.12	28 (22.2) B (0.002)	14 (8.9)	0.002	0 (0.0)	2 (4.1)	25 (21.4) B (0.04)	11 (31.4) B (0.004)	5 (10.9)	<0.001
Mechanical Thrombectomy; N (%)	10 (17.9)	10 (21.7)	0 (0.0)	1 (7.7)	2 (5.3)	0.07	24 (19.0)	0 (0.0)	<0.001	0 (0.0)	1 (2.0)	8 (6.8)	9 (25.7) B (0.01) C (0.01)	6 (13.0)	<0.001
National Institutes of Health Stroke Scale (NIHSS); Median [IQR]	9.0 [5.0 – 17.0	13.0 [8.0 -20.0]	4.0 [2.0 -8.0]	14.0 [6.0 – 18.0]	7.0 [3.0 – 17.0]	0.10	15.0 [8.0 – 21.0]	6.0 [3.0 – 12.0	<0.001	-	-	-	-	-	-
Imaging Patterns															
Embolic/Large Vessel athero-Thromboembolism; N (%)	54 (96.4) C (<0.001) D (0.017)	43 (93.5) C (<0.001)	4 (23.5)	9 (69.2)	34 (89.5) C (<0.001)		99 (88.4 B (0.003)	99 (73.3	<0.001	4 (57.1)	31 (73.8)	85 (81.7)	24 (75.0)	39 (92.9)	<0.001
Lacune; N (%)	0 (0.0)	1 (2.2)	13 (76.5) B (<0.001) D (0.001) E (<0.001)	1 (7.7)	4 (10.5)	<0.001	2 (1.8)	12 (17.0 B (<0.001		2 (28.6)	8 (19.0)	10 (9.6)	1 (3.1)	1 (2.4)	
Borderzone; N (%)	2 (3.6)	1 (2.2)	0 (0.0)	3 (23.1) A (0.044) B (0.024)	0 (0.0)		11 (9.8)	12 (8.9)		0 (0.0)	3 (7.1)	9 (8.7)	7 (21.9)	2 (4.8)	
Vasculitis Pattern; N (%)	0 (0.0)	1 (2.2)	0 (0.0)	0 (0.0)	0 (0.0)		0 (0.0)	1 (0.7)		1 (14.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	

Interval Between infection Onset to Index Event; Median [IQR]; Days	4.0 [1.0 – 10.0]	2.0 [0.0 – 11.0]	2.0 [0.0 – 6.0]	10.0 [4.0 – 17.0]	4.0 [0.0 – 12.0]	0.12	46 (58.2) A (<0.001)	8 (9.6)	<0.001	3.0 [0.0 – 12.0]	3.0 [0.0 – 10.0]	4.0 [1.0 – 8.0]	3.0 [0.0 – 10.0]	1.0 [0.0 – 10.0]	0.64
Mechanical Ventilatation; N (%)	22 (29.3)	10 (21.7)	4 (23.5)	9 (69.2)	11 (28.9)	0.02	36 (28.6)	42 (26.8)	0.73	2 (10.0)	8 (16.3)	25 (21.4)	12 (34.3)	29 (63.0) A (0.001) B (<0.001) C (<0.001)	<0.001
Disposition*							0 (0.0)	17 (20.5)							
Discharged Home; N (%)	24 (42.9)	16 (34.8)	11 (64.7)	8 (61.5)	9 (24.3)		46 (37.7)	66 (44.6)		5 (62.5)	28 (62.2) D (0.001) E (0.001)	52 (47.3) D (0.24) E (0.02)	6 (17.6)	9 (20.5)	
In Hospital Mortality; N (%)	14 (25.0)	13 (28.3)	3 (17.6)	2 (15.4)	9 (24.3)	0.12	40 (32.8)	35 (23.6)	0.24	1 (12.5)	5 (11.1)	18 (16.4)	19 (55.9) B (<0.001) C (<0.001)	23 (52.3) B (<0.001) C (<0.001)	<0.001
Still in Hospital/Subacute Care; N (%)	18 (32.1)	17 (37.0)	3 (17.6)	3 (23.1)	19 (54.1)		36 (29.5)	47 (31.8)		2 (25.0)	12 (26.7)	40 (36.4)	9 (26.5)	12 (27.3)	
Length of Hospital Stay; Median (IQR); Days	6.0 [4.0 – 15.0]	7.0 [5.0 - 14.0]	7.0 [5.0 – 16.0]	20.0 [4.0 – 35.0]	8.0 [6.0 – 26.0]	0.17	6.0 [4.0 – 15.0])	8.0 [5.0 – 17.0]	0.10	2 (10.0)	8 (16.3)	25 (21.4)	12 (34.3)	29 (63.0) A (0.001) B (<0.001) C (<0.001)	<0.001

* Data on patients' disposition were sparse.

Table 4. Localization and presenting symptoms regarding the SARS-CoV-2 infected patients with intracranial hemorrhage and cerebral sinus and venous thrombosis.

Localization	N (%)	Symptoms	N (%)
Intracranial Hemorrhage			
Subarachnoid Hemorrhage	23 (25.3)	Intracranial (Minus Subarachnoid) Hemorrhage	68 (15.7)
Middle Cerebral Artery Aneurysm	2 (8.6)	Altered Level of Consciousness	27 (39.7)
Posterior Cerebral Artery Aneurysm	1 (4.3)	Limb Paresis	25 (36.7)
Anterior Communicating Artery Aneurysm	2 (8.6)	Aphasia	3 (4.4)
Basilar Top Aneurysm	2 (8.6)	Facial Paresis	2 (2.9)
No Aneurysm Detected	16 (69.5)	Sensory Loss	3 (2.9)
Subarachnoid and Intraparenchymal Hemorrhages	3 (3.3)	Dysarthria	4 (2.9)
Intraventricular and Intraparenchymal Hemorrhages	4 (4.4)	Ataxia	5 (2.9)
Intraparenchymal Hemorrhage	61 (67)	Visual Field Defect	1 (1.5)
Cerebellar Hemorrhage	3 (7.0)	Subarachnoid Hemorrhage	23 (25.3)
Brain Stem Hemorrhage	4 (9.3)	Thunderclap headache	16 (69.5)
Basal Ganglia Hemorrhage	19 (43.1)	Decreased level of consciousness	5 (21.7)
Thalamic Hemorrhage	7 (16.3)	Cerebral herniation	1 (4.3)
Lobar Hemorrhage	20 (46.5)	Seizure	1 (4.3)
Missing	17 (28.3)	Missing	21 (30.9)
Cerebral Venous and Sinus Thrombosis			
Superior Sagittal Sinus	6 (33.3)	Headache	9 (50)
Sigmoid Sinus	6 (33.3)	Seizure	7 (38.9)
Transverse Sinus	5 (27.8)	Altered Level of Consciousness	5 (27.8)
Lateral Sinus	3 (16.7)	Increased Intracerebral Pressure	2 (11.1)
Straight Sinus	1 (5.6)	Concomitant Intracranial Hemorrhage	2 (11.1)
Cortical Veins	1 (5.6)		
Multiple Sinus or Venous Involvement	5 (27.8)		