Artificial Intelligence-Based Analytics for Diagnosis of Small Bowel Enteropathies and Black Box Feature Detection

Authors: Sana Syed, MD, MS[1,2*]; Lubaina Ehsan, MBBS[1]; Aman Shrivastava, MS[1,3]; Saurav Sengupta, MS[1,3]; Marium Khan, MD[1]; Kamran Kowsari, PhD[4,5]; Shan Guleria, MD[6]; Rasoul Sali, MS[4]; Karan Kant, MS[3]; Sung-Jun Kang, MS[3]; Kamran Sadiq, MD[2]; Najeeha T. Iqbal, PhD[2]; Lin Cheng, MD[7]; Christopher A. Moskaluk, MD, PhD[8]; Paul Kelly, MD, FRCP[9,10]; Beatrice C. Amadi, MD, MMed[9]; S. Asad Ali, MD, MPH[2]; Sean R. Moore, MD, MS[1]; Donald E. Brown, PhD[4*]

Author Affiliations:

[1]Department of Pediatrics, School of Medicine, University of Virginia, Charlottesville, VA, USA

[2]Department of Pediatrics and Child Health, Aga Khan University, Karachi, Pakistan

[3]Data Science Institute, University of Virginia, Charlottesville, VA

[4]Systems and Information Engineering, University of Virginia, Charlottesville, VA, USA

[5]University of California Los Angeles, Los Angeles, CA, USA

[6]School of Medicine, University of Virginia, Charlottesville, VA, USA

[7]Pathology Department, Rush University Medical Center, Chicago, IL, USA

[8]Deparment of Pathology, University of Virginia, Charlottesville, VA, USA

[9]Tropical Gastroenterology and Nutrition group, University of Zambia School of Medicine, Lusaka, Zambia

[10]Blizard Institute, Barts and the London School of Medicine, Queen Mary University of London, London, United Kingdom

*co-corresponding authors

Contact Information for Corresponding Authors:

Sana Syed, MD, MS

Assistant Professor

Division of Gastroenterology, Hepatology & Nutrition

Department of Pediatrics

University of Virginia School of Medicine

Office: Room 2035A, MR-4 Bldg, 409 Lane Rd., Charlottesville, VA 22908

Phone: 434-297-5064

Email: sana.syed@virginia.edu

Donald E. Brown, PhD

W.S. Calcott Professor

Engineering Systems and Environment

Office: Olsson Hall 102C P.O. Box 400747, Charlottesville, Virginia 22904

Phone: 434-982-2074

Email: deb@virginia.edu

Funding Information: Research reported in this manuscript was supported by National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health under award number K23DK117061-01A1 (Syed), Bill and Melinda Gates Foundation under award numbers OPP1066203 (Ali) and OPP1066118 (Kelly), University of Virginia Center for Engineering in Medicine Grant (Syed and Brown), and University of Virginia THRIV Scholar Career Development Award (Syed). The content is solely the responsibility of the authors and does not necessarily represent the official views of the funding agencies.

Word Count (Manuscript Body): 3,490

Abstract

Objectives: Striking histopathological overlap between distinct but related conditions poses a

significant disease diagnostic challenge. There is a major clinical need to develop computational

methods enabling clinicians to translate heterogeneous biomedical images into accurate and

quantitative diagnostics. This need is particularly salient with small bowel enteropathies;

Environmental Enteropathy (EE) and Celiac Disease (CD). We built upon our preliminary analysis

by developing an artificial intelligence (AI)-based image analysis platform utilizing deep learning

convolutional neural networks (CNNs) for these enteropathies.

Methods: Data for secondary analysis was obtained from three primary studies at different sites.

The image analysis platform for EE and CD was developed using convolutional neural networks

(CNNs: ResNet and custom Shallow CNN). Gradient-weighted Class Activation Mappings (Grad-

CAMs) were used to visualize the models' decision making process. A team of medical experts

simultaneously reviewed the stain color normalized images done for bias reduction and Grad-

CAM visualizations to confirm structural preservation and biological relevance, respectively.

Results: 461 high-resolution biopsy images from 150 children were acquired. Median age

(interquartile range) was 37·5 (19·0 to 121·5) months with a roughly equal sex distribution; 77

males (51·3%). ResNet50 and Shallow CNN demonstrated 98% and 96% case-detection

accuracy, respectively, which increased to 98·3% with an ensemble. Grad-CAMs demonstrated

models' ability to learn distinct microscopic morphological features.

Conclusion: Our AI-based image analysis platform demonstrated high classification accuracy for

small bowel enteropathies which was capable of identifying biologically relevant microscopic

features, emulating human pathologist decision making process, performing in the case of

suboptimal computational environment, and being modified for improving disease classification

accuracy. Grad-CAMs that were employed illuminated the otherwise 'black box' of deep learning

in medicine, allowing for increased physician confidence in adopting these new technologies in

clinical practice.

Keywords: Biopsy Image Analysis, Environmental Enteropathy, Global Health, Intestinal

Structure, Convolutional Neural Networks

What is known

• Striking histopathological overlap exists between distinct but related conditions which

poses a significant disease diagnostic challenge; such as for small bowel enteropathies

including Environmental Enteropathy (EE) and Celiac Disease (CD).

• There is a major clinical need to develop computational [including Artificial Intelligence (AI)

and deep learning] methods enabling clinicians to translate heterogeneous biomedical

images into accurate and quantitative diagnostics.

• A major issue plaguing the use of AI in medicine is the so-called 'black box' of deep

learning, an analogy which describes the lack of insight that humans have into how the

models arrive at their decision-making

What is new

• AI-based image analysis platform demonstrated high classification accuracy for small

bowel enteropathies (EE vs. CD vs. histologically normal controls).

• Gradient-weighted Class Activation Mappings (Grad-CAMs) illuminated the otherwise

'black box' of deep learning in medicine, allowing for increased physician confidence in

adopting these new technologies in clinical practice.

<u>Abbreviations</u>

EE: Environmental Enteropathy

CD: Celiac Disease

AI: Artificial Intelligence

CNN: Convolutional Neural Networks

Grad-CAMs: Gradient-weighted Class Activation Mappings

ResNet: Deep residual network

H&E: Hematoxylin and Eosin

PK: Pakistan

ZA: Zambia

UVA: University of Virginia

US: United States

EEDBI: Environmental Enteropathic Dysfunction Biopsy Initiative

IQR: inter-quartile range

LAZ/ HAZ: Length/ Height-for-Age Z score

ROC: Receiver Operating Curves

AUC: Area Under the Curve

<u>Introduction</u>

A major challenge of interpreting clinical biopsy images to diagnose disease is the often striking overlap in histopathology between distinct but related conditions.(1-3) Due to this significant clinical need, computational methods can pave the way for accurate and quantitative diagnostics(4). Computational modeling enhancements in medicine, particularly for image analysis, have shown the potential benefit of artificial intelligence (AI) for disease characterization.(5) There is an increasing interest in the utilization of deep learning,(6, 7) a subset of AI involving iterative optimization strategies based on pixel-by-pixel image evaluation.(8) These AI-based deep learning models, in particular Convolutional Neural Networks (CNNs), have been shown to be effective for medical image analysis.(5) CNNs have demonstrated potential for image feature extraction from diseases relying on radiological and histopathological diagnosis, particularly in the fields of ophthalmology and oncology.(9-13) Deep residual network (ResNet) is a CNN that has repeatedly shown success for image classification as it is optimized to gather fine grain attributes from regions of interest within the image.(14-16) It has also outperformed early deep learning models such as AlexNet(17) and VGG(18) and achieved superior performance on the ImageNet(19, 20) and COCO(21) image recognition benchmarks. While the deep learning models are noteworthy of their own accord, a major issue plaguing the use of AI in medicine is the so-called 'black box' of deep learning, an analogy which describes the lack of insight that humans have into how the models arrive at their decision-making.(22)

An accurate AI-based biopsy image analysis platform may enable efficient detection and differentiation of small bowel enteropathy damaged tissue architecture features which is challenging due to histopathological overlap.(2, 23-25) This will not only enable pathologists to filter and pre-populate scans, improving turn-around time, but also provide insight into previously

uncharacterized tissue features unique to complex small bowel enteropathies such as

Environmental Enteropathy (EE) and Celiac Disease (CD).

EE has been linked to poor sanitation and hygiene with dire consequences such as cognitive

decline, oral vaccine response, and growth failure.(3) EE assessment and its histopathological

differentiation from similar diseases (i.e. CD) is an essential task performed by pathologists.(26)

Even though an EE scoring system has been proposed, it will benefit from approaches narrowing

down specific cellular parameters important for an accurate EE diagnosis.(27) CD is an immune-

mediation condition with sensitivity to gluten leading to small bowel injury. Modified Marsh score

system(28, 29) has been used to classify the severity of CD but it does not account for complex

disease features, such as the role of goblet or enteroendocrine cells, that can potentially improve

diagnostic accuracy.

We have previously published a histopathological analysis model demonstrating 93.4%

classification accuracy for identifying and differentiating between duodenal biopsies from children

with EE and CD.(30) We also added a layer of explainability by using deconvolutions but they

lacked biopsy regions of interest being specifically highlighted and did not fully explain the deep

learning decision making process.(22) Despite increasing utilization of deep learning

architectures in medicine, they remain underutilized for improving diagnostic accuracy and

differentiation between histologically similar small bowel enteropathies. We now build upon our

prior work to address knowledge gaps in the approaches previously reported. We aimed to: 1)

optimize datasets by removing unavoidable bias acquired by archival data sourcing from multiple

sites by using a hematoxylin and eosin (H&E) color normalization method with structure-

preserving capabilities; 2) deploy an enteropathy focused deep learning CNN model with multiple

layers modified to gather fine grain attributes from image regions of interest; 3) mimic the

interpretation methodology of human pathologists by using a combination of multi-zoom and

reduced parameter approaches (shallow deep learning models); and, 4) improve visualization of

deep learning classification decision making.

Methods

Study Design and Archival Biopsy Image Dataset Acquisition

<u>Archival Specimen Sources</u>

Archival data for secondary analyses was sourced from: 1) primary Environmental Enteropathy

(EE) studies from Pakistan (PK; Aga Khan University) and Zambia (ZA; University of Zambia

School of Medicine, University Teaching Hospital), conducted 2013 to 2015,(31, 32) and 2) the

University of Virginia (UVA), United States (US). UVA biopsies were from archival specimens with

clinical diagnoses of CD and histologically normal controls (referred to as controls) from

participants who had undergone esophagogastroduodenoscopy in the past 25 years (data

accessed: 1992 to 2017) as part of an archival controls sub-study, methods reported

elsewhere.(33) Biopsies available from each site for this project were archival and had been

previously processed per local institutional H&E slide staining, tissue sectioning, and tissue

paraffin embedding protocols. Our timeline for secondary data collection and analysis was from

November 2017 to December 2019.

<u>Biopsy Digitization</u>

Biopsy slides for our secondary analysis had been previously digitized at high resolutions

(average 20,000 by 20,000 pixels) to allow visualization of microscopic cellular features up to 40x

(PK, US) and 20x (ZA) magnification. Digitization was done using Olympus VS 120 (Olympus

Corporation Inc., Center Valley, Pennsylvania), Aperio Scanscope CS scanners (Leica

Biosystems Division of Leica Microsystems Inc., Buffalo Grove, IL, United States), and Leica

SCN400 brightfield scanner (Leica Microsystems CMS GmbH, Germany) at PK, ZA, and US,

respectively. US archival biopsies were scanned using a prior detailed methods and work from

these datasets has been published elsewhere.(2, 30, 32)

<u>Archival Biopsy Image Data Acquisition</u>

For the biopsy images acquired, diagnosis of EE was as previously defined in primary studies. At

each site, clinical pathologists made diagnoses based on histological and clinical findings.(31, 32)

For the US archival images, duodenal biopsy slides were obtained from the Biorepository and

Tissue Research Facility at UVA with disease diagnoses as per clinical pathology reports.

Biopsies from participants reported as controls were only included if there was no disease in any

other part of the gastrointestinal tract (e.g., eosinophilic esophagitis, inflammatory bowel disease,

H. Pylori gastritis, post-transplant liver disease etc.) or overall (e.g., patients with solid organ

transplant, leukemia etc.).

Biopsy Image Analysis Model Design

<u>Dataset Pre-processing</u>

Biopsy Image Patch Creation

High resolution whole slide images are patched for deep learning models to account for

computational limitations.(34) Images were split into patches of 1000x1000 and 2000x2000 pixels

with an overlap in horizontal and vertical axes of 750 and 1000 pixels, respectively. Patches that

contained less than 50% tissue area were discarded and the rest were resized to 256x256 pixels

(details noted in supplemental methods Figure S1). Each biopsy whole slide image generated an

average of 250 1000x1000 and 40 2000x2000 patches.

Sample Size Augmentation, Balancing, and Justification

As there were more biopsy images for CD, EE and control images were up-sampled to balance

datasets. Extensive data augmentations were performed for better generalization. As biopsy

images exhibit both rotational and axial symmetry, a random combination of rotation (90, 180 or

270 degree angle), mirroring, and zoom (between 1x and 1·1x) was applied (Figure S2 in

supplement). Deep learning studies have focused on obtaining as much image data as possible

without any standard sample size recommendations.(34) There are limited studies focused on EE

deep learning models and the sample size of this model is larger compared to our preliminary

analysis.(30)

Stain Color Normalization

Stain color normalization using structure-preserving method as described by Vahadane et al.(35,

36) was used to eliminate bias due to color differences in the biopsies from different sites. It

involved empirically selecting a target patch to normalize color across all patches (Figure 1).

Three independent pathologists (LC, ZA, RI) completed a blind review of the color normalized

biopsy images from different sites to assess the structure-preserving ability of the method.

<u>Image Analysis Model: Deep Learning Computer-Aided Biopsy Disease Classification System</u>

Several deep learning Convolutional Neural Network (CNN) architectures were used to address

specific questions as outlined below:

1. *Need for the identification of microscopic features visible at high magnification (zoom)*

level: ResNet50 is a widely used deep CNN architecture with 50 layers for image

classification requiring identification of microscopic patterns.(14-16) Final decision layers

were modified to improve accuracy.(36) To combat data sparsity we used transfer

learning, an established methods used to improve training using limited datasets by pre-

training the model on the ImageNet dataset.(19, 37) Additional details of our modifications

to the ResNet50 architecture are described in the Appendix S1 in supplement.

2. *Emulate human pathologist decision making process for visualizing biopsies at multiple*

levels of magnification (zoom): we developed a framework incorporating multiple zoom

levels (Figure 2). Biopsy images were segmented into 2000x2000 pixels then further into

1000x1000 pixel patches with an overlap of 750 pixels (horizontal and vertical axes). Two

independent ResNet50 models were independently trained and the 1000x1000 and

2000x2000 corresponding patches were paired. Each pair was passed through the

respective trained ResNet50 model and features from the last fully connected layer of

each model were extracted and concatenated for an overall representation of specific

regions of the biopsy images. This concatenated vector was further passed through a set

of trainable linear layers for the final classification.

3. *Reduce computational model complexity used for disease classification:* Custom shallow

CNN architecture was designed to help reduce the number of parameters the model

optimized for disease classification. This Shallow CNN consisted of three convolutional

layers (Figure 2; detailed methods in Appendix S2 as part of the Supplement).

4. *Explore methods of improving disease classification by using a combined model:*

Ensemble models have been shown to generally improve the accuracy and robustness of

classification.(38) We combined Resnet50 and shallow CNN to improve model

classification accuracy. We obtained 3 values from the output layer of each CNN

architecture which were then passed into a softmax function to obtain prediction

probabilities for each patch being EE, CD or control. These predictions were then

aggregated to acquire the final and better prediction for each biopsy image.

Visualization of Model Decision-Making

CNN activation maps were generated using Gradient-weighted Class Activation Mappings (Grad-

CAMs)(39) to visualize the regions of interest utilized by the model for decision making (detailed

methods outlined in Appendix S3 in the Supplement). These were reviewed by a team of medical

professionals [pathologist specialized in gastroenterology (CM), pediatric gastroenterologist (SS)]

enabling corroboration of the model results with incumbent classification. Since human intuition

is required to assess if features highlighted can be biologically explained, we used the

Environmental Enteropathic Dysfunction Biopsy Initiative (EEDBI) scoring system for EE(27) and

the modified Marsh score classification for CD(28, 29) to inform human medical professional

intuition (Figure S3 and Table S1 in the Supplement).

<u>Base Case for Comparison Using Existing Computer Vision Approach</u>

An alternative method, CellProfiler,(40) using cellular feature extraction for explainability of the

models was also explored (Appendix S4 and Figure S4 in the Supplement). CellProfiler isolated

nucleated cells from the biopsy images and classified EE vs. CD vs. controls based on the cellular

feature differences, achieving 65% accuracy.

Ethical Considerations

The secondary analyses as part of this study were approved by University of Virginia Institutional

Review Board. Ethical approval for prior original primary studies was obtained from: 1) the Ethical

Review Committee of Aga Khan University, Karachi, PK (informed consent obtained from parents

and/ or guardians for EE cases), 2) the Biomedical Research Ethics Committee of the University

of Zambia School of Medicine, University Teaching Hospital, Lusaka, ZA (informed consent

obtained from caregivers for EE cases), and 3) University of Virginia Institutional Review Board

(waiver of consent granted).

This manuscript has been prepared in accordance with STARD guidelines.(41)

Results

Biopsy Image Dataset

We obtained 461 digitized biopsy images (171 H&E stained duodenal biopsy glass slides) from 150 participants (US: 124; PK: 10; ZA: 16). Our EE data consisted of 29 and 19 biopsy images from 10 PK and 16 ZA participants, respectively. Data from 124 UVA participants; 63 CD, 61 controls, was available for these analyses.

Population Clinical Characteristics

Of the 150 participants, 77 (51·3%) were male. Median (inter-quartile range: IQR) age and LAZ/ HAZ (Length/ Height-for-Age Z score) of the EE participants was 22·2 (IQR: 20·8 to 23·4) months and -2·8 (IQR: -3·6 to -2·3), respectively (Table 1). Participants with EE were overall younger as compared to the US CD and controls (median age: 25·0; IQR: 16·5 to 41·0 months).

Stain Color Normalization Assessment

Our panel of blinded independent pathologists confirmed that medically relevant cell types (lymphocytes, polymorphonuclear neutrophils, epithelial cells, eosinophils, goblet cells, paneth cells, and neuroendocrine cells) were preserved after color normalization. The panel also commented that features used for EE assessment via the published EE scoring system(27) were also visible. Although the granularity of eosinophilic cytoplasm and sharpness of paneth cell globules was less appreciated in post color-normalized images, these features were consistent throughout the biopsy images and were therefore hypothesized to not result in classification bias.

Disease Classification Accuracy and Performance

Patches were used for training the models and the predictions were then aggregated to identify classifications for their parent biopsy images. Modified ResNet50 and Shallow CNN exhibited

overall accuracies at the biopsy level of 98% (sensitivity: 93%, specificity: 94%) and 96%

(sensitivity: 80%, specificity: 88%), respectively. The accuracy increased to 98·3% (sensitivity:

95%, specificity: 96%) with the ensemble. Accuracy of the multi-zoom ResNet50 architecture was

98% (sensitivity: 96%, specificity: 97%). Confusion matrices of the models' classification for each

disease are shown in Figure 4. Receiver operating curves (ROC) with area under the curve (AUC)

were used to assess the certainty of the model classifications. Biopsy level AUC for modified

ResNet50, ResNet50 with multi-zoom architecture, and shallow CNN were 0·99, 0·99, and 0·96,

respectively (Figure 3). Error analysis and performance statistics for the models are included in

Table S2 in the supplement.

Grad-CAM Interpretations

Grad-CAMs were obtained for all the models (Figure 5). Modified ResNet50 and ResNet50 Multi-

zoom mainly focused on similar areas for classification of EE vs CD vs controls while Shallow

CNN focused on distinct, yet medically relevant cellular features (Table S3 in supplemental

material for detailed Grad-CAM findings).

<u>Discussion</u>

Diagnosis of overlapping histological diseases, such as small bowel enteropathies, lacks

computational approaches for histopathological assessment.(30) While scoring systems such as

the EE score and modified Marsh score for EE and CD, respectively, have the potential to

diagnose and classify the severity of these diseases, there is a need to more accurately define

the cellular parameters that play an important role in diagnosis.(27) We developed an AI-based

image analysis platform, based on our preliminary results,(30) for the diagnosis of small bowel

enteropathies. To the best of our knowledge, similar small bowel enteropathy image analysis

platforms that include EE do not exist. We now build upon our prior work to address knowledge

gaps in the approaches previously reported. Our modified ResNet50, ResNet50 with multi-zoom,

Shallow CNN, and an ensemble model (modified ResNet50 and Shallow CNN) depicted overall

>90% classification accuracies. These AI-bases analytics and black box feature detection findings

pave the way for standardizing and improving diagnoses for small bowel enteropathies.

Prior to training our deep learning models, dataset preparation included stain color normalization

since with studies obtaining data from multiple sites for further analysis, an important

consideration is to optimize datasets to remove site-specific bias.(35, 42) In our case, the biopsy

images displayed visible stain color differences and the method we used to counter this bias has

previously been proposed by Vahadane et al.(35) Along with stain color normalization, it

successfully preserved small bowel architecture as assessed by our team of pathologists.

Our modified ResNet50 deep learning model trained to classify biopsy images of EE vs CD vs

healthy controls was able to identify microscopic features of each disease. This work has been

shown to be successful for other medical fields with ResNet providing superior performance for

biopsy images.(14-16) This expands upon our preliminary model, as with the use of ResNet50 it

provides powerful analytics for images. Further, the input to the previously published model(30)

and other similar published studies for other clinical diseases consisted of small patches of whole

slide images(15, 34) that represent approximately 2% of the original biopsy image. Such patch

inputs do not take into account the fact that tissue features can be better visualized at multiple

magnifications. The multi-zoom ResNet50 model employed in this study more closely emulates

the process of human pathologists, who visualize biopsies at multiple magnification (zoom) levels

for diagnoses.(43) The multi-zoom model's Grad-CAMs focused on biologically similar areas as

the model without multi-zoom, suggesting that the multi-zoom architecture does not bias the

model significantly.

To further build upon our deep learning methods, we incorporated a shallow CNN and a final

ensemble of ResNet and shallow CNN. The architecture of the shallow CNN was less complex

than that of ResNet, a useful quality if an optimal computational environment is unavailable. A

less complex model would also demonstrate utility if model parameters need to be reduced for a

first-pass analysis of biopsy images to show proof of concept. Although the shallow CNN Grad-

CAMs focused on far too many features for EE to be clinical useful, it focused on distinct and

biologically informative features for CD. These Grad-CAM results led us to hypothesize that the

shallow CNN would have better utility when combined with a more complex model such as

ResNet50. This combination was manifested in the ensemble model, which showed an overall

accuracy of 98.3%, higher than either model alone (ResNet50-only accuracy: 98%, Shallow-CNN-

only accuracy: 96%).

While these results are noteworthy of their own accord, we further extracted Grad-CAMs to help

illuminate the 'black box'(22) of these deep learning models, to improve the insight into how the

models arrive at their decision-making and provide the much needed 'explainability' for clinicians

and researchers. Grad-CAMs supported the models' use of biologically relevant features for

defining each class: EE, CD or controls. The image features visualized via Grad-CAMs, if

confirmed via molecular methods such as immunohistochemistry or RNA in situ hybridization,

may pave the way for optimizing the EE and modified Marsh (for CD) scoring systems by drawing

attention to the parameters pertinent to disease diagnoses.

Our secondary analysis built upon our earlier work by expanding and addressing the limitations.

We computationally balanced the datasets and further augmented them to combat our prior

issues with imbalanced datasets. In our preliminary work, we also utilized γ correction for stain

color normalization without an assessment of the method's structure-preserving ability. We have

now utilized a more robust stain color normalization method that preserves structure, confirmed

by a team of pathologists. We have also built upon the single model that we reported earlier to

create a complex AI-based image analysis platform which utilizes multiple models to ensure a

robust computational approach for histopathological image analysis. Lastly, our prior

deconvolution-based approach to explaining the decision-making process of the preliminary

model left much to be desired. Our current approach using Grad-CAMs to explain the models'

decision-making is a significant improvement, one more easily understood by the average medical

professional.

Major strengths of our secondary analysis study include the addition of data from 48 new patients

(in addition to the data reported for preliminary results) which increased the data volume for patch-

level analysis. We also assessed the validity of our models with a wide variety of performance

statistics (accuracy, sensitivity/ specificity, positive predictive value/ negative predictive value,

precision, recall, and F1-score), many of which are often missing in the work done for deep

learning in medicine. Our stain color normalization method for eliminating bias preserved tissue

structure, an important quality for any successful normalization process. Our multi-zoom

architecture emulating human pathologist biopsy assessment process, brings us one step closer

to automated detection of biopsy disease features. Further, our use of Grad-CAMs for

visualization of the models' regions of interest will help both increase the confidence in AI-based

decision making and also improve biopsy based small bowel enteropathy diagnosis.

Despite these strengths, we experienced several limitations. Since the data was obtained for

secondary analysis, we were limited in terms of improving data quality due to differences in the

scanners used for digitization and staining methods. Due to this we were unable to further modify

the EE dataset to account for high classification accuracies. Despite this, we utilized an approach

to potentially eliminate stain color differences. We were also limited in our ability to benchmark

our findings due to limited literature for the use of deep learning models for small bowel

architecture, especially the use of ResNet vs. custom deep learning approaches. To improve on

our current efforts, more robust color normalization methods are underway. Finally, our

techniques remain subject to the limitations inherent to tissue-based diagnostic and management

approaches in clinical medicine, e.g., patchiness of findings and site specific workflows for tissue

acquisition and processing.

<u>Conclusion</u>

Artificial Intelligence-based analytics provide an exciting opportunity for improving small bowel

enteropathy diagnostics with histological overlap. Our work suggests that models incorporated in

our image analysis platform are capable of identifying biologically relevant microscopic features,

emulating human pathologist decision making process, performing in the case of suboptimal

computational environment, and being modified for improving disease classification accuracy.

This work was improved by structure-preserving stain color normalization of the image inputs

along with visualizations of the model outputs, which are imperative for determining the tissue

features used for decision making. Further work will advance the clinical use of deep learning

models for enteropathies and other histopathology based diseases, improving the effectiveness

and efficiency of clinical care in gastroenterology and beyond.

<u>References</u>

1.   Sullivan PB, Marsh MN, Mirakian R, et al. Chronic diarrhea and malnutrition--histology of
the small intestinal lesion. Journal of pediatric gastroenterology and nutrition. 1991;12(2):195-
203.

2.   Syed S, Yeruva S, Herrmann J, et al. Environmental enteropathy in undernourished
Pakistani children: clinical and histomorphometric analyses. The American journal of tropical
medicine and hygiene. 2018;98(6):1577-84.

3.   Syed S, Ali A, Duggan C. Environmental enteric dysfunction in children: a review. Journal
of pediatric gastroenterology and nutrition. 2016;63(1):6.

4.   Topol EJ. High-performance medicine: the convergence of human and artificial
intelligence. Nature medicine. 2019;25(1):44.

5.   Sharma G, Carter A. Artificial Intelligence and the Pathologist: Future Frenemies?
Archives of pathology & laboratory medicine. 2017;141(5):622-3.

6.   Patel V, Khan MN, Shrivastava A, et al. Artificial intelligence applied to gastrointestinal
diagnostics: a review. Journal of pediatric gastroenterology and nutrition. 2020;70(1):4-11.

7.   Yang YJ, Bang CS. Application of artificial intelligence in gastroenterology. World J
Gastroenterol. 2019;25(14):1666.

8.   LeCun Y, Bengio Y, Hinton G. Deep learning. nature. 2015;521(7553):436.

9.   Medeiros FA, Jammal AA, Thompson AC. From Machine to Machine: An OCT-Trained
Deep Learning Algorithm for Objective Quantification of Glaucomatous Damage in Fundus
Photographs. Ophthalmology. 2019;126(4):513-21.

10.   Li Z, He Y, Keel S, et al. Efficacy of a Deep Learning System for Detecting Glaucomatous
Optic Neuropathy Based on Color Fundus Photographs. Ophthalmology. 2018;125(8):1199-206.

11.   Zhang Z, Xie Y, Xing F, et al., editors. Mdnet: A semantically and visually interpretable

medical image diagnosis network. Proceedings of the IEEE conference on computer vision and

pattern recognition; 2017.

12.   Coudray N, Ocampo PS, Sakellaropoulos T, et al. Classification and mutation prediction

from non–small cell lung cancer histopathology images using deep learning. Nature medicine.

2018;24(10):1559.

13.   Beck AH, Sangoi AR, Leung S, et al. Systematic analysis of breast cancer morphology

uncovers stromal features associated with survival. Science translational medicine.

2011;3(108):108ra13-ra13.

14.   Bejnordi BE, Veta M, Van Diest PJ, et al. Diagnostic assessment of deep learning

algorithms for detection of lymph node metastases in women with breast cancer. Jama.

2017;318(22):2199-210.

15.   Wei JW, Wei JW, Jackson CR, et al. Automated detection of celiac disease on duodenal

biopsy slides: A deep learning approach. Journal of pathology informatics. 2019;10.

16.   Jiang Y, Chen L, Zhang H, et al. Breast cancer histopathological image classification using

convolutional neural networks with small SE-ResNet module. PloS one. 2019;14(3).

17.   Krizhevsky A, Sutskever I, Hinton GE, editors. Imagenet classification with deep

convolutional neural networks. Advances in neural information processing systems; 2012.

18.   Simonyan K, Zisserman A. Very deep convolutional networks for large-scale image

recognition. arXiv preprint arXiv:14091556. 2014.

19.   Deng J, Dong W, Socher R, et al., editors. Imagenet: A large-scale hierarchical image

database. 2009 IEEE conference on computer vision and pattern recognition; 2009: Ieee.

20.   Russakovsky O, Deng J, Su H, et al. Imagenet large scale visual recognition challenge.

International journal of computer vision. 2015;115(3):211-52.

21.   Lin T-Y, Maire M, Belongie S, et al., editors. Microsoft coco: Common objects in context.

European conference on computer vision; 2014: Springer.

22.   Castelvecchi D. Can we open the black box of AI? Nature News. 2016;538(7623):20.

23.   Ramakrishna BS, Venkataraman S, Mukhopadhya A. Tropical malabsorption. Postgrad

Med J. 2006;82(974):779-87.

24.   Sullivan PB, Marsh MN, Mirakian R, et al. Chronic diarrhea and malnutrition--histology of

the small intestinal lesion. J Pediatr Gastroenterol Nutr. 1991;12(2):195-203.

25.   Pai RK, editor A practical approach to small bowel biopsy interpretation: celiac disease

and its mimics. Seminars in diagnostic pathology; 2014: Elsevier.

26.   Liu T-C, VanBuskirk K, Ali SA, et al. A novel histological index for evaluation of

environmental enteric dysfunction identifies geographic-specific features of enteropathy among

children with suboptimal growth. PLOS Neglected Tropical Diseases. 2020;14(1):e0007975.

27.   Liu T-C, VanBuskirk K, Ali SA, et al. A novel histological index for evaluation of

environmental enteric dysfunction identifies geographic-specific features of enteropathy among

children with suboptimal growth. PLoS neglected tropical diseases. 2020;14(1):e0007975.

28.   Modified Marsh Classification of histologic findings in celiac disease (Oberhuber) Stanford

Medicine [Available from: http://surgpathcriteria.stanford.edu/gi/celiac-disease/marsh.html.

29.   Oberhuber G. Histopathology of celiac disease. Biomedicine & pharmacotherapy.

2000;54(7):368-72.

30.   Syed S, Al-Boni M, Khan MN, et al. Assessment of Machine Learning Detection of

Environmental Enteropathy and Celiac Disease in ChildrenAssessment of Machine Learning

Detection of Environmental Enteropathy and Celiac Disease in ChildrenAssessment of Machine

Learning Detection of Environmental Enteropathy and Celiac Disease in Children. JAMA Network

Open. 2019;2(6):e195822-e.

31.   Iqbal NT, Sadiq K, Syed S, et al. Promising biomarkers of environmental enteric

dysfunction: a prospective cohort study in Pakistani children. Scientific reports. 2018;8(1):2966.

32.   Amadi B, Besa E, Zyambo K, et al. Impaired barrier function and autoantibody generation

in malnutrition enteropathy in Zambia. EBioMedicine. 2017;22:191-9.

33. Iqbal NT, Syed S, Sadiq K, et al. Study of Environmental Enteropathy and Malnutrition
(SEEM) in Pakistan: protocols for biopsy based biomarker discovery and validation. BMC
pediatrics. 2019;19(1):247.

34. Iizuka O, Kanavati F, Kato K, et al. Deep Learning Models for Histopathological
Classification of Gastric and colonic epithelial tumours. Scientific Reports. 2020;10(1):1-11.

35. Vahadane A, Peng T, Sethi A, et al. Structure-preserving color normalization and sparse
stain separation for histological images. IEEE transactions on medical imaging. 2016;35(8):1962-
71.

36. Shrivastava A, Kant K, Sengupta S, et al., editors. Deep Learning for Visual Recognition
of Environmental Enteropathy and Celiac Disease. 2019 IEEE EMBS International Conference
on Biomedical & Health Informatics (BHI); 2019: IEEE.

37. Pan SJ, Yang Q. A survey on transfer learning. IEEE Transactions on knowledge and data
engineering. 2009;22(10):1345-59.

38. Liu B, Cui Q, Jiang T, et al. A combinational feature selection and ensemble neural
network method for classification of gene expression data. BMC bioinformatics. 2004;5(1):136.

39. Selvaraju RR, Cogswell M, Das A, et al., editors. Grad-cam: Visual explanations from deep
networks via gradient-based localization. Proceedings of the IEEE International Conference on
Computer Vision; 2017.

40. Yu K, Zhang C, Berry G, et al. Predicting non-small cell lung cancer prognosis by fully
automated microscopic pathology image features. Nat Commun. 2016; 7: 12474. Epub
2016/08/17. https://doi. org/10.1038/ncomms12474 PMID: 27527408.

41. STARD 2015: An Updated List of Essential Items for Reporting Diagnostic Accuracy
Studies: Enhancing the QUAlity and Transparency Of health Research;   [Available from:
https://www.equator-network.org/reporting-guidelines/stard/.

42. Khan AM, Rajpoot N, Treanor D, et al. A nonlinear mapping approach to stain normalization in digital histopathology images using image-specific color deconvolution. IEEE Transactions on Biomedical Engineering. 2014;61(6):1729-38.

43. Brunyé TT, Mercan E, Weaver DL, et al. Accuracy is in the eyes of the pathologist: the visual interpretive process and diagnostic accuracy with digital whole slide images. Journal of biomedical informatics. 2017;66:171-9.

Author Contributions

Concept and design: Syed, Brown, Shrivastava, Sengupta, Kant, Kang, Kelly, Amadi, Ali, Moore.

Acquisition, analysis, or interpretation of data: Syed, Ehsan, Khan, Guleria, Shrivastava, Sengupta, Kowsari, Cheng, Moskaluk, Kelly, Ali, Moore, Brown.

Drafting of the manuscript: Syed, Ehsan, Shrivastava, Sengupta.

Critical revision of the manuscript for important intellectual content: Syed, Guleria, Khan, Iqbal, Moskaluk, Kelly, Ali, Moore, Brown.

Statistical analysis: Syed, Shrivastava, Sengupta, Kowsari, Sali, Ehsan, Brown.

Obtained funding: Syed, Amadi, Ali, Moore, Brown.

Administrative, technical, or material support: Syed, Ehsan, Sadiq, Iqbal, Kelly, Ali.

Supervision: Syed, Kelly, Ali, Moore, Brown.

Acknowledgements

Additional Contributions: We acknowledge the following individuals from Aga Khan University, Karachi, Pakistan, for contributing to this manuscript: Drs. Romana Idrees, MBBS, FCPS, and Zubair Ahmad, MBBS, FCPS, FRCPath, for assessment of stain color normalized biopsy images, Saad Mallick, medical student, for organization of image activation mappings, Dr. Saman Siddiqui, MD, for coordinating with the pathologists at Aga Khan University, and field workers (community health workers, led by Sadaf Jakhro, MSc, and coordinated by Dr. Tauseef Akhund, MBBS), data management unit (Najeeb Rahman, BS) and laboratory staff (Aneeta Hotwani, BS) for contributing to the data collection in this work. Patcharin Pramoonjago, PhD, Biorepository and Tissue Research Facility, University of Virginia, Charlottesville, assisted in data collection for this work. All individuals were compensated for their time.

Conflicts of Interest: None

Figure Legends

Figure 1: Stain Color Normalization – images in the top row highlight the differences in stain colors before normalization while the bottom row shows images post color normalization which was applied using the specified target image.

Figure 2: Combined figure explaining the methods for development of an image analysis model for diagnosis of small bowel enteropathies and black box feature detection: (A) Digitized duodenal biopsy slides were obtained for secondary analysis; (B and C) dataset was pre-processed by creation of biopsy image patches, sample size augmentation, and stain color normalization [shown in figure 1]; (D) development of the image analysis model with (D1) ResNet50, (D2) ResNet50 Multi-zoom Architecture, and (D3) custom Shallow Convolutional Neural Network (CNN); and, (E) visualization of model decision-making [explained in detail in Figure 4] Frameworks.

Figure 3: Accuracies of the deep learning models. A1, B1, and C1 show patch-level while A2, B2, and C2 show biopsy level accuracies of ResNet50, ResNet50 Multi-zoom Architecture, and Shallow CNN, respectively.

Figure 4: Confusion matrices for patch-level disease classification models (top left: ResNet50; top right: ResNet50 multi-zoom architecture; bottom left: Shallow Convolutional Neural Network (CNN); bottom right: ensemble. The numbers within the matrices represent normalized data in the form of percentages and darker colors indicate higher percentages.

Figure 5: Gradient-weighted Class Activation Mappings for patches trained via modified ResNet50, ResNet50 Multi-zoom, and Shallow CNN are shown. Areas being highlighted are: A:

superficial epithelium with high number of intraepithelial lymphocytes (IELs) and mononuclear

cells in the lamina propria (LP); B: areas of white slit-like spaces representing artefactual

separation of tissue within the LP and ignoring areas of crypt cross sections; C: crypt cross

sections; D: areas with cells consistent with mononuclear cellular infiltrate; E: telescoping crypt;

F: lymphocytes stacked in a row simulating a linear epithelial architecture; G: superficial

epithelium with tall columnar cells and a brush border; H: base of crypts resembling superficial

epithelium; I: superficial epithelium with high number of IELs; J: several crypt cross sections

rather than 1 to 2 crypts when compared with ReNet50 only; K: superficial epithelium with

abundant cytoplasm; L: surface epithelium with IELs and goblet cells; M: inner lumen of crypt

cross sections; N: mononuclear cells within the LP; O: surface epithelium with epithelial cells

containing abundant cytoplasm and goblet cells. Key: EE: Environmental Enteropathy; CD:

Celiac Disease; CNN: Convolutional Neural Network

Figure 1: Stain Color Normalization – images in the top row highlight the differences in stain colors before normalization while the bottom row shows images post color normalization which was applied using the specified target image.

Figure 2: Combined figure explaining the methods for development of an image analysis model for diagnosis of small bowel enteropathies and black box feature detection: (A) Digitized duodenal biopsy slides were obtained for secondary analysis; (B and C) dataset was pre-processed by creation of biopsy image patches, sample size augmentation, and stain color normalization [shown in figure 1]; (D) development of the image analysis model with (D1) ResNet50, (D2) ResNet50 Multi-zoom Architecture, and (D3) custom Shallow Convolutional Neural Network (CNN); and, (E) visualization of model decision-making [explained in detail in Figure 4] Frameworks.

Figure 3

Figure 3: Accuracies of the deep learning models. A1, B1, and C1 show patch-level while A2, B2, and C2 show biopsy level accuracies of ResNet50, ResNet50 Multi-zoom Architecture, and Shallow CNN, respectively.

Figure 4: Confusion matrices for patch-level disease classification models (top left: ResNet50; top right: ResNet50 multi-zoom architecture; bottom left: Shallow Convolutional Neural Network (CNN); bottom right: ensemble. The numbers within the matrices represent normalized data in the form of percentages and darker colors indicate higher percentages.

Figure 5: Gradient-weighted Class Activation Mappings for patches trained via modified ResNet50, ResNet50 Multi-zoom, and Shallow CNN are shown. Areas being highlighted are: A: superficial epithelium with high number of intraepithelial lymphocytes (IELs) and mononuclear cells in the lamina propria (LP); B: areas of white slit-like spaces representing artefactual separation of tissue within the LP and ignoring areas of crypt cross sections; C: crypt cross sections; D: areas with cells consistent with mononuclear cellular infiltrate; E: telescoping crypt; F: lymphocytes stacked in a row simulating a linear epithelial architecture; G: superficial epithelium with tall columnar cells and a brush border; H: base of crypts resembling superficial epithelium; I: superficial epithelium with high number of IELs; J: several crypt cross sections rather than 1 to 2 crypts when compared with ReNet50 only; K: superficial epithelium with abundant cytoplasm; L: surface epithelium with IELs and goblet cells; M: inner lumen of crypt cross sections; N: mononuclear cells within the LP; O: surface epithelium with epithelial cells containing abundant cytoplasm and goblet cells.

Key: EE: Environmental Enteropathy; CD: Celiac Disease; CNN: Convolutional Neural Network

Tables

Table 1: Population characteristics of the children from which biopsy images were obtained.

Key: PK = Pakistan, Aga Khan University; Zambia = University of Zambia Medical Center, US = United States, University of Virginia; EE = Environmental Enteropathy; LAZ/HAZ = Length/Height-for-Age Z score; IQR = inter-quartile range (written as first quartile to third quartile)

a – For some patients 2-3 biopsy images were available

b – Height of 3 patients was not available for children with celiac disease (Z scores calculated for 60)

	Total Population	PK	Zambia	US	
	n = 150	EE (n = 10)	EE (n = 16)	Celiac (n = 63)[b]	Normal (n = 61)
Biopsy Images[a]	461	29	19	239	174
Age, median (IQR), months	37.5 (19.0 to 121.5)	22.2 (20.8 to 23.4)	16.5 (9.5 to 21.0)	130.0 (85.0 to 176.0)	25.0 (16.5 to 41.0)
Gender; Male, n (%)	77 (51.3)	5 (50.0)	10 (62.5)	29 (46.0)	33 (54.0)
LAZ/ HAZ, median (IQR)	-0.6 (-1.9 to 0.4)	-2.8 (-3.6 to -2.3)	-3.1 (-4.1 to -2.2)	-0.3 (-0.8 to 0.7)	-0.2 (-1.3 to 0.5)

Click here to access/download
Supplemental Data File (doc, pdf, etc.)
AI_Enteropathy_JPGN_Supplement_LE.docx